

# Commutative Algebra Notes

---

Notes By: Aryaman Maithani

Last Updated: June 5, 2023

# Contents

<b>1</b>	<b>Artinian and Noetherian rings</b>	<b>2</b>
1.1	Primary decompositions . . . . .	2
1.2	Artinian rings . . . . .	2
<b>2</b>	<b>Valuation rings</b>	<b>5</b>
2.1	Definitions . . . . .	5
2.2	Discrete Valuation Rings . . . . .	9
<b>3</b>	<b>Graded rings and modules</b>	<b>12</b>
3.1	Graded rings and modules . . . . .	12
3.2	Filtrations and Topology . . . . .	19
3.3	The Artin-Rees Lemma . . . . .	22
3.4	Krull's intersection theorems . . . . .	23
<b>4</b>	<b>Dimension Theory</b>	<b>26</b>
4.1	Integer valued polynomials . . . . .	26
4.2	Embedding dimension . . . . .	28
4.3	Dimension à la Hilbert polynomials . . . . .	28
4.4	Properties of dimension . . . . .	32
4.5	Characterisation of dimension . . . . .	37
4.6	Krull dimension . . . . .	38
4.7	Dimension via length and ideals of definition . . . . .	43
4.8	Height and minimal generators . . . . .	45
<b>5</b>	<b>Faithfully flat modules</b>	<b>47</b>
<b>6</b>	<b>Completion</b>	<b>49</b>
6.1	Completion via inverse limits . . . . .	49
6.2	Completion via Cauchy Sequences . . . . .	52
6.3	Exactness . . . . .	54
6.4	Properties of completion . . . . .	57
<b>7</b>	<b>Regular sequences and depth</b>	<b>59</b>
7.1	Regular sequences . . . . .	59
7.2	Grade and depth . . . . .	66
7.3	Depth and projective dimension . . . . .	75
7.4	Some linear algebra . . . . .	80
<b>8</b>	<b>Appendix</b>	<b>87</b>

## §1. Artinian and Noetherian rings

We assume basic results about Artinian and Noetherian rings. We prove some others.

### §§1.1. Primary decompositions

### §§1.2. Artinian rings

**Proposition 1.2.1.** Let  $R$  be a ring, and  $I \subseteq R$  an ideal. Let  $M$  be an  $R$ -module such that  $IM = 0$ . Equivalently,  $M$  is an  $R/I$ -module.

Then,  $M$  is Artinian (resp. Noetherian) as an  $R$ -module if it is so as an  $R/I$ -module.

*Proof.* Check that the family of submodules is the same in both cases. □

**Proposition 1.2.2.** Let  $0 \rightarrow N \rightarrow M \rightarrow L \rightarrow 0$  be an exact sequence of  $R$ -modules. The following are equivalent:

1.  $M$  is Artinian (resp. Noetherian).
2.  $N$  and  $L$  are Artinian (resp. Noetherian).

**Corollary 1.2.3.** Any quotient of an Artinian ring is Artinian (as a ring).

**Corollary 1.2.4.** Every prime ideal in an Artinian ring is maximal. In other words, the (Krull) dimension of an Artinian ring is zero.

Since it is convenient to refer to this, we make a special definition of dimension now. We have an elaborate discussion of dimension later.

**Definition 1.2.5.** A ring  $R$  is said to be of **dimension zero**, denoted  $\dim(R) = 0$ , if every prime ideal of  $R$  is maximal.

**Proposition 1.2.6.** An Artinian ring has finitely many prime (maximal) ideals.

*Proof.* Suppose not. Let  $\mathfrak{m}_1, \mathfrak{m}_2, \dots$  be a sequence of distinct prime (and hence, maximal) ideals. Note that the chain

$$\mathfrak{m}_1 \supseteq \mathfrak{m}_1 \cap \mathfrak{m}_2 \supseteq \dots$$

must stabilise. Thus, there exists  $n$  such that

$$\mathfrak{m}_1 \cap \cdots \cap \mathfrak{m}_n = \mathfrak{m}_1 \cap \cdots \cap \mathfrak{m}_n \cap \mathfrak{m}_{n+1}.$$

Note that the ideal on the right is contained in  $\mathfrak{m}_{n+1}$ . Thus,  $\mathfrak{m}_{n+1}$  contains the intersection on the left. But then, since  $\mathfrak{m}_{n+1}$  is prime, it must contain some  $\mathfrak{m}_i$  with  $i \leq n$ . This contradicts that  $\mathfrak{m}_i$  and  $\mathfrak{m}_{n+1}$  are distinct maximal ideals.  $\square$

**Proposition 1.2.7.** Let  $R$  be a field, and  $M$  an  $R$ -module, i.e., an  $R$ -vector space. Then,  $M$  is Noetherian iff  $M$  is Artinian (iff  $\dim_R(M) < \infty$ ).

**Proposition 1.2.8.** Let  $M$  be an  $R$ -module. Suppose there exists a filtration of submodules

$$0 = M_0 \subseteq M_1 \subseteq M_2 \subseteq \cdots \subseteq M_n = M$$

such that each  $M_{i+1}/M_i$  is Artinian (resp. Noetherian). Then,  $M$  is Artinian (resp. Noetherian).

Conversely, if  $M$  is an Artinian (resp. Noetherian) module and we have a filtration as above, then each  $M_{i+1}/M_i$  is Artinian (resp. Noetherian).

*Proof.* By hypothesis,  $M_1 \cong M_1/M_0$  is Artinian (resp. Noetherian). The exact sequence

$$0 \rightarrow M_1 \rightarrow M_2 \rightarrow M_2/M_1$$

shows the same for  $M_2$ . Induct.

The converse is again an easy consequence of Proposition 1.2.2.  $\square$

**Corollary 1.2.9.** Let  $R$  be any ring, and  $M$  an  $R$ -module. Let  $\mathfrak{m}_1, \dots, \mathfrak{m}_n$  be maximal ideals of  $R$  (not necessarily distinct) such that  $\mathfrak{m}_1 \cdots \mathfrak{m}_n M = 0$ .

Then,  $M$  is Noetherian iff  $M$  is Artinian.

*Proof.* Consider the filtration

$$0 = \mathfrak{m}_1 \cdots \mathfrak{m}_n M \subseteq \mathfrak{m}_1 \cdots \mathfrak{m}_{n-1} M \subseteq \cdots \subseteq \mathfrak{m}_1 M \subseteq M.$$

Note that  $\mathfrak{m}_1 \cdots \mathfrak{m}_i M / \mathfrak{m}_1 \cdots \mathfrak{m}_{i+1} M$  is an  $R/\mathfrak{m}_{i+1}$ -module.

Now use the previous two propositions.  $\square$

**Proposition 1.2.10.** Let  $R$  be an Artinian ring. The nilradical  $\mathcal{N}$  is nilpotent, i.e.,  $\mathcal{N}^k = 0$  for some  $k \geq 0$ .

*Proof.* The descending chain of powers of  $\mathcal{N}$  must stabilise to some  $\mathcal{N}^k$ . We claim  $\mathcal{N}^k = 0$ . Suppose not. Let  $I := \mathcal{N}^k$ . By hypothesis,  $I \neq 0$ . Thus, by the Artinian condition, we may pick  $J$  minimal with respect to the property that  $IJ \neq 0$ .

Thus, there exists  $z \in J$  with  $Iz \neq 0$ . But also,  $I(Iz) = Iz \neq 0$ .

Since  $Iz \subseteq (z) \subseteq J$ , minimality forces  $Iz = (z)$ . Thus, we can write  $z = xz$  for some  $x \in I$ . Thus, we have  $z = xz = x^2z = x^3z = \dots$ .

But  $x$  is nilpotent, since  $I \subseteq \mathcal{N}$ . Thus,  $z = 0$ , contradicting that  $Iz \neq 0$ .  $\square$

**Corollary 1.2.11.** If  $(R, \mathfrak{m})$  is an Artinian local ring, then  $\mathfrak{m}$  is nilpotent.

*Proof.* By Corollary 1.2.4, all primes of  $R$  are maximal and there is only one such maximal ideal. Thus, the nilradical of  $R$  is  $\mathfrak{m}$ .  $\square$

**Corollary 1.2.12.** Any Artinian ring is Noetherian.

*Proof.* Let  $R$  be an Artinian ring. Note that  $\mathcal{N} = \bigcap_{\mathfrak{p} \in \text{Spec } R} \mathfrak{p}$ . Since  $R$  is Artinian,  $\mathcal{N}$  is the intersection of all the finitely many maximal ideals of  $R$ .

But this is just the product  $\prod \mathfrak{m}_i$  since the ideals are comaximal. In turn,  $\mathcal{N}^k$  is a product of maximal ideals. Now use Corollary 1.2.9.  $\square$

**Corollary 1.2.13.** Let  $R$  be a ring. The following are equivalent:

- (i)  $R$  is Artinian.
- (ii)  $R$  is Noetherian and  $\dim(R) = 0$ .

*Proof.* (i)  $\Rightarrow$  (ii) follows from earlier.

(ii)  $\Rightarrow$  (i): Since  $R$  is Noetherian,  $R$  has finitely many minimal primes. Since  $\dim(R) = 0$ , all of these are maximal as well. Since every prime contains a minimal prime, it follows that there are only finitely many primes. Then, we have  $\mathcal{N} = \bigcap_i \mathfrak{p}_i = \prod_i \mathfrak{p}_i$ . But in a Noetherian ring, the nilradical is nilpotent and thus,  $\mathcal{N}^k = 0$  for some  $k$ . Using Corollary 1.2.9, we get that  $R$  is Artinian.  $\square$

## §2. Valuation rings

### §§2.1. Definitions

**Definition 2.1.1.** A **valuation ring** is an integral domain  $R$  such that for all  $a, b \in R$ : either  $a$  divides  $b$  or  $b$  divides  $a$ .

Equivalently, the set of principal ideals is totally ordered by inclusion.

Equivalently, if  $K = \text{Frac}(R)$ , then for every  $x \in K^\times$ , either  $x \in R$  or  $x^{-1} \in R$ .

Recall that a totally ordered abelian group is an abelian group  $(A, +)$  with a total order  $\leq$  such that  $x \leq y \Rightarrow x + z \leq y + z$  for all  $x, y, z \in A$ .

We often consider the ordered set  $A_\infty := A \sqcup \{\infty\}$ , this extends the order from  $A$  by defining  $a < \infty$  for all  $a \in A$ .

**Definition 2.1.2.** Let  $K$  be a field, and  $A$  a totally ordered abelian group. A **valuation** is a map

$$v : K \rightarrow A \sqcup \{\infty\}$$

such that

- $v(0) = \infty$ ,
- $v(K^\times) \subseteq A$ ,
- $v(xy) = v(x) + v(y)$  for all  $x, y \in K^\times$ ,
- $v(x + y) \geq \min(v(x), v(y))$  for all  $x, y \in K$ .

Note that the third point is simply stating that  $v$  is a group homomorphism when restricted to  $K^\times \rightarrow A$ . We often just specify valuations by defining them on  $K^\times$ .

**Observation 2.1.3.** Since  $v$  is a group homomorphism, we have  $v(1) = 0$  and  $v(x^{-1}) = -v(x)$  for  $x \in K^\times$ .

**Proposition 2.1.4.** Let  $R$  be an integral domain, and  $K = \text{Frac}(R)$ . The following are equivalent:

- (i)  $R$  is a valuation ring.
- (ii) There exists a totally ordered abelian group  $A$  and a valuation  $v : K \rightarrow A_\infty$  such that  $R = \{x \in K : v(x) \geq 0\}$ .

*Proof.* We prove the nontrivial direction, namely (i)  $\Rightarrow$  (ii).

Assume that  $R$  is a valuation ring. Define a preorder on  $K^\times$  as follows:  $x \leq y$  iff  $y/x \in R$ . By hypothesis, any two elements are comparable.

Define the equivalence relation  $\sim$  on  $K^\times$  as  $x \sim y$  iff  $x \leq y$  and  $y \leq x$ .

Let  $A := K^\times / \sim$ . Note that we can define multiplication on  $A$  by  $[x] \cdot [y] = [xy]$ . (Check that this is well-defined.)

The map  $\nu : K^\times \rightarrow A$  defined by  $x \mapsto [x]$  does the job.  $\square$

In view of the above proposition, whenever we talk about a valuation ring  $R$ , we always have an associated valuation with it. Proceeding forward,  $K$  will denote the fraction field of  $R$ . We have the relation  $R = \{x \in K : \nu(x) \geq 0\}$ .

For  $x, y \in R$ , we see that  $x \mid y$  is equivalent to  $\nu(x) \leq \nu(y)$ .

**Proposition 2.1.5.** Any valuation ring  $R$  is normal.

*Proof.* Let  $x \in K$  be integral over  $R$ . We wish to show  $\nu(x) \geq 0$ . Write

$$x^n = a_{n-1}x^{n-1} + \cdots + a_1x + a_0$$

for  $a_i \in R$ . Applying  $\nu$ , we get

$$n\nu(x) \geq \min(\nu(a_{n-1}) + (n-1)\nu(x), \dots, \nu(a_0)).$$

Thus, there exists some  $i \in \{0, \dots, n-1\}$  such that  $n\nu(x) \geq \nu(a_i) + i\nu(x)$ . Thus,  $(n-i)\nu(x) \geq \nu(a_i) \geq 0$ . This proves the result.  $\square$

**Proposition 2.1.6.** Any valuation ring  $R$  is normal. The unique maximal ideal is given by  $\{x \in R : \nu(x) > 0\} = \{x \in K : \nu(x) > 0\}$ .

*Proof.* Let  $\mathfrak{m} := \{x \in R : \nu(x) > 0\}$ . The properties of valuation imply that  $\mathfrak{m}$  is indeed an ideal.  $\mathfrak{m}$  is proper since  $\nu(1) = 0$  and thus,  $1 \notin \mathfrak{m}$ .

On the other hand, if  $x \in R \setminus \mathfrak{m}$ , then  $\nu(x) = 0$ . This means that  $x^{-1} \in K$  also satisfies  $\nu(x^{-1}) = -\nu(x) = 0$  and thus,  $x^{-1} \in R$ , i.e.,  $x$  is a unit.  $\square$

**Theorem 2.1.7.** Let  $R$  be a domain, not necessarily a valuation ring.

Let  $\mathfrak{p} \in \text{Spec } R$ , and  $K = \text{Frac}(R)$ .

Then, there exists a valuation ring  $V$  such that

- $R \subseteq V \subseteq K$ ,
- the maximal ideal of  $V$  contracts to  $\mathfrak{p}$ .

In particular, if  $\nu$  is the associated valuation to  $V$ , we have  $\mathfrak{p} = \{x \in R : \nu(x) > 0\}$ .



*Proof.* Throughout this proof, localisations of  $R$  (and of other subrings of  $K$ ) will be considered as subrings of  $K$  in the natural way.

Let  $\Sigma$  denote the collection of intermediate subrings  $R'$ :  $R_p \subseteq R' \subseteq K$  such that  $pR' \neq R'$ .

$\Sigma$  is nonempty since  $R_p \in \Sigma$ . Ordering  $\Sigma$  by inclusion, we note that  $\Sigma$  satisfies the hypothesis of Zorn's Lemma<sup>1</sup> and thus, there exists a maximal  $V$ . We show that  $V$  has the desired properties.

**Claim 1.**  $V$  is local.

*Proof.* By construction,  $pV \neq V$ . Thus,  $pV$  is contained in some maximal ideal  $\mathfrak{m} \subseteq V$ . In turn,  $pV_{\mathfrak{m}}$  is a proper subset of  $V_{\mathfrak{m}}$  and hence,  $V_{\mathfrak{m}} \in \Sigma$ . By maximality, we must have  $V = V_{\mathfrak{m}}$ , proving that  $V$  is a local ring.  $\square$

Going forth, let  $\mathfrak{m}$  denote the maximal ideal of  $V$ . Note that  $pV \subseteq \mathfrak{m}$ . In particular, intersecting with  $R_p$  shows that

$$pV \cap R_p = \mathfrak{m} \cap R_p.$$

But  $pR_p$  is clearly contained in the left ideal. Thus, we have

$$pR_p = \mathfrak{m} \cap R_p.$$

Further contracting to  $R$  gives us

$$p = \mathfrak{m} \cap R.$$

Thus, we now only need to prove that  $V$  is a valuation ring.

**Claim 2.**  $V$  is normal.

*Proof.* Let  $x \in K$  be integral over  $V$ . Showing  $x \in V$  is equivalent to saying  $V = V[x]$ . Using maximality of  $V$  in  $\Sigma$ , it suffices to prove that  $V[x] \in \Sigma$ .

Note that  $V \hookrightarrow V[x]$  is an integral extension. Thus, there is a prime  $\mathfrak{m}' \subseteq V[x]$  lying over  $V$ . In particular,  $p \subseteq \mathfrak{m}'$  showing that  $pV[x] \neq V[x]$ .  $\square$

We are now ready to show that  $V$  is a valuation ring. Let  $x \in K^\times$  with  $x \notin V$ . We need to show that  $x^{-1} \in V$ .

Consider the subring  $V[x]$ . This is strictly larger than  $V$ . By maximality, we must have that  $V[x] \notin \Sigma$  and thus,  $1 \in pV[x]$ . We can then write

$$1 = p_0 + p_1x + \cdots + p_tx^t$$

for some  $p_i \in pV$ .

---

<sup>1</sup>Note that  $1 \in pR'$  is a "finite condition".

Rearrange the above to get

$$1 - p_0 = p_1x + \cdots + p_tx^t.$$

Note that  $p_0 \in pV \subseteq \mathfrak{m}$  and thus,  $1 - p_0$  is a unit in  $V$ . Thus, we can write

$$\frac{1}{x^t} = \frac{1}{1 - p_0} \left( \frac{p_1}{x^{t-1}} + \cdots + p_t \right).$$

The above shows that  $x^{-1}$  is integral over  $V$ . By Claim 2, it follows that  $x^{-1} \in V$ , as desired.  $\square$

**Corollary 2.1.8.** Let  $R$  be an integral domain, and let  $\mathfrak{q} \subseteq \mathfrak{p}$  be primes. There exists a valuation ring  $V$  and a ring homomorphism  $f : R \rightarrow V$  such that  $f^{-1}(\mathfrak{m}_V) = \mathfrak{p}$  and  $f^{-1}(0) = \mathfrak{q}$ . ( $\mathfrak{m}_V$  denotes the maximal ideal of  $V$ .)

*Proof.* Applying the previous proposition to  $R/\mathfrak{q}$ , there exists a valuation ring  $V$  and an injection  $g : R/\mathfrak{q} \hookrightarrow V$  such that  $g^{-1}(\mathfrak{m}_V) = \mathfrak{p}/\mathfrak{q}$  (and necessarily  $g^{-1}(0) = 0$ ). Compose this with the projection  $R \twoheadrightarrow R/\mathfrak{q}$ .  $\square$

**Corollary 2.1.9.** If  $R$  is a normal domain, then

$$R = \bigcap_{\substack{R \subseteq V \subseteq K \\ V \text{ is a valuation ring}}} V.$$

More generally, if  $R$  is any integral domain, and  $\bar{R}$  is its integral closure in  $K$ , then

$$\bar{R} = \bigcap_{\substack{R \subseteq V \subseteq K \\ V \text{ is a valuation ring}}} V.$$

*Proof.* Note that since valuation rings are integrally closed, it follows that  $\bar{R}$  is contained in the intersection.

Conversely, suppose  $x$  belongs to the intersection. This means that  $v(x) \geq 0$  for every valuation  $v$  on  $K$ . We must show that  $x$  is integral over  $R$ .

Let  $R' = R \left[ \frac{1}{x} \right]$  and  $I = \left( \frac{1}{x} \right) R'$ .

**Claim.**  $I = R'$ .

*Proof.* If not, then  $I \subseteq \mathfrak{m}$  for some maximal ideal  $\mathfrak{m} \subseteq R'$ . By the previous proposition, there exists a valuation ring  $V \supseteq R'$  such that  $x^{-1} \in \mathfrak{m}_V$ . This implies that  $v(x) = -v(x^{-1}) > 0$ . A contradiction.  $\square$

Thus,  $I = R'$ . In other words,  $1/x$  is a unit in  $R'$  and hence,  $x \in R'$ . This means we can write

$$x = r_0 + \frac{r_1}{x} + \cdots + \frac{r_n}{x^n}$$

for some  $r_i \in R$ . Multiplying with  $x^n$  shows that  $x$  satisfies a monic polynomial over  $R$ , as desired.  $\square$

**Proposition 2.1.10.** Let  $R$  be an integral domain. The following are equivalent:

- (i)  $R$  is a valuation ring.
- (ii)  $R$  is a local Bézout domain. (That is,  $R$  is a local ring, where every finitely generated ideal is principal.)

*Proof.* (i)  $\Rightarrow$  (ii): Suppose  $R$  is a valuation ring. We have already seen that  $R$  is local. Let  $I = (f_1, \dots, f_n)$  be a finitely generated ideal. By definition of a valuation ring, we may pick a maximal ideal among the principal ideals  $(f_1), \dots, (f_n)$ . Then,  $I$  is equal to that principal ideal.

(ii)  $\Rightarrow$  (i): Let  $x, y \in R$  be nonzero. We wish to show that either  $x \mid y$  or  $y \mid x$ . By hypothesis, we can write  $(x, y) = (d)$  for some  $d \in R$ . In turn, we have the relations

$$x = dx', y = dy', d = ax + by,$$

for some  $a, b, x', y' \in R$ .

Plugging the first two relations in the last gives us

$$d = (ax' + by')d.$$

We may cancel  $d$  to see that  $(x', y') = (1)$ . Since  $R$  is a local ring, this implies that one of  $x'$  or  $y'$  is a unit. Without loss of generality,  $x'$  is a unit (in  $R$ ).

Then, we have

$$y = dy' = \frac{y'}{x'}x.$$

$\square$

## §§2.2. Discrete Valuation Rings

**Definition 2.2.1.** A valuation  $v : K \rightarrow A_\infty$  is said to be a **discrete valuation** if  $v(K^\times)$  is isomorphic to  $\mathbb{Z}$ .

A **discrete valuation ring (DVR)** is a ring that is of the form  $\{x \in K : v(x) \geq 0\}$  for some discrete valuation  $v : K \rightarrow A_\infty$ .

Note that the trivial valuation is not a discrete valuation. Equivalently, a field is not a discrete valuation ring.

Note that given a discrete valuation, we may always assume that  $A = \mathbb{Z}$  and that  $v(x) \neq 0$  for some  $x \in K$ .

**Proposition 2.2.2.** Let  $R$  be a valuation ring. The following are equivalent.

- (i)  $R$  is Noetherian.
- (ii)  $R$  is a DVR or a field.
- (iii)  $R$  is a local PID.

((ii) and (iii) are equivalent even without the a priori assumption that  $R$  is a valuation ring. See Corollary 2.2.4)

*Proof.* Let  $v : K \rightarrow A_\infty$  and  $\mathfrak{m} \subseteq R$  be as usual

(i)  $\Leftrightarrow$  (iii) is clear by Proposition 2.1.10.

(iii)  $\Rightarrow$  (ii): Assume  $R$  is a local PID and not a field.

By hypothesis, we have  $\mathfrak{m} = (x)$  with  $0 < v(x) < \infty$ . In turn,

$$v(x) = \min\{v(r) : r \in K, v(r) > 0\}.$$

(The above follows since  $v(r) > 0 \Rightarrow r \in \mathfrak{m} = (x) \Rightarrow x \mid r$ .)

In other words,  $v(x)$  is the smallest positive valuation.

We show that  $v(K^\times)$  is generated by  $v(x)$ . Let  $y \in K^\times$  be arbitrary. We may assume  $y$  is not a unit. By replacing  $y$  with  $y^{-1}$ , we may assume that  $v(y) > 0$  and hence,  $y \in R$ .

Consider the following subsets of  $K$ :

$$yR \subsetneq \frac{y}{x}R \subsetneq \frac{y}{x^2}R \subsetneq \cdots$$

Note that as long as  $y/x^n \in R$ , the set  $(y/x^n)R$  is an ideal of  $R$ . Since  $R$  is Noetherian, the chain must eventually escape  $R$ , i.e., there exists  $n \geq 0$  such that  $y/x^{n+1} \notin R$ . Choose the smallest such  $n$ . Then, we have

$$v(y/x^n) \geq 0 > v(y/x^{n+1}).$$

If  $v(y/x^n) = 0$ , then we are done since we have  $v(y) = nv(x)$ . If this is not the case, then we have strict inequalities above, which gives

$$v(y/x^n) > 0 > v(y/x^{n+1}).$$

Rearranging gives

$$v(x) > v(x^{n+1}/y) > 0.$$

But this contradicts that  $x$  had smallest positive valuation.

(ii)  $\Rightarrow$  (i): Clearly we may assume that  $R$  is a DVR. Let  $v : K \rightarrow \mathbb{Z}_\infty$  be the associated valuation. Consider the ideals  $I_n$  defined as

$$I_n := \{x \in R : v(x) \geq n\},$$

for  $n \geq 0$ . ( $I_n$  is an ideal by properties of valuation.)

We claim that any nonzero ideal in  $R$  is of the form. Indeed, given a nonzero ideal  $J$ , let  $n \geq 0$  and  $x \in J$  be such that  $v(x) = n$  is the smallest valuation among elements of  $J$ .

In particular,  $J \subseteq I_n$ . Conversely, if  $y \in I_n$ , then  $v(x) \leq v(y)$  shows that  $x \mid y$  in  $R$  and hence,  $y \in (x) \subseteq J$ .

Thus, the ideals of  $R$  all appear in the following chain:

$$I_0 \supseteq I_1 \supseteq I_2 \supseteq \cdots \supseteq 0.$$

It follows that  $R$  is Noetherian. □

Note that if  $v : K \rightarrow \mathbb{Z}_\infty$  is a valuation, then  $v(K^\times)$  is a nonzero subgroup of  $\mathbb{Z}$  and thus, of the form  $n\mathbb{Z}$  for some  $n > 0$ . This gives us a new valuation  $v' : K \rightarrow \mathbb{Z}_\infty$  defined by  $v'(x) = v(x)/n$ .

Both of these define the same valuation ring. Thus, we may always assume that a discrete valuation surjects onto  $\mathbb{Z}_\infty$ .

From the last part of the above proof, we extract the following.

**Porism 2.2.3** (Description of ideals in DVRs). Let  $R$  be a DVR, and  $v : K \rightarrow \mathbb{Z}_\infty$  the associated valuation.

Pick any  $t \in K$  with  $v(t) = 1$ . Then, the ideals

$$(1), (t), (t^2), (t^3), \dots, (0)$$

are all the distinct ideals of  $R$ .

In particular,  $R$  is a local PID.

**Corollary 2.2.4.** Let  $R$  be a ring which is not a field. The following are equivalent:

- (i)  $R$  is a local PID.
- (ii)  $R$  is a DVR.

*Proof.* In either case, we need to show that  $R$  is a valuation ring and then we can appeal to Proposition 2.2.2. If we assume (ii), then this follows tautologically. If we assume (i), then this follows from Proposition 2.1.10. □

### §3. Graded rings and modules

Going forth, we make the convention that  $0 \in \mathbb{N}$ .

#### §§3.1. Graded rings and modules

**Definition 3.1.1.** A  $(\mathbb{N})$ -graded ring is a ring  $R$  with a sequence of additive subgroups  $\langle R_n \rangle_{n \geq 0}$  such that

$$R = \bigoplus_{n \geq 0} R_n \quad \text{and} \quad R_m R_n \subseteq R_{m+n}$$

for all  $n, m \geq 0$ .

Elements of  $R_0 \cup R_1 \cup \cdots$  are called **homogeneous elements**. A nonzero homogeneous element  $x$  belongs to  $R_n$  for some unique  $n$ , this is called the **degree** of  $x$ .

$R_+ := \bigoplus_{n \geq 1} R_n$  is an ideal of  $R$ , called the **irrelevant ideal**.

Note that  $0 \in R_n$  for all  $n$ . There is some care that must be taken when defining the degree of 0. We do not bother about it and leave the edge cases to the reader.

Note that being graded is not a property of a ring. Rather, it consists of additional data given to a ring. (In contrast to something like how being a field is a property of a ring.) However, we still often say “ $R$  is a graded ring” and tacitly assume that we are given a *grading*  $\langle R_n \rangle_{n \in \mathbb{N}}$ .

**Definition 3.1.2.** Let  $R$  be a graded ring. A **graded  $R$ -module** is an  $R$ -module  $M$  with a sequence of additive subgroups  $\langle M_n \rangle_{n \in \mathbb{Z}}$  such that

$$M = \bigoplus_{n \in \mathbb{Z}} M_n \quad \text{and} \quad R_m M_n \subseteq M_{m+n}$$

for all  $n, m \geq 0$ .

A **homogeneous element** and its **degree** is defined as before.

**Proposition 3.1.3.** Let  $R$  be a graded ring, and  $M$  a graded  $R$ -module. Then,  $R_0$  is a subring of  $R$ . In particular,  $1 \in R$ . Moreover, each  $M_n$  is an  $R_0$ -module.

Note that  $M_n$  will typically *not* be an  $R$ -module.

*Proof.* The only nontrivial thing to check is that  $1 \in R$ . By hypothesis, we can write

$$1 = r_0 + r_1 + \cdots, \tag{3.1}$$

where  $r_i \in R_i$  and  $r_n = 0$  for  $n \gg 0$ .

We show that  $r_i = 0$  for all  $i > 0$  and hence,  $1 = r_0 \in R_0$ .

Multiplying (3.1) with  $r_0$  shows

$$r_0 = r_0^2 + r_0 r_1 + \cdots.$$

Comparing the homogeneous components on each side shows that  $r_0 r_i = 0$  for  $i > 0$ .

Now, fix  $i > 0$  and multiply (3.1) with  $r_i$  to get

$$r_i = r_0 r_i + r_1 r_i + \cdots.$$

Again, comparing the homogeneous components gives

$$r_i = r_0 r_i.$$

But by the earlier calculation, the element on the right is 0. □

**Example 3.1.4.** 1. If  $k$  is a field, and  $R = k[x_1, \dots, x_d]$ , then  $R$  has a natural grading with  $R_n$  consisting of the  $k$ -vector space generated by monomials of degree  $n$ .

2. Let  $R$  be a graded ring, and  $f \in R$  be a nonzero homogeneous element. Then,  $R_f = R[f^{-1}]$  has a natural  $R$ -module structure. (Note that unless  $f$  is nilpotent (in which case,  $R_f = 0$ ) or  $\deg(f) = 0$ ,  $R_f$  will have graded pieces in the negative component. Thus,  $R_f$  is not an  $\mathbb{N}$ -graded ring.)

3. More generally, let  $S \subseteq R$  be a multiplicative subset consisting of homogeneous elements. Then,  $S^{-1}R$  is a graded  $R$ -module:  $(S^{-1}R)_n$  consists of elements of the form  $r/s$  where  $r \in R$  and  $s \in S$  are homogeneous with  $\deg(r) - \deg(s) = n$ . (If  $r/s = r'/s'$  with  $r'$  and  $s'$  also homogeneous, it will follow that  $\deg(r') - \deg(s') = n$ .)

By the above definition, it follows that  $R_n(S^{-1}R)_m \subseteq (S^{-1}R)_{m+n}$ . In fact, it even follows that  $S^{-1}R$  is a  $\mathbb{Z}$ -graded ring (the definition is the obvious one).

Even in this case, the homogeneous elements of degree 0 form a subring of  $S^{-1}R$ .

If  $\mathfrak{p} \subseteq R$  is a graded homogeneous prime ideal (defined just below), then one can consider  $S$  to be the set of homogeneous elements not contained in  $\mathfrak{p}$ . This is a multiplicative subset. The 0-degree subring of  $S^{-1}R$  is sometimes denoted  $R_{(\mathfrak{p})}$ .

**Example 3.1.5** (Rees algebra). Given a ring  $R$  and an ideal  $I$ , we can construct the **Rees algebra** as follows:

Let  $S_0 := R$ ,  $S_n := I^n$  for  $n \geq 1$ . Then,

$$\bigoplus_{n \geq 0} S_n$$

has a natural ring structure.

There is a notational issue as to how one would write an element of  $S$ . One way is to write them as tuples, this is unwieldy. We would like to write them as sums, but then there's the issue of whether one interprets an element  $x \in I$  as sitting in  $S_1$  or  $S_0$ .

To combat this, we attach a dummy variable and define

$$R[It] := \bigoplus_{n \geq 0} I^n t^n \cong \bigoplus_{n \geq 0} S_n.$$

The upshot is that the variable  $t$  now succinctly acts as a bookkeeping device.

**Definition 3.1.6.** Let  $M, N$  be graded  $R$ -modules. A **graded module homomorphism**  $f : M \rightarrow N$  is a module homomorphism  $f$  such that  $f(M_n) \subseteq N_n$ .

A sequence of graded modules is defined similarly.

**Definition 3.1.7.** Given a graded  $R$ -module  $M$ , and an integer  $d$ , we defined the **shift**  $M(d)$  to be the graded  $R$ -module  $M$  with new grading

$$(M(d))_n = M_{d+n}.$$

**Observation 3.1.8.** Let  $M$  be a graded  $R$ -module, and  $x \in R$  a homogeneous element of degree  $d$ . Note that

$$M \xrightarrow{x} M$$

is a map of  $R$ -modules, but not a graded map (barring some trivial cases).

However, by shifting the domain, we do get a graded map as

$$M(-d) \xrightarrow{x} M.$$

Indeed, note that if  $m \in (M(-d))_n$ , then

$$xm \in R_d M_{-d+n} \subseteq M_n,$$

as desired.

**Definition 3.1.9.** Let  $M$  be a graded  $R$ -module. A submodule  $N \subseteq M$  is said to be a **graded (or homogeneous)  $R$ -submodule** if  $N$  satisfies any of the following equivalent properties:



1.  $N$  is generated by homogeneous elements.
2.  $N = \bigoplus_{n \in \mathbb{Z}} (N \cap M_n)$ .
3. Whenever  $x \in N$ , every homogeneous component (in  $M$ ) of  $x$  is an element of  $N$ .

**Observation 3.1.10.** If  $N \leq M$  is a graded submodule, then  $M/N$  has a natural grading given by

$$(M/N)_n := M_n / (N \cap M_n).$$

Note that by definition, we have  $N = \bigoplus_{n \in \mathbb{Z}} (N \cap M_n)$ , where this is an internal direct sum. Thus, one has the natural isomorphism  $M/N \cong \bigoplus_{n \in \mathbb{Z}} (M/N)_n$ .

Under this grading, we have an exact sequence of graded  $R$ -modules given as

$$0 \rightarrow N \rightarrow M \rightarrow M/N \rightarrow 0.$$

For homogeneous ideals, primality can be checked on homogeneous elements:

**Proposition 3.1.11.** Let  $I \subseteq R$  be a graded ideal with the property that if  $x, y$  are homogeneous elements with  $xy \in I$ , then either  $x \in I$  or  $y \in I$ . Then,  $I$  is prime.

**Proposition 3.1.12.** Let  $R$  be a graded ring. Let  $S \subseteq R_+$  be any subset. The following are equivalent:

- (i)  $S$  generates  $R_+$  as an  $R$ -ideal.
- (ii)  $S$  generates  $R$  as an  $R_0$ -algebra.

*Proof.* Only (i)  $\Rightarrow$  (ii) is nontrivial. Let  $T := R_0[S]$ . We show that  $R = T$  by proving  $R_n \subseteq R$  for all  $n \geq 0$ .

We prove this by induction.  $n = 0$  is clear. Let  $n > 0$ . By hypothesis, we have  $R_n \subseteq R_+ = RS$ . Thus,

$$\begin{aligned} R_n &= (R \cdot S) \cap R_n \\ &= (R_0 + \cdots + R_{n-1})S \cap R_n \\ &\subseteq TS \cap R_n \\ &\subseteq T, \end{aligned} \quad \left. \begin{array}{l} \text{If } m \geq n, \text{ then } R_m S \cap R_n = 0 \\ \text{by induction} \end{array} \right\}$$

as desired. □

**Corollary 3.1.13.** Let  $R$  be a graded ring.

$R$  is a Noetherian ring iff  $R_0$  is a Noetherian ring and  $R$  is a finitely generated  $R_0$ -algebra.

*Proof.* ( $\Leftarrow$ ) Follows from Hilbert's Basis Theorem since we can write  $R$  as a quotient of  $R[x_1, \dots, x_d]$ .

( $\Rightarrow$ )  $R_+$  is a finitely generated ideal and thus,  $R$  is finitely generated as an  $R_0$ -algebra.  $R_0$  is Noetherian since  $R_0 = R/R_+$ .  $\square$

**Definition 3.1.14.** Let  $R$  be a graded ring with grading  $\langle R_n \rangle_{n \geq 0}$ . Let  $d > 0$ . We define the **twisted** graded ring  $R^{(d)}$  to be the graded ring with

$$\left(R^{(d)}\right)_n = R_{dn}.$$

This is also called the  **$d$ -th Veronese subalgebra**.

$R^{(d)}$  can be visualised as follows: We have the additive subgroups  $R_0, R_d, R_{2d}, \dots$  of  $R$  such that their *internal* sum is direct. This realises  $R^{(d)}$  as a subset (subgroup even) of  $R$ . We now just scale the grading by  $d$ .

**Example 3.1.15.** Consider  $R = k[x, y]$  with the standard grading. Then,  $R^{(2)}$ , when considered as a subring of  $R$  consists of those polynomials such that each monomial has even degree.

This can be written as  $k[x^2, xy, y^2]$ , as a set. However,  $x^2$  now has degree 1 in  $R^{(2)}$ .

Similarly,  $R^{(3)} = k[x^3, x^2y, xy^2, y^3]$ .

**Definition 3.1.16.** Let  $M$  be a graded  $R$ -module, and  $d > 0$ . Let  $\ell \in \{0, \dots, d-1\}$ . Then, we define

$$M^{(d, \ell)} := \bigoplus_{k: k \equiv \ell (d)} M_k.$$

The above has the structure of a graded  $R^{(d)}$ -module.

A slightly more precise way of writing the above would be to specify that the  $n$ -th graded component of  $M^{(d, \ell)}$  is  $nd + \ell$ . Note that

$$R_m^{(d)} M_n^{(d, \ell)} = R_{md} M_{nd+\ell} \subseteq M_{(m+n)d+\ell} = M_{m+n}^{(d, \ell)}$$

and thus,  $M^{(d, \ell)}$  is indeed a graded  $R^{(d)}$ -module.

**Definition 3.1.17.** Given a graded ring  $R$ ,  $\text{Proj}(R)$  denotes the set of homogeneous primes of  $R$  not containing  $R_+$ .

Fact:  $\text{Proj}(R)$  is in bijection with  $\text{Proj}(R^{(d)})$  for all  $d > 0$ .

**Proposition 3.1.18.** Let  $R$  be a graded ring such that  $R$  is a finitely generated  $R_0$ -algebra. Let  $M$  be a finitely generated graded  $R$ -module. (For example, if  $R$  and  $M$  are both Noetherian.) Let  $d > 0$ .

Then,

- (i)  $M_i$  is a finitely generated  $R_0$ -module for all  $i$ . Moreover,  $M_i = 0$  for  $i \ll 0$ .
- (ii)  $M^{(d,\ell)}$  is a finitely generated  $R^{(d)}$ -module for all  $\ell$  and thus,

$$M \cong \bigoplus_{\ell=0}^{d-1} M^{(d,\ell)}$$

is so.

- (iii) In particular,  $R$  is a finitely generated  $R^{(d)}$ -module.
- (iv)  $R^{(d)}$  is a finitely generated  $R_0$ -algebra.

Note that in the above,  $M$  (and similarly  $R$ ) is an ordinary  $R^{(d)}$ -module, not a graded one. (The module structure is the one inherited by virtue of  $R^{(d)}$  being a subring of  $R$ , ignoring grading.)

*Proof.* We may assume that  $m_1, \dots, m_t$  are nonzero homogeneous and generate  $M$ . Let  $d = \min_i \deg(m_i)$ . Then,  $M_n$  must be zero for  $i < d$  since any  $R$ -linear combination of the  $m_i$  has degree at least  $d$ .<sup>2</sup>

Now, let  $r_1, \dots, r_p \in R_+ \setminus \{0\}$  be homogeneous elements that generate  $R$  as an  $R_0$ -algebra. Let  $d_i := \deg(r_i)$ .

Now, note that elements of  $M_i$  can be written as  $R_0$ -linear combination of elements of the form

$$r_1^{a_1} \cdots r_p^{a_p} m_s,$$

where  $s \in \{1, \dots, t\}$  and  $a_1, \dots, a_p \geq 0$  satisfy

$$\deg(m_s) + \sum_j a_j d_j = i.$$

But there are only finitely many such ways to pick  $s$  and  $a_1, \dots, a_p$  (once we fix  $i$ ). Thus, each  $M_i$  is a finite  $R_0$ -module. This proves (i).

<sup>2</sup>It was crucial here that  $R$  is  $\mathbb{N}$ -graded.

Note that each  $M^{(d,\ell)}$  is a quotient of  $M$  and hence it suffices to prove that  $M$  is finitely generated.  $M$  can be generated *over*  $R_0$  by elements of the form

$$r_1^{a_1} \cdots r_p^{a_p} m_s$$

where  $a_1, \dots, a_p \geq 0$  and  $s \in \{1, \dots, t\}$ .

Since all powers of the form  $r_i^{kd}$  are in  $R^{(d)}$ , we see that we have a finite generating set by restricting the  $a_i$  to be in  $[0, d-1]$ . This proves (ii) and (iii).

Since  $R$  is a finitely generated  $R_0$ -algebra,  $R_+$  is a finitely generated  $R$ -module (Proposition 3.1.12). By (ii), we see that  $R_+$  is a finitely generated  $R^{(d)}$ -module. But  $(R^{(d)})_+ = (R_+)^{(d,0)}$  and thus, part (ii) tells us that this is a finitely generated  $R^{(d)}$ -module. Using Proposition 3.1.12 again, we see that  $R^{(d)}$  is a finitely generated  $R_0$ -algebra. (Note that  $(R^{(d)})_0 = R_0$ .)

This proves (iv). □

**Corollary 3.1.19.** If  $R$  is a Noetherian graded ring, then so is  $R^{(d)}$  for all  $d > 0$ .

*Proof.* Use Corollary 3.1.13 and part (iv) of the previous proposition. □

**Proposition 3.1.20.** Let  $R$  be a graded ring, finitely generated as an  $R_0$ -algebra. Let  $M$  be a finitely generated graded  $R$ -module.

Then, there exist integers  $n_0, d \geq 0$  such that

$$R_d M_n = M_{d+n}$$

for all  $n \geq n_0$ .

Note that we always have  $R_d M_n \subseteq M_{d+n}$ .

*Proof.* Let notation be as in the earlier proof:  $m_1, \dots, m_t$  are nonzero homogeneous and generate  $M$  over  $R$ ;  $r_1, \dots, r_p \in R_+ \setminus \{0\}$  are homogeneous elements that generate  $R$  as an  $R_0$ -algebra, with  $d_i := \deg(r_i)$ .

Set  $d := \text{lcm}_i d_i$  and  $n_0 := 1 + pd + \max_i \deg(m_i)$ . For  $1 \leq i \leq p$ , define  $s_i := r_i^{d/d_i} \in R_d$ . We show that  $d$  and  $n_0$  have the properties as desired. To this end, let  $n \geq n_0$ . Pick  $m \in M_n$ . As seen earlier, we can write  $m$  as an  $R_0$ -linear combination of elements of the form

$$r_1^{a_1} \cdots r_p^{a_p} m_s \tag{3.2}$$

such that  $a_1 d_1 + \cdots + a_p d_p + \deg(m_s) = n$ . Note that

$$a_1 d_1 + \cdots + a_p d_p = n - \deg(m_s) > pd.$$

Thus, there is some  $i \in \{1, \dots, p\}$  such that  $a_i d_i > d$  and thus,  $a_i > d/d_i$ . Thus, we may write

$$r_i^{a_i} = r_i^{d/d_i} r_i^{a_i - d/d_i} = s_i r_i^{a_i - d/d_i}.$$

Thus, each term of the form in (3.2) can be written as an element of  $R_d M_{n-d}$ .  $\square$

Looking at the above proof, we may in fact extract a special result.

**Porism 3.1.21.** With the same hypothesis as earlier, assume further that  $R = R_0[r_1, \dots, r_p]$  with  $\deg(r_i) = 1$  for all  $i$ .

Then, we may take  $d = 1$  above, i.e.,  $R_1 M_n = M_{n+1}$  for  $n \gg 0$ .

**Corollary 3.1.22.** Suppose  $R$  is a graded ring and a finitely generated  $R_0$ -algebra. Then, there exists  $d > 0$  such that  $R^{(d)}$  is generated over  $R_0$  by  $(R^{(d)})_1 = R_d$ .

In other words, by taking a high enough Veronese subalgebra, we can ensure that it is generated in degree 1.

*Proof.* By Proposition 3.1.20, there exist  $d', n_0 > 0$  such that  $R_{d'} R_n = R_{n+d'}$  for all  $n \geq n_0$ . Let  $d$  be a multiply of  $d'$  which is greater than  $n_0$ . Then, for  $n \geq d$ , we have

$$R_{d+n} = R_{d'} R_{d-d'+n} = \dots = R_d R_n.$$

In particular, we have  $R_{kd} = (R_d)^k$  for all  $k \geq 1$ , as desired.  $\square$

## §§3.2. Filtrations and Topology

**Definition 3.2.1.** Let  $R$  be a ring. A **filtration** on  $R$  is a sequence of ideals  $\langle I_n \rangle_{n \in \mathbb{N}}$  satisfying

$$R = I_0 \supseteq I_1 \supseteq I_2 \supseteq \dots \quad \text{and} \quad I_n I_m \subseteq I_{n+m}$$

for all  $n, m \geq 0$ .

We say that  $(R, \langle I_n \rangle_{n \in \mathbb{N}})$  is a **filtered ring**.

Given such a filtered ring, and an  $R$ -module  $M$ , a **filtration** on  $M$  is a sequence of  $R$ -submodules  $\langle M_n \rangle_{n \in \mathbb{N}}$  satisfying

$$M = M_0 \supseteq M_1 \supseteq M_2 \supseteq \dots \quad \text{and} \quad I_n M_m \subseteq M_{n+m}$$

for all  $n, m \geq 0$ .

**Example 3.2.2.** Let  $R$  be any ring,  $I \subseteq R$  an ideal, and  $M$  an  $R$ -module. The **I-adic filtration** on  $R$  is the sequence  $\langle I^n \rangle_{n \geq 0}$ . Corresponding to this filtration, we have the filtration on  $M$  given by  $\langle I^n M \rangle_{n \geq 0}$ .

**Example 3.2.3.** Let  $R$  be a graded ring with grading  $\langle R_n \rangle_{n \geq 0}$ . Set  $I_n := R_{\geq n} = \bigoplus_{i \geq n} R_i$ . Then,  $\langle I_n \rangle_{n \geq 0}$  is a filtration on  $R$ .

**Definition 3.2.4.** Let  $(R, \langle I_n \rangle_{n \geq 0})$  be a filtered ring. The **associated graded ring**  $\text{gr}(R)$  is defined by

$$\text{gr}(R) := R/I_1 \oplus I_1/I_2 \oplus I_2/I_3 \oplus \cdots.$$

This has the natural structure of a graded ring.

Correspondingly, if we have a filtered module  $(M, \langle M_n \rangle_{n \geq 0})$ , then we define

$$\text{gr}(M) := M/M_1 \oplus M_1/M_2 \oplus M_2/M_3 \oplus \cdots.$$

$\text{gr}(M)$  is a graded  $\text{gr}(R)$ -module.

**Definition 3.2.5.** A **topological ring**  $R$  is a ring with a topology such that the following three maps are continuous:

- $+: R \times R \rightarrow R, (x, y) \mapsto x + y$ ;
- $\cdot: R \times R \rightarrow R, (x, y) \mapsto xy$ ;
- $-: R \rightarrow R, x \mapsto -x$ .

$R$  is said to have a **linear topology** if  $0$  has a neighbourhood basis of ideals (that is, there is a collection of ideals  $\mathcal{I}$  such that any neighbourhood of  $0$  contains an ideal  $I \in \mathcal{I}$ ).

Note for any fixed  $x \in R$ , the translation map  $T_x: R \rightarrow R$  defined by  $y \mapsto x + y$  is a homeomorphism which takes  $0$  to  $x$ . Thus, studying neighbourhoods of  $0$  is sufficient.

Note that if  $\langle I_n \rangle_{n \geq 0}$  is a filtration on a ring  $R$ , then the collection of all cosets  $\{x + I_n : x \in R, n \geq 0\}$  is a basis for a topology on  $R$ . Indeed, it is clear that the union is all of  $R$ . Suppose that  $z \in (x + I_n) \cap (y + I_m)$ . Then, check that

$$z \in (z + I_{n+m}) \subseteq (x + I_n) \cap (y + I_m).$$

**Definition 3.2.6.** Given a filtration  $\langle I_n \rangle_{n \geq 0}$  on  $R$ , the topology generated by the basis  $\{x + I_n : x \in R, n \geq 0\}$  is called the **topology induced** by  $\langle I_n \rangle_{n \geq 0}$ .

Similarly, given a filtration  $\langle M_n \rangle_{n \geq 0}$  on an  $R$ -module  $M$ , one gives  $M$  a topology.

This is the topology that we will be focusing on now.

**Proposition 3.2.7.** Let  $\langle I_n \rangle_{n \geq 0}$  be a filtration on  $R$ . Then,  $R$  is a topological ring under the induced topology. (Similarly,  $M$  is a topological  $R$ -module.)

*Proof.* For  $x, y \in R$  and  $n \geq 0$ , note that

$$\begin{aligned} (x + I_n) + (y + I_n) &\subseteq x + y + I_n, \\ (x + I_n) \cdot (y + I_n) &\subseteq xy + I_n, \\ -(x + I_n) &= -x + I_n. \end{aligned}$$

The continuity of the three operations now follows. The reader may formulate the definition of a topological module and prove the result similarly.  $\square$

**Observation 3.2.8.** Note that the  $I_n$  are clopen subsets of  $R$ . Indeed, each  $I_n$  by virtue of being a basis element. On the other hand, we can write the complement as a union of open sets as well:  $R \setminus I_n = \bigcup_{x \notin I_n} (x + I_n)$ .

Similarly, each  $x + I_n$  is a clopen subset.

**Example 3.2.9.** We consider the  $m$ -adic topology on  $R$  for the following cases.

1.  $R = \mathbb{Z}$ ,  $m = (p)$  for some prime  $p$ .  
In this topology,  $x$  and  $y$  are “close” if  $x - y$  is divisible by a “large” power of  $p$ .
2.  $R = k[x, y]$ ,  $m = (x, y)$ .  
Here,  $f$  and  $g$  are “close” if  $f - g \in m^i$  for “large”  $i$  (i.e., all monomials appearing in  $f - g$  have degree  $\geq i$ ).

**Proposition 3.2.10.**  $\langle I_n \rangle_{n \geq 0}$  induces a Hausdorff topology iff  $\bigcap_{n \geq 0} I_n = 0$ . Similarly, a filtered  $R$ -module  $M$  is Hausdorff iff  $\bigcap_{n \geq 0} M_n = 0$ .

*Proof.* We prove the statement about rings and leave the other one to the reader.

( $\Rightarrow$ ) Suppose  $\bigcap_{n \geq 0} I_n \neq 0$ . Pick a nonzero  $y$  in the intersection. Then,  $y$  and  $0$  cannot be separated.

( $\Leftarrow$ ) As noted, each  $I_n$  is closed. Thus, the intersection being  $0$  implies that  $\{0\}$  is closed. By continuity of multiplication, the diagonal is then closed in  $R \times R$ , which is equivalent to  $R$  being Hausdorff.  $\square$

An alternate proof for ( $\Leftarrow$ ) direction: Let  $x, y \in R$  be distinct. Pick  $N$  large such that  $x - y \notin I_N$ . Then,  $x + I_N$  and  $y + I_N$  are disjoint neighbourhoods.

**Proposition 3.2.11.** Let  $M$  be a filtered  $R$ -module, and  $N \leq M$  a submodule. Then,

$$\overline{N} = \bigcap_{n \geq 0} (N + M_n).$$

*Proof.* Let  $x \in M$ .  $x \in \overline{N}$  iff every basic neighbourhood of  $x$  intersects  $N$  iff  $(x + M_n) \cap N \neq \emptyset$  for all  $n$ . This happens iff  $x \in N + M_n$  for all  $n$ .  $\square$

### §§3.3. The Artin-Rees Lemma

We wish to prove the Artin-Rees Lemma. Loosely speaking, it tells us the following (under Noetherian hypotheses): Suppose  $N \subseteq M$  is a submodule, and  $I \subseteq R$  an ideal. Give  $M$  the  $I$ -adic topology. Now, we can give  $N$  a topology in two ways: either the subspace topology, or the  $I$ -adic topology. The Artin-Rees lemma gives us that the two are the same.

We will assume the following setup for this subsection.

**Setup.**  $R$  is a Noetherian ring.  $I \subseteq R$  is an ideal, and  $R$  is given the  $I$ -adic filtration.  $M$  is a finitely generated  $R$ -module with some filtration  $\langle M_n \rangle_{n \geq 0}$ .  $N \subseteq M$  is an  $R$ -submodule.  
 $S := R[It]$  is the Rees algebra (as defined in Example 3.1.5).

**Definition 3.3.1.** The filtration  $\langle M_n \rangle_{n \geq 0}$  is said to be  $I$ -good if  $IM_n = M_{n+1}$  for  $n \gg 0$ .

**Lemma 3.3.2.**  $\langle M_n \rangle_{n \geq 0}$  is  $I$ -good iff  $F = \bigoplus_{n \geq 0} M_n t^n$  is a finitely generated  $S$ -module.

As before, the variable  $t^n$  in the direct sum above is for bookkeeping. This definition makes  $F$  a graded  $S$ -module. (Indeed, the definition of filtration implies that  $I_n t^n M_m t^m \subseteq M_{n+m} t^{n+m}$ .)

*Proof.* ( $\Leftarrow$ ) Suppose  $F$  is a finitely generated  $S$ -module. Note that  $S$  is finitely generated in degree 1. Thus, Porism 3.1.21 tells us that  $S_1 M_n = M_{n+1}$  for  $n \gg 0$ . This translates to  $IM_n = M_{n+1}$  for  $n \gg 0$ , which is precisely being  $I$ -good.

( $\Rightarrow$ ) Let  $n_0$  be such that  $IM_n = M_{n+1}$  holds for all  $n \geq n_0$ .

Now, each of  $M_0, \dots, M_{n_0}$  is finitely generated (over  $R = S_0$ ) and the above equation shows that using generators for these is enough to generate everything.  $\square$



**Lemma 3.3.3.**  $S$  is Noetherian.

*Proof.* Note that  $S_0 = R$  is Noetherian, and  $S$  is a finitely generated  $R$ -algebra. (Indeed, if  $x_1, \dots, x_n$  generate  $I$  as an  $R$ -ideal, then  $x_1 t, \dots, x_n t$  generate  $S$  as an  $R_0$ -algebra.)  $\square$

**Theorem 3.3.4** (Artin-Rees Lemma). Let  $R$  be a Noetherian ring,  $I \subseteq R$  an ideal,  $M$  a finitely generated  $R$ -module, and  $N \subseteq M$  a submodule.

Then,

- (i) there exists  $c \geq 1$  such that  $I^{n+c}M \cap N = I^n(I^c M \cap N)$  for all  $n \geq 1$ ;
- (ii) there exists  $c \geq 1$  such that  $I^{n+c}M \cap N \subseteq I^n N$  for all  $n \geq 1$ ;
- (iii) the subspace topology on  $N$  (when  $M$  is given the  $I$ -adic topology) agrees with the  $I$ -adic topology on  $N$ .

In the statement of (ii), we can add some more obvious inclusions:

$$I^{n+c}M \subseteq I^{n+c}M \cap N \subseteq I^n N \subseteq I^n M.$$

This shows how (ii) implies (iii). It is also clear that (i) implies (ii). So we shall only prove (i).

*Proof.* Note that (i) is equivalent to proving the filtration  $\langle I^n M \cap N \rangle_{n \geq 0}$  on  $N$  is  $I$ -good. Clearly, the  $I$ -adic filtration on  $M$  is  $I$ -good. Thus,  $\bigoplus_{n \geq 0} I^n M$  is a finitely generated  $S$ -module (Lemma 3.3.2).

Since  $S$  is Noetherian (Lemma 3.3.3), it follows that the submodule  $\bigoplus_{n \geq 0} (I^n M \cap N)$  is finitely generated.

Now, using Lemma 3.3.2 again shows that  $\langle I^n M \cap N \rangle_{n \geq 0}$  is  $I$ -good, as desired.  $\square$

### §§3.4. Krull's intersection theorems

We now use the Artin-Rees lemma to deduce that certain infinite intersections are zero. In view of Proposition 3.2.10, it is saying that a certain module is Hausdorff.

**Theorem 3.4.1** (Krull's Intersection Theorem). Let  $(R, \mathfrak{m})$  be a local Noetherian ring, and  $I \subsetneq R$  a proper ideal. If  $M$  is a finitely generated  $R$ -module, then

$$\bigcap_{n \geq 0} I^n M = 0.$$

That is, the  $I$ -adic topology on  $M$  is Hausdorff.

In particular,  $\bigcap_{n \geq 0} I^n = 0$ .

A special case is  $I = \mathfrak{m}$ .

*Proof.* Let  $N := \bigcap_{n \geq 0} \mathfrak{m}^n M$ . We wish to show that  $N = 0$ . Let  $c$  be as given by the **Artin-Rees Lemma 3.3.4**. Note that  $N \subseteq \mathfrak{m}^{1+c} M$ , by definition of  $N$ . Thus, we have

$$N \subseteq \mathfrak{m}^{1+c} M \cap N \subseteq \mathfrak{m} N.$$

Now, using NAK, we see that  $N = 0$ , as desired.  $\square$

*Aliter.* We use topological language in this proof. As before, let  $N$  be the intersection.

**Artin-Rees Lemma 3.3.4** tells us that the  $\mathfrak{m}$ -adic topology on  $N$  is the restriction of the  $\mathfrak{m}$ -adic topology on  $M$ . Note that  $N$  is contained in every  $\mathfrak{m}$ -adic neighbourhood of  $0$  in  $M$ . Thus, only neighbourhood of  $0$  in  $N$  is  $N$  itself.

It follows that  $N$  has the indiscrete topology. But  $\mathfrak{m}N$  is a nonempty open subset in the  $\mathfrak{m}$ -adic topology. Thus,  $N = \mathfrak{m}N$  and hence,  $N = 0$  by NAK.  $\square$

**Porism 3.4.2.** If  $R$  is a commutative ring,  $M$  a finitely generated  $R$ -module, and  $I$  is contained in the Jacobson radical of  $R$ , then  $\bigcap_{n \geq 0} I^n M = 0$ .

**Corollary 3.4.3.** Let  $R$  be a Noetherian ring,  $M$  a finitely generated  $R$ -module, and  $I$  be an ideal contained in the Jacobson radical. Then, every submodule  $N \leq M$  is closed in the  $I$ -adic topology. In particular, every ideal of  $R$  is closed in the  $I$ -adic topology.

*Proof.*  $M/N$  is a Hausdorff space in the  $I$ -adic topology. In particular,  $\{0\}$  is closed. The map  $\pi : M \rightarrow M/N$  is continuous (both have the  $I$ -adic topology). Thus,  $N = \pi^{-1}(0)$  is closed.  $\square$

**Theorem 3.4.4.** Let  $R$  be a Noetherian integral domain (not necessarily local).

Let  $I \subsetneq R$  be a proper ideal. Then,

$$\bigcap_{n \geq 0} I^n = 0.$$

*Proof.* Let  $\mathfrak{m} \supseteq I$  be maximal. Since  $R$  is a domain, have the inclusion  $R \hookrightarrow R_{\mathfrak{m}}$ . Thinking of  $R$  and  $R_{\mathfrak{m}}$  as subrings of the fraction field, we note the containments

$$\begin{aligned} \bigcap_{n \geq 0} I^n &\subseteq \left( \bigcap_{n \geq 0} I^n \right) R_{\mathfrak{m}} \\ &\subseteq \bigcap_{n \geq 0} (IR_{\mathfrak{m}})^n = 0, \end{aligned}$$

where the last equality follows from Theorem 3.4.1, since  $R_{\mathfrak{m}}$  is local.  $\square$

**Example 3.4.5.** The theorems breaks down if drop both the hypotheses of being local and integral domain.

Indeed, take any ring  $R$  with a nontrivial idempotent  $e$ . Then, the ideal  $(e)$  is an idempotent ideal and hence  $\bigcap_{n \geq 0} (e)^n = (e) \neq 0$ . (Note that this is possible when  $R$  is Noetherian, finite even.)

If  $R$  is not assumed Noetherian, then one cannot conclude even with the integral domain and local hypothesis. Consider  $R = \mathbb{Z}_{(p)}[p^{1/p}, p^{1/p^2}, \dots]$ .

Then,  $R$  is a local domain with maximal ideal  $\mathfrak{m} = (p^{1/p}, p^{1/p^2}, \dots)$ . But  $\mathfrak{m}$  is idempotent.

## §4. Dimension Theory

We may tacitly be assuming that  $M \neq 0$  in many places, when we talk about dimension. It may be safe to assume that we are defining  $\dim(M) = -1$  when  $M = 0$ .

### §§4.1. Integer valued polynomials

As before, we continue with the notation that  $0 \in \mathbb{N}$ . We use  $\mathbb{N}_{\geq 1}$  to denote the set of positive integers.

**Definition 4.1.1.** A function  $f : \mathbb{Z} \rightarrow \mathbb{Z}$  is an **eventual polynomial** if there exists a polynomial  $P(t) \in \mathbb{Q}[t]$  such that

$$f(n) = P(n) \quad \text{for all } n \gg 0.$$

The **degree** of  $f$  is defined to be the degree of  $P$ .

Note that the polynomial  $P$  is uniquely determined. For if  $Q$  is another such polynomial, then  $P - Q$  vanishes at infinitely many points. Thus, the degree is well-defined.

**Definition 4.1.2.** For a function  $f : \mathbb{Z} \rightarrow \mathbb{Z}$ , we define the **difference function**  $\Delta f : \mathbb{N}_{\geq 1} \rightarrow \mathbb{Z}$  by

$$(\Delta f)(n) := f(n) - f(n-1).$$

**Proposition 4.1.3.** Let  $f : \mathbb{Z} \rightarrow \mathbb{Z}$  be any function. The following are equivalent:

1.  $f$  is eventually polynomial.
2.  $\Delta f$  is eventually polynomial.

In this case, we have  $\deg(f) = \deg(\Delta f) + 1$ .

We are not worrying about degree of the zero polynomial. The reader can take care of that.

*Proof.* If  $f$  agrees with  $P$  eventually, then  $\Delta f$  agrees with  $\Delta P$  eventually. In this case, we clearly have  $\deg(f) = \deg(\Delta f)$ .

Now suppose that  $\Delta f$  is eventually a polynomial. Assume  $Q(t) \in \mathbb{Q}[t]$  and  $N > 1$  are such that

$$\Delta f(n) = Q(n)$$

for  $n \geq N$ .

For  $n > N$ , note that we have  $f(n) = f(n-1) + \Delta f(n)$ . Adding such equalities gives us

$$f(n) = f(N) + \sum_{k=N}^{n-1} \Delta f(k) = f(N) + \sum_{k=N}^{n-1} Q(k).$$

Thus, for a constant  $C \in \mathbb{Z}$ , we can write

$$f(n) = C + \sum_{k=0}^{n-1} Q(k).$$

But now note that the sum is itself a polynomial of degree  $\deg(Q) + 1$ . This finishes the proof.  $\square$

**Definition 4.1.4.** For  $n \geq 0$ , define the rational polynomial  $\binom{t}{n} \in \mathbb{Q}[t]$  as

$$\binom{t}{n} := \frac{t(t-1) \cdots (t-(n-1))}{n!}.$$

Note that  $\binom{t}{n}$  is a polynomial with (non-integer) rational coefficients, but takes integer values at all integers. We show that these polynomials are essentially all.

**Proposition 4.1.5.** Let  $P(t) \in \mathbb{Q}[t]$ . The following are equivalent:

- (i)  $P(t) = \sum_n c_n \binom{t}{n}$  for some sequence of integers  $\langle c_n \rangle_{n \geq 0}$  eventually zero.
- (ii)  $P(n) \in \mathbb{Z}$  for all  $n \in \mathbb{Z}$ .
- (iii)  $P(n) \in \mathbb{Z}$  for all  $n \gg 0$ .

*Proof.* (i)  $\Rightarrow$  (ii)  $\Rightarrow$  (iii) is clear.

(iii)  $\Rightarrow$  (ii): Write  $P(x) = Q(x)/M$  for  $Q(x) \in \mathbb{Z}[x]$  and  $M \in \mathbb{N}_{\geq 1}$ .

If there exists  $n \in \mathbb{Z}$  such that  $M$  does not divide  $Q(n)$ , then  $M$  does not divide  $Q(n+M)$  either.

In other words, if  $P(n) \notin \mathbb{Z}$ , then  $P(n+M), P(n+2M), \dots$  are also not in  $\mathbb{Z}$ .

(ii)  $\Rightarrow$  (i): Let  $d := \deg(P)$ . By linear algebra, we can write

$$P(t) = \sum_{k=0}^d a_k \binom{t}{k},$$

for some choice of rationals  $a_k \in \mathbb{Q}$ .

Now, evaluating the above sequentially at  $t = 0, \dots, d$  shows that each  $a_k$  is an integer. (Note that for such  $t$ ,  $\binom{t}{k}$  vanishes when  $k > t$  and equals 1 when  $k = t$ .)  $\square$

**Remark 4.1.6.** The above holds even if we started with  $P(t) \in \mathbb{R}[t]$  or  $\mathbb{C}[t]$ . Lagrange interpolation immediately tells us that such a  $P$  must have rational coefficients.

## §§4.2. Embedding dimension

**Definition 4.2.1.** Let  $(R, \mathfrak{m}, k)$  be a local Noetherian ring. The **embedding dimension** of  $R$  is defined as

$$\text{emb. dim}(R) := \dim_k(\mathfrak{m}/\mathfrak{m}^2).$$

By Nakayama's lemma, the above is equal to the minimum number of generators needed to generate  $\mathfrak{m}$  (as an  $R$ -ideal). This is not a very good notion. We try capture information by looking at all quotients of the form  $\mathfrak{m}^k/\mathfrak{m}^{k+1}$ .

## §§4.3. Dimension à la Hilbert polynomials

We now define some functions, which we show are eventually polynomial. Then we define dimension using the degree.

**Definition 4.3.1.** Let  $(R, \mathfrak{m}, k)$  be a local Noetherian ring. The function

$$n \mapsto \ell(R/\mathfrak{m}^n)$$

is eventually polynomial. This function is called the **Hilbert-Samuel function**.

The corresponding polynomial is called the **Hilbert-Samuel polynomial**. Its degree is defined to be the **dimension** of  $R$ , denoted  $\dim(R)$ .

More generally, if  $M \neq 0$  is a finitely generated  $R$ -module, then

$$n \mapsto \ell(M/\mathfrak{m}^n M)$$

is eventually polynomial, whose degree defines  $\dim(M)$ . We define the **Hilbert-Samuel polynomial**  $H_M(t) \in \mathbb{Q}[t]$  to be the polynomial such that  $H_M(n) = \ell(M/\mathfrak{m}^n M)$  for  $n \gg 0$ .

Moreover,  $\dim(M) \leq \text{emb. dim}(R)$ . (In particular,  $\dim(R) \leq \text{emb. dim}(R)$ .)

These assertions are Proposition 4.3.7.

In the above,  $\ell$  denotes the length of the  $R$ -module. Note that  $M/\mathfrak{m}^n M$  is annihilated by  $\mathfrak{m}^n$  and is hence, Artinian (Corollary 1.2.9) and therefore has finite length. We also have

$$\ell(M/\mathfrak{m}^n) = \sum_{i=0}^{n-1} \dim_k(\mathfrak{m}^i M / \mathfrak{m}^{i+1} M).$$

**Definition 4.3.2.** Let  $k$  be a field, and  $M \neq 0$  be a finitely generated graded module over the polynomial ring  $k[x_1, \dots, x_r]$ . Then, the **Hilbert function**, defined by

$$n \mapsto \sum_{i \leq n} \dim_k(M_i)$$

is eventually a polynomial of degree  $\leq r$  (Proposition 4.3.5).

The corresponding polynomial is denoted by  $f_M^+(t) \in \mathbb{Q}[t]$ , and is called the **Hilbert polynomial**. We also define  $f_M := \Delta f_M^+$ .

Note that in the above, it makes sense to talk about  $\dim_k(M_i)$  since  $k$  is in fact a subring of the polynomial ring. (As opposed to earlier, when we cannot talk about  $\dim_k(R/\mathfrak{m}^n)$ .) Moreover, these dimensions are finite, by Proposition 3.1.18.

**Example 4.3.3** (Dimension of a field). Let  $k$  be a field (and hence, a local ring), and  $M \neq 0$  be a finitely generated  $k$ -module. Then,  $\dim(M) = 0$ . In particular,  $\dim(k) = 0$ . (Note the contrast from the usual vector space dimension.)

To see this, note that the maximal ideal here is 0. Thus,  $\ell(M/\mathfrak{m}^n M) = \ell(M)$  for all  $n$ . Thus, the Hilbert-Samuel polynomial is constant, i.e., its degree is zero.

**Example 4.3.4** (Dimension of a polynomial ring). Let  $k$  be a field, and  $R = k[x_1, \dots, x_d]_{(x_1, \dots, x_d)}$ .

Note that  $R$  is a local ring with maximal ideal  $\mathfrak{m} = (x_1, \dots, x_d)$ . We claim that  $\dim(R) = d$ .

To see this, note that  $\mathfrak{m}^n/\mathfrak{m}^{n+1}$  has a  $k$ -basis given by (the images of) the monomials of degree  $n$ . (Here we are looking at  $k$  as the quotient  $R/\mathfrak{m}$  and not the subring  $k \subseteq R$ .)

Thus,

$$\dim_k(\mathfrak{m}^n/\mathfrak{m}^{n+1}) = \binom{d+n-1}{d-1}.$$

Thus, the above is a polynomial (in  $n$ ) of degree  $d-1$ . In turn,  $\ell(M/\mathfrak{m}^n)$  is a polynomial of degree  $d$ .

**Proposition 4.3.5.** The assertions made in Definition 4.3.2 hold.

*Proof.* Define  $g_M : \mathbb{Z} \rightarrow \mathbb{Z}$  to be  $g_M(n) = \dim_k(M_n)$ . Note that this is the difference function of the one given in the definition. Thus, by Proposition 4.1.3, it suffices to show that  $g_M$  is eventually polynomial.

We repeatedly use the following fact: If

$$0 \rightarrow M' \rightarrow M \rightarrow M'' \rightarrow 0$$

is an exact sequence of *graded*  $R$ -modules, then  $g_M = g_{M'} + g_{M''}$ .

Note that as a consequence, we get that if there is a filtration

$$0 = M_0 \subseteq M_1 \subseteq \cdots \subseteq M_t,$$

then

$$g_{M_t} = g_{M_t/M_{t-1}} + \cdots + g_{M_1/M_0}.$$

In particular, if all the functions on the right are eventually polynomials of degree  $d$ , then the same is true for  $g_{M_t}$ . ( $t$  here is fixed.)

Another trick that we will use is that we will consider modules annihilated by  $x_r$ , so that we can view them as modules over a smaller polynomial ring. Note that the function  $g_M$  does not change.

We prove the statement by induction on  $r$ .

If  $r = 0$ , then the statement is clear since  $g_M$  is eventually zero. Thus,  $f_M^+$  is a constant polynomial.

Since  $x_r$  is homogeneous, we have a chain of graded submodules of  $M$  given by

$$0 \subseteq \text{ann}_M(x_r) \subseteq \text{ann}_M(x_r^2) \subseteq \cdots$$

Since  $M$  is Noetherian, then above stabilises to some  $M'$ .

Note that each  $\text{ann}_M(x_r^i) / \text{ann}_M(x_r^{i+1})$  is a finitely generated module over the polynomial ring in  $r - 1$  variables. By induction, each of these modules have a Hilbert polynomial<sup>3</sup> of degree  $\leq r - 1$ . Thus, by our observation about filtrations, the same is true for  $M'$ .

Thus, we have shown that  $g_{M'}$  is eventually a polynomial of degree  $\leq r - 1$ . Since we have the graded exact sequence

$$0 \rightarrow M' \rightarrow M \rightarrow M/M' \rightarrow 0,$$

it suffices to prove that  $g_{M/M'}$  is eventually a polynomial of degree  $\leq r$ . Thus, we may replace  $M$  with  $M/M'$ . The gain is that  $x_r$  is a nonzerodivisor on  $M/M'$ .

Now, note that we have a graded exact sequence

$$0 \rightarrow M(-1) \xrightarrow{x_r} M \rightarrow M/(x_r) \rightarrow 0.$$

Thus, we have

$$g_M - g_{M(-1)} = g_{M/(x_r)}.$$

Again, by induction, the right function is eventually polynomial of degree  $\leq r - 1$ . Moreover the left function is exactly  $\Delta g_M$ . Thus, Proposition 4.1.3 tells us that  $g_M$  is eventually a polynomial of degree  $\leq r$ .  $\square$

<sup>3</sup>By this, we mean that the corresponding function is eventually polynomial for them.



**Remark 4.3.6.** Note that in the above proof, we did not really use that  $k[x_1, \dots, x_r]$  is a polynomial ring. Rather, all we needed was a finitely generated graded  $k$ -algebra, with generators in degree 1. (The proof would have to be slightly modified if the generators are homogeneous of different degrees.)

**Proposition 4.3.7.** The assertions made in Definition 4.3.1 hold.

*Proof.* We prove the result for modules. Consider the associated graded ring  $\text{gr}(R) = \bigoplus_{n \geq 0} \mathfrak{m}^n / \mathfrak{m}^{n+1}$  and the associated graded module  $\text{gr}(M) = \bigoplus_{n \geq 0} \mathfrak{m}^n M / \mathfrak{m}^{n+1} M$ .

Let  $r = \text{emb. dim}(R)$ . Then, we can generate  $\mathfrak{m}$  by some  $t_1, \dots, t_r \in \mathfrak{m}$ . Note that  $\text{gr}(R)_0 = R/\mathfrak{m} = k$ . We have a surjective map of graded  $k$ -algebras

$$k[x_1, \dots, x_r] \rightarrow \text{gr}(R)$$

given by

$$x_i \mapsto \bar{t}_i.$$

(Note that  $x_i$  and  $\bar{t}_i \in \mathfrak{m}/\mathfrak{m}^2$  both do have degree 1.)

Thus,  $\text{gr}(M)$  is a finitely generated graded  $k[x_1, \dots, x_r]$ -module. Now, by Proposition 4.3.5, the function

$$n \mapsto \dim_k(\mathfrak{m}^n M / \mathfrak{m}^{n+1} M)$$

is eventually polynomial, of degree  $\leq r - 1$ .

In turn, the Hilbert-Samuel function

$$n \mapsto \sum_{i=0}^{n-1} \dim_k(\mathfrak{m}^i M / \mathfrak{m}^{i+1} M)$$

is eventually a polynomial of degree  $\leq r$ . □

**Example 4.3.8.** Let  $(R, \mathfrak{m})$  be a local Noetherian ring and  $M$  a finitely generated  $R$ -module. Then,  $\dim(M) = 0$  iff  $M$  is of finite length (iff  $M$  is Artinian).

Indeed, if  $\dim(M) = 0$ , then the Hilbert-Samuel polynomial is constant. This means that  $\mathfrak{m}^n M = \mathfrak{m}^{n+1} M$  for  $n \gg 0$ . Nakayama's lemma now implies that  $\mathfrak{m}^n M = 0$  for  $n \gg 0$ . In turn, Corollary 1.2.9 tells us that  $M$  is Artinian.

Conversely, if  $M$  is Artinian, then the chain

$$M \supseteq \mathfrak{m}M \supseteq \mathfrak{m}^2 M \supseteq \dots$$

eventually stabilises. Thus,  $\ell(\mathfrak{m}^n M / \mathfrak{m}^{n+1} M) = 0$  for  $n \gg 0$ .

### §§4.4. Properties of dimension

To summarise, we have shown the following:

- If  $M$  is a finitely generated module over  $(R, \mathfrak{m})$ , then we have a polynomial  $H_M$  such that  $H_M(n) = \ell(M/\mathfrak{m}^n M)$  for large  $n$ .
- If  $M$  is a finitely generated graded module over  $k[x_1, \dots, x_r]$ , then there are polynomials  $f_M, f_M^+$  such that

$$f_M(n) = \dim_k(M_n) \quad \text{and} \quad f_M^+(n) = \sum_{i \leq n} \dim_k(M_i)$$

for large  $n$ .

We now wish to see how the degrees of these polynomials interact with exact sequences. It is direct for the graded case. The local case requires more work.

**Proposition 4.4.1.** Suppose we have an exact sequence

$$0 \rightarrow M' \rightarrow M \rightarrow M'' \rightarrow 0$$

of graded modules over  $k[x_1, \dots, x_r]$ . Then,

$$f_M = f_{M'} + f_{M''} \quad \text{and} \quad f_M^+ = f_{M'}^+ + f_{M''}^+.$$

Consequently,  $\deg(f_M) = \max(\deg(f_{M'}), \deg(f_{M''}))$ .

*Proof.* The first result about the sum is clear since the exact sequence is graded, giving an exact sequence of  $k$ -vector space.

The statement about degree follows since the leading coefficients of the concerned polynomials are positive.  $\square$

**Proposition 4.4.2.** Let  $(R, \mathfrak{m})$  be a Noetherian local ring. Let

$$0 \rightarrow M' \rightarrow M \rightarrow M'' \rightarrow 0$$

be an exact sequence of finite  $R$ -modules. Then,  $\dim(M) = \max(\dim(M'), \dim(M''))$ .

*Proof.* Note that since  $\otimes_R R/\mathfrak{m}^n$  is a right-exact functor, we have the exact sequence

$$M'/\mathfrak{m}^n M' \rightarrow M/\mathfrak{m}^n M \rightarrow M''/\mathfrak{m}^n M'' \rightarrow 0.$$

We show that the genuine kernel is a “good approximation” for the leftmost term there. To this end, let  $M'_n := M'/\mathfrak{m}^n M'$ , and let  $K_n$  be the genuine kernel, i.e., the sequence

$$0 \rightarrow K_n \rightarrow M/\mathfrak{m}^n M \rightarrow M''/\mathfrak{m}^n M'' \rightarrow 0 \tag{4.1}$$

is exact.

Note that  $K_n \cong (M' + \mathfrak{m}^n M) / \mathfrak{m}^n M \cong M' / (M' \cap \mathfrak{m}^n M)$ . In particular, we have a surjection  $M'_n \twoheadrightarrow K_n$  and thus,  $\ell(K_n) \leq \ell(M'_n)$ .

On the other hand, **Artin-Rees Lemma 3.3.4** implies the existence of  $c \geq 1$  such that

$$M' \cap \mathfrak{m}^n M \subseteq \mathfrak{m}^{n-c} M'$$

for all  $n \geq c$ .

Thus, for  $n \geq c$ , we have

$$\ell(M'_{n-c}) \leq \ell(K_n) \leq \ell(M'_n). \quad (4.2)$$

Note that (4.1) tells us that the Hilbert polynomials satisfy

$$H_M(n) = H_{M''}(n) + \ell(K_n) \quad (4.3)$$

for all  $n$ .

This implies that  $n \mapsto \ell(K_n)$  is eventually a polynomial. (4.2) shows that  $H_{M'}$  has the same degree as this. The result now follows from (4.3) since the leading coefficients of the polynomials in questions are positive.  $\square$

**Corollary 4.4.3.** If  $M$  is a Noetherian module over a Noetherian local ring, then  $\dim(M^{\oplus n}) = \dim(M)$  for all  $n \geq 1$ .

**Theorem 4.4.4.** Let  $R$  be a local Noetherian ring. Then,  $\dim(R) = \dim(R/\text{nil}(R))$ .

$\text{nil}(R)$  above denotes the ideal of nilpotents.

*Proof.* Let  $I = \text{nil}(R)$ . Since  $R$  is Noetherian, we have  $I^n = 0$  for  $n \gg 0$ . Thus, it suffices to prove that

$$\dim(R/I^k) = \dim(R/I^{k+1})$$

for all  $k \geq 1$ .

Note that we have an exact sequence

$$0 \rightarrow I^k/I^{k+1} \rightarrow R/I^{k+1} \rightarrow R/I^k \rightarrow 0$$

of  $R$ -modules.

In view of Proposition 4.4.2, it suffices to show that  $\dim(I^k/I^{k+1}) \leq \dim(R/I^{k+1})$ .

Note that  $I^k/I^{k+1}$  is a (finitely generated)  $R/I^k$ -module. Thus, we have a surjection of the form

$$\bigoplus_{i=1}^N R/I^k \rightarrow I^k/I^{k+1} \rightarrow 0.$$

Corollary 4.4.3 and Proposition 4.4.2 now imply that  $\dim(R/I^k) \geq \dim(I^k/I^{k+1})$ .  $\square$

**Theorem 4.4.5** (Dimension only depends on support). Let  $(R, \mathfrak{m})$  be a local Noetherian, and  $M \neq 0$  a finitely generated  $R$ -module. Then,

$$\dim(M) = \max_{\substack{\mathfrak{p} \in \text{Supp}(M) \\ \mathfrak{p} \text{ minimal}}} \dim(R/\mathfrak{p}).$$

In particular,  $\dim(R) = \max_{\mathfrak{p}} \dim(R/\mathfrak{p})$ , where  $\mathfrak{p}$  runs over all minimal primes.

*Proof.* There exists a filtration of  $M$  of the form

$$0 = M_0 \subseteq M_1 \subseteq \cdots \subseteq M_m = M$$

such that each  $M_{i+1}/M_i$  is isomorphic to some  $R/\mathfrak{p}_i$ , with  $\mathfrak{p}_i \in \text{Supp}(M)$ .

Using Proposition 4.4.2 successively on exact sequences of the form  $0 \rightarrow M_i \rightarrow M_{i+1} \rightarrow M_{i+1}/M_i \rightarrow 0$ , we see that

$$\dim(M) = \max_i \dim(R/\mathfrak{p}_i) \leq \sup_{\mathfrak{p} \in \text{Supp } M} \dim(R/\mathfrak{p}).$$

To prove the reverse inequality, we fix  $\mathfrak{p} \in \text{Supp}(M)$  and show that  $\dim(R/\mathfrak{p}) \leq \dim(M)$ .

Since  $M_{\mathfrak{p}} \neq 0$  and localisation is exact, we must have that  $(R/\mathfrak{p}_i)_{\mathfrak{p}} \neq 0$  for some  $i$ <sup>4</sup> and hence,  $\mathfrak{p}_i \subseteq \mathfrak{p}$ . This gives us  $\dim(R/\mathfrak{p}) \leq \dim(R/\mathfrak{p}_i)$ .

This also shows that the supremum is indeed a maximum.

To see that the maximum can be taken over minimal primes, first note that we clearly have

$$\max_{\mathfrak{p} \in \text{Supp } M} \dim(R/\mathfrak{p}) \geq \max_{\substack{\mathfrak{p} \in \text{Supp}(M) \\ \mathfrak{p} \text{ minimal}}} \dim(R/\mathfrak{p}).$$

For the other direction, note that if  $\mathfrak{p} \in \text{Supp } M$ , then  $\mathfrak{p} \supseteq \mathfrak{q}$  for some minimal  $\mathfrak{q} \in \text{Supp } M$  and  $\dim(R/\mathfrak{p}) \leq \dim(R/\mathfrak{q})$ .  $\square$

**Corollary 4.4.6.** With the same hypothesis,  $\dim(M) = \dim(R/\text{ann}(M))$ . In particular,  $\dim(M) \leq \dim(R)$ .

*Proof.* It suffices to show that  $\text{Supp}(M) = \text{Supp}(R/\text{ann}(M))$ . But note that  $\text{Supp}(M) = V(\text{ann}(M))$ , since  $M$  is finitely generated. But  $V(\text{ann}(M)) = \text{Supp}(R/\text{ann}(M))$ .  $\square$

<sup>4</sup>Indeed, localise the filtration  $M_0 \subseteq \cdots \subseteq M_m$  at  $\mathfrak{p}$ . Since  $(M_m)_{\mathfrak{p}} \neq 0$ , there is some  $i$  with  $(M_i)_{\mathfrak{p}} \neq (M_{i+1})_{\mathfrak{p}}$ .

**Theorem 4.4.7.** Let  $(R, \mathfrak{m})$  be local, and  $M \neq 0$  be a finitely generated  $R$ -module. Let  $x \in \mathfrak{m}$  be a nonzerodivisor on  $M$ . Then,

$$\dim(M/xM) = \dim(M) - 1.$$

*Proof.* We have the exact sequence

$$0 \rightarrow xM \rightarrow M \rightarrow M/xM \rightarrow 0.$$

Thus,

$$0 \rightarrow xM/(xM \cap \mathfrak{m}^n M) \rightarrow M/\mathfrak{m}^n M \rightarrow M/(xM + \mathfrak{m}^n M) \rightarrow 0$$

is an exact sequence for all  $n$ . In terms of the Hilbert polynomials, we see that

$$H_M(n) = H_{M/xM}(n) + \ell(xM/(xM \cap \mathfrak{m}^n M)) \quad (4.4)$$

for  $n \gg 0$ . In particular, the rightmost term is eventually polynomial.

Note that since  $x$  is a nonzerodivisor, we have  $xM \cong M$  and hence,

$$xM/(xM \cap \mathfrak{m}^n M) \cong M/N_n,$$

where  $N_n = (\mathfrak{m}^n M :_M x) = \{a \in M : xa \in \mathfrak{m}^n M\}$ . Indeed,  $N_n$  is the preimage of  $xM \cap \mathfrak{m}^n M$  under the isomorphism  $M \xrightarrow{x} xM$ .

Note that since  $x \in \mathfrak{m}$ , we have  $\mathfrak{m}^{n-1}M \subseteq N_n$  and hence,

$$\ell(M/N_n) \leq \ell(M/\mathfrak{m}^{n-1}M) \leq H_M(n-1)$$

for  $n \gg 0$ . Combining this with (4.4) gives

$$H_M(n) \leq H_{M/xM}(n) + H_M(n-1)$$

for  $n \gg 0$ . Thus,  $H_{M/xM} \geq \Delta H_M$  eventually and hence,  $\dim(M/xM) \geq \dim(M) - 1$ .

We now need to prove the other direction of the inequality.

The **Artin-Rees Lemma 3.3.4** tells us that there exists  $c$  such that  $N_{n+c} \subseteq \mathfrak{m}^n M$  for all  $n$ .<sup>5</sup> Thus,

$$\ell(M/N_n) \geq \ell(M/\mathfrak{m}^{n-c}M) = H_M(n-c)$$

for  $n \gg 0$ . Since  $M/N_n \cong xM/(xM \cap \mathfrak{m}^n M)$ , (4.4) now tells us that

$$H_M(n) \geq H_{M/xM}(n) + H_M(n-c)$$

for  $n \gg 0$ .

Thus,  $H_{M/xM}(n) \leq H_M(n) - H_M(n-c)$  and thus, the degree drops in the desired way giving us  $\dim(M/xM) \leq \dim(M) - 1$ .  $\square$

<sup>5</sup>Indeed, it tells us that there exists  $c$  such that  $\mathfrak{m}^{n+c}M \cap xM \subseteq \mathfrak{m}^n(xM)$  for all  $n$ . Now, take preimages under the isomorphism  $M \xrightarrow{x} xM$ .

Using the above, we can prove a weaker inequality when  $x$  is possibly a zerodivisor.

**Corollary 4.4.8.** Let  $M$  be a finitely generated module over the local Noetherian ring  $(R, \mathfrak{m})$ .

If  $x \in \mathfrak{m}$ , then

$$\dim(M) - 1 \leq \dim(M/xM) \leq \dim(M).$$

In words, the dimension can drop by at most 1.

*Proof.*  $\dim(M/xM) \leq \dim(M)$  is clear since  $M$  surjects onto  $M/xM$ .

Let  $N$  denote the  $x$ -torsion submodule of  $M$ , i.e.,  $N = \{a \in M : x^n a = 0 \text{ for some } n \geq 1\}$ . Let  $M'' = M/N$ . Then, we have an exact sequence

$$0 \rightarrow N \rightarrow M \rightarrow M'' \rightarrow 0.$$

Tensoring with  $R/x$  gives us an exact sequence

$$0 \rightarrow N/xN \rightarrow M/xM \rightarrow M''/xM'' \rightarrow 0.$$

We only need to check that  $N/xN \rightarrow M/xM$  is indeed an inclusion. This is equivalent to showing  $N \cap (xM) \subseteq xN$ . But this is clear.<sup>6</sup>

Now, note that  $\dim(M''/xM'') = \dim(M'') - 1$  since  $x$  is a nonzerodivisor on  $M''$ , by construction.

On the other hand,  $\dim(N/xN) = \dim(N)$  since dimension only depends on the support (Theorem 4.4.5), and  $\text{Supp}(N/xN) = \text{Supp}(N)$  since  $N$  is  $x$ -torsion.<sup>7</sup> We now apply Proposition 4.4.2 twice. We have

$$\begin{aligned} \dim(M/xM) &= \max(\dim(M''/xM''), \dim(N/xN)) \\ &= \max(\dim(M'') - 1, \dim(N)) \\ &\geq \max(\dim(M'') - 1, \dim(N) - 1) \\ &= \max(\dim(M''), \dim(N)) - 1 = \dim(M). \end{aligned} \quad \square$$

**Corollary 4.4.9.** Let  $(R, \mathfrak{m})$  be a (nonzero) local Noetherian ring of dimension zero. Then, every element of  $\mathfrak{m}$  is a zerodivisor. In particular, if  $R$  is a domain, then  $R$  is a field.

*Proof.* Let  $x \in \mathfrak{m}$  be arbitrary. If  $x$  is a nonzerodivisor, then

$$\dim(R) = \dim(R/xR) + 1 \geq 1,$$

a contradiction. □

<sup>6</sup>If  $a \in M$  and  $xa \in N$ , then  $x^n(xa) = 0$  for some  $n$  and hence,  $a \in N$ .

<sup>7</sup>In general,  $\text{Supp}(N/xN) = \text{Supp}(N) \cap V(x)$ . Now, since  $N$  is  $x$ -torsion, if  $N_p \neq 0$ , we must have  $x \in \mathfrak{p}$ . Thus,  $\text{Supp}(N) \subseteq V(x)$ .

Note that the above is a converse of sorts to Example 4.3.3. However, note that there do exist rings of dimension zero which are not fields. Indeed, Artinian local rings are precisely these zero dimensional Noetherian local rings.

We have already shown most of this in Corollary 1.2.13; we only need to show that the notion of  $\dim(R)$  is the same.

### §§4.5. Characterisation of dimension

For this discussion, assume that  $(R, \mathfrak{m}, k)$  is local Noetherian. We showed that the function  $n \mapsto \ell(R/\mathfrak{m}^n)$  is eventually polynomial, and defined  $\dim(R)$  to be the degree of this (unique) polynomial.

We also showed the following properties.

- (P1)  $\dim(R) = \max_{\mathfrak{p} \text{ a minimal prime}} \dim(R/\mathfrak{p})$ .  
(Note that  $R$  has only finitely many minimal primes.)
- (P2)  $\dim(R) = 0$  if  $R$  is a field (Example 4.3.3).
- (P3) If  $R$  is a domain, and  $x \in \mathfrak{m} \setminus \{0\}$ , then  $\dim(R/(x)) = \dim(R) - 1$ .

Note that (P3) above implies the following.

- (P3') If  $R$  is a domain which is not a field, then  $\dim(R) = \sup_{x \in \mathfrak{m} \setminus \{0\}} \dim(R/xR) + 1$ .

**Theorem 4.5.1.** (P1)-(P3) uniquely characterise the dimension function. In other words, if we are given a function

$$d : \{\text{local Noetherian rings}\} \rightarrow \mathbb{Z}_{\geq 0}$$

satisfying (P1)-(P3), then  $d = \dim$ .

The statement is true even if we replace (P3) with (P3').

(We are ignoring any set-theoretic issues and using the term “function”.)

*Proof.* Note that if  $d$  satisfies (P3), then it satisfies (P3'). Thus, we may assume that  $d$  satisfies (P1), (P2), (P3'), and show that  $d = \dim$ .

It suffices to show  $d(R) = \dim(R)$  whenever  $R$  is a domain. It then follows for a general ring by using (P1).

We prove this by induction on  $\dim(R)$ .

$\dim(R) = 0$ : In this case,  $R$  is a field, by Corollary 4.4.9. But then  $d(R) = 0$ , by (P2).

$\dim(R) > 0$ : Then,  $R$  is not a field (by (P2)). By (P3'), there exists  $x \in \mathfrak{m} \setminus \{0\}$  such that  $d(R) = d(R/xR) + 1$ .

Note that the ring  $R' = R/xR$  has dimension equal to  $\dim(R) - 1$ .<sup>8</sup> We wish to use the inductive hypothesis, but  $R'$  need not be a domain. But note that  $R'/\mathfrak{p}$  is, for every  $\mathfrak{p} \in \text{Spec}(R')$ .

Moreover,  $\dim(R'/\mathfrak{p}) \leq \dim(R') < \dim(R)$  due to (P1). Thus, the induction hypothesis applies to all such quotients. Now using (P1) again, we see

$$d(R') = \max_{\mathfrak{p} \text{ minimal}} d(R'/\mathfrak{p}) = \max_{\mathfrak{p} \text{ minimal}} \dim(R'/\mathfrak{p}) = \dim(R').$$

Thus,  $d(R) = d(R') + 1 = \dim(R') + 1 = \dim(R)$ . □

### §§4.6. Krull dimension

We now define a new notion of dimension that makes sense for any commutative ring (with the additional possibility of it being  $\infty$  in some cases). We will then show that when restricted to local Noetherian rings, this dimension function is finite and satisfies (P1), (P2), (P3'). This will show that the newly defined dimension agrees with the earlier definition.

**Definition 4.6.1.** Let  $R$  be any commutative ring. A **chain of prime ideals** in  $R$  is a finite sequence of prime ideals

$$\mathfrak{p}_0 \subsetneq \mathfrak{p}_1 \subsetneq \cdots \subsetneq \mathfrak{p}_n.$$

The above chain is said to have **length**  $n$ .

The **Krull dimension** of  $R$  is defined as

$$\text{krdim}(R) := \sup\{n : \text{there exists a chain of prime ideals of length } n\}.$$

**Example 4.6.2.** Corollary 1.2.13 told us that Artinian rings are precisely Noetherian rings with Krull dimension zero.

**Theorem 4.6.3.** If  $(R, \mathfrak{m})$  is a Noetherian local ring, then  $\text{krdim}(R) = \dim(R)$ .

*Proof.* We refer to (P1), (P2), (P3') from the previous section.

(P1): Any chain in  $R/\mathfrak{p}$  can be lifted back to  $R$  showing that  $\dim(R/\mathfrak{p}) \leq \dim(R)$  for all  $\mathfrak{p} \in \text{Spec}(R)$ .

Conversely, any chain of primes in  $R$  contains a minimal prime and then continues to be a chain in the quotient by that prime.

(P2): Obvious.

---

<sup>8</sup>We are allowed to use (P3) for  $\dim$ !



(P3'): Let  $R$  be a domain and not a field.

Let  $x \in \mathfrak{m} \setminus \{0\}$  be arbitrary. A chain of length  $n$  in  $R/xR$  can be lifted to a chain of length  $n$  in  $R$ . Moreover,  $(0)$  is a prime strictly contained in this chain. Thus, we have a chain of length  $n + 1$  in  $R$ . This shows

$$\text{krdim}(R) \geq \sup_{x \in \mathfrak{m} \setminus \{0\}} \text{krdim}(R/xR) + 1. \quad (4.5)$$

Conversely, if

$$0 = \mathfrak{p}_0 \subsetneq \mathfrak{p}_1 \subsetneq \cdots \subsetneq \mathfrak{p}_{m+1} = \mathfrak{m}$$

is a chain of length  $m + 1$  in  $R$ , then pick any nonzero  $x \in \mathfrak{p}_1$ . Then,

$$\mathfrak{p}_1/(x) \subsetneq \cdots \subsetneq \mathfrak{p}_{m+1}/(x)$$

is a chain of length  $m$  in  $R/xR$ , proving equality in (4.5).

To finish concluding, we must show that  $\text{krdim}(R) < \infty$  for all local Noetherian rings  $R$ . We do this by showing that  $\text{krdim}(R) \leq \dim(R)$  for all Noetherian local *domains*  $R$ . (The general case then follows by (P1).)

We prove this by induction on  $\dim(R)$ .

$\dim(R) = 0$ : In this case,  $R$  must be a field, for which the statement is known.

Now, assume that  $d := \dim(R) \geq 1$  and that the statement is known whenever  $\dim < R$ . For the sake of contradiction, assume that  $\text{krdim}(R) > d$ . Thus, there exists a chain

$$0 = \mathfrak{p}_0 \subsetneq \cdots \subsetneq \mathfrak{p}_{d+1} = \mathfrak{m}.$$

Pick  $x \in \mathfrak{p}_1 \setminus \{0\}$ . Then,  $\dim(R/xR) = \dim(R) - 1$ . Since we have a surjection  $R/x \twoheadrightarrow R/\mathfrak{p}_1$ , we get  $\dim(R/\mathfrak{p}_1) \leq d - 1$  as well.

But  $R/\mathfrak{p}_1$  is a domain in which we have the chain

$$0 \subseteq \mathfrak{p}_2/\mathfrak{p}_1 \subseteq \cdots \subseteq \mathfrak{p}_{d+1}/\mathfrak{p}_1,$$

of length  $d$ , contradicting the inductive hypothesis. □

A related concept is the height of a prime.

**Definition 4.6.4.** Let  $R$  be a commutative ring. The **height** of a prime  $\mathfrak{p} \in R$  is defined to be

$$\text{ht}(\mathfrak{p}) := \sup\{n : \text{there exists a chain of prime ideals of length } n \text{ ending at } \mathfrak{p}\}.$$

Equivalently,  $\text{ht}(\mathfrak{p}) = \text{krdim}(R_{\mathfrak{p}})$ .

For an arbitrary ideal  $I \subseteq R$ , one defines **height** as

$$\text{ht}(I) := \inf\{\dim R_{\mathfrak{p}} : \mathfrak{p} \in V(I)\}.$$

Note that under this definition, we have

$$\text{krdim}(R) = \sup_{\mathfrak{p} \in \text{Spec}(R)} \text{ht}(\mathfrak{p}).$$

Note that  $\text{ht}(\mathfrak{p})$  is always finite if  $R$  is Noetherian, since then  $\text{ht}(\mathfrak{p}) = \text{krdim}(R_{\mathfrak{p}}) = \dim(R_{\mathfrak{p}}) < \infty$ .

Moreover, for a local ring  $(R, \mathfrak{m})$ , we have  $\dim(R) = \text{ht}(\mathfrak{m})$ .

#### 4.6.1 Whacky behaviour of Krull dimension

To begin with,  $\text{krdim}(R)$  can be infinite even when  $R$  is Noetherian (and necessarily not local).

Note that in the local case, the (Krull) dimension being finite tells us that there exists a maximal chain of primes (i.e., a chain of primes which cannot be extended to a strictly larger chain of primes) with length equal to the dimension. (Thus, this chain is maximal in terms of inclusion and in terms of length.)

But even in the local case, there can exist maximal chains of different length.

##### Maximal chains of different lengths.

Consider the local (!) Noetherian ring  $R = k[x, y, z]_{(x, y, z)} / (xy, xz)$ .

$R$  has two minimal primes:  $(\bar{x})$ ,  $(\bar{y}, \bar{z})$ . Computing dimension using (P1) shows that  $\dim(R) = 2$ . (Example 4.3.4 tells us that the dimensions of  $k[x]_x$  and  $k[y, z]_{(y, z)}$  are 1 and 2 respectively.)

$R$  has the following two maximal chains, of different lengths:

$$\begin{aligned} (\bar{y}, \bar{z}) &\subsetneq (\bar{x}, \bar{y}, \bar{z}), \\ (\bar{x}) &\subsetneq (\bar{x}, \bar{y}) \subsetneq (\bar{x}, \bar{y}, \bar{z}). \end{aligned}$$

Note that here the chains did start at different minimal primes. However, there are more complicated examples of local Noetherian domains with maximal chains of different lengths (these maximal chains are necessarily between the zero ideal and the maximal ideal).

##### Maximal ideals of different heights.

Let  $R = \mathbb{Q}[[t]]$  be the power series ring over  $\mathbb{Q}$  in one variable, and  $\mathfrak{p} = (t)$  the maximal ideal.

Let  $S = R[x]$ .

Consider the ideals  $\mathfrak{m}_1 := (tx - 1)$  and  $\mathfrak{m}_2 = (t, x)$ .

Note that  $S/\mathfrak{m}_1 = R[x]/(tx - 1) \cong R[t^{-1}]$  is the field of Laurent series. In particular,  $\mathfrak{m}_1$  is maximal. Moreover, since  $tx - 1$  is prime, it follows that  $\text{ht}(\mathfrak{m}_1) = 1$ .<sup>9</sup>

On the other hand, note that we have the chain  $0 \subsetneq (x) \subsetneq (t, x)$  of primes showing that  $\text{ht}(\mathfrak{m}_2) \geq 2$ . Thus,  $\text{ht}(\mathfrak{m}_1) \neq \text{ht}(\mathfrak{m}_2)$ .

<sup>9</sup>Note that  $R[x]$  is a unique factorisation domain since  $R$  is a PID. Consequently, if  $0 \subsetneq \mathfrak{p} \subseteq (tx - 1)$ , then we can pick a nonzero element  $f \in \mathfrak{p}$ . By factoring and using the fact that  $\mathfrak{p}$  is prime, we may assume that  $f$  is irreducible. But  $f \in (tx - 1)$  implies that  $tx - 1 \mid f$ . Thus,  $(tx - 1) = (f)$ .

More generally, one can replace  $R$  with a DVR. In that case,  $\mathfrak{p}$  will again be principally generated (Corollary 2.2.4). Moreover,  $R[t^{-1}] \cong \text{Frac}(R)$ , showing that  $\mathfrak{m}_1$  is still maximal. As we shall later see,  $\dim(R[x]) = \dim(R) + 1 = 2$  and so we can in fact conclude  $\text{ht}(\mathfrak{m}_2) = 2$  and not just  $\geq 2$ .

#### 4.6.2 Dimension theory for $k$ -algebras

We introduce some terminology for ease of reference.

**Definition 4.6.5.** Let  $k$  be a ring. By a  **$k$ -affine ring**, we mean a  $k$ -algebra of finite type, i.e., a ring of the form  $k[x_1, \dots, x_n]/I$ .

**Definition 4.6.6.** Given a ring  $R$  and a prime ideal  $\mathfrak{p} \in \text{Spec } R$ , we define  $\dim(\mathfrak{p}) := \text{krdim}(R/\mathfrak{p})$ .

Note that  $\dim(\mathfrak{p})$  is dual to  $\text{ht}(\mathfrak{p})$ . The former calculates the length of chain of primes *starting* at  $\mathfrak{p}$ , whereas the latter calculates the length of those *ending* at  $\mathfrak{p}$ . This immediately gives us

$$\dim(\mathfrak{p}) + \text{ht}(\mathfrak{p}) \leq \dim(R)$$

for any ring  $R$  and any prime  $\mathfrak{p} \in \text{Spec}(R)$ . Note that the above inequality can be strict. We will examine some cases where we can conclude better results.

**Definition 4.6.7.** A ring  $R$  is **catenary** if given any two primes  $\mathfrak{p} \subsetneq \mathfrak{p}'$ , any two maximal prime chains from  $\mathfrak{p}$  to  $\mathfrak{p}'$  have the same length.

As noted in the previous section (without proof), there do exist Noetherian local domains which are not catenary. However, as we shall show,  $k$ -affine rings are always catenary.

**Definition 4.6.8.** Let  $k$  be a field, and  $R$  a  $k$ -affine domain. Then, we define

$$\text{trdeg}_k(R) := \text{trdeg}_k(\text{Frac}(R)).$$

**Lemma 4.6.9.** Let  $R$  be a  $k$ -affine domain with  $\text{trdeg}_k(R) = d$ , and let  $\mathfrak{p} \in \text{Spec}(R)$  be of height one. Then,  $\text{trdeg}_k(R/\mathfrak{p}) = d - 1$ .

*Proof.* **Case 1.**  $R = k[x_1, \dots, x_d]$ .

Suppose  $\mathfrak{p} \subseteq R$  is a height one prime. Then,  $\mathfrak{p} = (f)$  for some irreducible  $f$ .<sup>10</sup> Without loss

<sup>10</sup>Indeed, pick any nonzero  $f \in \mathfrak{p}$ . By primality, we may replace  $f$  by an irreducible factor. Then,  $(0) \subsetneq$

of generality,  $f$  has positive degree in  $x_d$ . We may then write

$$f = g_0(x_1, \dots, x_{d-1}) + \dots + g_n(x_1, \dots, x_{d-1})x_d^n,$$

where  $n \geq 1$ .

Note that a degree comparison shows that  $k[x_1, \dots, x_{d-1}] \cap (f) = (0)$ .

Thus, we have an inclusion  $k[x_1, \dots, x_{d-1}] \hookrightarrow R/(f)$  and hence, the images of  $x_1, \dots, x_{d-1}$  in  $R/\mathfrak{p}$  are algebraically independent over  $k$ . But  $\overline{x_d}$  is algebraic over  $\text{Frac}(k[\overline{x_1}, \dots, \overline{x_{d-1}}])$ , as witnessed by  $f$ .

Thus,  $\text{trdeg}_k(\text{Frac}(R/\mathfrak{p})) = d - 1$ .

**Case 2.** The general case.

Consider a Noetherian normalisation  $A = k[x_1, \dots, x_n] \subseteq R$ . Necessarily,  $n \leq d$  since  $\text{Frac}(A) \subseteq \text{Frac}(R)$ . On the other hand  $A \subseteq R$  is an integral extension, and hence  $\text{Frac}(R)$  is algebraic over  $\text{Frac}(A)$ . This gives us  $n = d$ .

Now, given a height one prime  $\mathfrak{p} \subseteq R$ , consider its contraction  $\mathfrak{p}_0 = \mathfrak{p} \cap A$ . Since  $A$  is a UFD and  $R$  a domain, the Going down theorem applies. This gives us that  $\text{ht}(\mathfrak{p}_0) = 1$ . Case 1 now gives us that  $\text{trdeg}_k(A/\mathfrak{p}_0) = d - 1$ . Now, since again  $R/\mathfrak{p}$  is integral over  $A/\mathfrak{p}_0$ , we get  $\text{trdeg}_k(R/\mathfrak{p}) = d - 1$  as well.  $\square$

**Theorem 4.6.10.** Let  $R$  be a  $k$ -affine ring, not necessarily a domain. Then,  $R$  is catenary.

More precisely, any maximal chain from  $\mathfrak{p}$  to  $\mathfrak{p}'$  has length  $\dim(\mathfrak{p}) - \dim(\mathfrak{p}')$ .

If  $R$  is a domain, then all maximal ideals have the same length, which is equal to  $\dim(R) = \text{trdeg}_k(R)$ .

*Proof.* Let

$$\mathfrak{p}_0 \subsetneq \dots \subsetneq \mathfrak{p}_r$$

be a maximal chain. This gives us a sequence of ring maps

$$R/\mathfrak{p}_0 \twoheadrightarrow \dots \twoheadrightarrow R/\mathfrak{p}_r.$$

Each prime ideal  $\mathfrak{p}_i/\mathfrak{p}_{i-1}$  has height one in  $R/\mathfrak{p}_{i-1}$ . Thus, each arrow above decreases the transcendence degree by exactly one.

This gives

$$\text{trdeg}_k(R/\mathfrak{p}_r) = \text{trdeg}_k(R/\mathfrak{p}_0) - r.$$

By the previous lemma, we can replace  $\text{trdeg}_k$  above with  $\dim$ , since  $R/\mathfrak{p}_i$  are domains. Rearranging gives

$$r = \dim(R/\mathfrak{p}_0) - \dim(R/\mathfrak{p}_r) = \dim(\mathfrak{p}_0) - \dim(\mathfrak{p}_r).$$

$(f) \subseteq \mathfrak{p}$ . Being height one forces equality  $(f) = \mathfrak{p}$ .

To see the last statement, if  $\mathfrak{m}$  is a maximal ideal, we can take  $\mathfrak{p}_0 = 0$  and  $\mathfrak{p}_r = \mathfrak{m}$ . We then get

$$r = \dim(0) - \dim(\mathfrak{m}) = \dim(R) - 0 = \text{trdeg}_k(R). \quad \square$$

Note that the affine ring  $k[x, y, z]/(xy, xz)$  has maximal ideals of different height. (Thus, being a domain is not an unnecessary condition.)

**Corollary 4.6.11.** Let  $R$  be a  $k$ -affine domain, and  $\mathfrak{p} \in \text{Spec}(R)$ . Then,

$$\text{ht}(\mathfrak{p}) + \dim(\mathfrak{p}) = \dim(R).$$

*Proof.* Pick a maximal chain of primes ending at  $\mathfrak{p}$ . This has length  $\text{ht}(\mathfrak{p})$ .

Pick a maximal chain of primes starting at  $\mathfrak{p}$ . This has length  $\dim(\mathfrak{p})$ .

Join these together to get a maximal chain (ending at some maximal ideal). This total length is  $\dim(R)$ .  $\square$

## §§4.7. Dimension via length and ideals of definition

We now see yet another definition of dimension for local Noetherian rings. Recall the following.

**Proposition 4.7.1.** Let  $(R, \mathfrak{m})$  be a local Noetherian ring, and  $J \subsetneq R$  a proper ideal. The following are equivalent.

1.  $J$  contains a power of  $\mathfrak{m}$ .
2.  $\sqrt{J} = \mathfrak{m}$ .
3.  $J$  is  $\mathfrak{m}$ -primary.
4.  $R/J$  is Artinian.

**Definition 4.7.2.** Let  $(R, \mathfrak{m})$  be a local Noetherian ring. A proper ideal  $J \subseteq \mathfrak{m}$  is said to be an **ideal of definition** of  $R$  if  $J$  is  $\mathfrak{m}$ -primary.

The above proposition tells us alternate definitions of an ideal of definition.

**Proposition 4.7.3.** Let  $(R, \mathfrak{m})$  be a local Noetherian ring. Define

$$d'(R) := \min\{n : \text{there exist an ideal of definition with } n \text{ generators}\}.$$

In other words,  $d'(R)$  is the minimal  $n$  such that there exist elements  $x_1, \dots, x_n \in \mathfrak{m}$  with  $\sqrt{(x_1, \dots, x_n)} = \mathfrak{m}$ .

Then,  $d'(R) = \dim(R)$ .

The above will follow from the more general theorem about dimension of modules below.

**Theorem 4.7.4.** Let  $(R, \mathfrak{m})$  be local Noetherian, and  $M$  a finitely generated  $R$ -module. Then,

$$\dim(M) = \min\{n : \text{there exist } x_1, \dots, x_n \in \mathfrak{m} \text{ such that } M/(x_1, \dots, x_n)M \text{ has finite length}\}.$$

Equivalently, the minimal  $n$  such that  $M/(x_1, \dots, x_n)M$  is Artinian.

Note that one can always take generators for the maximal ideals to get a finitely generated module over a field. Thus, the set is indeed nonempty.

*Proof.* Let  $d'(M)$  denote the minimal  $n$  such that there exist  $x_1, \dots, x_n$  with  $M/(x_1, \dots, x_n)M$  Artinian. We wish to show  $\dim(M) = d'(M)$ .

The inequality  $\leq$  is clear. Indeed, suppose that we have  $x_1, \dots, x_n \in \mathfrak{m}$ . By Corollary 4.4.8, we know that

$$\dim(M/(x_1, \dots, x_n)M) \geq \dim(M) - n.$$

Thus, if the dimension of the quotient is zero, then  $n \geq \dim(M)$ .

We prove  $d'(M) \leq \dim(M)$  by induction on  $\dim(M)$ . If  $\dim(M) = 0$ , then  $M$  is already of finite length (Example 4.3.8).

Now suppose that  $d = \dim(M) > 0$ . We first construct  $x_1, \dots, x_d$  such that the desired quotient is Artinian.

To this end, let  $M' \subseteq M$  be a maximal submodule of finite length. There is an exact sequence

$$0 \rightarrow M' \rightarrow M \rightarrow M'' \rightarrow 0$$

where  $M''$  has no finite length submodules. Note that  $\dim(M') = 0$  by Example 4.3.8 and consequently,  $\dim(M) = \dim(M'')$  by Proposition 4.4.2.

On the other hand, for any  $x_1, \dots, x_n \in \mathfrak{m}$ , we see that  $M/(x_1, \dots, x_n)M$  has finite length iff  $M''/(x_1, \dots, x_n)M''$  has finite length.<sup>11</sup> Thus,  $d'(M) = d'(M'')$  and we may assume that  $M = M''$ .

Thus,  $M$  does not have any submodule of finite length. In particular,  $R/\mathfrak{m}$  does not inject into  $M$ , i.e.,  $\mathfrak{m} \notin \text{Ass}(M)$ . By prime avoidance, there exists  $x_1 \in \mathfrak{m}$  which is not in any associated prime. Thus,  $x_1$  is a nonzerodivisor on  $M$ . By Theorem 4.4.7, we have  $\dim(M/x_1M) = \dim(M) - 1$ .

By inductive hypothesis, there exist  $x_2, \dots, x_d$  with

$$(M/x_1M)/(x_2, \dots, x_d)(M/x_1M) \cong M/(x_1, \dots, x_d)M$$

<sup>11</sup>We have the exact sequence  $0 \rightarrow K \rightarrow M/(x_1, \dots, x_n)M \rightarrow M''/(x_1, \dots, x_n)M'' \rightarrow 0$ , where  $K$  is a quotient of  $M'$  and hence of finite length.

of finite length. □

**Definition 4.7.5.** Let  $(R, \mathfrak{m})$  be a noetherian local ring, and  $M \neq 0$  a finite  $R$ -module of dimension  $n$ . A **system of parameters for  $M$**  is a sequence  $x_1, \dots, x_n \in \mathfrak{m}$  such that  $M/(x_1, \dots, x_n)M$  has finite length.

The preceding theorem tells us that systems of parameters always exist.

### §§4.8. Height and minimal generators

**Definition 4.8.1.** Given an ideal  $I$  in a ring  $R$ , and a prime ideal  $\mathfrak{p}$  containing  $I$ , we say that  $\mathfrak{p}$  is **minimal over  $I$**  if there does not exist a prime ideal  $\mathfrak{q}$  such that

$$I \subseteq \mathfrak{q} \subsetneq \mathfrak{p}.$$

In other words,  $\mathfrak{p}$  is minimal in the set of all primes containing  $I$ .

Note that a prime ideal is always minimal over itself.

**Proposition 4.8.2.** Let  $R$  be a Noetherian ring. Let  $x \in R$ .

- (i) If  $\mathfrak{p}$  is minimal over  $(x)$ , then  $\text{ht}(\mathfrak{p}) \leq 1$ . If  $x$  is a nonzerodivisor, then  $\text{ht}(\mathfrak{p}) = 1$ .
- (ii) If  $\mathfrak{p}, \mathfrak{q} \in \text{Spec}(R)$  and  $\mathfrak{q}$  is minimal over  $(\mathfrak{p}, x)$ , then there is no prime strictly between  $\mathfrak{p}$  and  $\mathfrak{q}$ .

*Proof.* (i): Assume  $\mathfrak{p}$  is minimal over  $(x)$ . Then,  $\mathfrak{p}R_{\mathfrak{p}}$  is the only prime in  $R_{\mathfrak{p}}$  that contains  $(\frac{x}{1})$ . Thus,  $\sqrt{(\frac{x}{1})} = \mathfrak{p}R_{\mathfrak{p}}$ , i.e.,  $(\frac{x}{1})$  is an ideal of definition generated by 1 element. Thus,  $\dim(R_{\mathfrak{p}}) \leq 1$  or  $\text{ht}(\mathfrak{p}) \leq 1$ .

Moreover, note that  $\text{ht}(\mathfrak{p}) = 0$  iff  $R_{\mathfrak{p}}$  is already Artinian. This happens iff  $\mathfrak{p}R_{\mathfrak{p}}$  is nilpotent (see Corollary 1.2.11 and Corollary 1.2.9). This happens iff  $\frac{x}{1}$  is nilpotent in  $R_{\mathfrak{p}}$ .

However, if  $x$  is not a zerodivisor, then neither is  $\frac{x}{1}$  and hence,  $\text{ht}(\mathfrak{p}) = 1$ .

(ii): By part (i), we see that  $\text{ht}(\mathfrak{q}/\mathfrak{p}) \leq 1$  in  $R/\mathfrak{p}$ . Thus, there cannot be a prime strictly in between  $\mathfrak{q}$  and  $\mathfrak{p}$ . □

The same proof gives the following more general result.

**Proposition 4.8.3.** Let  $R$  be a Noetherian ring, and  $x_1, \dots, x_r \in R$ .

- 1. If  $\mathfrak{p}$  is minimal over  $(x_1, \dots, x_r)$ , then  $\text{ht}(\mathfrak{p}) \leq r$ .  
In particular, if  $\mathfrak{p}$  is generated by  $r$  elements, then  $\text{ht}(\mathfrak{p}) \leq r$ .

2. If  $\mathfrak{p}, \mathfrak{q} \in \operatorname{Spec}(R)$  and  $\mathfrak{q}$  is minimal over  $(\mathfrak{p}, x_1, \dots, x_r)$ , then every chain between  $\mathfrak{p}$  and  $\mathfrak{q}$  has length at most  $r$ .



## §5. Faithfully flat modules

**Definition 5.0.1.** Let  $R$  be a ring. An  $R$ -module is said to be **flat** if  $- \otimes_R M$  is exact. A flat module is said to be **faithfully flat (fflat)** if  $- \otimes_R M$  is a faithful functor, i.e., the map

$$\mathrm{Hom}_R(N, N') \rightarrow \mathrm{Hom}_R(N \otimes_R M, N' \otimes_R M)$$

is an injection for all  $R$ -modules  $N, N'$ .

A map of rings  $f : R \rightarrow S$  is said to be **flat** (resp. **faithfully flat**) if  $S$  is a flat (resp. faithfully flat)  $R$ -module via  $f$ .

We leave the following proposition as an exercise.

**Proposition 5.0.2.** The following are equivalent for an  $R$ -module  $M$ .

1.  $M$  is fflat.
2.  $N' \rightarrow N \rightarrow N''$  is exact iff  $N' \otimes M \rightarrow N \otimes M \rightarrow N'' \otimes M$  is exact (for all sequences  $N' \rightarrow N \rightarrow N''$ ).
3.  $M$  is flat, and  $M \otimes_R N \neq 0$  for all nonzero  $R$ -modules  $N$ .
4.  $M$  is flat, and  $M \otimes k(\mathfrak{p}) \neq 0$  for all prime ideals  $\mathfrak{p} \subseteq R$ .  
(Here,  $k(\mathfrak{p}) = R_{\mathfrak{p}}/\mathfrak{p}R_{\mathfrak{p}} \cong \mathrm{Frac}(R/\mathfrak{p})$ .)
5.  $M$  is flat, and  $M \otimes k(\mathfrak{m}) \neq 0$  for all maximal ideals  $\mathfrak{m} \subseteq R$ .

**Example 5.0.3.**  $0$  is flat but not fflat (unless  $R = 0$ ).

$\mathbb{Q}$  is a flat  $\mathbb{Z}$ -module, but not fflat. (Flatness follows since  $\mathbb{Q}$  is a localisation. On the other hand, the nonzero map  $\mathbb{Z} \twoheadrightarrow \mathbb{Z}/2$  becomes the zero map after tensoring.)

$R^{\oplus n}$  is fflat.

**Lemma 5.0.4.** Suppose  $f : (R, \mathfrak{m}) \rightarrow (S, \mathfrak{n})$  is a ring homomorphism of local rings such that  $f^{-1}(\mathfrak{n}) = \mathfrak{m}$ .

If  $S$  is a flat  $R$ -module (via  $f$ ), then  $S$  is fflat.

*Proof.* We use Item 5 of Proposition 5.0.2. We only need to show that  $S \otimes_R k(\mathfrak{m})$  is nonzero. Note that

$$\begin{aligned} S \otimes_R k(\mathfrak{m}) &\cong S \otimes_R R/\mathfrak{m} \\ &\cong S/\mathfrak{m}S. \end{aligned}$$

By hypothesis,  $\mathfrak{m}S \subseteq \mathfrak{n} \neq S$ . Thus, the quotient  $S/\mathfrak{m}S$  is nonzero. □

**Theorem 5.0.5** (Lying-over theorem for fflat maps). Suppose  $f : R \rightarrow S$  is fflat. Then,  $\text{Spec}(S) \rightarrow \text{Spec}(R)$  is surjective.

*Proof.* Let  $\mathfrak{p} \subseteq R$  be prime. Then, we have a map

$$k(\mathfrak{p}) \xrightarrow{\bar{f}} S \otimes_R k(\mathfrak{p})$$

of rings.

Since  $S$  is fflat, the ring on the right is nonzero. Thus, there exists a prime  $Q \in \text{Spec}(S \otimes_R k(\mathfrak{p}))$ . Moreover,  $\bar{f}^{-1}(Q)$  is a prime in  $k(\mathfrak{p})$  and hence, is zero.

Now, note that

$$S \otimes_R k(\mathfrak{p}) \cong S_{\mathfrak{p}} / \mathfrak{p}S_{\mathfrak{p}}.$$

Primes of the above ring corresponding to primes of  $S$  that contain  $\mathfrak{p}S$  and do not intersect  $f(R \setminus \mathfrak{p})$ . Let  $\mathfrak{q} \in \text{Spec}(S)$  be the prime corresponding to  $\mathfrak{p}$ .

Then,  $f^{-1}(\mathfrak{q}) = \mathfrak{p}$ . □

**Theorem 5.0.6** (Going-down theorem for flat maps). If  $f : R \rightarrow S$  is flat, then it has the going-down property.

More explicitly: if  $\mathfrak{p}_1, \mathfrak{p}_2 \in \text{Spec}(R)$  and  $\mathfrak{q}_2 \in \text{Spec}(S)$  satisfy

$$\mathfrak{p}_1 \subseteq \mathfrak{p}_2 = f^{-1}(\mathfrak{q}_2),$$

then there exists a prime  $\mathfrak{q}_1 \in \text{Spec}(S)$  such that

$$f^{-1}(\mathfrak{q}_1) = \mathfrak{p}_1 \quad \text{and} \quad \mathfrak{q}_1 \subseteq \mathfrak{q}_2.$$

Note that  $\mathbb{Z} \rightarrow \mathbb{Q}$  is flat and hence, has the going-down property. However, it does not have the lying-over property. There is no lift of  $2\mathbb{Z}$ .

*Proof.* Note that

$$R_{\mathfrak{p}_2} \rightarrow S_{\mathfrak{q}_2}$$

is flat. Moreover, this is a map of local rings since  $f^{-1}(\mathfrak{q}_2) = \mathfrak{p}_2$ . Thus, Lemma 5.0.4 tells us that the map is fflat. Now use the lying-over theorem. □

## §6. Completion

### §§6.1. Completion via inverse limits

**Definition 6.1.1.** Let  $R$  be a ring, and  $\langle I_n \rangle_{n \geq 1}$  a collection of ideals such that  $I_{n+1} \subseteq I_n$ . Then, we have a inverse system  $R/I_{n+1} \rightarrow R/I_n$  and we define

$$\widehat{R}^{\langle I_n \rangle_n} := \varprojlim_n R/I_n.$$

If  $I_n = I^n$  for some ideal  $I$ , then the above inverse limit is denoted by  $\widehat{R}^I$ . In this case, we call  $\widehat{R}^I$  the **I-adic completion** of  $R$ .

We will typically be interested in the case where  $(R, \mathfrak{m})$  is a local ring and  $I_n = \mathfrak{m}^n$ . We may also typically write  $\widehat{R}$  if the system of ideals is clear.

**Definition 6.1.2.** More generally, if  $M$  is an  $R$ -module and  $\langle M_n \rangle_{n \geq 1}$  a sequence of submodules with  $M_n \subseteq M_{n-1}$ , then we define

$$\widehat{M}^{\langle M_n \rangle_n} := \varprojlim_n M/M_n.$$

If  $M_n = I^n M$  for some ideal  $I$ , then the above inverse limit is denoted by  $\widehat{M}^I$ . In this case, we call  $\widehat{M}^I$  the **I-adic completion** of  $M$ .

We will typically be interested in  $I$ -adic completions, and often write  $\widehat{M}$  if the inverse system is clear.

Note that the limit  $\varprojlim_n R/I_n$  is the subring of  $\prod_n R/I_n$  consisting of the elements of the form

$$(\overline{a_1}, \overline{a_2}, \overline{a_3}, \dots)$$

such that  $\overline{a_{n+1}} \mapsto \overline{a_n}$  under the map  $R/I_{n+1} \rightarrow R/I_n$ . (We may sometimes use the more elaborate notation  $a_n + I_n$  instead of  $\overline{a_n}$  for clarity.)

In particular, there is a map  $R \rightarrow \widehat{R}$  given by

$$a \mapsto (a + I_1, a + I_2, a + I_3, \dots).$$

The description of  $\widehat{M}$  is analogous. Similarly, we have a map  $M \rightarrow \widehat{M}$ .

**Remark 6.1.3.** From the discussion, we note that  $\widehat{R}$  is naturally a ring. Similarly,  $\widehat{M}$  is an  $R$ -module. If both of these are  $I$ -adic completions, then  $\widehat{M}$  is also an  $\widehat{R}$ -module.

**Observation 6.1.4.** Let  $R$  be a ring, and  $I \subseteq R$  an ideal. The description above should also make it clear that  $I$ -adic completion is a functor from the category of  $R$ -modules to itself.

Indeed, if  $f : M \rightarrow N$  is an  $R$ -module homomorphism, then  $f(J^n M) \subseteq J^n f(M) \subseteq J^n N$ .

Thus, we have compatible maps  $M/J^n M \rightarrow N/J^n N$  inducing a map  $\widehat{M} \rightarrow \widehat{N}$ .

Explicitly, the map is given by

$$(m_1 + IM, m_2 + I^2M, \dots) \mapsto (f(m_1) + IN, f(m_2) + I^2N, \dots).$$

This also shows that  $\widehat{\cdot}$  is an additive functor, i.e.,  $\widehat{f+g} = \widehat{f} + \widehat{g}$  for maps  $f, g : M \rightarrow N$ .

Consequently,  $\widehat{M \oplus N} \cong \widehat{M} \oplus \widehat{N}$ . In particular,  $\widehat{R^{\oplus n}} \cong \widehat{R}^{\oplus n}$ . (In general, any additive functor preserves finite direct sums.)

**Proposition 6.1.5.** Let  $R$  be a ring, and  $\langle I_n \rangle_{n \geq 1}$  a filtration on  $R$ .

$R \rightarrow \widehat{R}$  is injective iff  $\bigcap_{n \geq 1} I_n = 0$  iff  $R$  is Hausdorff in the topology induced by  $\langle I_n \rangle_n$ .

*Proof.* Clearly, the kernel of  $R \rightarrow \widehat{R}$  is  $\bigcap_n I_n$ . This gives the first equivalence. The last equivalence is just Proposition 3.2.10.  $\square$

**Example 6.1.6.** Let  $R$  be a ring. Consider the polynomial ring  $S = R[x_1, \dots, x_n]$  and the ideal  $\mathfrak{m} = (x_1, \dots, x_n)$ . (Note that  $\mathfrak{m}$  need not be maximal.)

Then, the  $\mathfrak{m}$ -adic completion  $\widehat{S}$  is isomorphic to  $R[[x_1, \dots, x_n]]$ .

To see this, note that we have a map

$$R[[x_1, \dots, x_n]] \rightarrow \varprojlim_t R[x_1, \dots, x_n]/(x_1, \dots, x_n)^t$$

given by truncation. It is easy to see that this is an isomorphism.

We also recall the following.

**Theorem 6.1.7.** If  $R$  is Noetherian, then so is the power series ring  $R[[x_1, \dots, x_n]]$ .

Note that the power series has the analogous universal property of polynomial rings, in the following way: Suppose  $J \subseteq R$  is an ideal generated by  $y_1, \dots, y_n$ .

Then, we have a map

$$R[x_1, \dots, x_n] \rightarrow R$$

given by  $x_i \mapsto y_i$ . Note this maps  $(x_1, \dots, x_n)$  onto  $J$ . More generally, this maps  $(x_1, \dots, x_n)^t$  onto  $J^t$  for all  $t \geq 1$ .

For all  $t \geq 1$ , this induces maps

$$R[x_1, \dots, x_n]/(x_1, \dots, x_n)^t \rightarrow R/J^t.$$

The universal property of limits now gives us a map

$$R[[x_1, \dots, x_n]] \rightarrow \widehat{R}^J.$$

Note that under this map, we see that

$$x_i \mapsto (\overline{y_i}, \overline{y_i}, \overline{y_i}, \dots).$$

(Of course, in the first coordinate, we have  $\overline{y_i} = 0$ .)

**Theorem 6.1.8.** Let  $R$  be a Noetherian ring,  $J \subseteq R$  an ideal. Then, the  $J$ -adic completion  $\widehat{R}$  is Noetherian.

*Proof.* Let  $J = (y_1, \dots, y_n)$ . Let  $\widetilde{y_i}$  denote the image of  $y_i$  in  $\widehat{R}$  via  $R \rightarrow \widehat{R}$ .

As noted, we have a (ring) map  $\Phi : R[[x_1, \dots, x_n]] \rightarrow \widehat{R}$  given by

$$x_i \mapsto \widetilde{y_i}.$$

We claim that this map is onto. In particular,  $\widehat{R}$  is Noetherian since  $R[[x_1, \dots, x_n]]$  is so.

Let  $\mathbf{a} = (\overline{a_1}, \overline{a_2}, \overline{a_3}, \dots) \in \widehat{R}$ , and let  $a_1, \dots \in R$  be lifts.

By hypothesis, we must have  $a_{t+1} - a_t \in J^t$ . Thus, we can write

$$a_{t+1} - a_t = F_t(y_1, \dots, y_n),$$

where  $F_t$  is a homogeneous polynomial of degree  $t$ .

Then,  $F = \sum_{t=1}^{\infty} F_t(x_1, \dots, x_n) \in R[[x_1, \dots, x_n]]$ . Moreover,  $\Phi(F) = \mathbf{a}$ . □

**Example 6.1.9.** Let  $J \subseteq R$  be an ideal, and consider the  $R$ -module  $M = R/J$ . We claim that the map  $M \rightarrow \widehat{M}^J$  is an isomorphism.

This follows immediately since  $JM = 0$  and thus,  $M/J^t M \cong M$  for all  $t$  and the maps  $M/J^{t+1} M \rightarrow M/J^t M$  are just the identity maps.

More generally, if  $n \geq 1$  and  $M = R/J^n$ , then we have  $M \cong \widehat{M}^J$  via the natural map. Explicitly, the map is given by

$$r + J^n \mapsto (r + J, r + J^2, \dots, r + J^n, r + J^n, r + J^n, \dots).$$

On the right, we have identified  $M/J^t M \cong (R/J^n)/(J^t/J^n) \cong R/J^t$  for  $t \leq n$ . For  $t > n$ , we have  $J^t M = 0$  and  $M/J^t M \cong M = R/J^n$ .

## §§6.2. Completion via Cauchy Sequences

We quickly see an alternate description of  $\widehat{R}$  to justify the term “completion”. Throughout this discussion, we fix a ring  $R$  and a system of ideals  $\langle I_n \rangle_{n \geq 1}$  such that  $I_{n+1} \subseteq I_n$  for all  $n$ .

**Definition 6.2.1.** A sequence  $(a_n)_{n \geq 1}$  in  $R$  is said to be **Cauchy** if for every  $t \geq 0$ , there exists  $N = N(t) \geq 1$  such that

$$a_n - a_m \in I_t$$

for all  $n, m \geq N$ .

Thus, these ideals replace the  $\varepsilon$ -balls.<sup>12</sup>

For the moment, we now forget the notation  $\widehat{R}$  from earlier and define it in a new way.

First, define  $\mathcal{R} = \prod_{\mathbb{N}} R$  to be the direct product of  $\mathbb{N}$ -many copies of  $R$ . In other words,  $\mathcal{R}$  is the ring of all sequences in  $R$ .

Let  $\mathcal{C} \subseteq \mathcal{R}$  be the set of all Cauchy sequences.

**Exercise 6.2.2.** Check that  $\mathcal{C}$  is a subring of  $\mathcal{R}$ .

Now, define  $\mathcal{I} \subseteq \mathcal{C}$  to be the set of Cauchy sequences converging to 0. ( $(r_n)_{n \geq 0}$  is said to converge to 0 if for all  $t \geq 1$ , there exists  $N = N(t)$  such that  $r_n \in I_t$  for all  $N \geq t$ . Uniqueness of limit is not assumed.)

**Exercise 6.2.3.** Check that  $\mathcal{I}$  is an ideal of  $\mathcal{C}$ .

Finally, define  $\widehat{R} = \mathcal{C} / \mathcal{I}$ . The description of  $\widehat{R}$  is given as follows: Its elements are equivalence classes of Cauchy sequences  $(r_n)_{n \geq 1}$ , where the equivalence relation is

$$(r_n)_{n \geq 1} \sim (s_n)_{n \geq 1} \Leftrightarrow \text{for all } t, \text{ there exists } N \text{ such that } r_n - s_n \in I_t \text{ for all } n \geq N.$$

The ring operations are defined in the obvious manner. The previous exercises ensure that this is well-defined.

Note that we have an obvious ring map  $R \rightarrow \widehat{R}$ , mapping  $r \in R$  to the equivalence of the constant sequence  $(r)_{n \geq 1}$ .

**Theorem 6.2.4.**  $\varprojlim_t R/I_t \cong \widehat{R}$ .

<sup>12</sup>If  $R$  is Hausdorff, i.e., if  $\bigcap_{t \geq 0} I_t = 0$ , then the topology is indeed metrisable with  $d(x, y) = \inf\{2^{-t} : x - y \in I^t\}$ . In this case, the definition above coincides with the usual notion of a Cauchy sequence. Note that if  $R$  is not Hausdorff, then  $d$  is only a pseudometric.

More precisely, there is an isomorphism  $\varphi$  making the following diagram commute

$$\begin{array}{ccc}
 & \varprojlim_t R/I_t & \\
 R & \nearrow & \downarrow \cong \varphi \\
 & \searrow & \widehat{R}
 \end{array}$$

*Proof.* Define  $\varphi$  as

$$(r_1 + I_1, r_2 + I_2, \dots) \mapsto [(r_1, r_2, \dots)].$$

We need to check that  $\varphi$  is well-defined. Note that the above formula makes it clear that the desired diagram does commute.

Firstly, note that by definition of inverse limit, we have  $r_{t+1} - r_t \in I_t$  for all  $t$ . Since the ideals are decreasing, we get  $r_n - r_m \in I_t$  for all  $n, m \geq t$ . This proves that the sequence  $(r_n)_{n \geq 1}$  is indeed Cauchy.

Second, note that if  $(r'_t)_{t \geq 1}$  is another sequence representing the same element of the inverse limit, i.e., if  $r_t - r'_t \in I_t$  for all  $t$ , then  $r_n - r'_n \in I_t$  for all  $n \geq t$ , proving that  $(r_t - r'_t)_{t \geq 1}$  converges to 0.

Thus,  $\varphi$  is well-defined. It is clearly a ring map. We now check that it is bijection.

**Injection:** Suppose that  $(r_1 + I_1, r_2 + I_2, \dots) \in \ker(\varphi)$ . We wish to show that  $r_t \in I_t$  for all  $t$ .

Fix  $t$ . By hypothesis, there exists  $N \geq t$  such that  $r_N \in I_t$ . By definition of an inverse limit, we must have  $r_N - r_t \in I_t$ , showing that  $r_t \in I_t$ .

**Surjection:** Let  $(r_1, r_2, \dots)$  be a Cauchy sequence.

Pick  $N_1 < N_2 < \dots$  such that

$$r_n - r_m \in I_t$$

for all  $n, m \geq N_t$ .

Then,

$$\mathbf{r} = (r_{N_1} + I_1, r_{N_2} + I_2, \dots)$$

is an element of the inverse limit.

We claim  $\varphi(\mathbf{r}) = [(r_1, r_2, \dots)]$  proving that  $\varphi$  is onto.

To see this, note that

$$\varphi(\mathbf{r}) = (r_{N_1}, r_{N_2}, \dots).$$

We wish to show that

$$\lim_{t \rightarrow \infty} r_{N_t} - r_t = 0.$$

(We are not claiming uniqueness of limit.)

To see this, note that  $N_n \geq n$  for all  $n$ . Thus,

$$r_{N_n} - r_n \in I_t$$

for all  $t$  and all  $n \geq N_t$ . □

### §§6.3. Exactness

Recall that one has the obvious definition of a morphism of inverse systems. In particular, a short exact sequence

$$0 \rightarrow \langle A_n \rangle_{n \geq 1} \rightarrow \langle B_n \rangle_{n \geq 1} \rightarrow \langle C_n \rangle_{n \geq 1} \rightarrow 0$$

of inverse systems of  $R$ -modules corresponds to a commutative diagram of the form

$$\begin{array}{ccccccc} & \vdots & & \vdots & & \vdots & \\ & \downarrow & & \downarrow & & \downarrow & \\ 0 & \longrightarrow & A_2 & \longrightarrow & B_2 & \longrightarrow & C_2 \longrightarrow 0 \\ & & \downarrow & & \downarrow & & \downarrow \\ 0 & \longrightarrow & A_1 & \longrightarrow & B_1 & \longrightarrow & C_1 \longrightarrow 0 \\ & & \downarrow & & \downarrow & & \downarrow \\ 0 & \longrightarrow & A_0 & \longrightarrow & B_0 & \longrightarrow & C_0 \longrightarrow 0, \end{array}$$

where each row is exact.

Corresponding to this, we can form the sequence

$$0 \rightarrow \varprojlim_n A_n \rightarrow \varprojlim_n B_n \rightarrow \varprojlim_n C_n \rightarrow 0.$$

**Proposition 6.3.1.** The sequence above is exact, except possibly at  $\varprojlim_n C_n$ .

If  $\langle A_n \rangle_{n \geq 1}$  is a surjective system, i.e., the map  $A_{n+1} \rightarrow A_n$  is a surjection for all  $n$ , then the sequence above is exact.

Note that in the case that we are interested in, the inverse systems *are* surjective.

*Proof.* Diagram chase. Left as an exercise. □



**Theorem 6.3.2.** Let  $R$  be a Noetherian ring,  $J \subseteq R$  an ideal. Let

$$0 \rightarrow M' \rightarrow M \rightarrow M'' \rightarrow 0$$

be a short exact sequence of finitely generated  $R$ -modules.

Then,

$$0 \rightarrow \widehat{M'}^J \rightarrow \widehat{M}^J \rightarrow \widehat{M''}^J \rightarrow 0$$

is exact.

*Sketch.* For each  $n$ , the sequence

$$0 \rightarrow M'/(J^n M \cap M') \rightarrow M/J^n M \rightarrow M''/J^n M'' \rightarrow 0$$

is exact.

Thus,

$$0 \rightarrow \varprojlim_n M'/(J^n M \cap M') \rightarrow \widehat{M} \rightarrow \widehat{M''} \rightarrow 0$$

is exact.

Use [Artin-Rees Lemma 3.3.4](#) to conclude that the limit above is isomorphic to  $\widehat{M'}$  (and that the appropriate diagram commutes).  $\square$

**Corollary 6.3.3.** Let  $R$  be a Noetherian ring,  $J \subseteq R$  an ideal, and  $M$  a finitely generated  $R$ -module. Then,

$$\widehat{M}^J \cong M \otimes_R \widehat{R}^J.$$

*Proof.* Since  $R$  is Noetherian and  $M$  finitely generated,  $M$  is also finitely presented, i.e., there is an exact sequence

$$R^{\oplus m} \rightarrow R^{\oplus n} \rightarrow M \rightarrow 0.$$

Now, consider the following commutative diagram:

$$\begin{array}{ccccccc} \widehat{R^{\oplus m}} & \longrightarrow & \widehat{R^{\oplus n}} & \longrightarrow & \widehat{M} & \longrightarrow & 0 \\ \downarrow \cong & & \downarrow \cong & & & & \\ \widehat{R}^{\oplus m} & \longrightarrow & \widehat{R}^{\oplus n} & & & & \\ \downarrow \cong & & \downarrow \cong & & & & \\ R^{\oplus m} \otimes_R \widehat{R} & \longrightarrow & R^{\oplus n} \otimes_R \widehat{R} & \longrightarrow & M \otimes_R \widehat{R} & \longrightarrow & 0 \end{array}$$

The top row is exact by the previous result. The bottom row is exact since tensoring is right exact. The vertical arrows are the natural ones that follow by virtue of completion and tensoring being additive functors (and therefore preserving finite direct sums).

In turn, there is an induced isomorphism  $\widehat{M} \rightarrow M \otimes_R \widehat{R}$ .  $\square$

**Corollary 6.3.4.** If  $R$  is a Noetherian ring, and  $J \subseteq R$  an ideal, then the  $J$ -adic completion  $\widehat{R}$  is a flat  $R$ -module.

*Proof.* In general, if  $K$  is an  $R$ -module such that  $\otimes_R K$  preserves short exact sequences of finitely generated modules, then  $K$  is flat.  $\square$

**Corollary 6.3.5.** Let  $R$  be a Noetherian ring, and  $J \subseteq R$  an ideal. Let  $\widehat{R}$  denote the  $J$ -adic completion.

For each  $n$ , the natural projection map  $\widehat{R} \twoheadrightarrow R/J^n$  has kernel  $J^n \widehat{R} \cong \widehat{J^n}$ .

In particular,  $\widehat{R}/J^n \widehat{R} \cong R/J^n$ .

Consequently,  $\widehat{R} \cong \widehat{\widehat{R}}$ , where the second completion is with respect to  $J\widehat{R}$ .

*Proof.* Fix  $n$ . We have the short exact sequence

$$0 \rightarrow J^n \rightarrow R \rightarrow R/J^n \rightarrow 0$$

of  $R$ -modules.

Note that by Example 6.1.9, we have  $\widehat{R/J^n} \cong R/J^n$ . Thus, applying  $\widehat{\phantom{x}}$  to the above short exact sequence yields

$$0 \rightarrow \widehat{J^n} \rightarrow \widehat{R} \rightarrow R/J^n \rightarrow 0.$$

(Check that the map  $\widehat{R} \rightarrow R/J^n$  is indeed the natural projection mentioned in the statement of the result.)

Moreover, note that  $\widehat{J^n} \cong J^n \otimes_R \widehat{R} \cong J^n \widehat{R}$ , where the last isomorphism follows since  $\widehat{R}$  is flat. (Again, one has to check that the maps are the natural ones.)

Thus, we have shown that  $J^n \widehat{R}$  is the kernel of  $\widehat{R} \rightarrow R/J^n$ .

The other statements follow as well.  $\square$

**Exercise 6.3.6.** Check that in the above setup, we have  $(J\widehat{R})^n = J^n \widehat{R}$ .

### §§6.4. Properties of completion

For this subsection, we shall assume that  $R$  is a Noetherian ring, and  $J \subseteq R$  is an ideal,  $\widehat{R}$  will denote the  $J$ -adic completion.

**Definition 6.4.1.**  $R$  is said to be **complete** in the  $J$ -adic topology if the map  $R \rightarrow \widehat{R}$  is an isomorphism.

**Remark 6.4.2.** Note that  $R$  is complete  $\Rightarrow R$  is Hausdorff in the  $J$ -adic topology, by Proposition 6.1.5.

In terms of Cauchy sequences, one can note the following.

**Proposition 6.4.3.** The following are equivalent:

- (i)  $R$  is complete.
- (ii) Every Cauchy sequence in  $R$  has a unique limit.
- (iii) Every Cauchy sequence in  $R$  converges, and  $R$  is Hausdorff.
- (iv) Every Cauchy sequence in  $R$  converges, and  $\bigcap_{n \geq 1} J^n = 0$ .

For the above proposition, one could work in the full generality of  $R$  being an arbitrary ring and  $\langle I_t \rangle_{t \geq 1}$  an arbitrary sequence of decreasing ideals.

*Proof.* (iii)  $\Leftrightarrow$  (iv) is clear from earlier discussions (Proposition 3.2.10).

(iii)  $\Rightarrow$  (ii) is clear from general topology.

(ii)  $\Rightarrow$  (iv): If there is a nonzero element  $y \in \bigcap_{n \geq 1} J^n$ , then the constant sequence  $(0)_{n \geq 0}$  has converges to both 0 and  $y$ .

Thus, it suffices to now show (i)  $\Leftrightarrow$  (iv). The proof is very similar to that of Theorem 6.2.4, and is left to the reader.  $\square$

Note that Corollary 6.3.5 told us that  $\widehat{R}$  is complete in the  $J\widehat{R}$ -adic topology, in the sense that  $\widehat{R} \rightarrow \widehat{\widehat{R}}$  is an isomorphism. In particular, this tells us that  $\widehat{R}$  is Hausdorff and that every Cauchy sequence in  $\widehat{R}$  converges (in the  $J\widehat{R}$ -adic topology).

**Corollary 6.4.4.** If  $u \in \widehat{R}$  is a unit, and  $x \in J\widehat{R}$ , then  $u + x$  is a unit.

Consequently,  $I\widehat{R}$  is contained in the Jacobson radical of  $R$ . (Equivalently, every maximal ideal of  $\widehat{R}$  contains  $I\widehat{R}$ .)

*Proof.* It suffices to show that  $1 - x$  is a unit. Consider the following sequence in  $\widehat{R}$ :

$$1, 1 + x, 1 + x + x^2, \dots$$

Note that if  $m > n \geq M$ , then

$$(1 + x + \dots + x^m) - (1 + x + \dots + x^n) = x^{n+1} + \dots + x^{m+1} \in (J\widehat{R})^M.$$

Thus, the sequence is Cauchy and has a limit, which we denote by  $1 + x + x^2 + \dots$ .

Note that

$$(1 - x)(1 + x + \dots + x^n) = 1 - x^{n+1}.$$

Thus,

$$1 - (1 - x)(1 + x + \dots + x^n) = -x^{n+1} \in (J\widehat{R})^{n+1}.$$

From the above, conclude that  $1 + x + x^2 + \dots$  is the desired inverse. □

**Corollary 6.4.5.** Suppose  $(R, \mathfrak{m})$  is a local Noetherian ring. Then,  $(\widehat{R}, \mathfrak{m}\widehat{R})$  is a local Noetherian ring.

Moreover,

$$R \rightarrow \widehat{R}^{\mathfrak{m}}$$

is a local and faithfully flat ring map, with  $R/\mathfrak{m} \cong \widehat{R}/\mathfrak{m}\widehat{R}$ .

## §7. Regular sequences and depth

### §§7.1. Regular sequences

**Definition 7.1.1.** Let  $M$  be an  $R$ -module, and  $x \in R$ . We say that  $x \in R$  is an  **$M$ -regular element** if  $x$  is a nonzerodivisor on  $M$ .

A sequence  $\mathbf{x} = x_1, \dots, x_n$  of elements in  $R$  is called an  **$M$ -regular sequence** or simply an  **$M$ -sequence** if the following conditions are met:

- (i)  $x_i$  is an  $M/(x_1, \dots, x_{i-1})M$ -regular element for  $i \in \{1, \dots, n\}$ ,
- (ii)  $M/\mathbf{x}M \neq 0$ .

A **regular sequence** is an  $R$ -sequence.

A **weak  $M$ -sequence** is only required satisfy (i).

Note that if  $(R, \mathfrak{m})$  is a local ring,  $M \neq 0$  a finitely generated  $R$ -module, and  $\mathbf{x} \subseteq M$ , then condition (ii) is automatically satisfied due to Nakayama's lemma.

Note that condition (i) can be rephrased as saying that the  $R$ -module map

$$M/(\mathbf{x}_{<i})M \xrightarrow{x_i} M/(\mathbf{x}_{<i})M$$

is injective for all  $i$ .

**Proposition 7.1.2.** Let  $R$  be a ring,  $M$  an  $R$ -module, and  $\mathbf{x} \subseteq R$  a weak  $M$ -sequence. Suppose  $\varphi : R \rightarrow S$  is a ring homomorphism, and  $N$  an  $S$ -module which is flat as an  $R$ -module. Then,  $\mathbf{x} \subseteq R$  and  $\varphi(\mathbf{x}) \subseteq S$  are weak  $(M \otimes_R N)$ -sequences. If  $\mathbf{x}(M \otimes_R N) \neq M \otimes_R N$ , then  $\mathbf{x}$  and  $\varphi(\mathbf{x})$  are  $(M \otimes_R N)$ -sequences.

*Proof.* For the first part, note that by hypothesis,

$$M/(\mathbf{x}_{<i})M \xrightarrow{x_i} M/(\mathbf{x}_{<i})M$$

is injective for all  $i$ . Since  $N$  is a flat  $R$ -module, we get that

$$M/(\mathbf{x}_{<i})M \otimes_R N \xrightarrow{x_i} M/(\mathbf{x}_{<i})M \otimes_R N$$

is injective for all  $i$ . Furthermore, by exactness again, we have a natural isomorphism

$$M/(\mathbf{x}_{<i})M \otimes_R N \cong (M \otimes_R N)/((\mathbf{x}_{<i})(M \otimes_R N)).$$

This shows that  $\mathbf{x} \subseteq R$  is weakly regular. The same is true for  $\varphi(\mathbf{x}) \subseteq S$  because the  $R$ - $S$  bimodule structure on  $N$  gives us

$$\mathbf{x}(m \otimes n) = m \otimes (\mathbf{x} \cdot n) = m \otimes (\varphi(\mathbf{x})n) = \varphi(\mathbf{x})(m \otimes n).$$

The second statement now follows by definition of being regular. □

**Corollary 7.1.3.** Let  $R$  be a noetherian ring,  $M$  a finite  $R$ -module, and  $\mathbf{x}$  an  $M$ -sequence.

- Suppose  $\mathfrak{p} \in \text{Supp } M$  is such that  $\mathbf{x} \subseteq \mathfrak{p}$ . Then,  $\mathbf{x} \subseteq R_{\mathfrak{p}}$  is an  $M_{\mathfrak{p}}$ -sequence.
- Suppose that  $R$  is local with maximal ideal  $\mathfrak{m}$ . Then,  $\mathbf{x} \subseteq \widehat{R}$  is an  $\widehat{M}$ -sequence, where  $\widehat{(-)}$  denotes completion with respect to  $\mathfrak{m}$ .

*Proof.* Both the extensions  $R \rightarrow R_{\mathfrak{p}}$  and  $R \rightarrow \widehat{R}$  are flat. We just need to ensure that quotient-ing by  $\mathbf{x}$  in either case remains nonzero.

For the first, note that  $M_{\mathfrak{p}}$  is a finite  $R_{\mathfrak{p}}$  module, which is nonzero by the hypothesis that  $\mathfrak{p} \in \text{Supp } M$ . Thus, Nakayama's lemma now gives the result.

Similarly,  $\widehat{M} = M \otimes_R \widehat{R}$  is a nonzero finite  $\widehat{R}$ -module. □

**Proposition 7.1.4.** Let  $R$  be a ring,  $M$  an  $R$ -module, and  $\mathbf{x}$  a weak  $M$ -sequence. Then an exact sequence

$$N_2 \xrightarrow{\varphi_2} N_1 \xrightarrow{\varphi_1} N_0 \xrightarrow{\varphi_0} M \rightarrow 0$$

of  $R$ -modules induces an exact sequence

$$N_2/\mathbf{x}N_2 \xrightarrow{\varphi_2} N_1/\mathbf{x}N_1 \xrightarrow{\varphi_1} N_0/\mathbf{x}N_0 \xrightarrow{\varphi_0} M/\mathbf{x}M \rightarrow 0.$$

*Proof.* By induction on length of  $\mathbf{x}$ , it suffices to prove it in the case that  $\mathbf{x}$  is a single element, say  $x$ . By right-exactness of  $- \otimes_R R/x$ , exactness only needs to be checked at  $N_1/\mathbf{x}N_1$ .

This is now a simple diagram chase: let  $\overline{n_1} \in \ker(\overline{\varphi_1})$ .

- $\overline{\varphi_1(n_1)} = 0$ ; hence,  $\varphi_1(n_1) \in \mathbf{x}N_0$ .
- Write  $\varphi_1(n_1) = \mathbf{x}n_0$  for some  $n_0 \in N_0$ .
- $\mathbf{x}\varphi_0(n_0) = \varphi_0(\mathbf{x}n_0) = \varphi_0(\varphi_1(n_1)) = 0$ , since  $\varphi_0 \circ \varphi_1 = 0$ .
- Since  $x$  is a nonzerodivisor on  $M$ , we get  $\varphi_0(n_0) = 0$ .
- By exactness of original sequence, we can write  $n_0 = \varphi_1(n'_1)$  for  $n'_1 \in N_1$ .
- Thus,

$$\varphi_1(n_1) = \mathbf{x}n_0 = \mathbf{x}\varphi_1(n'_1)$$

or  $n_1 - \mathbf{x}n'_1 \in \ker(\varphi_1)$ .

- By exactness, we can write  $n_1 = \mathbf{x}n'_1 + \varphi_2(n_2)$  for some  $n_2 \in N_2$ .
- Going modulo  $\mathbf{x}N_1$ , we are done. □

**Proposition 7.1.5.** Let  $R$  be a ring and

$$N_{\bullet} : \cdots \rightarrow N_m \xrightarrow{\varphi_m} N_{m-1} \rightarrow \cdots \rightarrow N_0 \xrightarrow{\varphi_0} N_{-1} \rightarrow 0$$

an exact complex of  $R$ -complexes. If  $\mathbf{x} \subseteq R$  is weakly  $N_i$ -regular for all  $i \geq -1$ , then  $N_{\bullet} \otimes_R R/(\mathbf{x})$  is exact again.

*Proof.* Adapt the previous proof. □

**Proposition 7.1.6.** Let  $(R, \mathfrak{m})$  be a noetherian local ring,  $M$  a finite  $R$ -module, and  $\mathbf{x}$  and  $M$ -sequence (necessarily,  $\mathbf{x} \subseteq \mathfrak{m}$ ). Then, every permutation of  $\mathbf{x}$  is a regular sequence.

*Proof.* It suffices to prove the theorem for transpositions that switch consecutive elements. Thus, it suffices to prove it for the case  $\mathbf{x} = x_1, x_2$ .

**Step 1.**  $x_2$  is  $M$ -regular.

Let  $K := (0 :_M x_2)$ . We wish to show  $K = 0$ . By Nakayama's lemma, it suffices to prove  $K \subseteq x_1 K$ .

Let  $m \in K$ . Thus,  $x_2 m = 0$ . Going modulo  $x_1 M$  and using the regularity hypothesis, we get  $\overline{m} = 0$  in  $M/x_1 M$ . Write  $m = x_1 m'$ . We show that  $m' \in K$ , i.e.,  $x_2 m' = 0$ . To this end, note

$$x_1(x_2 m') = x_2(x_1 m') = x_2 m = 0.$$

Since  $x_1$  is  $M$ -regular, we are done.

**Step 2.**  $x_1$  is  $M/x_2 M$ -regular.

Let  $m \in M$  be such that  $x_1 m \in x_2 M$ . We wish to show  $m \in x_2 M$ .

Write  $x_1 m = x_2 m'$ . Going modulo  $x_1 M$  and using regularity shows  $m' \in x_1 M$ . Write  $m' = x_1 m''$ . Then,

$$x_1 m = x_1 x_2 m''.$$

Cancel  $x_1$  to get the result. □

Let  $R$  be a ring,  $M$  an  $R$ -module, and  $\mathbf{X} = X_1, \dots, X_n$  be indeterminates over  $R$ . We define  $M[\mathbf{X}] := M \otimes_R R[\mathbf{X}]$  and call its elements **polynomials with coefficients in  $M$** .

If  $\mathbf{x} = x_1, \dots, x_n$  is a sequence of elements in  $R$ , then the substitution  $X_i \mapsto x_i$  induces an  $R$ -algebra map  $R[\mathbf{X}] \rightarrow R$  and an  $R$ -module map  $M[\mathbf{X}] \rightarrow M$ . We write  $F(\mathbf{x})$  for the image of  $F \in M[\mathbf{X}]$  under this map.

Since the monomials form a free  $R$ -basis for  $R[\mathbf{X}]$ , we may speak of the coefficients and the degree of an element of  $M[\mathbf{X}]$ . For example, in the one variable case, elements of  $M[X]$  are of the form

$$m_0 + m_1 X + \cdots + m_n X^n$$

for  $m_0, \dots, m_n$ . The action of  $r_0 + \dots + r_n X^n \in R[X]$  on the above element is given by the obvious definition.

**Theorem 7.1.7** (Rees). Let  $R$  be a ring,  $M$  an  $R$ -module,  $\mathbf{x} = x_1, \dots, x_n$  an  $M$ -sequence, and  $I = (\mathbf{x})$ . Let  $\mathbf{X} = X_1, \dots, X_n$  be indeterminates over  $R$ . If  $F \in M[\mathbf{X}]$  is homogeneous of (total) degree  $d$  and  $F(\mathbf{x}) \in I^{d+1}M$ , then the coefficients of  $F$  are in  $IM$ .

**Remark 7.1.8.** The above can be rephrased as following: Let  $R$  be a ring,  $M$  an  $R$ -module,  $\mathbf{x} = x_1, \dots, x_n$  an  $M$ -sequence, and  $I = (\mathbf{x})$ . Suppose we have an equation of the form

$$\sum_{|\alpha|=d} \mathbf{x}^\alpha m_\alpha = 0,$$

where  $m_\alpha \in M$ . (Notational clarification:  $\alpha \in \mathbb{N}_0^n$  varies over multi-indices with  $\sum \alpha_i = d$ .  $\mathbf{x}^\alpha := x_1^{\alpha_1} \dots x_n^{\alpha_n}$ .)

Then,  $m_\alpha \in IM$  for all  $\alpha$ .

*Proof.* ??

□

Recall Definition 3.2.4: given an ideal  $I \subseteq R$ , we define

$$\mathrm{gr}_I(R) = \bigoplus_{i \geq 0} I^i / I^{i+1},$$

and given an  $R$ -module  $M$ , we put

$$\mathrm{gr}_I(M) = \bigoplus_{i \geq 0} I^i M / I^{i+1} M,$$

making  $\mathrm{gr}_I(R)$  a graded ring, and  $\mathrm{gr}_I(M)$  a graded  $\mathrm{gr}_I(R)$ -module.

If  $I = (x_1, \dots, x_n)$ , then we have a natural surjective graded ring homomorphism

$$R[X_1, \dots, X_n] \twoheadrightarrow \mathrm{gr}_I(R) \tag{7.1}$$

given by  $X_i \mapsto \overline{x_i} \in I/I^2$ . (Elements of  $R$  mapped via the natural map  $R \rightarrow R/I$ .) To see that this is a surjection, note that  $\mathrm{gr}_I(R)$  is generated (as an  $R/I$ -algebra) by  $I/I^2$ .

Similarly, there is an epimorphism (of graded abelian groups)

$$\psi : M[X_1, \dots, X_n] \twoheadrightarrow \mathrm{gr}_I(M)$$

defined as following: a homogeneous polynomial  $F \in M[\mathbf{X}]$  of degree  $d$  is mapped to  $\overline{F(\mathbf{x})} \in I^d M / I^{d+1} M$ .  $\psi$  is then extended additively. Surjectivity is clear since any element of  $I^d M$  is an additive linear combination of elements of the form  $\mathbf{x}^\alpha m$ , where  $m \in M$  and  $|\alpha| = d$ .



Now, note that (7.1) makes  $\text{gr}_I(R)$  into an  $R[X]$ -algebra. Thus,  $\text{gr}_I(M)$  has the structure of a graded  $R[X]$ -module as well. With this structure, one checks  $\psi$  is a map of graded  $R[X]$ -modules.

Note that  $IM[X] \subseteq \ker \psi$ . Thus, we get an induced epimorphism

$$\varphi : (M/IM)[X] \twoheadrightarrow \text{gr}_I(M),$$

where we have used the isomorphism  $M[X]/IM[X] \cong (M/IM)[X]$  since  $R[X]$  is a flat (free even)  $R$ -module. Note that  $\varphi$  is an isomorphism iff  $IM[X] = \ker \psi$ . Since  $\psi$  is a graded map,  $\ker \psi$  is determined by its homogeneous components. Note that  $F \in M[X]$  of degree  $d$  is in the kernel iff  $F(\mathbf{x}) \in I^{d+1}M$ . Thus, a reformulation of Theorem 7.1.7, we get the following.

**Theorem 7.1.9.** Let  $R$  be a ring,  $M$  an  $R$ -module,  $\mathbf{x} = x_1, \dots, x_n$  an  $M$ -sequence, and  $I = (\mathbf{x})$ . Then, the map

$$(M/IM)[X_1, \dots, X_n] \rightarrow \text{gr}_I(M)$$

induced by  $X_i \mapsto \overline{x_i} \in I/I^2$  is an isomorphism.

**Definition 7.1.10.** Let  $R$  be a ring,  $M$  an  $R$ -module, and  $\mathbf{x} = x_1, \dots, x_n \subseteq R$  a sequence of elements of  $R$ .

We say that  $\mathbf{x}$  is  **$M$ -quasi-regular** if  $xM \neq M$  and the conclusion of Theorem 7.1.9.

### Exercises.

**Exercise 7.1.11.** Let  $0 \rightarrow U \rightarrow M \rightarrow N \rightarrow 0$  be an exact sequence of  $R$ -modules, and  $\mathbf{x}$  a sequence which is weakly  $U$ -regular and (weakly)  $N$ -regular. Prove that  $\mathbf{x}$  is (weakly)  $M$ -regular too.

*Solution.* Write  $\mathbf{x} = x_1, \dots, x_n$ . We have a commutative diagram with exact rows:

$$\begin{array}{ccccccccc} 0 & \longrightarrow & U & \longrightarrow & M & \longrightarrow & N & \longrightarrow & 0 \\ & & \downarrow x_1 & & \downarrow x_1 & & \downarrow x_1 & & \\ 0 & \longrightarrow & U & \longrightarrow & M & \longrightarrow & N & \longrightarrow & 0 \end{array}$$

The outer vertical arrows are injections by hypothesis. The Five Lemma now implies the same for the middle arrow. By Proposition 7.1.5, we see that the sequence

$$0 \rightarrow U/x_1U \rightarrow M/x_1M \rightarrow N/x_1N \rightarrow 0$$

is again exact. Induction now proves the statement involving “weakly”.

Now, if  $xM = M$ , then it follows that  $xN = N$ . This finishes the proof.  $\square$

**Exercise 7.1.12.** (a) Let  $x_1, \dots, x_i, \dots, x_n$  and  $x_1, \dots, x_i x'_i, \dots, x_n$  be (weakly)  $M$ -regular. Show that  $x_1, \dots, x_i x'_i, \dots, x_n$  is (weakly)  $M$ -regular.

(b) Prove that  $x_1^{e_1}, \dots, x_n^{e_n}$  is (weakly)  $M$ -regular for all  $e_i \geq 1$ .

*Solution.* It is clear that (b) follows from (a). We prove (a). It suffices to assume  $i = 1$ . If  $x_1$  and  $x'_1$  are nonzerodivisors, then so is  $x_1 x'_1$ . We now wish to prove that  $x_{>1}$  is a (weakly) regular sequence on  $M/x_1 x'_1 M$ .

We have a short sequence

$$0 \rightarrow M/x'_1 M \xrightarrow{\iota} M/x_1 x'_1 M \xrightarrow{\pi} M/x_1 M \rightarrow 0$$

where  $\pi$  is the natural surjection and  $\iota$  is given by  $\overline{m} \mapsto \overline{x_1 m}$ . It is not difficult to check that  $\iota$  is well-defined and that  $\ker \pi = \text{im } \iota$ .

To see that  $\iota$  is injective, one uses that  $x_1$  is a nonzerodivisor.

Now, by the previous exercise, the statement follows.  $\square$

**Exercise 7.1.13.** Prove that the converse of Proposition 7.1.2 holds if, in the situation of 7.1.2,  $N$  is faithfully flat over  $R$ .

*Solution.* We wish to prove the following statement:

Let  $R$  be a ring,  $M$  an  $R$ -module, and  $\mathbf{x} \subseteq R$ . Suppose  $\varphi : R \rightarrow S$  is a ring homomorphism, and  $N$  an  $S$ -module which is faithfully flat as an  $R$ -module. If  $\mathbf{x} \subseteq R$  (and  $\varphi(\mathbf{x}) \subseteq S$ ) are (weak)  $(M \otimes_R N)$ -sequences, then  $\mathbf{x} \subseteq R$  is a (weak)  $M$ -sequence.

Let  $\mathbf{x} = x_1, \dots, x_n$ . First, we wish to show that  $M \xrightarrow{x_1} M$  is injective. We are given that the map is injective after tensoring with  $N$ . Since  $N$  is faithfully flat, it must have been injective to begin with.

Now, note that by flatness, we have

$$(M \otimes_R N)/(x_1(M \otimes_R N)) \cong (M/x_1 M) \otimes_R N.$$

Thus, injectivity of  $M/x_1 M \xrightarrow{x_2} M/x_1 M$  also follows similarly and so on.

Lastly, if  $\mathbf{x}M = M$ , then we must have  $\mathbf{x}(M \otimes_R N) = M \otimes_R N$ .  $\square$

**Exercise 7.1.14.** 1. Prove that if  $\mathbf{x}$  is a weak  $M$ -sequence, then  $\text{Tor}_1^R(M, R/(\mathbf{x})) = 0$ .

2. Prove that if, in addition,  $\mathbf{x}$  is a weak  $R$ -sequence, then  $\mathrm{Tor}_i^R(M, R/(\mathbf{x})) = 0$  for all  $i \geq 1$ .

*Solution.* ?? Need Koszul. □

**Exercise 7.1.15.** Let  $R = k[X, Y, Z]$ ,  $k$  a field. Show that  $X, Y(1 - X), Z(1 - X)$  is an  $R$ -sequence, but  $Y(1 - X), Z(1 - X), X$  is not.

*Solution.* Direct. □

**Exercise 7.1.16.** Prove that  $x_1, \dots, x_n$  is  $M$ -quasi-regular if and only if  $\bar{x}_1, \dots, \bar{x}_n \in I/I^2$  is a  $\mathrm{gr}_I(M)$ -regular sequence where  $I = (x_1, \dots, x_n)$ .

*Solution.* Let  $\varphi$  denote the map

$$\varphi : (M/IM)[\mathbf{X}] \rightarrow \mathrm{gr}_I(M)$$

( $\Rightarrow$ ) Suppose  $\bar{\mathbf{x}}$  is  $M$ -quasi-regular, i.e.,  $\varphi$  is an isomorphism, and  $\bar{\mathbf{x}} \mathrm{gr}_I(M) \neq \mathrm{gr}_I(M)$ . Note that  $\bar{\mathbf{x}} \subseteq \mathrm{gr}_I(R)$  acts on  $\mathrm{gr}_I(M)$  via  $\mathbf{X}$ . Since  $\varphi$  is an isomorphism, it suffices to show that  $\mathbf{X}$  is  $(M/IM)[\mathbf{X}]$ -regular. But this is clear.

( $\Leftarrow$ ) We just need to show that  $\varphi$  is one-one. Since  $\varphi$  is graded, it suffices to show that there is no nonzero homogeneous element in the kernel. To this end, let  $F \in \ker \varphi$  be homogeneous of degree  $d$ . Then, we can write

$$F = \sum_{|\alpha|=d} \bar{m}_\alpha \mathbf{X}^\alpha,$$

for  $\bar{m}_\alpha \in M/IM$ . We wish to show that  $\bar{m}_\alpha = 0 \in M/IM$  for all  $\alpha$ . Note that  $F$  being in the kernel gives us that

$$\sum_{|\alpha|=d} \bar{\mathbf{x}}^\alpha \bar{m}_\alpha = 0.$$

Now, Theorem 7.1.7 tells us that  $\bar{m}_\alpha \in J \mathrm{gr}_I(M)$  for all  $\alpha$ , where  $J := (\bar{\mathbf{x}}) \mathrm{gr}_I(R)$ . But note that  $\bar{m}_\alpha$  (as an element of  $\mathrm{gr}_I(M)$ ) has degree zero, whereas every nonzero element of  $J \mathrm{gr}_I(M)$  has positive degree. Thus, it follows that  $\bar{m}_\alpha = 0$  in  $M/IM$ , as desired. □

**Exercise 7.1.17.** Suppose that  $\mathbf{x}$  is  $M$ -quasi-regular, and let  $I = (x_1, \dots, x_n)$ . Prove

- (a) if  $x_1 z \in I^{i+1}M$  for  $z \in M$ , then  $z \in I^i M$ ,

- (b)  $x_2, \dots, x_n$  is  $(M/x_1M)$ -quasi-regular,  
(c) if  $R$  is noetherian local and  $M$  is finite, then  $x$  is an  $M$ -sequence.

*Solution.* As before, we let  $\varphi$  denote the map

$$\varphi : (M/IM)[X] \xrightarrow{\cong} \text{gr}_I(M)$$

- (a) Suppose  $z \notin I^i M$ . Pick the largest  $j$  such that  $z \in I^j M$ .<sup>13</sup> By hypothesis,  $j \leq i-1$  and  $\bar{z} \in I^j M/I^{j+1} M$  is nonzero. Let  $G := \varphi^{-1}(\bar{z}) \neq 0$ .

Then,

$$\varphi(X_1 \cdot G) = X_1 \cdot \varphi(G) = \bar{x}_1 \bar{z} = 0.$$

The last equality follows since  $x_1 z \in I^{i+1} M \subseteq I^{j+2} M$ , giving us  $\bar{x}_1 \bar{z} = 0 \in I^{j+1} M/I^{j+2} M$ .

But this is a contradiction because  $\varphi$  is an isomorphism and  $X_1 G \neq 0$  in  $(M/IM)[X]$ .

- (b) Let  $J := (x_2, \dots, x_n)$ . By the previous exercise, we know that  $\bar{x}_1, \dots, \bar{x}_n \in \text{gr}_I(R)$  is regular on  $\text{gr}_I(M)$ , and that it suffices to show that  $\bar{x}_2, \dots, \bar{x}_n \in \text{gr}_J(R)$  is regular on  $\text{gr}_J(M/x_1 M)$ .

??

- (c) By part (b), it suffices to prove this in the case  $x = x$  is a single element. Necessarily,  $I = (x) \subseteq m$ .

Suppose that  $m \in M$  is such that  $xm = 0$ . Then,  $xm \in I^{i+1} M$  for all  $i$ . By (a), this implies that  $m \in \bigcap_{i \geq 0} I^i M$ . But by **Krull's Intersection Theorem 3.4.1**, we get that  $m = 0$ , as desired.  $\square$

## §§7.2. Grade and depth

**Definition 7.2.1.** Let  $R$  be a ring,  $M$  an  $R$ -module, and  $I \subseteq R$  an ideal. An  $M$ -sequence  $x_1, \dots, x_n$  (contained in  $I$ ) is **maximal (in  $I$ )**, if  $x_1, \dots, x_{n+1}$  is not an  $M$ -sequence for any  $x_{n+1} \in R$  ( $x_{n+1} \in I$ ).

Note that if  $x$  is an  $M$ -sequence, then the sequence  $(x_1) \subsetneq (x_1, x_2) \cdots (x_1, \dots, x_n)$  strictly increases. Thus, every  $M$ -sequence (contained in  $I$ ) can be extended to a maximal such sequence if the ring is noetherian.

We recall the following fact regarding associated primes for noetherian rings and modules.

<sup>13</sup>Such a  $j$  exists since  $z \in I^0 M$ .

**Proposition 7.2.2.** Let  $R$  be a noetherian ring, and  $M \neq 0$  a finite  $R$ -module. If an ideal  $I \subseteq R$  consists of zerodivisors of  $M$ , then  $I \subseteq \mathfrak{p}$  for some  $\mathfrak{p} \in \text{Ass}(M)$ .

*Proof.* Recall  $\mathcal{Z}(M) = \bigcup_{\mathfrak{p} \in \text{Ass } M} \mathfrak{p}$ , and that  $\text{Ass}(M)$  is nonempty and finite. By prime avoidance, the result now follows.  $\square$

**Proposition 7.2.3.** Let  $R$  be a ring, and  $M \neq 0$ ,  $N$  be  $R$ -modules. Set  $I := \text{ann}_R(N)$ .

- (a) If  $I$  contains an  $M$ -regular element, then  $\text{Hom}_R(N, M) = 0$ .
- (b) Conversely, if  $R$  is noetherian, and  $M, N$  are finite,  $\text{Hom}_R(N, M) = 0$  implies that  $I$  contains an  $M$ -regular element.

*Proof.* (a) Let  $\varphi \in \text{Hom}_R(N, M) = 0$ , and  $r \in I$  be  $M$ -regular. We wish to show that  $\varphi(n) = 0$  for all  $n$ . To this end, note that

$$r\varphi(n) = \varphi(rn) = \varphi(0) = 0$$

for all  $n \in N$ . Since  $r$  is  $M$ -regular, we may cancel  $r$  to get the result.

- (b) Suppose that  $I$  does not contain any  $M$ -regular element. Then,  $I \subseteq \mathfrak{p}$  for some  $\mathfrak{p} \in \text{Ass } M$ . Since  $N$  is finite, we get that  $\mathfrak{p} \in \text{Supp } N$  and that  $N_{\mathfrak{p}} \otimes_{R_{\mathfrak{p}}} k(\mathfrak{p})$  is nonzero. Since we are working over a field, we have an epimorphism

$$N_{\mathfrak{p}} \otimes_{R_{\mathfrak{p}}} k(\mathfrak{p}) \twoheadrightarrow k(\mathfrak{p}).$$

Note that  $\mathfrak{p}R_{\mathfrak{p}} \in \text{Ass}(M_{\mathfrak{p}})$ . This means that we have an inclusion  $k(\mathfrak{p}) \hookrightarrow M_{\mathfrak{p}}$ . Composing with the map above gives us a nonzero map  $N_{\mathfrak{p}} \rightarrow M_{\mathfrak{p}}$ . Since

$$\text{Hom}_{R_{\mathfrak{p}}}(M_{\mathfrak{p}}, N_{\mathfrak{p}}) \cong \text{Hom}_R(M, N)_{\mathfrak{p}},$$

we are done.  $\square$

**Lemma 7.2.4.** Let  $R$  be a ring,  $M, N$  be  $R$ -modules, and  $\mathbf{x} = x_1, \dots, x_n$  a weak  $M$ -sequence in  $\text{ann}_R(N)$ . Then,

$$\text{Hom}_R(N, M/\mathbf{x}M) \cong \text{Ext}_R^n(N, M).$$

*Proof.* We induct on  $n$ . The case  $n = 0$  is clear. Now, assume  $n \geq 1$ . By induction, we have

$$\text{Ext}_R^{n-1}(N, M) \cong \text{Hom}_R(N, M/\mathbf{x}_{<n}M).$$

Since  $x_n \in \text{ann}_R(N)$  is regular on  $M/\mathbf{x}_{<n}M$ , the above object is zero, by the previous result.

Note that we have the short exact sequence

$$0 \rightarrow M \xrightarrow{x_1} M \rightarrow M/x_1M \rightarrow 0.$$

Applying (the derived)  $\text{Hom}_R(N, -)$  to the above gives the long exact sequence

$$0 \rightarrow \text{Ext}_R^{n-1}(N, M/x_1M) \xrightarrow{\psi} \text{Ext}_R^n(N, M) \xrightarrow{x_1} \text{Ext}_R^n(N, M).$$

However, multiplication by  $x_1$  above is zero since  $x_1 \in \text{ann}_R(N)$ . Thus,  $\psi$  is an isomorphism. Using induction again proves the result.  $\square$

Let  $R$  be noetherian,  $I$  an ideal,  $M$  a finite  $R$ -module with  $M \neq IM$ , and  $\mathbf{x} = x_1, \dots, x_n$  a maximal  $M$ -sequence contained in  $I$ . Using Proposition 7.2.3 and Lemma 7.2.4, we get that

$$\text{Ext}_R^{i-1}(R/I, M) \cong \text{Hom}_R(R/I, M/(x_1, \dots, x_{i-1})M) = 0 \quad \text{for all } i = 1, \dots, n,$$

since  $I$  contains an  $M/(x_{<i})M$ -regular sequence (and  $\text{ann}_R(R/I) = I$ ). On the other hand, since  $\mathbf{x}$  is a maximal  $M$ -sequence in  $I$  and  $IM \neq I$ ,  $I$  must consist of zerodivisors of  $M/\mathbf{x}M$ .<sup>14</sup> Thus,  $\text{Ext}_R^n(R/I, M/\mathbf{x}M) \cong \text{Hom}_R(R/I, M/\mathbf{x}M) \neq 0$ .

We have therefore proved the following.

**Theorem 7.2.5 (Rees).** Let  $R$  be a noetherian ring,  $M$  a finite  $R$ -module, and  $I$  an ideal such that  $IM \neq M$ . Then, all maximal  $M$ -sequences in  $I$  have the same length  $n$  given by

$$n := \inf \text{Ext}_R^*(R/I, M).$$

Notation: If  $\langle V^i \rangle_{i \in \mathbb{Z}}$  is a sequence of  $R$ -modules, we set

$$\inf V^* := \inf\{i : V^i \neq 0\}.$$

Note that  $\inf V^* = \infty$  iff  $V^i = 0$  for all  $i$ .

**Definition 7.2.6.** Let  $R$  be a noetherian ring,  $M$  a finite  $R$ -module, and  $I$  an ideal such that  $IM \neq M$ . Then, the common length of the maximal  $M$ -sequences in  $I$  is called the **grade of  $I$  on  $M$** , denoted by  $\text{grade}(I, M)$ .

If  $IM = M$ , we define  $\text{grade}(I, M) := \infty$ . (Including the possibility  $M = 0$ .)

**Lemma 7.2.7.** If  $R$  is noetherian,  $M$  a finite  $R$ -module, and  $I$  an ideal, then  $IM = M$  iff  $\text{Ext}_R^i(R/I, M) = 0$  for all  $i$ . Thus, the convention for grade is consistent with  $\inf \text{Ext}_R^*(R/I, M)$ .

<sup>14</sup>Note that a different reason for  $\mathbf{x}$  not being maximal could have been that  $(\mathbf{x}, y)M = M$  for all  $y \in I$ . But  $IM \neq M$  ensures that that is not the case.

*Proof.* ( $\Rightarrow$ ) Assume  $IM = M$ . Then,  $\text{Supp}(M) \cap \text{Supp}(R/I) = \emptyset$  by Nakayama's lemma,<sup>15</sup> hence

$$\text{Supp Ext}_R^i(R/I, M) \subseteq \text{Supp } M \cap \text{Supp } R/I = \emptyset$$

for all  $i$ . The inclusion above follows from the natural isomorphism

$$\text{Ext}_{R_p}^i((R/I)_p, M_p) \cong \text{Ext}_R^i(R/I, M)_p.$$

( $\Leftarrow$ ) If  $IM \neq M$ , then we have already seen  $\text{grade}(I, M)$  to be finite.  $\square$

**Definition 7.2.8.** Let  $(R, \mathfrak{m})$  be a noetherian local ring, and  $M$  a finite  $R$ -module. Then the **depth of  $M$**  is defined as

$$\text{depth}(M) := \text{grade}(\mathfrak{m}, M).$$

We restate Theorem 7.2.5 in term of depth.

**Theorem 7.2.9.** Let  $(R, \mathfrak{m}, k)$  be a noetherian local ring, and  $M$  a finite module. Then,

$$\text{depth } M = \inf \text{Ext}_R^*(k, M).$$

The following also follows from the Ext characterisation.

**Corollary 7.2.10.** Let  $(R, \mathfrak{m}, k)$  be a noetherian local ring, and  $M, M'$  finite  $R$ -modules. Then,

$$\text{depth}(M \oplus M') = \min(\text{depth}(M), \text{depth}(M')).$$

In particular, if  $F$  is a nonzero finite free  $R$ -module, then  $\text{depth}(F) = \text{depth}(R)$ .

**Proposition 7.2.11.** Let  $R$  be a noetherian ring,  $I \subseteq R$  an ideal, and  $0 \rightarrow U \rightarrow M \rightarrow N \rightarrow 0$  an exact sequence of finite  $R$ -modules. Then,

$$\begin{aligned} \text{grade}(I, M) &\geq \min\{\text{grade}(I, U), \text{grade}(I, N)\}, \\ \text{grade}(I, U) &\geq \min\{\text{grade}(I, M), \text{grade}(I, N) + 1\}, \\ \text{grade}(I, N) &\geq \min\{\text{grade}(I, U) - 1, \text{grade}(I, M)\}. \end{aligned}$$

*Proof.* Follows from the long exact sequence of  $\text{Ext}_R^i(R/I, -)$ . For the first one, we have

<sup>15</sup>Indeed, if not then there exists  $\mathfrak{p} \in \text{Supp } M$  such that  $I \subseteq \mathfrak{p}$ . Then,  $IM = M$  implies  $(IR_{\mathfrak{p}})M_{\mathfrak{p}} = M_{\mathfrak{p}}$ , a contradiction since  $M_{\mathfrak{p}} \neq 0$ .

the exact sequence

$$\operatorname{Ext}_R^i(R/I, U) \rightarrow \operatorname{Ext}_R^i(R/I, M) \rightarrow \operatorname{Ext}_R^i(R/I, N).$$

Thus, if the outer two terms are zero, then so is the middle one. This proves the inequality.  $\square$

**Remark 7.2.12.** We can sometimes comment about equality if we know some more information. For example, if  $\operatorname{grade}(I, U) < \operatorname{grade}(I, N)$ , then for  $i = \operatorname{depth}(I, U)$ , we note that we have the exact sequence

$$0 = \operatorname{Ext}_R^{i-1}(R/I, N) \rightarrow \operatorname{Ext}_R^i(R/I, U) \rightarrow \operatorname{Ext}_R^i(R/I, M) \rightarrow \operatorname{Ext}_R^i(R/I, N) = 0.$$

Thus,  $\operatorname{Ext}_R^i(R/I, M) \cong \operatorname{Ext}_R^i(R/I, U) \neq 0$ .

This shows that  $\operatorname{grade}(I, M) = \operatorname{grade}(I, U) = \min(\operatorname{grade}(I, U), \operatorname{grade}(I, N))$ .

Similarly, if  $\operatorname{grade}(I, N) < \operatorname{grade}(I, M)$ , then for  $i = \operatorname{grade}(I, N)$ , we have

$$0 = \operatorname{Ext}_R^i(R/I, M) \rightarrow \operatorname{Ext}_R^i(R/I, N) \hookrightarrow \operatorname{Ext}_R^{i+1}(R/I, U),$$

showing  $\operatorname{Ext}_R^{i+1}(R/I, U) \neq 0$ . Coupled with

$$0 = \operatorname{Ext}_R^{i-1}(R/I, N) \rightarrow \operatorname{Ext}_R^i(R/I, U) \rightarrow \operatorname{Ext}_R^i(R/I, M) = 0,$$

we get that  $\operatorname{grade}(I, U) = \operatorname{grade}(I, N) + 1$ .

**Proposition 7.2.13.** Let  $R$  be a noetherian ring,  $I, J$  ideals of  $R$ , and  $M$  a finite  $R$ -module. Then,

- (a)  $\operatorname{grade}(I, M) = \inf\{\operatorname{depth}_{R_p}(M_p) : p \in V(I)\}$ ,
- (b)  $\operatorname{grade}(I, M) = \operatorname{grade}(\sqrt{I}, M)$ ,
- (c)  $\operatorname{grade}(I \cap J, M) = \min\{\operatorname{grade}(I, M), \operatorname{grade}(J, M)\}$ ,
- (d) if  $\mathbf{x} = x_1, \dots, x_n$  is an  $M$ -sequence in  $I$ , then  $\operatorname{grade}(I/(\mathbf{x}), M/\mathbf{x}M) = \operatorname{grade}(I, M/\mathbf{x}M) = \operatorname{grade}(I, M) - n$ ,
- (e) if  $N$  is a finite  $R$ -module with  $\operatorname{Supp} N = V(I)$ , then

$$\operatorname{grade}(I, M) = \inf \operatorname{Ext}_R^*(N, M).$$

The last property generalises the equation  $\operatorname{grade}(I, M) = \inf \operatorname{Ext}_R^*(N, M)$ .

*Proof.* (a): we may assume that  $I \neq R$ . Note that if  $IM = M$ , then the same is true for all primes containing  $I$  and both sides are  $\infty$ . Thus, we may assume  $IM \neq M$ . In this



case, grade is given by the length of the longest  $M$ -sequence in the ideal. This gives us  $\text{grade}(I, M) \leq \inf\{\text{depth}(M_p) : p \in V(I)\}$  (using Corollary 7.1.3).

We now show that equality is achieved: pick  $x \subseteq I$  to be a maximal  $M$ -sequence. Since  $IM \neq M$ , this tells us that  $I$  consists of zerodivisors on  $M/xM$ . Thus,  $I \subseteq p$  for some  $p \in \text{Ass}(M)$ . By definition, we have  $p \in V(I)$ . Moreover,  $x \subseteq pR_p$  is a regular sequence on  $M_p$ . But  $pR_p$  is now an associated prime of  $(M/xM)_p = M_p/xM_p$  showing that  $pR_p$  consists of zerodivisors on  $M_p/xM_p$  and thus,  $x$  is a maximal  $M_p$ -sequence.

(b) and (c): Follow from (a) since  $V(I) = V(\sqrt{I})$  and  $V(I \cap J) = V(I) \cup V(J)$ .

(d): Let  $\bar{\cdot}$  denote going modulo  $x$ . Note that

$$IM = M \Leftrightarrow \overline{IM} = \overline{M} \Leftrightarrow \overline{I}\overline{M} = \overline{M}.$$

Thus, we may only concern ourselves with the case of finite grade. The second equality follows since any regular sequence can be extended to a maximal one, whose length is the grade (and that  $y$  is regular on  $M/xM$  iff  $x, y$  is regular on  $M$ ). For the first equality, one notes that  $y \subseteq I$  is an  $\overline{M}$ -sequence iff  $\overline{y} \subseteq \overline{I}$  is so.

(e): The hypothesis tells us that  $V(\text{ann}_R(N)) = V(I)$ . By (a), we may assume that  $I = \text{ann}_R(N)$ . Now, note that in the proof of Theorem 7.2.5, we only used that  $\text{ann}_R(R/I) = I$ .  $\square$

**Remark 7.2.14.** The name ‘grade’ was originally used by Rees for a different invariant: given a noetherian ring  $R$ , and  $M$  a finite  $R$ -module, the **grade of  $M$**  is given by

$$\text{grade}(M) := \inf \text{Ext}_R^i(M, R).$$

(Thus, grade of the zero module is  $\infty$ .)

Now, by Proposition 7.2.13 (e), we know that  $\inf \text{Ext}_R^i(M, R) = \text{grade}(\text{ann}_R(M), R)$ . Thus,  $\text{grade}(M) = \text{grade}(\text{ann}_R(M), R)$ .

It is customary to set

$$\text{grade}(I) := \text{grade}(R/I) = \text{grade}(I, R),$$

for an ideal  $I \subseteq R$ . (grade( $I$ ) now has two meanings, but we shall never use it to denote the grade of  $I$  as an  $R$ -module.)

Note that if  $I \neq R$ , then  $\text{grade}(I)$  is the length of the longest ( $R$ -)regular sequence in  $I$ .

### 7.2.1 Depth and dimension

Let  $(R, \mathfrak{m})$  be noetherian local and  $M \neq 0$  a finite  $R$ -module. By Theorem 4.4.7, we see that if  $x$  is an  $M$ -sequence (necessarily contained in  $\mathfrak{m}$ ), then  $\dim(M/xM) = \dim(M) - n$ . Since a regular sequence cuts down the dimension by the correct amount, we get the following.

**Proposition 7.2.15.** Let  $(R, \mathfrak{m})$  be a noetherian local ring and  $M \neq 0$  a finite  $R$ -module. Then every  $M$ -sequence is a part of a system of parameters of  $M$ . In particular,  $\text{depth } M \leq \dim M$ .

In fact, something stronger can be proven.

**Proposition 7.2.16.** With the same notation as above, one has  $\text{depth } M \leq \dim R/\mathfrak{p}$  for all  $\mathfrak{p} \in \text{Ass } M$ .

*Proof.* We induct on  $\text{depth } M$ . The base case  $\text{depth } M = 0$  is trivial.

If  $\text{depth } M > 0$ , there exists  $x \in \mathfrak{m}$  which is  $M$ -regular. By Proposition 7.2.13, we have  $\text{depth}(M/xM) = \text{depth}(M) - 1$  and thus, we may apply the inductive hypothesis.

Now, let  $\mathfrak{p} \in \text{Ass } M$  be arbitrary. We wish to show that  $\text{depth } M \leq \dim(R/\mathfrak{p})$ . Choose  $z \in M$  such that  $Rz$  is maximal among the cyclic submodules of  $M$  annihilated by  $\mathfrak{p}$ .

**Claim.**  $\mathfrak{p}$  consists of zerodivisors of  $M/xM$ .

*Proof.* Note that  $\mathfrak{p}$  kills  $\bar{z} \in M/xM$ . We just need to show that  $\bar{z} \neq 0 \in M/xM$ , i.e.,  $z \notin xM$ .

To the contrary, assume that  $z = xm$  for some  $m \in M$ . Then,

$$\mathfrak{p}z = x(\mathfrak{p}m) = 0$$

implies that  $\mathfrak{p}m = 0$  since  $x$  is  $M$ -regular.

But then,  $\mathfrak{p}$  annihilates  $Rm \supsetneq Rz$ , a contradiction. □

Thus,  $\mathfrak{p} \subseteq \mathfrak{q}$  for some  $\mathfrak{q} \in \text{Ass}(M/xM)$ .

Note that  $x \notin \mathfrak{p}$  since  $\mathfrak{p}$  is associated to  $M$  whereas  $x$  is regular on  $M$ . Consequently,  $(M/xM)_{\mathfrak{p}} = 0$ . Thus,  $\mathfrak{p} \neq \mathfrak{q}$  since the latter is actually in the support of  $M/xM$ .

Thus, we get

$$\dim(R/\mathfrak{p}) - 1 \geq \dim(R/\mathfrak{q}) \geq \text{depth}(M/xM) = \text{depth}(M) - 1,$$

as desired. □

The global analogue of the above says the following.

**Proposition 7.2.17.** Let  $R$  be a noetherian ring, and  $I \subseteq R$  an ideal. Then,  $\text{grade } I \leq \text{ht } I$ .

*Proof.* We have  $\text{grade}(I) = \inf\{\text{depth } R_{\mathfrak{p}} : \mathfrak{p} \in V(I)\}$  by Proposition 7.2.13, and  $\text{ht}(I) := \inf\{\dim R_{\mathfrak{p}} : \mathfrak{p} \in V(I)\}$  by definition. The assertion now follows from Proposition 7.2.15. □

### 7.2.2 Depth, type, and flat extensions

**Definition 7.2.18.** Let  $(R, \mathfrak{m}, k)$  be a noetherian local ring, and  $M \neq 0$  a finite  $R$ -module of depth  $t$ . The number

$$\text{type}(M) := \dim_k \text{Ext}_R^t(k, M)$$

is called the **type** of  $M$ .

**Proposition 7.2.19.** Let  $\varphi : (R, \mathfrak{m}, k) \rightarrow (S, \mathfrak{n}, \ell)$  be a homomorphism of noetherian local rings. Suppose  $M$  is a finite  $R$ -module, and  $N$  is a finite  $S$ -module which is flat over  $R$ ; both nonzero. Then,

- (a)  $\text{depth}_S(M \otimes_R N) = \text{depth}_R(M) + \text{depth}_S(N/\mathfrak{m}N)$ ,
- (b)  $\text{type}_R(M \otimes_R N) = \text{type}_R(M) \cdot \text{type}_S(N/\mathfrak{m}N)$ .

Note: the hypothesis above ensures  $M \otimes_R N \neq 0$ . Recall that  $\varphi$  being a local map means  $\varphi(\mathfrak{m}) \subseteq \mathfrak{n}$  (equivalently,  $\varphi^{-1}(\mathfrak{n}) = \mathfrak{m}$ ).

**Lemma 7.2.20.** Under the same setup as above, the following hold:

- (a)  $\dim_\ell \text{Hom}_S(\ell, M \otimes_R N) = \dim_k \text{Hom}_R(k, M) \cdot \dim_\ell \text{Hom}_S(\ell, N/\mathfrak{m}N)$ ,
- (b) if  $\mathbf{y}$  is an  $(N/\mathfrak{m}N)$ -sequence in  $S$ , then  $\mathbf{y}$  is an  $(M \otimes_R N)$ -sequence, and  $N/\mathbf{y}N$  is flat over  $R$ .

*Proof.* Set  $T := S/\mathfrak{m}S = k \otimes_R S$ . There is a natural isomorphism

$$\text{Hom}_S(\ell, M \otimes_R N) \cong \text{Hom}_S(\ell, \text{Hom}_S(T, M \otimes_R N)), \quad (7.2)$$

since both sides can be identified with the submodule of  $M \otimes_R N$  consisting of elements annihilated by  $\mathfrak{n}$ .

As  $N$  is flat over  $R$ , we have

$$\begin{aligned} \text{Hom}_S(T, M \otimes_R N) &= \text{Hom}_S(k \otimes_R S, M \otimes_R N) \\ &\cong \text{Hom}_R(k, M \otimes_R N) \\ &\cong \text{Hom}_R(k, M) \otimes_R N. \end{aligned} \quad \left. \vphantom{\begin{aligned} \text{Hom}_S(T, M \otimes_R N) &= \text{Hom}_S(k \otimes_R S, M \otimes_R N) \\ &\cong \text{Hom}_R(k, M \otimes_R N) \\ &\cong \text{Hom}_R(k, M) \otimes_R N. \end{aligned}} \right\} N \text{ flat, } k \text{ finite}$$

Now,  $\text{Hom}_R(k, M) \cong k^s$  for some finite  $s \geq 0$ , and so  $\text{Hom}_R(k, M) \otimes_R N \cong (N/\mathfrak{m}N)^s$ . Combined with (7.2), this proves (a).

One has a natural isomorphism  $(M \otimes_R N)/J(M \otimes_R N) \cong M \otimes_R (N/JN)$  for an arbitrary ideal  $J \subseteq S$ .<sup>16</sup> Thus, it suffices to consider the case that  $\mathbf{y} = y$  is a single element in  $S$ .

<sup>16</sup>Both modules are isomorphic to  $M \otimes_R N \otimes_S (S/J)$ .

For the sake of contradiction, assume that  $y$  is a zerodivisor on  $M \otimes_R N$ . Then, there exists nonzero  $z \in M \otimes_R N$  such that  $yz = 0$ . By **Krull's Intersection Theorem 3.4.1**, there exists  $i \geq 0$  such that  $z \in \mathfrak{m}^i(M \otimes_R N) \setminus \mathfrak{m}^{i+1}(M \otimes_R N)$ , but then  $y$  would be a zerodivisor on

$$\mathfrak{m}^i(M \otimes_R N) / \mathfrak{m}^{i+1}(M \otimes_R N) \cong (\mathfrak{m}^i M / \mathfrak{m}^{i+1} M) \otimes_R N \cong k^t \otimes_R N \cong (N / \mathfrak{m} N)^t,$$

for some  $t \geq 1$ , contradicting the fact that  $y$  is a zerodivisor on  $N / \mathfrak{m} N$ . (The first isomorphism above follows by flatness of  $N$ .)

In order to test flatness for  $N / yN$ , it suffices to consider exact sequences

$$0 \rightarrow M_1 \rightarrow M_2 \rightarrow M_3 \rightarrow 0$$

of finite  $R$ -modules. By flatness of  $N$ , the sequence

$$0 \rightarrow M_1 \otimes_R N \rightarrow M_2 \otimes_R N \rightarrow M_3 \otimes_R N \rightarrow 0$$

is exact. By what we just showed,  $y$  is a regular sequence on  $M_3 \otimes_R N$ . By **Proposition 7.1.4**, the sequence above is exact even after tensoring with  $S / yS$ . But the obtained sequence is precisely

$$0 \rightarrow M_1 \otimes_R (N / yN) \rightarrow M_2 \otimes_R (N / yN) \rightarrow M_3 \otimes_R (N / yN) \rightarrow 0. \quad \square$$

*Proof of Proposition 7.2.19.* Let  $x_1, \dots, x_m$  be a maximal  $M$ -sequence, and  $y_1, \dots, y_n$  a maximal  $(N / \mathfrak{m} N)$ -sequence. We will show that  $\varphi(\mathbf{x}), \mathbf{y}$  is an  $M \otimes_R N$ -sequence. Moreover, showing  $\text{Ext}_S^{m+n}(\ell, M \otimes_R N) \neq 0$  will then prove (a).

By **Proposition 7.1.2**, we already know that  $\varphi(\mathbf{x})$  is an  $(M \otimes_R N)$ -sequence. Now, by part (b) of the previous lemma, we get that  $\mathbf{y}$  is an  $(\overline{M} \otimes_R N)$ -sequence, where  $\overline{M} := M / \mathbf{x}M$ . Since

$$\overline{M} \otimes_R N \cong (M \otimes_R N) / \varphi(\mathbf{x})(M \otimes_R N),$$

this proves that  $\varphi(\mathbf{x}), \mathbf{y}$  is an  $M \otimes_R N$ -sequence.

Next, set  $N' := N / yN$ . We then see that going modulo the above regular sequence, we get

$$(M \otimes_R N) / (\varphi(\mathbf{x}), \mathbf{y})(M \otimes_R N) \cong \overline{M} \otimes_R N'.$$

Now, using **Lemma 7.2.4**, we get

$$\begin{aligned} \text{Hom}_R(k, \overline{M}) &\cong \text{Ext}_R^m(k, M), & \text{Hom}_S(\ell, N' / \mathfrak{m} N') &\cong \text{Ext}_S^n(\ell, N / \mathfrak{m} N), \\ \text{Hom}_S(\ell, \overline{M} \otimes_R N') &\cong \text{Ext}_R^{m+n}(\ell, M \otimes_R N). \end{aligned}$$

Now, by the previous lemma, we get that

$$\begin{aligned} \dim_\ell \text{Ext}_R^{m+n}(\ell, M \otimes_R N) &= \dim_\ell \text{Hom}_S(\ell, \overline{M} \otimes_R N') \\ &= \dim_k \text{Hom}_R(k, \overline{M}) \cdot \dim_\ell \text{Hom}_S(\ell, N') \\ &= \text{type}_R(M) \cdot \text{type}_S(N). \end{aligned}$$

In particular, the above  $\text{Ext}_R^{m+n}$  is nonzero. Thus, the depth is the desired one and the above quantity is the desired type.  $\square$

**Definition 7.2.21.** Let  $M$  be a module over a local ring  $(R, \mathfrak{m}, k)$ . Then the **socle of  $M$**  is defined as

$$\text{soc}(M) := (0 :_M \mathfrak{m}) = \text{Hom}_R(k, M).$$

The socle is the largest  $R$ -submodule of  $M$  which has the structure of a  $k$ -module. The proof above showed us the following intermediate calculation.

**Porism 7.2.22.** Let  $(R, \mathfrak{m}, k)$  be a noetherian local ring,  $M$  a finite  $R$ -module, and  $\mathbf{x}$  a maximal  $M$ -sequence. Then,  $\text{type}(M) = \dim_k \text{soc}(M/\mathbf{x}M)$ .

### Exercises.

**Exercise 7.2.23.** Let  $k$  be a field and  $R = k[[X]][Y]$ . Deduce that  $X, Y$  and  $1 - XY$  are maximal  $R$ -sequences. (This example shows that  $IM \neq M$  in Theorem 7.2.5 is relevant.)

*Solution.* It is clear that both sequences written are actually regular. Moreover, the ideal generated is maximal in either case. Thus, these sequences cannot be extended.  $\square$

??

## §§7.3. Depth and projective dimension

Let  $R$  be a ring,  $M$  an  $R$ -module;  $M$  has a(n augmented) projective resolution

$$\cdots \rightarrow P_n \xrightarrow{\varphi_n} P_{n-1} \rightarrow \cdots \rightarrow P_1 \xrightarrow{\varphi_1} P_0 \xrightarrow{\varphi_0} M \rightarrow 0.$$

Set  $M_0 := M$  and  $M_i := \ker(\varphi_{i-1})$  for  $i \geq 1$ . The modules  $M_i$  are determined up to projective equivalence.<sup>17</sup>  $M_i$  is called the  **$i$ -th syzygy** of  $M$ . Since being projective is preserved by projective equivalence, the following definition makes sense.

**Definition 7.3.1.** The **projective dimension of  $M$** , abbreviated  $\text{projdim } M$ , is infinity if no syzygy of  $M$  is projective; otherwise, it is the least integer  $n$  for which the  $n$ -th syzygy is projective.

If  $n := \text{projdim } M$ , then replacing  $P_n$  by  $M_n$ , we get a finite projective resolution as follows:

$$0 \rightarrow M_n \rightarrow P_{n-1} \rightarrow \cdots \rightarrow P_0 \rightarrow M \rightarrow 0.$$

<sup>17</sup>Modules  $M$  and  $N$  are said to be **projectively equivalent** if there exist projectives  $P$  and  $Q$  with  $A \oplus P \cong B \oplus Q$ .

Thus, one could equivalently define  $\text{projdim } M$  as the infimum of the lengths of all of its projective resolutions.

Now, if  $(R, \mathfrak{m}, k)$  is a noetherian local ring, and  $M$  a finitely generated  $R$ -module, then the quantity

$$\mu(M) := \dim_k(M/\mathfrak{m}M)$$

is finite. This is the number of minimal generators of  $M$ . Set  $\beta_0 := \mu(M)$ . Let  $x_1, \dots, x_{\beta_0} \in M$  be a system of generators of  $M$ . Then, we get a surjection  $\varphi_0 : R^{\beta_0} \rightarrow M$  by mapping  $e_i \mapsto x_i$ . Repeating this procedure with  $\ker(\varphi_0)$  and so on, we get a resolution as

$$F_\bullet : \dots \rightarrow R^{\beta_n} \xrightarrow{\varphi_n} R^{\beta_{n-1}} \rightarrow \dots \rightarrow R^{\beta_1} \xrightarrow{\varphi_1} R^{\beta_0} \rightarrow M \rightarrow 0.$$

**Exercise 7.3.2.**  $F_\bullet$  is determined up to isomorphism of complexes.

*Solution.* Suppose that  $G_\bullet$  is another resolution built in the same way. Note that necessarily,  $G_0 \cong R^{\beta_0}$ .

Let  $x_1, \dots, x_{\beta_0}$  be as before, and let  $y_1, \dots, y_{\beta_0}$  be a minimal generating set corresponding to  $G_0$ . Then, there exists an  $n \times n$  matrix  $A \in M_n(R)$  relating  $x$  and  $y$ . Moreover, going modulo  $\mathfrak{m}$  shows that  $\bar{A}$  is related two bases of  $k^n$ . Thus,  $\det(\bar{A}) \neq 0$  in  $k$ . Consequently,  $\det(A)$  is a unit in  $R$ . This gives us an invertible map making the following diagram commute

$$\begin{array}{ccccc} F_0 & \xrightarrow{\varphi_0} & M & \longrightarrow & 0 \\ \eta_0 \downarrow \cong & & \parallel & & \\ G_0 & \xrightarrow{\psi_0} & M & \longrightarrow & 0. \end{array}$$

Now, a diagram chase shows that we have an induced map as shown

$$\begin{array}{ccccccc} \ker(\varphi_0) & \longrightarrow & F_0 & \xrightarrow{\varphi_0} & M & \longrightarrow & 0 \\ \downarrow \text{dotted} & & \downarrow \eta_0 & & \parallel & & \\ \ker(\psi_0) & \longrightarrow & G_0 & \xrightarrow{\psi_0} & M & \longrightarrow & 0. \end{array}$$

The map above is just the restriction of  $\eta_0$ . By symmetry,  $\eta_0^{-1}$  also restricts to a map in the other direction. Since the compositions are identity, we see that the kernels are isomorphic.

Now,  $F_1$  and  $G_1$  were constructed by mapping onto these kernels. The same argument as before shows that a dotted arrow below exists, making the diagram commute:

$$\begin{array}{ccccc} F_1 & \longrightarrow & \ker(\varphi_0) & \longrightarrow & 0 \\ \downarrow \text{dotted } \eta_1 & & \downarrow \eta_0|_{\ker} & & \\ G_1 & \longrightarrow & \ker(\psi_0) & \longrightarrow & 0. \end{array}$$

Induction finishes the job. □

**Definition 7.3.3.** The resolution built in the above way is called the **minimal free resolution of  $M$**  (and is well-defined up to isomorphism of  $R$ -complexes). The number  $\beta_i$  as above is called the  **$i$ -th Betti number** of  $M$ , denoted  $\beta_i(M)$ .

Note that projectives are free over a local ring.

**Proposition 7.3.4.** Let  $(R, \mathfrak{m}, k)$  be a local ring,  $M$  a finite  $R$ -module, and

$$F_\bullet : \cdots \rightarrow F_n \xrightarrow{\varphi_n} F_{n-1} \rightarrow \cdots \rightarrow F_1 \xrightarrow{\varphi_1} F_0 \rightarrow 0$$

a free resolution of  $M$ . Then the following are equivalent:

- (a)  $F_\bullet$  is minimal,
- (b)  $\varphi_i(F_i) \subseteq \mathfrak{m}F_{i-1}$  for all  $i \geq 1$ ,
- (c)  $\text{rank } F_i = \dim_k \text{Tor}_i^R(M, k)$  for all  $i \geq 0$ ,
- (d)  $\text{rank } F_i = \dim_k \text{Ext}_R^i(M, k)$  for all  $i \geq 0$ .

*Proof.* (a)  $\Leftrightarrow$  (b) follows at once from Proposition 8.0.2, using our definition (/construction) of the minimal free resolution.

Note that the above free resolution comes with a surjection  $F_0 \xrightarrow{\varphi_0} M$  such that the augmented complex is exact.

By definition, we have  $\text{Tor}_i^R(M, k) \cong H_i(F_\bullet \otimes_R k)$ .

(b)  $\Rightarrow$  (c), (d): By the assumption of (b), we see that

$$F_\bullet : \cdots \rightarrow F_n/\mathfrak{m}F_n \xrightarrow{0} F_{n-1}/\mathfrak{m}F_{n-1} \rightarrow \cdots \rightarrow F_1/\mathfrak{m}F_1 \xrightarrow{0} F_0/\mathfrak{m}F_0 \rightarrow 0.$$

Thus, the  $i$ -th homology is simply  $F_i/\mathfrak{m}F_i$ . Its rank as a  $k$ -vector space is precisely the same as  $\text{rank } F_i$ . This proves (c). (d) follows in the same way.

(c)  $\Rightarrow$  (b): After tensoring with  $k$ , we note that we have

$$\cdots \xrightarrow{\overline{\varphi_{i+1}}} \overline{F_i} \xrightarrow{\overline{\varphi_i}} \cdots,$$

where  $\overline{\phantom{x}}$  denotes going mod  $\mathfrak{m}$ .

Note that

$$\dim_k \text{Tor}_i^R(M, k) = \dim_k(\ker(\overline{\varphi_i})/\text{im}(\overline{\varphi_{i+1}})) \leq \dim_k(\overline{F_i}) = \text{rank } F_i.$$

Thus, for equality to hold, we must have  $\ker(\overline{\varphi}_i) = \overline{F}_i$  and  $\operatorname{im}(\overline{\varphi}_{i+1}) = 0$ . Thus,  $\overline{\varphi}_i = 0$  for all  $i$ , giving us (b).

(d)  $\Rightarrow$  (b): Follows the same way as above. Note that  $\operatorname{Hom}_R(F_i, k) \cong k^{\operatorname{rank} F_i}$ , as  $R$ -modules (and  $k$ -modules).  $\square$

**Corollary 7.3.5.** Let  $(R, \mathfrak{m}, k)$  be a noetherian local ring, and  $M$  a finite  $R$ -module. Then,  $\beta_i(M) = \operatorname{Tor}_i^R(M, k)$  for all  $i$  and

$$\operatorname{projdim} M = \sup \operatorname{Tor}_*^R(M, k).$$

**Corollary 7.3.6.** Let  $(R, \mathfrak{m})$  be a noetherian local ring, and  $M$  a finite  $R$ -module. If  $x \in \mathfrak{m}$  is  $M$ -regular and  $R$ -regular, then

$$\operatorname{projdim}_R M = \operatorname{projdim}_{R/(x)} M/xM.$$

*Proof.* Let  $F_\bullet \rightarrow M$  be a minimal free resolution. Note that  $x$  is regular on each  $F_i$ . By Proposition 7.1.5, it follows that we still have a resolution after going mod  $(x)$ . This continues to be minimal by Proposition 7.3.4. Using that again with Corollary 7.3.5, we get the result.  $\square$

**Lemma 7.3.7.** Let  $(R, \mathfrak{m}, k)$  be a local ring, and  $\varphi : F \rightarrow G$  a homomorphism of finite  $R$ -modules. Suppose that  $F$  is free, and let  $M$  be an  $R$ -module with  $\mathfrak{m} \in \operatorname{Ass} M$ . Suppose that  $\varphi \otimes_R M$  is injective. Then,

- (a)  $\varphi \otimes_R k$  is injective,
- (b) if  $G$  is a free  $R$ -module, then  $\varphi$  is injective, and  $\varphi(F)$  is a direct summand of  $G$ .

*Proof.*

- (a) Since  $\mathfrak{m} \in \operatorname{Ass} M$ , there exists an embedding  $\iota : k \hookrightarrow M$ .  $F$  being free implies that  $F \otimes_R \iota$  is also injective. We have a commutative diagram as

$$\begin{array}{ccc} F \otimes_R k & \xrightarrow{F \otimes \iota} & F \otimes_R M \\ \varphi \otimes k \downarrow & & \downarrow \varphi \otimes M \\ G \otimes_R k & \xrightarrow{G \otimes \iota} & G \otimes_R M, \end{array}$$

proving that the left map is an injection.



- (b) We are given that  $\varphi : R^n \rightarrow R^m$  is a map such that  $\overline{\varphi} : k^n \rightarrow k^m$  is an injection. We wish to show that  $\varphi$  is an injection and splits.

Let  $e_1, \dots, e_n$  be the standard basis of  $R^n$ , and put  $x_i := \varphi(e_i)$  for  $i \in [n]$ . Then, we are given that  $\overline{x}_1, \dots, \overline{x}_n$  are  $k$ -linearly independent. Thus, we may extend it to a  $k$ -basis  $\overline{x}_1, \dots, \overline{x}_m$  of  $k^m$ . Then,  $x_1, \dots, x_m$  is an  $R$ -basis for  $R^m$ . One can now define an inverse to  $\varphi$  in the obvious way.  $\square$

**Theorem 7.3.8** (Auslander-Buchsbaum). Let  $(R, \mathfrak{m})$  be a noetherian local ring, and  $M \neq 0$  a finite  $R$ -module. If  $\text{projdim } M < \infty$ , then

$$\text{projdim } M + \text{depth } M = \text{depth } R.$$

*Proof.* We proceed by induction on  $\text{depth } R$ .

First, let  $\text{depth } R = 0$ . We wish to show  $\text{projdim } M = \text{depth } M = 0$ . Suppose  $n := \text{projdim } M > 0$ . Then,  $M$  has a minimal free resolution

$$0 \rightarrow F_n \xrightarrow{\varphi_n} F_{n-1} \rightarrow \cdots \rightarrow F_0 \xrightarrow{\varphi_0} M \rightarrow 0.$$

Now,  $\text{depth } R = 0$  tells us that  $\mathfrak{m}$  consists of zero divisors, i.e.,  $\mathfrak{m} \in \text{Ass } R$ . Applying Lemma 7.3.7 to  $M = R$ , we see that  $\varphi_n$  maps  $F_n$  isomorphically onto a free summand of  $F_{n-1}$ , contradicting minimality of the above resolution.

Thus,  $n = 0$ , proving that  $M$  is projective and hence free. Consequently,  $\text{depth } M = \text{depth } R = 0$ .

Now, assume  $\text{depth } R > 0$ . Suppose first that  $\text{depth } M = 0$ . Let  $M_1$  be the first syzygy of  $M$ . Note that  $M_1 \neq 0$  since  $M$  is not free (since  $\text{depth } M \neq \text{depth } R$ ). Now, by Remark 7.2.12, we see that  $\text{depth } M' = 1$ . Since  $\text{projdim } M' = \text{projdim } M - 1$ , it is enough to prove the result for  $M'$ . Thus, we may assume  $\text{depth } M > 0$ .

Now, this means that  $\mathfrak{m} \notin \text{Ass } M$  and  $\mathfrak{m} \in \text{Ass } R$ . By prime avoidance, we may pick  $x \in \mathfrak{m}$  which is both  $\mathfrak{m}$ -regular and  $R$ -regular.

Using Proposition 7.2.13 and Corollary 7.3.6, induction now finishes the job.  $\square$

### Exercises.

**Example 7.3.9.** Let  $R$  be a noetherian local ring,  $M$  a finite  $R$ -module, and  $x$  a  $M$ -sequence of length  $n$ . Show  $\text{projdim}_R(M/xM) = \text{projdim}_R(M) + n$ .

*Proof.* Note that  $\text{depth}(M/xM) = \text{depth}(M) - n$  by Proposition 7.2.13. Thus, Auslander-Buchsbaum 7.3.8 gives the result if both  $M$  and  $M/xM$  have finite projective dimensions.

??

$\square$

**Example 7.3.10.** Let  $R$  be a noetherian local ring, and  $N$  an  $n$ -th syzygy of a finite  $R$ -module in a finite free resolution. Prove that  $\text{depth } N \geq \min(n, \text{depth } R)$ .

*Proof.* Let  $F_\bullet \rightarrow M \rightarrow 0$  be the finite free resolution mentioned. Let  $M_n$  denote the  $n$ -th syzygy (with  $M_0 = M$ ). We prove by induction on  $n$ , that  $\text{depth } M_n \geq \min(n, \text{depth } R)$ . The statement is clear for  $n = 0$ .

We have the exact sequence

$$0 \rightarrow M_n \rightarrow F_{n-1} \rightarrow M_{n-1} \rightarrow 0.$$

Note that  $\text{depth}(F_{n-1}) = \text{depth}(R)$  since  $F_{n-1}$  is a (nonzero) finite free module (Corollary 7.2.10). Now, by Proposition 7.2.11, we see that

$$\text{depth } M_n \geq \min(\text{depth}(F_{n-1}), \text{depth}(M_{n-1}) + 1) = \min(\text{depth}(R), n). \quad \square$$

## §§7.4. Some linear algebra

### 7.4.1 Torsionfree and reflexive modules

Recall that  $\mathcal{Z}(R)$  is the set of zerodivisors of  $R$ .

**Definition 7.4.1.** Let  $R$  be a ring,  $M$  an  $R$ -module. An element  $x \in M$  is said to be **torsion** if there exists  $r \in R \setminus \mathcal{Z}(R)$  such that  $rx = 0$ .

$M$  is said to be **torsionfree** if  $M$  has no nonzero torsion elements.

$M$  is said to be **torsion** if every element of  $M$  is torsion.

The zero module is both torsion and torsionfree.

Recall that the **total ring of fractions** of a ring  $R$ , is the ring  $W^{-1}R$ , where  $W$  consists of nonzerodivisors of  $R$ .

**Proposition 7.4.2.** Let  $R$  be a ring,  $Q$  its total ring of fractions, and  $M$  an  $R$ -module.

- (a)  $M$  is torsionfree iff the natural map  $M \rightarrow M \otimes_R Q$  is injective.
- (b)  $M$  is torsion iff  $M \otimes_R Q = 0$ .

*Proof.* Let  $W$  denote the set of nonzerodivisors of  $R$ . Note that  $M \otimes_R Q = W^{-1}M$  and the map  $M \rightarrow M \otimes_R Q$  is simply  $m \mapsto \frac{m}{1}$ .

This map is injective iff no nonzero element of  $M$  is killed by an element of  $W$ . (a) follows.

Similarly,  $W^{-1}M = 0$  iff every element of  $M$  is killed by an element of  $W$ . (b) follows.  $\square$

<sup>17</sup>If  $F_{n-1} = 0$ , then  $\text{depth} = \infty$  and there is nothing to prove.

**Definition 7.4.3.** Let  $R$  be a ring,  $M$  a module.

The **dual** of  $M$  is the module  $\text{Hom}_R(M, R)$ , denoted  $M^*$ ; the **bidual** is  $M^{**}$ .

The bilinear map  $M \times M^* \rightarrow R$  given by  $(x, \varphi) \mapsto \varphi(x)$  induces a natural (evaluation) map  $h : M \rightarrow M^{**}$ .

$M$  is **torsionless** if  $h$  is injective;  $M$  is **reflexive** if  $h$  is an isomorphism.

Note that  $(-)^*$  is in fact a functor, with the obvious definition for homomorphisms.

Note that if  $R$  is noetherian,  $M$  a finite  $R$ -module,  $\mathfrak{p} \in \text{Spec } R$ , then

$$\text{Hom}_R(M, R)_{\mathfrak{p}} \cong \text{Hom}_{R_{\mathfrak{p}}}(M_{\mathfrak{p}}, R_{\mathfrak{p}}).$$

Thus,  $(M^*)_{\mathfrak{p}} \cong (M_{\mathfrak{p}})^*$ , where the latter dual is considered with respect to  $R_{\mathfrak{p}}$ .

Similarly,  $(M^{**})_{\mathfrak{p}} \cong (M_{\mathfrak{p}})^{**}$ . Now, as  $R_{\mathfrak{p}}$ -modules, we have a map

$$M_{\mathfrak{p}} \rightarrow M_{\mathfrak{p}}^{**}.$$

We claim that the above is just the localisation of

$$h : M \rightarrow M^{**},$$

under the identification mentioned earlier.

**Observation 7.4.4.** Let  $(R, \mathfrak{m})$  be a local ring,  $M$  a finite  $R$ -module.

Note that  $\text{depth } M = 0$  iff  $\mathfrak{m}$  consists of zerodivisors on  $M$  iff  $\mathfrak{m} \in \text{Ass } M$ .

Thus,

$$\text{depth } M = 0 \Leftrightarrow \mathfrak{m} \in \text{Ass } M; \quad \text{depth } M \geq 1 \Leftrightarrow \mathfrak{m} \notin \text{Ass } M.$$

Now, let  $R$  be an arbitrary noetherian ring, and  $M$  a finite  $R$ -module. Since localisation behaves well with associated primes, we see that given any  $\mathfrak{p} \in \text{Spec } R$ , we have

$$\text{depth}_{R_{\mathfrak{p}}} M_{\mathfrak{p}} = 0 \Leftrightarrow \mathfrak{p} \in \text{Ass } M; \quad \text{depth}_{R_{\mathfrak{p}}} M_{\mathfrak{p}} \geq 1 \Leftrightarrow \mathfrak{p} \notin \text{Ass } M.$$

(We have used that  $\mathfrak{p} \in \text{Ass } M \Leftrightarrow \mathfrak{p}R_{\mathfrak{p}} \in \text{Ass } M_{\mathfrak{p}}$ .)

**Proposition 7.4.5.** Let  $R$  be a noetherian ring, and  $M$  a finite  $R$ -module. Then:

- (a)  $M$  is torsionless if and only if
  - (i)  $M_{\mathfrak{p}}$  is torsionless for all  $\mathfrak{p} \in \text{Ass } R$ , and
  - (ii)  $\text{depth } M_{\mathfrak{p}} \geq 1$  for  $\mathfrak{p} \in \text{Spec } R$  with  $\text{depth } R_{\mathfrak{p}} \geq 1$ .

- (b)  $M$  is reflexive if and only if
- (i)  $M_p$  is reflexive for all  $p$  with  $\text{depth } R_p \leq 1$ , and
  - (ii)  $\text{depth } M_p \geq 2$  for  $p \in \text{Spec } R$  with  $\text{depth } R_p \geq 2$ .

In points (i) and (ii),  $M_p$  and  $R_p$  are being considered as  $R_p$ -modules.

*Proof.* Let  $h : M \rightarrow M^{**}$  be the evaluation map. Put  $K := \ker(h)$ , and  $C := \text{coker}(h)$ . Note that  $M$  ( $M_p$ ) is torsionless iff  $K$  ( $K_p$ ) is zero; furthermore, reflexive iff  $C$  ( $C_p$ ) is zero. As noticed,  $h$  behaves well with localisation. This proves that points (i) in (a) and (b) are necessary.

By Exercise 7.4.13, we

$$\text{depth } M_p^{**} \geq \min(2, \text{depth } R_p). \quad (7.3)$$

This proves necessity of (b) (ii). Next, if  $M$  is torsionless, then  $M_p$  is a submodule of  $M_p^{**}$ . Proposition 7.2.11 then tells us that

$$\text{depth}(M_p) \geq \min(\text{depth}(M_p^{**}), 1).$$

Combined with (7.3), this proves the necessity of (a) (ii).

We now need to prove the sufficiency directions of (a) and (b). We will make use of Observation 7.4.4 repeatedly.

(a): It suffices to prove that  $\text{Ass } K = \emptyset$ . Let  $p \in \text{Spec } R$ . If  $\text{depth } R_p = 0$ , then  $p \in \text{Ass } R$ . For such  $p$ , (a) (i) tells us that  $K_p = 0$  and thus,  $p \notin \text{Ass } K$ .

Now, if  $\text{depth}(R_p) \geq 1$ , then  $\text{depth}(M_p) \geq 1$ , proving that  $p \notin \text{Ass}(M) \supseteq \text{Ass}(K)$ .

(b): Suppose (i) and (ii) hold. Check that this implies that (a) is true, i.e.,  $K = 0$ . Thus, we have a short exact sequence

$$0 \rightarrow M \rightarrow M^{**} \rightarrow C \rightarrow 0.$$

As before, we show that  $\text{Ass } C = \emptyset$ .

Let  $p \in \text{Spec } R$ . If  $\text{depth } R_p \leq 1$ , then we are done by (i). If  $\text{depth } R_p \geq 2$ , then  $\text{depth } M_p \geq 2$ . By (7.3),  $\text{depth } M_p^{**} \geq 2$  as well. By Proposition 7.2.11, we get  $\text{depth}(C_p) \geq 1$ . Thus,  $p \notin \text{Ass } C$ .  $\square$

### 7.4.2 Rank

**Definition 7.4.6.** Let  $R$  be a ring,  $Q$  its total ring of fractions, and  $M$  an  $R$ -module. We say  $M$  **has a rank** if  $M \otimes_R Q$  is a free  $Q$ -module. In this case, we define the **rank** of  $M$  to be  $\text{rank}_Q(M \otimes_R Q)$ .

If  $\varphi : M \rightarrow N$  is a homomorphism of  $R$ -modules, then  $\varphi$  has **rank**  $r$  if  $\text{im}(\varphi)$  has rank  $r$ .

We shall often just say “ $M$  has rank  $r$ ” to mean that  $M$  has a rank, and that this rank is  $r$ . Note that not all modules have a rank. On the other hand, any free module has a rank, which coincides with its usual notion of a rank. At the same time, there are non-free modules which have a rank in the above sense. For example, the  $\mathbb{Z}$ -module  $\mathbb{Z}/(2)$  has rank 0.

Recall that a ring  $R$  is said to be **semilocal** if  $R$  has finitely many maximal ideals. (The zero ring is also a semilocal ring.)

**Lemma 7.4.7.** Let  $R$  be a semilocal ring,  $M$  a finite projective  $R$ -module. Then,  $M$  is free of rank  $r$  iff the localisations  $M_{\mathfrak{m}}$  (are free) have the same rank  $r$  for all maximal ideals  $\mathfrak{m}$  of  $R$ .

Note that  $M$  is assumed to be projective to begin with, so the localisations  $M_{\mathfrak{m}}$  are automatically free since projective modules over local rings are free.

*Proof.* Only the direction ( $\Leftarrow$ ) is to be proven. We prove this by induction. If  $r = 0$ , then  $M = 0$  and the result is true.

Now, assume  $r > 0$ . In particular  $M \neq 0$  and thus,  $M_{\mathfrak{m}} \not\subseteq \mathfrak{m}M_{\mathfrak{m}}$  for all the maximal ideals  $\mathfrak{m}$ . By **Generalised Prime Avoidance 8.0.3**, there exists  $x \in M$  such that  $x \notin \mathfrak{m}M_{\mathfrak{m}}$  for all  $\mathfrak{m}$ . Thus,  $x$  is a part of a basis of  $M_{\mathfrak{m}}$ , for each  $\mathfrak{m}$ . In particular,  $(M/Rx)_{\mathfrak{m}}$  is free of rank  $r - 1$ , for every  $\mathfrak{m}$ . By Theorem 8.0.4,  $M/Rx$  is again projective.

Thus, the inductive hypothesis now tells us that  $M/Rx$  is free of rank  $r - 1$ . In particular,  $M \rightarrow M/Rx$  splits, giving us  $M \cong (M/Rx) \oplus Rx$ . It now suffices to prove that  $Rx$  is free of rank 1.

Note that the natural map  $R \rightarrow Rx$  is injective since it is injective after localisation at every maximal ideal.  $\square$

**Proposition 7.4.8.** Let  $R$  be a noetherian ring,  $M$  an  $R$ -module with a finite free presentation  $F_1 \xrightarrow{\varphi} F_0 \rightarrow M \rightarrow 0$ . Then the following are equivalent:

- (a)  $M$  has rank  $r$ ;
- (b)  $M$  has a free submodule  $N$  of rank  $r$  such that  $M/N$  is a torsion module;
- (c) for all prime ideals  $\mathfrak{p} \in \text{Ass } R$  the  $R_{\mathfrak{p}}$ -module  $M_{\mathfrak{p}}$  is free of rank  $r$ ;
- (d)  $\text{rank } \varphi = \text{rank } F_0 - r$ .

*Proof.* (a)  $\Rightarrow$  (b): By clearing denominators, there is a  $Q$ -basis  $\frac{x_1}{1}, \dots, \frac{x_r}{1}$  of  $M \otimes_R Q$ , where  $x_i \in M$ . Put  $N := \sum Rx_i$ . We claim that  $N$  has the desired properties.

Note that  $(M/N) \otimes_R Q = (M \otimes Q)/(N \otimes Q) = 0$  by construction. Thus,  $M/N$  is indeed torsion.

We claim that  $x_1, \dots, x_r$  are  $R$ -linearly independent. Suppose

$$a_1 x_1 + \dots + a_r x_r = 0.$$

Viewing that as an equation in  $M \otimes Q$ , we get that  $a_i = 0$  for all  $i$ . (Note that  $a_i = 0$  in  $Q$  implies  $a_i = 0$  in  $R$ .)

(b)  $\Rightarrow$  (a): We have the short exact sequence

$$0 \rightarrow N \rightarrow M \rightarrow M/N \rightarrow 0.$$

Since  $Q$  is a flat  $R$ -module, tensoring with  $Q$  is exact. By hypothesis, the tensored sequence looks like

$$0 \rightarrow Q^r \rightarrow M \otimes_R Q \rightarrow 0 \rightarrow 0.$$

Thus,  $M$  has rank  $r$ .

(a)  $\Rightarrow$  (c): Note that  $\mathcal{Z}(R) = \bigcup_{\mathfrak{p} \in \text{Ass } R} \mathfrak{p}$ . Thus, if  $\mathfrak{p} \in \text{Ass } R$ , then  $M_{\mathfrak{p}}$  is just a further localisation of  $M \otimes_R Q$ .

(c)  $\Rightarrow$  (a):  $Q$  is a semilocal ring with  $\text{MaxSpec } Q = \text{Max}(\text{Ass } R)$ . By Theorem 8.0.4, we see that  $M \otimes_R Q$  is  $Q$ -projective. Now, by Lemma 7.4.7, this module is free of rank  $r$ .

(c)  $\Leftrightarrow$  (d): In view of (a)  $\Leftrightarrow$  (c), we can replace (d) by the condition that  $(\text{im } \varphi)_{\mathfrak{p}}$  is free and  $\text{rank}(\text{im } \varphi)_{\mathfrak{p}} = \text{rank } F_0 - r$  for all  $\mathfrak{p} \in \text{Ass } R$ . We have the exact sequence

$$0 \rightarrow (\text{im } \varphi)_{\mathfrak{p}} \rightarrow (F_0)_{\mathfrak{p}} \rightarrow M_{\mathfrak{p}} \rightarrow 0.$$

If  $M_{\mathfrak{p}}$  is free, then the sequence splits, showing that  $(\text{im } \varphi)_{\mathfrak{p}}$  is projective and hence, free. Since  $\mathfrak{p} \in \text{Ass } R$ . Conversely, if  $(\text{im } \varphi)_{\mathfrak{p}}$  is free, then too the sequence splits since  $\mathfrak{p} \in \text{Ass } R$ , by Lemma 7.3.7.  $\square$

**Proposition 7.4.9.** Let  $R$  be a noetherian ring, and  $0 \rightarrow U \rightarrow M \rightarrow N \rightarrow 0$  an exact sequence of finite  $R$ -modules. If two of  $U, M, N$  have a rank, then so does the third, and  $\text{rank } M = \text{rank } U + \text{rank } N$ .

*Proof.* In view of Proposition 7.4.8 (and Observation 7.4.4), we may assume  $(R, \mathfrak{m})$  is local and  $\text{depth } R = 0$ .

- If  $N$  is free then,  $M \cong U \oplus N$ . Now freeness of  $M$  implies that of  $U$  and conversely (since projectives are free over local rings).
- If  $M$  and  $U$  are free, then again  $M \cong U \oplus N$  by Lemma 7.3.7 ( $\mathfrak{m} \in \text{Ass } R$  since  $\text{depth } R = 0$ ). Again,  $N$  is free.

The result now follows.  $\square$

**Corollary 7.4.10.** Let  $R$  be a noetherian ring,  $M$  an  $R$ -module with a finite free resolution

$$F_{\bullet} : 0 \rightarrow F_s \rightarrow F_{s-1} \rightarrow \cdots \rightarrow F_0.$$

Then,  $M$  has a rank and  $\text{rank } M = \sum_{j=0}^s (-1)^j \text{rank } F_j$ .

*Proof.* Use the previous result and induction on  $s$ . □

Note that the hypothesis above assumes that  $M$  has a finite length finite free resolution. This need not always be true. For example, consider  $R = k[x]/(x^2)$ , with  $k$  a field. This is a local artinian ring. In particular, every nonzerodivisor is already a unit. Thus, its total field of fraction is  $R$  itself. However, not every finitely generated  $R$ -module is free. Indeed,  $k = R/(x)$  is not. Consequently,  $k$  has no rank and no finite length finite free resolution either.

**Corollary 7.4.11.** Let  $R$  be a noetherian ring, and  $I \neq 0$  an ideal such that  $I$  has a finite free resolution. Then,  $I$  contains an  $R$ -regular element.

*Proof.* We have a short exact sequence  $0 \rightarrow I \rightarrow R \rightarrow R/I \rightarrow 0$ . By the previous result,  $I$  has a rank. By Proposition 7.4.9, so does  $R/I$  and  $\text{rank } I + \text{rank } R/I = 1$ . Note that  $I$  is torsionfree and nonzero; thus,  $\text{rank } I = 1$  and  $\text{rank } R/I = 0$ . Thus,  $R/I$  is torsion, i.e., annihilated by an  $R$ -regular element. This element is necessarily in  $I$ . □

### Exercises.

**Exercise 7.4.12.** Let  $R$  be a ring, and  $M$  a finite torsionfree module. Prove that if  $M$  has a rank, then  $M$  is isomorphic to a submodule of a finite free  $R$ -module of the same rank.

*Proof.* Let  $W := R \setminus \mathcal{Z}(R)$ . Then, the total ring of fractions is  $W^{-1}R =: Q$ .

Since  $M$  is torsionfree, the map  $\varphi : M \rightarrow W^{-1}M$  is injective. Since  $M$  has a rank (say  $r$ ), we know that  $W^{-1}M \cong Q^r$ .

By clearing denominators, we may assume that  $\frac{x_1}{1}, \dots, \frac{x_r}{1}$  is a  $Q$ -basis for  $W^{-1}M$ , with  $x_i \in M$ .

Then, it is clear that  $F := \sum R \frac{x_i}{1} \subseteq Q^r$  is a free  $R$ -module of rank  $r$ . We show that there exists  $w \in W$  such that  $\varphi(M) \subseteq \frac{1}{w}F$ . Note that  $\frac{1}{w}F$  is again a free  $R$ -module of rank  $r$ , which would prove the result.

**Claim.** For every  $x \in M$ , there exists  $w \in W$  such that  $\frac{x}{1} \in \frac{1}{w}F$ .

*Proof.* By hypothesis, we can write

$$\frac{x}{1} = \sum \frac{a_i x_i}{w_i}$$

with  $a_i \in R$  and  $w_i \in W$ . By clearing denominators, the result follows.  $\square$

The desired result now follows from the claim using finite generation of  $M$ .  $\square$

**Exercise 7.4.13.** Let  $R$  be a noetherian ring,  $I$  an ideal,  $M, N$  finite  $R$ -modules. Show that  $\text{grade}(I, \text{Hom}_R(M, N)) \geq \min(2, \text{grade}(I, N))$ .

*Solution.* Put  $L := \text{Hom}_R(M, N)$ . This is a finite  $R$ -module.

Edge cases:

- (a) If  $IL = L$ , then there is nothing to prove.
- (b)  $IN = N$ . In this case, there exists  $i \in I$  such that  $in = n$  for all  $n \in N$ .  
Consequently,  $i\varphi = \varphi$  for all  $\varphi \in L$ . We are back in case (a).
- (c) If  $\text{grade}(I, N) = 0$ , then there is nothing to prove.

Thus, we may now assume  $IN \neq N$  and  $IL \neq L$ . Thus, both the relevant grades are given in terms of lengths of regular sequences.

**Case 1.**  $\text{grade}(I, N) = 1$ . We wish to prove  $\text{grade}(I, L) \geq 1$ .

By hypothesis, there exists  $x \in I$  which is regular on  $N$ . Consequently,  $x$  is regular on  $L$ . Thus,  $\text{grade}(I, L) \geq 1$ .

**Case 2.**  $\text{grade}(I, N) \geq 2$ . We wish to prove  $\text{grade}(I, L) \geq 2$ .

Let  $x, y \in I$  be regular on  $N$ . We show that  $x, y$  is an  $L$ -sequence. As noted earlier,  $x$  is regular on  $L$ . Suppose  $\varphi \in L$  is such that  $y\varphi \in xL$ . We wish to show that  $\varphi \in xL$ .

Writing  $y\varphi = x\psi$ , we see that

$$y\varphi(m) = x\psi(m)$$

for all  $m \in M$ . By hypothesis of regularity,  $\varphi(m) \in xN$  for all  $m \in M$ .

Thus, there is some function  $\eta : M \rightarrow N$  such that  $\varphi(m) = x\eta(m)$  for all  $m$ .

In fact, there is a unique such  $\eta$  since  $x$  is  $N$ -regular. Using this, one gets that  $\eta$  is actually  $R$ -linear, i.e.,  $\eta \in L$ .

Thus,  $\varphi = x\eta \in xL$ , as desired.  $\square$



## §8. Appendix

Miscellaneous facts that I haven't found a better place to put yet.

**Theorem 8.0.1.** Let  $R$  be any local ring. Every projective  $R$ -module is free.

**Proposition 8.0.2.** Let  $(R, \mathfrak{m}, k)$  be a noetherian local ring,  $N$  a finite  $R$ -module,  $F = R^{\oplus I}$  a free  $R$ -module. Let  $\{e_i\}_{i \in I}$  be an  $R$ -basis for  $F$ , and  $\varphi : F \rightarrow N$  be an epimorphism. The following are equivalent:

- (a)  $\langle \varphi(e_i) \rangle_{i \in I}$  is a minimal generating set of  $N$  (consisting of distinct elements),
- (b)  $\ker(\varphi) \subseteq \mathfrak{m}F$ .

*Proof.* (a)  $\Rightarrow$  (b): Assume (a). Let  $\sum a_i e_i \in \ker(\varphi)$ . If there is some  $a_{i_0} \in R \setminus \mathfrak{m}$ , then we may divide by it to get an equation

$$\varphi(e_{i_0}) = - \sum_{i \neq i_0} \frac{a_i}{a_{i_0}} \varphi(e_i),$$

contradicting the assumption about being a minimal generating set.

(b)  $\Rightarrow$  (a): By assumption,  $\langle \varphi(e_i) \rangle_{i \in I}$  is a spanning set. Suppose it is not minimal. Then, we may pick  $i_0 \in I$  such that there is an equation

$$\varphi(e_{i_0}) = \sum_{i \neq i_0} a_i \varphi(e_i).$$

But then,  $e_{i_0} - \sum a_i e_i \in \ker \varphi \setminus \mathfrak{m}F$ . □

**Proposition 8.0.3** (Generalised Prime Avoidance). Let  $R$  be a ring,  $\mathfrak{p}_1, \dots, \mathfrak{p}_m$  prime ideals,  $M$  an  $R$ -module, and  $x_1, \dots, x_n \in M$ . Set  $N = \sum R x_i$ . If  $N_{\mathfrak{p}_j} \not\subseteq \mathfrak{p}_j M_{\mathfrak{p}_j}$  for  $j = 1, \dots, m$ , then there exist  $a_2, \dots, a_n \in R$  such that

$$x_1 + \sum_{i=2}^n a_i x_i \notin \mathfrak{p}_j M_{\mathfrak{p}_j}$$

for  $j = 1, \dots, m$ .

In words: if  $N_{\mathfrak{p}} \not\subseteq \mathfrak{p} M_{\mathfrak{p}}$  for a finite collection of primes, then there exists  $x \in N$  such that  $x \notin \mathfrak{p} M_{\mathfrak{p}}$  for those primes. Note that the only assumption was that  $N$  is a finite  $R$ -module.

*Proof.* We use induction on  $m$ . The case  $m = 1$  is vacuous. Now, suppose  $m > 1$  and that we have elements  $a'_2, \dots, a'_n$  such that  $x'_1 := x_1 + \sum_{i=2}^n a'_i x_i \notin \mathfrak{p}_j M_{\mathfrak{p}_j}$  for  $1 \leq j < m$ . If  $x'_1 \notin \mathfrak{p}_m N_{\mathfrak{p}_m}$ , then we are done already. Assume now that  $x'_1 \in \mathfrak{p}_m N_{\mathfrak{p}_m}$ .

We may also assume that  $\mathfrak{p}_j$  are pairwise distinct and  $\mathfrak{p}_m$  is minimal among the  $\mathfrak{p}_j$ . Thus, we may pick  $r \in (\bigcap_{j=1}^{m-1} \mathfrak{p}_j) \setminus \mathfrak{p}_m$ .

Put  $x'_i := rx_i$  for  $2 \leq i \leq n$ , and  $N' := \sum_{i=1}^n Rx'_i \leq N$ . Since  $r \notin \mathfrak{p}_m$ ,  $N'_{\mathfrak{p}_m} = N_{\mathfrak{p}_m}$ . On the other hand, since  $r \in \mathfrak{p}_j$  for  $1 \leq j < m$ , we have

$$x'_1 + x'_i \notin \mathfrak{p}_j M_{\mathfrak{p}_j}$$

for  $1 < i \leq n$  and  $1 \leq j < m$ . Now, since  $x'_1 \in \mathfrak{p}_m N_{\mathfrak{p}_m}$  and  $N'_{\mathfrak{p}_m} \not\subseteq \mathfrak{p}_m M_{\mathfrak{p}_m}$ , there is some  $i \in \{2, \dots, n\}$  such that  $x'_1 + x'_i \notin \mathfrak{p}_m M_{\mathfrak{p}_m}$ . This is the desired element.  $\square$

**Theorem 8.0.4** (Projective iff locally free). Let  $R$  be a ring,  $M$  a finitely presented  $R$ -module.  $M$  is projective iff  $M_{\mathfrak{p}}$  is free (equivalently projective) for all  $\mathfrak{p} \in \operatorname{Spec} R$  iff  $M_{\mathfrak{m}}$  is free for all  $\mathfrak{m} \in \operatorname{MaxSpec} R$ .

*Proof.*  $M$  projective implies that each  $M_{\mathfrak{p}}$  is projective and hence free, since  $R_{\mathfrak{p}}$  is local. The only nontrivial implication is that third implies the first.

We show that  $\operatorname{Hom}_R(M, -)$  preserves surjections. Let  $N \twoheadrightarrow N'$  be a surjection. It suffices to show that the induced map

$$\operatorname{Hom}_R(M, N)_{\mathfrak{m}} \rightarrow \operatorname{Hom}_R(M, N')_{\mathfrak{m}}$$

is surjective for every  $\mathfrak{m} \in \operatorname{MaxSpec} R$ . Since  $M$  is finitely presented, the above is equivalent to showing that

$$\operatorname{Hom}_{R_{\mathfrak{m}}}(M_{\mathfrak{m}}, N_{\mathfrak{m}}) \rightarrow \operatorname{Hom}_{R_{\mathfrak{m}}}(M_{\mathfrak{m}}, N'_{\mathfrak{m}})$$

is surjective for all  $\mathfrak{m}$ . But now, for a fixed  $\mathfrak{m}$ , we note that the above map is also induced by the surjection  $N_{\mathfrak{m}} \twoheadrightarrow N'_{\mathfrak{m}}$ . Thus, the hypothesis of  $M_{\mathfrak{m}}$  being projective finishes the proof.  $\square$