

# Fourier Inversion for $L^1$ Functions

Aryaman Maithani

Department of Mathematics  
IIT Bombay

March 17, 2021

- 1 Recap
- 2 Notations and Setup
- 3 Proof of the Main Theorem
- 4 The Stronger Theorem

- 1 Recap
- 2 Notations and Setup
- 3 Proof of the Main Theorem
- 4 The Stronger Theorem

Recall how we had proven Fourier inversion for  $L^1$  functions in class.

Recall how we had proven Fourier inversion for  $L^1$  functions in class. Fix  $n \in \mathbb{N}$  for the entirety of this discussion.

## Definition (Heat Kernel)

Recall how we had proven Fourier inversion for  $L^1$  functions in class. Fix  $n \in \mathbb{N}$  for the entirety of this discussion.

## Definition (Heat Kernel)

For  $t > 0$ , define  $h_t : \mathbb{R}^n \rightarrow \mathbb{R}$  by

Recall how we had proven Fourier inversion for  $L^1$  functions in class. Fix  $n \in \mathbb{N}$  for the entirety of this discussion.

## Definition (Heat Kernel)

For  $t > 0$ , define  $h_t : \mathbb{R}^n \rightarrow \mathbb{R}$  by

$$h_t(x) = \frac{1}{t^{n/2}} \exp\left(-\frac{\pi}{t} \|x\|^2\right).$$

# The proof

We had seen that  $\{h_t\}_{t>0}$  constitutes a (continuous) approximate identity.



# The proof

We had seen that  $\{h_t\}_{t>0}$  constitutes a (continuous) approximate identity. We thus concluded that  $f * h_t \xrightarrow{t \rightarrow 0} f$  in  $L^1$  for any  $f \in L^1$ .

# The proof

We had seen that  $\{h_t\}_{t>0}$  constitutes a (continuous) approximate identity. We thus concluded that  $f * h_t \xrightarrow{t \rightarrow 0} f$  in  $L^1$  for any  $f \in L^1$ . In turn, that gave us that

$$\lim_{t \rightarrow 0} (f * h_t)(x) = f(x) \quad (\star)$$

for almost all  $x$ .

# The proof

We had seen that  $\{h_t\}_{t>0}$  constitutes a (continuous) approximate identity. We thus concluded that  $f * h_t \xrightarrow{t \rightarrow 0} f$  in  $L^1$  for any  $f \in L^1$ . In turn, that gave us that

$$\lim_{t \rightarrow 0} (f * h_t)(x) = f(x) \quad (\star)$$

for almost all  $x$ .

We then showed the following assuming  $\hat{f} \in L^1$ .

# The proof

We had seen that  $\{h_t\}_{t>0}$  constitutes a (continuous) approximate identity. We thus concluded that  $f * h_t \xrightarrow{t \rightarrow 0} f$  in  $L^1$  for any  $f \in L^1$ . In turn, that gave us that

$$\lim_{t \rightarrow 0} (f * h_t)(x) = f(x) \quad (\star)$$

for almost all  $x$ .

We then showed the following assuming  $\hat{f} \in L^1$ .

$$\textcircled{1} \quad (f * h_t)(x) = \int_{\mathbb{R}^n} \hat{f}(\xi) e^{-\pi t \|\xi\|^2} e^{2\pi i x \cdot \xi} d\xi \text{ for } \underline{\text{all}} \ x \in \mathbb{R}^n.$$

# The proof

We had seen that  $\{h_t\}_{t>0}$  constitutes a (continuous) approximate identity. We thus concluded that  $f * h_t \xrightarrow{t \rightarrow 0} f$  in  $L^1$  for any  $f \in L^1$ . In turn, that gave us that

$$\lim_{t \rightarrow 0} (f * h_t)(x) = f(x) \quad (\star)$$

for almost all  $x$ .

We then showed the following assuming  $\hat{f} \in L^1$ .

- ①  $(f * h_t)(x) = \int_{\mathbb{R}^n} \hat{f}(\xi) e^{-\pi t \|\xi\|^2} e^{2\pi i x \cdot \xi} d\xi$  for all  $x \in \mathbb{R}^n$ .
- ② Using DCT, we let  $t \rightarrow 0$  in the above to conclude that

$$f(x) = \int_{\mathbb{R}^n} \hat{f}(\xi) e^{2\pi i x \cdot \xi} d\xi$$

# The proof

We had seen that  $\{h_t\}_{t>0}$  constitutes a (continuous) approximate identity. We thus concluded that  $f * h_t \xrightarrow{t \rightarrow 0} f$  in  $L^1$  for any  $f \in L^1$ . In turn, that gave us that

$$\lim_{t \rightarrow 0} (f * h_t)(x) = f(x) \quad (\star)$$

for almost all  $x$ .

We then showed the following assuming  $\hat{f} \in L^1$ .

- ①  $(f * h_t)(x) = \int_{\mathbb{R}^n} \hat{f}(\xi) e^{-\pi t \|\xi\|^2} e^{2\pi i x \cdot \xi} d\xi$  for all  $x \in \mathbb{R}^n$ .
- ② Using DCT, we let  $t \rightarrow 0$  in the above to conclude that

$$f(x) = \int_{\mathbb{R}^n} \hat{f}(\xi) e^{2\pi i x \cdot \xi} d\xi$$

for those  $x \in \mathbb{R}^n$  for which  $(\star)$  holds.

# Conclusion

Thus, we have actually proven that the Fourier inversion holds for those points  $x$



Thus, we have actually proven that the Fourier inversion holds for those points  $x$  for which the equality

$$\lim_{t \rightarrow 0} (f * h_t)(x) = f(x) \quad (\star)$$

holds.

Thus, we have actually proven that the Fourier inversion holds for those points  $x$  for which the equality

$$\lim_{t \rightarrow 0} (f * h_t)(x) = f(x) \quad (\star)$$

holds.

Our aim now is to show that  $(\star)$  holds for all  $x$  in the *Lebesgue set* of  $f$ .

Thus, we have actually proven that the Fourier inversion holds for those points  $x$  for which the equality

$$\lim_{t \rightarrow 0} (f * h_t)(x) = f(x) \quad (\star)$$

holds.

Our aim now is to show that  $(\star)$  holds for all  $x$  in the *Lebesgue set* of  $f$ .

We will actually prove the result for a broader class of approximate identities.

- 1 Recap
- 2 **Notations and Setup**
- 3 Proof of the Main Theorem
- 4 The Stronger Theorem

Note that if  $f \in L^1$ , then  $f$  is finite a.e. Thus, we may assume that  $f$  is finite everywhere by changing it on a null set.

## Definition (Lebesgue set)

Note that if  $f \in L^1$ , then  $f$  is finite a.e. Thus, we may assume that  $f$  is finite everywhere by changing it on a null set.

## Definition (Lebesgue set)

Suppose  $f \in L^1(\mathbb{R}^n)$ .

Note that if  $f \in L^1$ , then  $f$  is finite a.e. Thus, we may assume that  $f$  is finite everywhere by changing it on a null set.

## Definition (Lebesgue set)

Suppose  $f \in L^1(\mathbb{R}^n)$ . The Lebesgue set of  $f$ , denoted  $\text{Leb}(f)$

Note that if  $f \in L^1$ , then  $f$  is finite a.e. Thus, we may assume that  $f$  is finite everywhere by changing it on a null set.

## Definition (Lebesgue set)

Suppose  $f \in L^1(\mathbb{R}^n)$ . The Lebesgue set of  $f$ , denoted  $\text{Leb}(f)$  is defined to be the set of all  $x \in \mathbb{R}^n$  for which

holds.



Note that if  $f \in L^1$ , then  $f$  is finite a.e. Thus, we may assume that  $f$  is finite everywhere by changing it on a null set.

## Definition (Lebesgue set)

Suppose  $f \in L^1(\mathbb{R}^n)$ . The Lebesgue set of  $f$ , denoted  $\text{Leb}(f)$  is defined to be the set of all  $x \in \mathbb{R}^n$  for which

$$\lim_{r \rightarrow 0} \frac{1}{r^n} \int_{|u| < r} |f(x - u) - f(x)| \, du = 0$$

holds.

Note that if  $f \in L^1$ , then  $f$  is finite a.e. Thus, we may assume that  $f$  is finite everywhere by changing it on a null set.

## Definition (Lebesgue set)

Suppose  $f \in L^1(\mathbb{R}^n)$ . The Lebesgue set of  $f$ , denoted  $\text{Leb}(f)$  is defined to be the set of all  $x \in \mathbb{R}^n$  for which

$$\lim_{r \rightarrow 0} \frac{1}{r^n} \int_{|u| < r} |f(x - u) - f(x)| \, du = 0$$

holds.

Note that the above  $\text{Leb}(f)$  is actually a superset of the  $\text{Leb}(f)$  we defined it in class.

Note that if  $f \in L^1$ , then  $f$  is finite a.e. Thus, we may assume that  $f$  is finite everywhere by changing it on a null set.

## Definition (Lebesgue set)

Suppose  $f \in L^1(\mathbb{R}^n)$ . The Lebesgue set of  $f$ , denoted  $\text{Leb}(f)$  is defined to be the set of all  $x \in \mathbb{R}^n$  for which

$$\lim_{r \rightarrow 0} \frac{1}{r^n} \int_{|u| < r} |f(x - u) - f(x)| \, du = 0$$

holds.

Note that the above  $\text{Leb}(f)$  is actually a superset of the  $\text{Leb}(f)$  we defined in class. So, we shall prove a stronger result.

# The identity

Let  $\varphi \in L^1(\mathbb{R}^n)$  be a radial function with  $\|\varphi\|_1 = 1$ .

# The identity

Let  $\varphi \in L^1(\mathbb{R}^n)$  be a radial function with  $\|\varphi\|_1 = 1$ . Let  $\psi_0 : \mathbb{R} \rightarrow \mathbb{R}$  be defined as

# The identity

Let  $\varphi \in L^1(\mathbb{R}^n)$  be a radial function with  $\|\varphi\|_1 = 1$ . Let  $\psi_0 : \mathbb{R} \rightarrow \mathbb{R}$  be defined as

$$\psi_0(\|x\|) = \varphi(x)$$

for  $x \in \mathbb{R}^n$ .

# The identity

Let  $\varphi \in L^1(\mathbb{R}^n)$  be a radial function with  $\|\varphi\|_1 = 1$ . Let  $\psi_0 : \mathbb{R} \rightarrow \mathbb{R}$  be defined as

$$\psi_0(\|x\|) = \varphi(x)$$

for  $x \in \mathbb{R}^n$ . (Well-defined by radial assumption.)

# The identity

Let  $\varphi \in L^1(\mathbb{R}^n)$  be a radial function with  $\|\varphi\|_1 = 1$ . Let  $\psi_0 : \mathbb{R} \rightarrow \mathbb{R}$  be defined as

$$\psi_0(\|x\|) = \varphi(x)$$

for  $x \in \mathbb{R}^n$ . (Well-defined by radial assumption.)

For  $t > 0$ , define  $\varphi_t \in L^1(\mathbb{R}^n)$  by



# The identity

Let  $\varphi \in L^1(\mathbb{R}^n)$  be a radial function with  $\|\varphi\|_1 = 1$ . Let  $\psi_0 : \mathbb{R} \rightarrow \mathbb{R}$  be defined as

$$\psi_0(\|x\|) = \varphi(x)$$

for  $x \in \mathbb{R}^n$ . (Well-defined by radial assumption.)

For  $t > 0$ , define  $\varphi_t \in L^1(\mathbb{R}^n)$  by

$$\varphi_t(x) = t^{-n} \varphi(x/t).$$

# The identity

Let  $\varphi \in L^1(\mathbb{R}^n)$  be a radial function with  $\|\varphi\|_1 = 1$ . Let  $\psi_0 : \mathbb{R} \rightarrow \mathbb{R}$  be defined as

$$\psi_0(\|x\|) = \varphi(x)$$

for  $x \in \mathbb{R}^n$ . (Well-defined by radial assumption.)

For  $t > 0$ , define  $\varphi_t \in L^1(\mathbb{R}^n)$  by

$$\varphi_t(x) = t^{-n} \varphi(x/t).$$

( $\varphi_t$  is in  $L^1$  since  $\varphi$  is, as can be seen by a change of variables.)

# The identity

Let  $\varphi \in L^1(\mathbb{R}^n)$  be a radial function with  $\|\varphi\|_1 = 1$ . Let  $\psi_0 : \mathbb{R} \rightarrow \mathbb{R}$  be defined as

$$\psi_0(\|x\|) = \varphi(x)$$

for  $x \in \mathbb{R}^n$ . (Well-defined by radial assumption.)

For  $t > 0$ , define  $\varphi_t \in L^1(\mathbb{R}^n)$  by

$$\varphi_t(x) = t^{-n} \varphi(x/t).$$

( $\varphi_t$  is in  $L^1$  since  $\varphi$  is, as can be seen by a change of variables.)

Recall that we had seen that  $\{\varphi_t\}_{t>0}$  constitutes an approximate identity.

# The identity theorem

## Theorem (Main theorem)

# The identity theorem

## Theorem (Main theorem)

Suppose that  $\varphi$  is non-negative.

## Theorem (Main theorem)

Suppose that  $\varphi$  is non-negative. Further assume that  $\psi_0$  is in  $L^1(\mathbb{R}^n)$  and is decreasing.

## Theorem (Main theorem)

Suppose that  $\varphi$  is non-negative. Further assume that  $\psi_0$  is in  $L^1(\mathbb{R}^n)$  and is decreasing. Let  $f \in L^1(\mathbb{R}^n)$ .

## Theorem (Main theorem)

Suppose that  $\varphi$  is non-negative. Further assume that  $\psi_0$  is in  $L^1(\mathbb{R}^n)$  and is decreasing. Let  $f \in L^1(\mathbb{R}^n)$ . Then,

$$\lim_{t \rightarrow 0} (f * \varphi_t)(x) = f(x)$$



## Theorem (Main theorem)

Suppose that  $\varphi$  is non-negative. Further assume that  $\psi_0$  is in  $L^1(\mathbb{R}^n)$  and is decreasing. Let  $f \in L^1(\mathbb{R}^n)$ . Then,

$$\lim_{t \rightarrow 0} (f * \varphi_t)(x) = f(x)$$

for all  $x \in \text{Leb}(f)$ .

# Less abstract, more concrete

# Less abstract, more concrete

For the remainder of the discussion, one may forget the general setup and consider the following example for concreteness' sake.

# Less abstract, more concrete

For the remainder of the discussion, one may forget the general setup and consider the following example for concreteness' sake. Let  $\varphi$  be defined as

For the remainder of the discussion, one may forget the general setup and consider the following example for concreteness' sake. Let  $\varphi$  be defined as

$$\varphi(x) = e^{-\pi\|x\|^2}.$$

For the remainder of the discussion, one may forget the general setup and consider the following example for concreteness' sake. Let  $\varphi$  be defined as

$$\varphi(x) = e^{-\pi\|x\|^2}.$$

This is radial, non-negative, radially decreasing and we have  $\|\varphi\|_1 = 1$ .

For the remainder of the discussion, one may forget the general setup and consider the following example for concreteness' sake. Let  $\varphi$  be defined as

$$\varphi(x) = e^{-\pi\|x\|^2}.$$

This is radial, non-negative, radially decreasing and we have  $\|\varphi\|_1 = 1$ . The above is just the kernel  $h_1$ .

For the remainder of the discussion, one may forget the general setup and consider the following example for concreteness' sake. Let  $\varphi$  be defined as

$$\varphi(x) = e^{-\pi\|x\|^2}.$$

This is radial, non-negative, radially decreasing and we have  $\|\varphi\|_1 = 1$ . The above is just the kernel  $h_1$ .

For  $t > 0$ , we have

$$\varphi_t(x) = t^{-n} \exp\left(-\frac{\pi}{t^2}\|x\|^2\right).$$



# Less abstract, more concrete

For the remainder of the discussion, one may forget the general setup and consider the following example for concreteness' sake. Let  $\varphi$  be defined as

$$\varphi(x) = e^{-\pi\|x\|^2}.$$

This is radial, non-negative, radially decreasing and we have  $\|\varphi\|_1 = 1$ . The above is just the kernel  $h_1$ .

For  $t > 0$ , we have

$$\varphi_t(x) = t^{-n} \exp\left(-\frac{\pi}{t^2}\|x\|^2\right).$$

The above isn't exactly the heat kernel  $h_t$ .

For the remainder of the discussion, one may forget the general setup and consider the following example for concreteness' sake. Let  $\varphi$  be defined as

$$\varphi(x) = e^{-\pi\|x\|^2}.$$

This is radial, non-negative, radially decreasing and we have  $\|\varphi\|_1 = 1$ . The above is just the kernel  $h_1$ .

For  $t > 0$ , we have

$$\varphi_t(x) = t^{-n} \exp\left(-\frac{\pi}{t^2}\|x\|^2\right).$$

The above isn't exactly the heat kernel  $h_t$ . Rather, it is  $h_{t^2}$ .

# Less abstract, more concrete

For the remainder of the discussion, one may forget the general setup and consider the following example for concreteness' sake. Let  $\varphi$  be defined as

$$\varphi(x) = e^{-\pi\|x\|^2}.$$

This is radial, non-negative, radially decreasing and we have  $\|\varphi\|_1 = 1$ . The above is just the kernel  $h_1$ .

For  $t > 0$ , we have

$$\varphi_t(x) = t^{-n} \exp\left(-\frac{\pi}{t^2}\|x\|^2\right).$$

The above isn't exactly the heat kernel  $h_t$ . Rather, it is  $h_{t^2}$ .

It is now clear that proving the Main Theorem will show that  $(\star)$  holds for  $x \in \text{Leb}(f)$ .

# Some final notation

$S^{n-1}$  will denote the  $n - 1$  sphere in  $\mathbb{R}^n$ .

$S^{n-1}$  will denote the  $n - 1$  sphere in  $\mathbb{R}^n$ . More explicitly,

$$S^{n-1} = \{x \in \mathbb{R}^n : \|x\| = 1\}.$$

$S^{n-1}$  will denote the  $n - 1$  sphere in  $\mathbb{R}^n$ . More explicitly,

$$S^{n-1} = \{x \in \mathbb{R}^n : \|x\| = 1\}.$$

For  $x \in \mathbb{R}^n$  and  $r > 0$ ,  $B(x, r) = \{y \in \mathbb{R}^n : \|y - x\| < r\}$ .

$S^{n-1}$  will denote the  $n - 1$  sphere in  $\mathbb{R}^n$ . More explicitly,

$$S^{n-1} = \{x \in \mathbb{R}^n : \|x\| = 1\}.$$

For  $x \in \mathbb{R}^n$  and  $r > 0$ ,  $B(x, r) = \{y \in \mathbb{R}^n : \|y - x\| < r\}$ .

$V_n$  is the volume of the unit ball  $B(0, 1)$ .



$S^{n-1}$  will denote the  $n - 1$  sphere in  $\mathbb{R}^n$ . More explicitly,

$$S^{n-1} = \{x \in \mathbb{R}^n : \|x\| = 1\}.$$

For  $x \in \mathbb{R}^n$  and  $r > 0$ ,  $B(x, r) = \{y \in \mathbb{R}^n : \|y - x\| < r\}$ .

$V_n$  is the volume of the unit ball  $B(0, 1)$ . Thus, we have

$$\int_{B(x, r)} 1 = V_n r^n$$

for  $x \in \mathbb{R}^n$  and  $r > 0$ .

# Recap on Polar Coordinates

Recall that given  $f \in L^1(\mathbb{R}^n)$ , we may compute its integral as

# Recap on Polar Coordinates

Recall that given  $f \in L^1(\mathbb{R}^n)$ , we may compute its integral as

$$\int_{\mathbb{R}^n} f(x) \, dx = \int_0^\infty \int_{S^{n-1}} r^{n-1} f(r\omega) \, d\omega \, dr.$$

Recall that given  $f \in L^1(\mathbb{R}^n)$ , we may compute its integral as

$$\int_{\mathbb{R}^n} f(x) \, dx = \int_0^\infty \int_{S^{n-1}} r^{n-1} f(r\omega) \, d\omega \, dr.$$

In particular, if  $f$  is a radial function

Recall that given  $f \in L^1(\mathbb{R}^n)$ , we may compute its integral as

$$\int_{\mathbb{R}^n} f(x) \, dx = \int_0^\infty \int_{S^{n-1}} r^{n-1} f(r\omega) \, d\omega \, dr.$$

In particular, if  $f$  is a radial function and  $g$  is such that  $f(x) = g(\|x\|)$ , then

# Recap on Polar Coordinates

Recall that given  $f \in L^1(\mathbb{R}^n)$ , we may compute its integral as

$$\int_{\mathbb{R}^n} f(x) \, dx = \int_0^\infty \int_{S^{n-1}} r^{n-1} f(r\omega) \, d\omega \, dr.$$

In particular, if  $f$  is a radial function and  $g$  is such that  $f(x) = g(\|x\|)$ , then

$$\int_{\mathbb{R}^n} f(x) \, dx = \omega(S^{n-1}) \int_0^\infty r^{n-1} g(r) \, dr.$$

# Proof of the Main Theorem

- 1 Recap
- 2 Notations and Setup
- 3 Proof of the Main Theorem
- 4 The Stronger Theorem



# Proof of the Main Theorem

Fix a point  $x \in \text{Leb}(f)$  and let  $\epsilon > 0$  be arbitrary.

# Proof of the Main Theorem

Fix a point  $x \in \text{Leb}(f)$  and let  $\epsilon > 0$  be arbitrary.  
By definition of Lebesgue set, there exists  $\delta > 0$  such that

# Proof of the Main Theorem

Fix a point  $x \in \text{Leb}(f)$  and let  $\epsilon > 0$  be arbitrary.

By definition of Lebesgue set, there exists  $\delta > 0$  such that

$$\int_{|t|<r} |f(x-u) - f(x)| \, du < \epsilon r^n \quad (L)$$

# Proof of the Main Theorem

Fix a point  $x \in \text{Leb}(f)$  and let  $\epsilon > 0$  be arbitrary.  
By definition of Lebesgue set, there exists  $\delta > 0$  such that

$$\int_{|t| < r} |f(x - u) - f(x)| \, du < \epsilon r^n \quad (L)$$

for all  $0 < r \leq \delta$ .

# Proof of the Main Theorem

Fix a point  $x \in \text{Leb}(f)$  and let  $\epsilon > 0$  be arbitrary.

By definition of Lebesgue set, there exists  $\delta > 0$  such that

$$\int_{|t| < r} |f(x - u) - f(x)| \, du < \epsilon r^n \quad (L)$$

for all  $0 < r \leq \delta$ .

Note that for all  $t > 0$ , we have  $\int_{\mathbb{R}^n} \varphi_t = \int_{\mathbb{R}^n} \varphi = \int_{\mathbb{R}^n} |\varphi| = 1$ .

# Still proving the main theorem

# Still proving the main theorem

Thus, for all  $t > 0$ , we have

# Still proving the main theorem

Thus, for all  $t > 0$ , we have

$$|(f * \varphi_t)(x) - f(x)| = \left| \int_{\mathbb{R}^n} f(x - u) \varphi_t(u) \, du - f(x) \right|$$



# Still proving the main theorem

Thus, for all  $t > 0$ , we have

$$\begin{aligned}|(f * \varphi_t)(x) - f(x)| &= \left| \int_{\mathbb{R}^n} f(x-u)\varphi_t(u) \, du - f(x) \right| \\ &= \left| \int_{\mathbb{R}^n} f(x-u)\varphi_t(u) \, du - f(x) \cdot 1 \right|\end{aligned}$$

# Still proving the main theorem

Thus, for all  $t > 0$ , we have

$$\begin{aligned}|(f * \varphi_t)(x) - f(x)| &= \left| \int_{\mathbb{R}^n} f(x - u) \varphi_t(u) \, du - f(x) \right| \\ &= \left| \int_{\mathbb{R}^n} f(x - u) \varphi_t(u) \, du - f(x) \int_{\mathbb{R}^n} \varphi_t(u) \, du \right|\end{aligned}$$

# Still proving the main theorem

Thus, for all  $t > 0$ , we have

$$\begin{aligned}|(f * \varphi_t)(x) - f(x)| &= \left| \int_{\mathbb{R}^n} f(x-u)\varphi_t(u) \, du - f(x) \right| \\&= \left| \int_{\mathbb{R}^n} f(x-u)\varphi_t(u) \, du - f(x) \int_{\mathbb{R}^n} \varphi_t(u) \, du \right| \\&= \left| \int_{\mathbb{R}^n} [f(x-u) - f(x)]\varphi_t(u) \, du \right|\end{aligned}$$

# Still proving the main theorem

Thus, for all  $t > 0$ , we have

$$\begin{aligned} |(f * \varphi_t)(x) - f(x)| &= \left| \int_{\mathbb{R}^n} f(x-u) \varphi_t(u) \, du - f(x) \right| \\ &= \left| \int_{\mathbb{R}^n} f(x-u) \varphi_t(u) \, du - f(x) \int_{\mathbb{R}^n} \varphi_t(u) \, du \right| \\ &= \left| \int_{\mathbb{R}^n} [f(x-u) - f(x)] \varphi_t(u) \, du \right| \\ &\leq \left| \int_{|u| < \delta} [f(x-u) - f(x)] \varphi_t(u) \, du \right| \\ &\quad + \left| \int_{|u| \geq \delta} [f(x-u) - f(x)] \varphi_t(u) \, du \right| \end{aligned}$$

# Still proving the main theorem

Thus, for all  $t > 0$ , we have

$$\begin{aligned} |(f * \varphi_t)(x) - f(x)| &= \left| \int_{\mathbb{R}^n} f(x-u) \varphi_t(u) \, du - f(x) \right| \\ &= \left| \int_{\mathbb{R}^n} f(x-u) \varphi_t(u) \, du - f(x) \int_{\mathbb{R}^n} \varphi_t(u) \, du \right| \\ &= \left| \int_{\mathbb{R}^n} [f(x-u) - f(x)] \varphi_t(u) \, du \right| \\ &\leq \left| \int_{|u| < \delta} [f(x-u) - f(x)] \varphi_t(u) \, du \right| \rightsquigarrow I_1 \\ &\quad + \left| \int_{|u| \geq \delta} [f(x-u) - f(x)] \varphi_t(u) \, du \right| \rightsquigarrow I_2 \end{aligned}$$

# Still proving the main theorem

Thus, for all  $t > 0$ , we have

$$\begin{aligned} |(f * \varphi_t)(x) - f(x)| &= \left| \int_{\mathbb{R}^n} f(x-u) \varphi_t(u) \, du - f(x) \right| \\ &= \left| \int_{\mathbb{R}^n} f(x-u) \varphi_t(u) \, du - f(x) \int_{\mathbb{R}^n} \varphi_t(u) \, du \right| \\ &= \left| \int_{\mathbb{R}^n} [f(x-u) - f(x)] \varphi_t(u) \, du \right| \\ &\leq \left| \int_{|u| < \delta} [f(x-u) - f(x)] \varphi_t(u) \, du \right| \rightsquigarrow I_1(t) \\ &\quad + \left| \int_{|u| \geq \delta} [f(x-u) - f(x)] \varphi_t(u) \, du \right| \rightsquigarrow I_2(t) \end{aligned}$$

First, we note that

First, we note that

$$\int_{r/2 \leq |u| \leq r} \varphi(u) \, du$$



First, we note that

$$\int_{r/2 \leq |x| \leq r} \psi_0(r) \, du \leq \int_{r/2 \leq |u| \leq r} \varphi(u) \, du$$

First, we note that

$$\left( \int_{r/2 \leq |x| \leq r} 1 \right) \psi_0(r) \leq \int_{r/2 \leq |u| \leq r} \varphi(u) \, du$$

First, we note that

$$V_n \left(1 - \frac{1}{2^n}\right) r^n \psi_0(r) \leq \int_{r/2 \leq |u| \leq r} \varphi(u) \, du$$

First, we note that

$$V_n \left(1 - \frac{1}{2^n}\right) r^n \psi_0(r) \leq \int_{r/2 \leq |u| \leq r} \varphi(u) \, du \rightarrow 0,$$

First, we note that

$$V_n \left(1 - \frac{1}{2^n}\right) r^n \psi_0(r) \leq \int_{r/2 \leq |u| \leq r} \varphi(u) \, du \rightarrow 0,$$

as  $r \rightarrow 0$

First, we note that

$$V_n \left(1 - \frac{1}{2^n}\right) r^n \psi_0(r) \leq \int_{r/2 \leq |u| \leq r} \varphi(u) \, du \rightarrow 0,$$

as  $r \rightarrow 0$  or as  $r \rightarrow \infty$ .

First, we note that

$$V_n \left(1 - \frac{1}{2^n}\right) r^n \psi_0(r) \leq \int_{r/2 \leq |u| \leq r} \varphi(u) \, du \rightarrow 0,$$

as  $r \rightarrow 0$  or as  $r \rightarrow \infty$ .

Thus,  $r^n \psi_0(r) \rightarrow 0$

First, we note that

$$V_n \left(1 - \frac{1}{2^n}\right) r^n \psi_0(r) \leq \int_{r/2 \leq |u| \leq r} \varphi(u) \, du \rightarrow 0,$$

as  $r \rightarrow 0$  or as  $r \rightarrow \infty$ .

Thus,  $r^n \psi_0(r) \rightarrow 0$  as  $r$  tends to 0 or  $\infty$ .



First, we note that

$$V_n \left(1 - \frac{1}{2^n}\right) r^n \psi_0(r) \leq \int_{r/2 \leq |u| \leq r} \varphi(u) \, du \rightarrow 0,$$

as  $r \rightarrow 0$  or as  $r \rightarrow \infty$ .

Thus,  $r^n \psi_0(r) \rightarrow 0$  as  $r$  tends to 0 or  $\infty$ .

Hence, there exists  $A > 0$  such that

First, we note that

$$V_n \left(1 - \frac{1}{2^n}\right) r^n \psi_0(r) \leq \int_{r/2 \leq |u| \leq r} \varphi(u) \, du \rightarrow 0,$$

as  $r \rightarrow 0$  or as  $r \rightarrow \infty$ .

Thus,  $r^n \psi_0(r) \rightarrow 0$  as  $r$  tends to 0 or  $\infty$ .

Hence, there exists  $A > 0$  such that  $r^n \psi_0(r) \leq A$

First, we note that

$$V_n \left(1 - \frac{1}{2^n}\right) r^n \psi_0(r) \leq \int_{r/2 \leq |u| \leq r} \varphi(u) \, du \rightarrow 0,$$

as  $r \rightarrow 0$  or as  $r \rightarrow \infty$ .

Thus,  $r^n \psi_0(r) \rightarrow 0$  as  $r$  tends to 0 or  $\infty$ .

Hence, there exists  $A > 0$  such that  $r^n \psi_0(r) \leq A$  for  $r \in (0, \infty)$ .

First, we note that

$$V_n \left(1 - \frac{1}{2^n}\right) r^n \psi_0(r) \leq \int_{r/2 \leq |u| \leq r} \varphi(u) \, du \rightarrow 0,$$

as  $r \rightarrow 0$  or as  $r \rightarrow \infty$ .

Thus,  $r^n \psi_0(r) \rightarrow 0$  as  $r$  tends to 0 or  $\infty$ .

Hence, there exists  $A > 0$  such that  $r^n \psi_0(r) \leq A$  for  $r \in (0, \infty)$ .

Using this, we first show that  $I_2(t) \xrightarrow{t \rightarrow 0} 0$ .

$$l_2(t)$$

$$I_2(t) = \left| \int_{|u| \geq \delta} [f(x-u) - f(x)] \varphi_t(u) \, du \right|$$

$$I_2(t) \leq \int_{|u| \geq \delta} |f(x-u)| \varphi_t(u) \, du + |f(x)| \int_{|u| \geq \delta} \varphi_t(u) \, du.$$

$$l_2(t) \leq \int_{|u| \geq \delta} |f(x-u)| \varphi_t(u) \, du + |f(x)| \int_{|u| \geq \delta} \varphi_t(u) \, du.$$

The second term goes to 0 as  $t \rightarrow 0$



$$I_2(t) \leq \int_{|u| \geq \delta} |f(x-u)| \varphi_t(u) \, du + |f(x)| \int_{|u| \geq \delta} \varphi_t(u) \, du.$$

The second term goes to 0 as  $t \rightarrow 0$  since  $\{\varphi_t\}_{t>0}$  is an approximate identity.

$$I_2(t) \leq \int_{|u| \geq \delta} |f(x-u)| \varphi_t(u) \, du + |f(x)| \int_{|u| \geq \delta} \varphi_t(u) \, du.$$

The second term goes to 0 as  $t \rightarrow 0$  since  $\{\varphi_t\}_{t>0}$  is an approximate identity.

Now, let  $\chi_\delta$  denote the characteristic function of  $\{u \in \mathbb{R}^n : |u| \geq \delta\}$ .

$$I_2(t) \leq \int_{|u| \geq \delta} |f(x-u)| \varphi_t(u) \, du + |f(x)| \int_{|u| \geq \delta} \varphi_t(u) \, du.$$

The second term goes to 0 as  $t \rightarrow 0$  since  $\{\varphi_t\}_{t>0}$  is an approximate identity.

Now, let  $\chi_\delta$  denote the characteristic function of  $\{u \in \mathbb{R}^n : |u| \geq \delta\}$ .

We see that the first term is at most  $\|f\|_1 \|\chi_\delta \varphi_t\|_\infty$ .

$$I_2(t) \leq \int_{|u| \geq \delta} |f(x-u)| \varphi_t(u) \, du + |f(x)| \int_{|u| \geq \delta} \varphi_t(u) \, du.$$

The second term goes to 0 as  $t \rightarrow 0$  since  $\{\varphi_t\}_{t>0}$  is an approximate identity.

Now, let  $\chi_\delta$  denote the characteristic function of  $\{u \in \mathbb{R}^n : |u| \geq \delta\}$ .

We see that the first term is at most  $\|f\|_1 \|\chi_\delta \varphi_t\|_\infty$ . Since  $\varphi$  is radially decreasing, we see that

$$I_2(t) \leq \int_{|u| \geq \delta} |f(x-u)| \varphi_t(u) \, du + |f(x)| \int_{|u| \geq \delta} \varphi_t(u) \, du.$$

The second term goes to 0 as  $t \rightarrow 0$  since  $\{\varphi_t\}_{t>0}$  is an approximate identity.

Now, let  $\chi_\delta$  denote the characteristic function of  $\{u \in \mathbb{R}^n : |u| \geq \delta\}$ .

We see that the first term is at most  $\|f\|_1 \|\chi_\delta \varphi_t\|_\infty$ . Since  $\varphi$  is radially decreasing, we see that

$$\|\chi_\delta \varphi_t\|_\infty = \sup_{\|u\| \geq \delta} \varphi_t(u)$$

$$I_2(t) \leq \int_{|u| \geq \delta} |f(x-u)| \varphi_t(u) \, du + |f(x)| \int_{|u| \geq \delta} \varphi_t(u) \, du.$$

The second term goes to 0 as  $t \rightarrow 0$  since  $\{\varphi_t\}_{t>0}$  is an approximate identity.

Now, let  $\chi_\delta$  denote the characteristic function of  $\{u \in \mathbb{R}^n : |u| \geq \delta\}$ .

We see that the first term is at most  $\|f\|_1 \|\chi_\delta \varphi_t\|_\infty$ . Since  $\varphi$  is radially decreasing, we see that

$$\|\chi_\delta \varphi_t\|_\infty = \sup_{|u| \geq \delta} t^{-n} \varphi(u/t)$$

$$I_2(t) \leq \int_{|u| \geq \delta} |f(x-u)| \varphi_t(u) \, du + |f(x)| \int_{|u| \geq \delta} \varphi_t(u) \, du.$$

The second term goes to 0 as  $t \rightarrow 0$  since  $\{\varphi_t\}_{t>0}$  is an approximate identity.

Now, let  $\chi_\delta$  denote the characteristic function of  $\{u \in \mathbb{R}^n : |u| \geq \delta\}$ .

We see that the first term is at most  $\|f\|_1 \|\chi_\delta \varphi_t\|_\infty$ . Since  $\varphi$  is radially decreasing, we see that

$$\|\chi_\delta \varphi_t\|_\infty = \sup_{\|u\| \geq \delta} t^{-n} \varphi(u/t) = t^{-n} \psi_0(\delta/t)$$

$$I_2(t) \leq \int_{|u| \geq \delta} |f(x-u)| \varphi_t(u) \, du + |f(x)| \int_{|u| \geq \delta} \varphi_t(u) \, du.$$

The second term goes to 0 as  $t \rightarrow 0$  since  $\{\varphi_t\}_{t>0}$  is an approximate identity.

Now, let  $\chi_\delta$  denote the characteristic function of  $\{u \in \mathbb{R}^n : |u| \geq \delta\}$ .

We see that the first term is at most  $\|f\|_1 \|\chi_\delta \varphi_t\|_\infty$ . Since  $\varphi$  is radially decreasing, we see that

$$\|\chi_\delta \varphi_t\|_\infty = \sup_{\|u\| \geq \delta} t^{-n} \varphi(u/t) = \delta^{-n} (\delta/t)^n \psi_0(\delta/t)$$



$$I_2(t) \leq \int_{|u| \geq \delta} |f(x-u)| \varphi_t(u) \, du + |f(x)| \int_{|u| \geq \delta} \varphi_t(u) \, du.$$

The second term goes to 0 as  $t \rightarrow 0$  since  $\{\varphi_t\}_{t>0}$  is an approximate identity.

Now, let  $\chi_\delta$  denote the characteristic function of  $\{u \in \mathbb{R}^n : |u| \geq \delta\}$ .

We see that the first term is at most  $\|f\|_1 \|\chi_\delta \varphi_t\|_\infty$ . Since  $\varphi$  is radially decreasing, we see that

$$\|\chi_\delta \varphi_t\|_\infty = \sup_{\|u\| \geq \delta} t^{-n} \varphi(u/t) = \delta^{-n} (\delta/t)^n \psi_0(\delta/t) \rightarrow 0,$$

$$l_2(t) \leq \int_{|u| \geq \delta} |f(x-u)| \varphi_t(u) du + |f(x)| \int_{|u| \geq \delta} \varphi_t(u) du.$$

The second term goes to 0 as  $t \rightarrow 0$  since  $\{\varphi_t\}_{t>0}$  is an approximate identity.

Now, let  $\chi_\delta$  denote the characteristic function of  $\{u \in \mathbb{R}^n : |u| \geq \delta\}$ .

We see that the first term is at most  $\|f\|_1 \|\chi_\delta \varphi_t\|_\infty$ . Since  $\varphi$  is radially decreasing, we see that

$$\|\chi_\delta \varphi_t\|_\infty = \sup_{\|u\| \geq \delta} t^{-n} \varphi(u/t) = \delta^{-n} (\delta/t)^n \psi_0(\delta/t) \rightarrow 0,$$

as  $t \rightarrow 0$ .

$$I_2(t) \leq \int_{|u| \geq \delta} |f(x-u)| \varphi_t(u) \, du + |f(x)| \int_{|u| \geq \delta} \varphi_t(u) \, du.$$

The second term goes to 0 as  $t \rightarrow 0$  since  $\{\varphi_t\}_{t>0}$  is an approximate identity.

Now, let  $\chi_\delta$  denote the characteristic function of  $\{u \in \mathbb{R}^n : |u| \geq \delta\}$ .

We see that the first term is at most  $\|f\|_1 \|\chi_\delta \varphi_t\|_\infty$ . Since  $\varphi$  is radially decreasing, we see that

$$\|\chi_\delta \varphi_t\|_\infty = \sup_{|u| \geq \delta} t^{-n} \varphi(u/t) = \delta^{-n} (\delta/t)^n \psi_0(\delta/t) \rightarrow 0,$$

as  $t \rightarrow 0$ . Thus,  $I_2(t) \xrightarrow{t \rightarrow 0} 0$ .

$l_2$  down,  $l_1$  to go

Let us now define

$$g(r) = \int_{S^{n-1}} |f(x - r\omega) - f(x)| \, d\omega$$

Let us now define

$$g(r) = \int_{S^{n-1}} |f(x - r\omega) - f(x)| \, d\omega$$

and

$$G(r) = \int_0^r s^{n-1} g(s) \, ds.$$

Let us now define

$$g(r) = \int_{S^{n-1}} |f(x - r\omega) - f(x)| \, d\omega$$

and

$$G(r) = \int_0^r s^{n-1} g(s) \, ds.$$

Thus, the Lebesgue set condition ( $L$ ) from earlier translates to

Let us now define

$$g(r) = \int_{S^{n-1}} |f(x - r\omega) - f(x)| \, d\omega$$

and

$$G(r) = \int_0^r s^{n-1} g(s) \, ds.$$

Thus, the Lebesgue set condition ( $L$ ) from earlier translates to

$$G(r) \leq \epsilon r^n$$



Let us now define

$$g(r) = \int_{S^{n-1}} |f(x - r\omega) - f(x)| \, d\omega$$

and

$$G(r) = \int_0^r s^{n-1} g(s) \, ds.$$

Thus, the Lebesgue set condition ( $L$ ) from earlier translates to

$$G(r) \leq \epsilon r^n \quad \text{for } r \leq \delta.$$

Note that  $G(0) = 0$ .

Let us now define

$$g(r) = \int_{S^{n-1}} |f(x - r\omega) - f(x)| \, d\omega$$

and

$$G(r) = \int_0^r s^{n-1} g(s) \, ds.$$

Thus, the Lebesgue set condition ( $L$ ) from earlier translates to

$$G(r) \leq \epsilon r^n \quad \text{for } r \leq \delta.$$

Note that  $G(0) = 0$ .

With these notations, we do some more calculations.

# Some more calculations

We have

# Some more calculations

We have

$$I_1(t) = \left| \int_{|u| < \delta} [f(x-u) - f(x)] \varphi_t(u) \, du \right|$$

# Some more calculations

We have

$$I_1(t) \leq \int_{|u| < \delta} |f(x-u) - f(x)| \varphi_t(u) \, du$$

# Some more calculations

We have

$$I_1(t) \leq \int_{|u| < \delta} |f(x-u) - f(x)| t^{-n} \varphi(u/t) \, du$$

# Some more calculations

We have

$$\begin{aligned} I_1(t) &\leq \int_{|u| < \delta} |f(x-u) - f(x)| t^{-n} \varphi(u/t) \, du \\ &= \int_0^\delta r^{n-1} g(r) t^{-n} \psi_0(r/t) \, dr \end{aligned}$$

# Some more calculations

We have

$$\begin{aligned} I_1(t) &\leq \int_{|u|<\delta} |f(x-u) - f(x)| t^{-n} \varphi(u/t) \, du \\ &= \int_0^\delta r^{n-1} g(r) t^{-n} \psi_0(r/t) \, dr \end{aligned}$$

Integrate by parts



# Some more calculations

We have

$$\begin{aligned} I_1(t) &\leq \int_{|u| < \delta} |f(x-u) - f(x)| t^{-n} \varphi(u/t) \, du \\ &= \int_0^\delta r^{n-1} g(r) t^{-n} \psi_0(r/t) \, dr \\ &= G(r) t^{-n} \psi_0(r/t) \Big|_0^\delta - \int_0^\delta G(r) \, d(t^{-n} \psi_0(r/t)) \end{aligned}$$

# Some more calculations

We have

$$\begin{aligned} I_1(t) &\leq \int_{|u| < \delta} |f(x-u) - f(x)| t^{-n} \varphi(u/t) \, du \\ &= \int_0^\delta r^{n-1} g(r) t^{-n} \psi_0(r/t) \, dr \\ &= G(\delta) t^{-n} \psi_0(\delta/t) - \int_0^\delta G(r) \, d(t^{-n} \psi_0(r/t)) \end{aligned}$$

# Some more calculations

We have

$$\begin{aligned} I_1(t) &\leq \int_{|u|<\delta} |f(x-u) - f(x)| t^{-n} \varphi(u/t) \, du \\ &= \int_0^\delta r^{n-1} g(r) t^{-n} \psi_0(r/t) \, dr \\ &= G(\delta) t^{-n} \psi_0(\delta/t) - \int_0^\delta G(r) \, d(t^{-n} \psi_0(r/t)) \\ &\leq \epsilon \delta^n t^{-n} \psi_0(\delta/t) - \int_0^{\delta/t} G(ts) t^{-n} \, d(\psi_0(s)) \end{aligned}$$

# Some more calculations

We have

$$\begin{aligned} I_1(t) &\leq \int_{|u|<\delta} |f(x-u) - f(x)| t^{-n} \varphi(u/t) \, du \\ &= \int_0^\delta r^{n-1} g(r) t^{-n} \psi_0(r/t) \, dr \\ &= G(\delta) t^{-n} \psi_0(\delta/t) - \int_0^\delta G(r) \, d(t^{-n} \psi_0(r/t)) \\ &\leq \epsilon(\delta/t)^n \psi_0(\delta/t) - \int_0^{\delta/t} G(ts) t^{-n} \, d(\psi_0(s)) \end{aligned}$$

# Some more calculations

We have

$$\begin{aligned} I_1(t) &\leq \int_{|u|<\delta} |f(x-u) - f(x)| t^{-n} \varphi(u/t) \, du \\ &= \int_0^\delta r^{n-1} g(r) t^{-n} \psi_0(r/t) \, dr \\ &= G(\delta) t^{-n} \psi_0(\delta/t) - \int_0^\delta G(r) \, d(t^{-n} \psi_0(r/t)) \\ &\leq \epsilon A - \int_0^{\delta/t} G(ts) t^{-n} \, d(\psi_0(s)) \end{aligned}$$

# Some more calculations

We have

$$\begin{aligned} I_1(t) &\leq \int_{|u|<\delta} |f(x-u) - f(x)| t^{-n} \varphi(u/t) \, du \\ &= \int_0^\delta r^{n-1} g(r) t^{-n} \psi_0(r/t) \, dr \\ &= G(\delta) t^{-n} \psi_0(\delta/t) - \int_0^\delta G(r) \, d(t^{-n} \psi_0(r/t)) \\ &\leq \epsilon A - \int_0^{\delta/t} G(ts) t^{-n} \, d(\psi_0(s)) \\ &\leq \epsilon A - \int_0^{\delta/t} \epsilon (ts)^n t^{-n} \, d(\psi_0(s)) \end{aligned}$$

Note that  $d\psi_0 \leq 0$ .

# Some more calculations

We have

$$\begin{aligned} I_1(t) &\leq \int_{|u|<\delta} |f(x-u) - f(x)| t^{-n} \varphi(u/t) \, du \\ &= \int_0^\delta r^{n-1} g(r) t^{-n} \psi_0(r/t) \, dr \\ &= G(\delta) t^{-n} \psi_0(\delta/t) - \int_0^\delta G(r) \, d(t^{-n} \psi_0(r/t)) \\ &\leq \epsilon A - \int_0^{\delta/t} G(ts) t^{-n} \, d(\psi_0(s)) \\ &\leq \epsilon A - \int_0^{\delta/t} \epsilon s^n \, d(\psi_0(s)) \end{aligned}$$

Note that  $d\psi_0 \leq 0$ .

# Some more calculations

We have

$$\begin{aligned} I_1(t) &\leq \int_{|u|<\delta} |f(x-u) - f(x)| t^{-n} \varphi(u/t) \, du \\ &= \int_0^\delta r^{n-1} g(r) t^{-n} \psi_0(r/t) \, dr \\ &= G(\delta) t^{-n} \psi_0(\delta/t) - \int_0^\delta G(r) \, d(t^{-n} \psi_0(r/t)) \\ &\leq \epsilon A - \int_0^{\delta/t} G(ts) t^{-n} \, d(\psi_0(s)) \\ &\leq \epsilon A - \int_0^{\delta/t} \epsilon s^n \, d(\psi_0(s)) \leq \epsilon \left( A - \int_0^\infty s^n \, d(\psi_0(s)) \right) \end{aligned}$$

Note that  $d\psi_0 \leq 0$ .



# Some more calculations

We have

$$\begin{aligned} I_1(t) &\leq \int_{|u|<\delta} |f(x-u) - f(x)| t^{-n} \varphi(u/t) \, du \\ &= \int_0^\delta r^{n-1} g(r) t^{-n} \psi_0(r/t) \, dr \\ &= G(\delta) t^{-n} \psi_0(\delta/t) - \int_0^\delta G(r) \, d(t^{-n} \psi_0(r/t)) \\ &\leq \epsilon A - \int_0^{\delta/t} G(ts) t^{-n} \, d(\psi_0(s)) \\ &\leq \epsilon A - \int_0^{\delta/t} \epsilon s^n \, d(\psi_0(s)) \leq \epsilon \left( A - \int_0^\infty s^n \, d(\psi_0(s)) \right) \end{aligned}$$

Note that  $d\psi_0 \leq 0$ .

The green integral can be calculated exactly quite simply.

The green integral can be calculated exactly quite simply.

$$- \int_0^\infty s^n d(\psi_0(s))$$

The green integral can be calculated exactly quite simply.

$$-\int_0^\infty s^n d(\psi_0(s)) = -s^n \psi_0(s) \Big|_0^\infty + n \int_0^\infty s^{n-1} \psi_0(s) ds$$

The green integral can be calculated exactly quite simply.

$$\begin{aligned} - \int_0^\infty s^n d(\psi_0(s)) &= -s^n \psi_0(s) \Big|_0^\infty + n \int_0^\infty s^{n-1} \psi_0(s) ds \\ &= 0 + \frac{n}{\omega(S^{n-1})} \int_{\mathbb{R}^n} \varphi \end{aligned}$$

The green integral can be calculated exactly quite simply.

$$\begin{aligned} - \int_0^\infty s^n d(\psi_0(s)) &= -s^n \psi_0(s) \Big|_0^\infty + n \int_0^\infty s^{n-1} \psi_0(s) ds \\ &= \frac{n}{\omega(S^{n-1})}. \end{aligned}$$

The green integral can be calculated exactly quite simply.

$$\begin{aligned} - \int_0^\infty s^n d(\psi_0(s)) &= -s^n \psi_0(s) \Big|_0^\infty + n \int_0^\infty s^{n-1} \psi_0(s) ds \\ &= \frac{n}{\omega(S^{n-1})}. \end{aligned}$$

Putting this back, we get

The green integral can be calculated exactly quite simply.

$$\begin{aligned} - \int_0^\infty s^n d(\psi_0(s)) &= -s^n \psi_0(s) \Big|_0^\infty + n \int_0^\infty s^{n-1} \psi_0(s) ds \\ &= \frac{n}{\omega(S^{n-1})}. \end{aligned}$$

Putting this back, we get

$$I_1(t) \leq \epsilon \left( A + \frac{n}{\omega(S^{n-1})} \right)$$



The green integral can be calculated exactly quite simply.

$$\begin{aligned} - \int_0^\infty s^n d(\psi_0(s)) &= -s^n \psi_0(s) \Big|_0^\infty + n \int_0^\infty s^{n-1} \psi_0(s) ds \\ &= \frac{n}{\omega(S^{n-1})}. \end{aligned}$$

Putting this back, we get

$$I_1(t) \leq \epsilon \left( A + \frac{n}{\omega(S^{n-1})} \right) = \epsilon B.$$

The green integral can be calculated exactly quite simply.

$$\begin{aligned} - \int_0^\infty s^n d(\psi_0(s)) &= -s^n \psi_0(s) \Big|_0^\infty + n \int_0^\infty s^{n-1} \psi_0(s) ds \\ &= \frac{n}{\omega(S^{n-1})}. \end{aligned}$$

Putting this back, we get

$$I_1(t) \leq \epsilon \left( A + \frac{n}{\omega(S^{n-1})} \right) = \epsilon B.$$

Thus, we have bounded  $I_1$

The green integral can be calculated exactly quite simply.

$$\begin{aligned} - \int_0^\infty s^n d(\psi_0(s)) &= -s^n \psi_0(s) \Big|_0^\infty + n \int_0^\infty s^{n-1} \psi_0(s) ds \\ &= \frac{n}{\omega(S^{n-1})}. \end{aligned}$$

Putting this back, we get

$$I_1(t) \leq \epsilon \left( A + \frac{n}{\omega(S^{n-1})} \right) = \epsilon B.$$

Thus, we have bounded  $I_1$  independent of  $t$

The green integral can be calculated exactly quite simply.

$$\begin{aligned} - \int_0^\infty s^n d(\psi_0(s)) &= -s^n \psi_0(s) \Big|_0^\infty + n \int_0^\infty s^{n-1} \psi_0(s) ds \\ &= \frac{n}{\omega(S^{n-1})}. \end{aligned}$$

Putting this back, we get

$$I_1(t) \leq \epsilon \left( A + \frac{n}{\omega(S^{n-1})} \right) = \epsilon B.$$

Thus, we have bounded  $I_1$  independent of  $t$  and of  $f$ .

# Putting it back

We had

We had

$$|(f * \varphi_t)(x) - f(x)| \leq I_1(t) + I_2(t).$$

We had

$$|(f * \varphi_t)(x) - f(x)| \leq I_1(t) + I_2(t).$$

We showed that  $I_1$  is bounded independent of  $t$ .

We had

$$|(f * \varphi_t)(x) - f(x)| \leq I_1(t) + I_2(t) \leq \epsilon B + I_2(t).$$

We showed that  $I_1$  is bounded independent of  $t$ .



We had

$$|(f * \varphi_t)(x) - f(x)| \leq I_1(t) + I_2(t) \leq \epsilon B + I_2(t).$$

We showed that  $I_1$  is bounded independent of  $t$ . We also showed that  $I_2(t) \xrightarrow{t \rightarrow 0} 0$ .

We had

$$|(f * \varphi_t)(x) - f(x)| \leq I_1(t) + I_2(t) \leq \epsilon B + I_2(t).$$

We showed that  $I_1$  is bounded independent of  $t$ . We also showed that  $I_2(t) \xrightarrow{t \rightarrow 0} 0$ . Thus, for  $t$  sufficiently small, we have

We had

$$|(f * \varphi_t)(x) - f(x)| \leq I_1(t) + I_2(t) \leq \epsilon B + I_2(t).$$

We showed that  $I_1$  is bounded independent of  $t$ . We also showed that  $I_2(t) \xrightarrow{t \rightarrow 0} 0$ . Thus, for  $t$  sufficiently small, we have

$$|(f * \varphi_t)(x) - f(x)| < \epsilon(B + 1).$$

We had

$$|(f * \varphi_t)(x) - f(x)| \leq I_1(t) + I_2(t) \leq \epsilon B + I_2(t).$$

We showed that  $I_1$  is bounded independent of  $t$ . We also showed that  $I_2(t) \xrightarrow{t \rightarrow 0} 0$ . Thus, for  $t$  sufficiently small, we have

$$|(f * \varphi_t)(x) - f(x)| < \epsilon(B + 1).$$

This completes the proof.

# The Stronger Theorem

- 1 Recap
- 2 Notations and Setup
- 3 Proof of the Main Theorem
- 4 The Stronger Theorem

# Concluding Remark

The theorem which I have proven is actually a weaker version of something more general.

# Concluding Remark

The theorem which I have proven is actually a weaker version of something more general. Forget all the notation and hypothesis that we had so far.

# Concluding Remark

The theorem which I have proven is actually a weaker version of something more general. Forget all the notation and hypothesis that we had so far.

## Theorem (General Theorem)

Suppose  $\varphi \in L^1(\mathbb{R}^n)$ .



# Concluding Remark

The theorem which I have proven is actually a weaker version of something more general. Forget all the notation and hypothesis that we had so far.

## Theorem (General Theorem)

Suppose  $\varphi \in L^1(\mathbb{R}^n)$ . Let  $\psi(y) = \operatorname{ess\,sup}_{\|z\| \geq \|y\|} \varphi(z)$

# Concluding Remark

The theorem which I have proven is actually a weaker version of something more general. Forget all the notation and hypothesis that we had so far.

## Theorem (General Theorem)

Suppose  $\varphi \in L^1(\mathbb{R}^n)$ . Let  $\psi(y) = \operatorname{ess\,sup}_{\|z\| \geq \|y\|} \varphi(z)$  and for  $t > 0$ , let  $\varphi_t(y) = t^{-n} \varphi(y/t)$ .

# Concluding Remark

The theorem which I have proven is actually a weaker version of something more general. Forget all the notation and hypothesis that we had so far.

## Theorem (General Theorem)

Suppose  $\varphi \in L^1(\mathbb{R}^n)$ . Let  $\psi(y) = \operatorname{ess\,sup}_{\|z\| \geq \|y\|} \varphi(z)$  and for  $t > 0$ , let  $\varphi_t(y) = t^{-n}\varphi(y/t)$ . If  $\psi \in L^1(\mathbb{R}^n)$  and  $f \in L^p(\mathbb{R}^n)$ ,  $1 \leq p \leq \infty$ ,

The theorem which I have proven is actually a weaker version of something more general. Forget all the notation and hypothesis that we had so far.

## Theorem (General Theorem)

Suppose  $\varphi \in L^1(\mathbb{R}^n)$ . Let  $\psi(y) = \operatorname{ess\,sup}_{\|z\| \geq \|y\|} \varphi(z)$  and for  $t > 0$ , let  $\varphi_t(y) = t^{-n} \varphi(y/t)$ . If  $\psi \in L^1(\mathbb{R}^n)$  and  $f \in L^p(\mathbb{R}^n)$ ,  $1 \leq p \leq \infty$ , then  $\lim_{t \rightarrow 0} (f * \varphi_t)(x) = f(x) \int_{\mathbb{R}^n} \varphi(t) \, dt$

# Concluding Remark

The theorem which I have proven is actually a weaker version of something more general. Forget all the notation and hypothesis that we had so far.

## Theorem (General Theorem)

Suppose  $\varphi \in L^1(\mathbb{R}^n)$ . Let  $\psi(y) = \operatorname{ess\,sup}_{\|z\| \geq \|y\|} \varphi(z)$  and for  $t > 0$ , let  $\varphi_t(y) = t^{-n} \varphi(y/t)$ . If  $\psi \in L^1(\mathbb{R}^n)$  and  $f \in L^p(\mathbb{R}^n)$ ,  $1 \leq p \leq \infty$ , then  $\lim_{t \rightarrow 0} (f * \varphi_t)(x) = f(x) \int_{\mathbb{R}^n} \varphi(t) \, dt$  whenever  $x \in \operatorname{Leb}(f)$ .

# Concluding Remark

The theorem which I have proven is actually a weaker version of something more general. Forget all the notation and hypothesis that we had so far.

## Theorem (General Theorem)

Suppose  $\varphi \in L^1(\mathbb{R}^n)$ . Let  $\psi(y) = \operatorname{ess\,sup}_{\|z\| \geq \|y\|} \varphi(z)$  and for  $t > 0$ , let  $\varphi_t(y) = t^{-n} \varphi(y/t)$ . If  $\psi \in L^1(\mathbb{R}^n)$  and  $f \in L^p(\mathbb{R}^n)$ ,  $1 \leq p \leq \infty$ , then  $\lim_{t \rightarrow 0} (f * \varphi_t)(x) = f(x) \int_{\mathbb{R}^n} \varphi(t) \, dt$  whenever  $x \in \operatorname{Leb}(f)$ .

Reference: *Introduction to Fourier Analysis on Euclidean Spaces* by Stein and Weiss