

Fourier Inversion for L^1 Functions

Aryaman Maithani

Department of Mathematics
IIT Bombay

March 17, 2021

- 1 Recap
- 2 Notations and Setup
- 3 Proof of the Main Theorem
- 4 The Stronger Theorem

- 1 Recap
- 2 Notations and Setup
- 3 Proof of the Main Theorem
- 4 The Stronger Theorem

Recall how we had proven Fourier inversion for L^1 functions in class.

Recall how we had proven Fourier inversion for L^1 functions in class. Fix $n \in \mathbb{N}$ for the entirety of this discussion.

Definition (Heat Kernel)

Recall how we had proven Fourier inversion for L^1 functions in class. Fix $n \in \mathbb{N}$ for the entirety of this discussion.

Definition (Heat Kernel)

For $t > 0$, define $h_t : \mathbb{R}^n \rightarrow \mathbb{R}$ by

Recall how we had proven Fourier inversion for L^1 functions in class. Fix $n \in \mathbb{N}$ for the entirety of this discussion.

Definition (Heat Kernel)

For $t > 0$, define $h_t : \mathbb{R}^n \rightarrow \mathbb{R}$ by

$$h_t(x) = \frac{1}{t^{n/2}} \exp\left(-\frac{\pi}{t} \|x\|^2\right).$$

The proof from class

We had seen that $\{h_t\}_{t>0}$ constitutes a (continuous) approximate identity.

The proof from class

We had seen that $\{h_t\}_{t>0}$ constitutes a (continuous) approximate identity. We thus concluded that $f * h_t \xrightarrow{t \rightarrow 0} f$ in L^1 for any $f \in L^1$.

The proof from class

We had seen that $\{h_t\}_{t>0}$ constitutes a (continuous) approximate identity. We thus concluded that $f * h_t \xrightarrow{t \rightarrow 0} f$ in L^1 for any $f \in L^1$. In turn, that gave us that there is a subsequence $\{t_n\}$ such that

$$\lim_{t_n \rightarrow 0} (f * h_{t_n})(x) = f(x) \quad (\star)$$

for almost all x .

The proof from class

We had seen that $\{h_t\}_{t>0}$ constitutes a (continuous) approximate identity. We thus concluded that $f * h_t \xrightarrow{t \rightarrow 0} f$ in L^1 for any $f \in L^1$. In turn, that gave us that there is a subsequence $\{t_n\}$ such that

$$\lim_{t_n \rightarrow 0} (f * h_{t_n})(x) = f(x) \quad (\star)$$

for almost all x .

We then showed the following assuming $\hat{f} \in L^1$.

The proof from class

We had seen that $\{h_t\}_{t>0}$ constitutes a (continuous) approximate identity. We thus concluded that $f * h_t \xrightarrow{t \rightarrow 0} f$ in L^1 for any $f \in L^1$. In turn, that gave us that there is a subsequence $\{t_n\}$ such that

$$\lim_{t_n \rightarrow 0} (f * h_{t_n})(x) = f(x) \quad (\star)$$

for almost all x .

We then showed the following assuming $\hat{f} \in L^1$.

$$\textcircled{1} \quad (f * h_t)(x) = \int_{\mathbb{R}^n} \hat{f}(\xi) e^{-\pi t \|\xi\|^2} e^{2\pi i x \cdot \xi} d\xi \text{ for } \underline{\text{all}} \ x \in \mathbb{R}^n.$$

The proof from class

We had seen that $\{h_t\}_{t>0}$ constitutes a (continuous) approximate identity. We thus concluded that $f * h_t \xrightarrow{t \rightarrow 0} f$ in L^1 for any $f \in L^1$. In turn, that gave us that there is a subsequence $\{t_n\}$ such that

$$\lim_{t_n \rightarrow 0} (f * h_{t_n})(x) = f(x) \quad (\star)$$

for almost all x .

We then showed the following assuming $\hat{f} \in L^1$.

❶ $(f * h_t)(x) = \int_{\mathbb{R}^n} \hat{f}(\xi) e^{-\pi t \|\xi\|^2} e^{2\pi i x \cdot \xi} d\xi$ for all $x \in \mathbb{R}^n$.

❷ Using DCT, we let $t \rightarrow 0$ in the above via $\{t_n\}$ to conclude that

$$f(x) = \int_{\mathbb{R}^n} \hat{f}(\xi) e^{2\pi i x \cdot \xi} d\xi$$

The proof from class

We had seen that $\{h_t\}_{t>0}$ constitutes a (continuous) approximate identity. We thus concluded that $f * h_t \xrightarrow{t \rightarrow 0} f$ in L^1 for any $f \in L^1$. In turn, that gave us that there is a subsequence $\{t_n\}$ such that

$$\lim_{t_n \rightarrow 0} (f * h_{t_n})(x) = f(x) \quad (\star)$$

for almost all x .

We then showed the following assuming $\hat{f} \in L^1$.

- ① $(f * h_t)(x) = \int_{\mathbb{R}^n} \hat{f}(\xi) e^{-\pi t \|\xi\|^2} e^{2\pi i x \cdot \xi} d\xi$ for all $x \in \mathbb{R}^n$.
- ② Using DCT, we let $t \rightarrow 0$ in the above via $\{t_n\}$ to conclude that

$$f(x) = \int_{\mathbb{R}^n} \hat{f}(\xi) e^{2\pi i x \cdot \xi} d\xi$$

for those $x \in \mathbb{R}^n$ for which (\star) holds.

Conclusion

Thus, we have actually proven that the Fourier inversion holds for those points x

Thus, we have actually proven that the Fourier inversion holds for those points x for which the equality

$$\lim_{t \rightarrow 0} (f * h_t)(x) = f(x) \quad (\star)$$

holds.

Thus, we have actually proven that the Fourier inversion holds for those points x for which the equality

$$\lim_{t \rightarrow 0} (f * h_t)(x) = f(x) \quad (\star)$$

holds. (In fact, we have something stronger since we allow $t \rightarrow 0$ via a subsequence.)

Thus, we have actually proven that the Fourier inversion holds for those points x for which the equality

$$\lim_{t \rightarrow 0} (f * h_t)(x) = f(x) \quad (\star)$$

holds. (In fact, we have something stronger since we allow $t \rightarrow 0$ via a subsequence.)

Our aim now is to show that (\star) holds for all x in the *Lebesgue set* of f .

Thus, we have actually proven that the Fourier inversion holds for those points x for which the equality

$$\lim_{t \rightarrow 0} (f * h_t)(x) = f(x) \quad (\star)$$

holds. (In fact, we have something stronger since we allow $t \rightarrow 0$ via a subsequence.)

Our aim now is to show that (\star) holds for all x in the *Lebesgue set* of f . (Even without passing to a subsequence.)

Thus, we have actually proven that the Fourier inversion holds for those points x for which the equality

$$\lim_{t \rightarrow 0} (f * h_t)(x) = f(x) \quad (\star)$$

holds. (In fact, we have something stronger since we allow $t \rightarrow 0$ via a subsequence.)

Our aim now is to show that (\star) holds for all x in the *Lebesgue set* of f . (Even without passing to a subsequence.)

We will actually prove the result for a broader class of approximate identities.

- 1 Recap
- 2 **Notations and Setup**
- 3 Proof of the Main Theorem
- 4 The Stronger Theorem

Note that if $f \in L^1$, then f is finite a.e. Thus, we may assume that f is finite everywhere by changing it on a null set.

Definition (Lebesgue set)

Note that if $f \in L^1$, then f is finite a.e. Thus, we may assume that f is finite everywhere by changing it on a null set.

Definition (Lebesgue set)

Suppose $f \in L^1(\mathbb{R}^n)$.

Note that if $f \in L^1$, then f is finite a.e. Thus, we may assume that f is finite everywhere by changing it on a null set.

Definition (Lebesgue set)

Suppose $f \in L^1(\mathbb{R}^n)$. The Lebesgue set of f , denoted $\text{Leb}(f)$

Note that if $f \in L^1$, then f is finite a.e. Thus, we may assume that f is finite everywhere by changing it on a null set.

Definition (Lebesgue set)

Suppose $f \in L^1(\mathbb{R}^n)$. The Lebesgue set of f , denoted $\text{Leb}(f)$ is defined to be the set of all $x \in \mathbb{R}^n$ for which

holds.

Note that if $f \in L^1$, then f is finite a.e. Thus, we may assume that f is finite everywhere by changing it on a null set.

Definition (Lebesgue set)

Suppose $f \in L^1(\mathbb{R}^n)$. The Lebesgue set of f , denoted $\text{Leb}(f)$ is defined to be the set of all $x \in \mathbb{R}^n$ for which

$$\lim_{r \rightarrow 0} \frac{1}{r^n} \int_{\|u\| < r} |f(x - u) - f(x)| \, du = 0$$

holds.

Note that if $f \in L^1$, then f is finite a.e. Thus, we may assume that f is finite everywhere by changing it on a null set.

Definition (Lebesgue set)

Suppose $f \in L^1(\mathbb{R}^n)$. The Lebesgue set of f , denoted $\text{Leb}(f)$ is defined to be the set of all $x \in \mathbb{R}^n$ for which

$$\lim_{r \rightarrow 0} \frac{1}{r^n} \int_{\|u\| < r} |f(x - u) - f(x)| \, du = 0$$

holds.

Note that the above $\text{Leb}(f)$ is actually a superset of the $\text{Leb}(f)$ we defined it in class.

Note that if $f \in L^1$, then f is finite a.e. Thus, we may assume that f is finite everywhere by changing it on a null set.

Definition (Lebesgue set)

Suppose $f \in L^1(\mathbb{R}^n)$. The Lebesgue set of f , denoted $\text{Leb}(f)$ is defined to be the set of all $x \in \mathbb{R}^n$ for which

$$\lim_{r \rightarrow 0} \frac{1}{r^n} \int_{\|u\| < r} |f(x - u) - f(x)| \, du = 0$$

holds.

Note that the above $\text{Leb}(f)$ is actually a superset of the $\text{Leb}(f)$ we defined in class. So, we shall prove a stronger result.

Let $\varphi \in L^1(\mathbb{R}^n)$ be a radial function with $\|\varphi\|_1 = 1$.

The identity

Let $\varphi \in L^1(\mathbb{R}^n)$ be a radial function with $\|\varphi\|_1 = 1$. Let $\psi_0 : [0, \infty) \rightarrow \mathbb{R}$ be defined as

The identity

Let $\varphi \in L^1(\mathbb{R}^n)$ be a radial function with $\|\varphi\|_1 = 1$. Let $\psi_0 : [0, \infty) \rightarrow \mathbb{R}$ be defined as

$$\psi_0(\|x\|) = \varphi(x)$$

for $x \in \mathbb{R}^n$.

The identity

Let $\varphi \in L^1(\mathbb{R}^n)$ be a radial function with $\|\varphi\|_1 = 1$. Let $\psi_0 : [0, \infty) \rightarrow \mathbb{R}$ be defined as

$$\psi_0(\|x\|) = \varphi(x)$$

for $x \in \mathbb{R}^n$. (Well-defined by radial assumption.)

The identity

Let $\varphi \in L^1(\mathbb{R}^n)$ be a radial function with $\|\varphi\|_1 = 1$. Let $\psi_0 : [0, \infty) \rightarrow \mathbb{R}$ be defined as

$$\psi_0(\|x\|) = \varphi(x)$$

for $x \in \mathbb{R}^n$. (Well-defined by radial assumption.)

For $t > 0$, define $\varphi_t \in L^1(\mathbb{R}^n)$ by

The identity

Let $\varphi \in L^1(\mathbb{R}^n)$ be a radial function with $\|\varphi\|_1 = 1$. Let $\psi_0 : [0, \infty) \rightarrow \mathbb{R}$ be defined as

$$\psi_0(\|x\|) = \varphi(x)$$

for $x \in \mathbb{R}^n$. (Well-defined by radial assumption.)

For $t > 0$, define $\varphi_t \in L^1(\mathbb{R}^n)$ by

$$\varphi_t(x) = t^{-n} \varphi(x/t).$$

The identity

Let $\varphi \in L^1(\mathbb{R}^n)$ be a radial function with $\|\varphi\|_1 = 1$. Let $\psi_0 : [0, \infty) \rightarrow \mathbb{R}$ be defined as

$$\psi_0(\|x\|) = \varphi(x)$$

for $x \in \mathbb{R}^n$. (Well-defined by radial assumption.)

For $t > 0$, define $\varphi_t \in L^1(\mathbb{R}^n)$ by

$$\varphi_t(x) = t^{-n} \varphi(x/t).$$

(φ_t is in L^1 since φ is, as can be seen by a change of variables.)

The identity

Let $\varphi \in L^1(\mathbb{R}^n)$ be a radial function with $\|\varphi\|_1 = 1$. Let $\psi_0 : [0, \infty) \rightarrow \mathbb{R}$ be defined as

$$\psi_0(\|x\|) = \varphi(x)$$

for $x \in \mathbb{R}^n$. (Well-defined by radial assumption.)

For $t > 0$, define $\varphi_t \in L^1(\mathbb{R}^n)$ by

$$\varphi_t(x) = t^{-n} \varphi(x/t).$$

(φ_t is in L^1 since φ is, as can be seen by a change of variables.)

Recall that we had seen that $\{\varphi_t\}_{t>0}$ constitutes an approximate identity.

The Main Theorem

Theorem (Main theorem)

Theorem (Main theorem)

Suppose that φ is non-negative.

Theorem (Main theorem)

Suppose that φ is non-negative. Further assume that ψ_0 is in $L^1(\mathbb{R}^n)$ and is decreasing.

Theorem (Main theorem)

Suppose that φ is non-negative. Further assume that ψ_0 is in $L^1(\mathbb{R}^n)$ and is decreasing. Let $f \in L^1(\mathbb{R}^n)$.

Theorem (Main theorem)

Suppose that φ is non-negative. Further assume that ψ_0 is in $L^1(\mathbb{R}^n)$ and is decreasing. Let $f \in L^1(\mathbb{R}^n)$. Then,

$$\lim_{t \rightarrow 0} (f * \varphi_t)(x) = f(x)$$

Theorem (Main theorem)

Suppose that φ is non-negative. Further assume that ψ_0 is in $L^1(\mathbb{R}^n)$ and is decreasing. Let $f \in L^1(\mathbb{R}^n)$. Then,

$$\lim_{t \rightarrow 0} (f * \varphi_t)(x) = f(x)$$

for all $x \in \text{Leb}(f)$.

Less abstract, more concrete

Less abstract, more concrete

For the remainder of the discussion, one may forget the general setup and consider the following example for concreteness' sake.

Less abstract, more concrete

For the remainder of the discussion, one may forget the general setup and consider the following example for concreteness' sake. Let φ be defined as

For the remainder of the discussion, one may forget the general setup and consider the following example for concreteness' sake. Let φ be defined as

$$\varphi(x) = e^{-\pi\|x\|^2}.$$

For the remainder of the discussion, one may forget the general setup and consider the following example for concreteness' sake. Let φ be defined as

$$\varphi(x) = e^{-\pi\|x\|^2}.$$

This is radial, non-negative, radially decreasing and we have $\|\varphi\|_1 = 1$.

For the remainder of the discussion, one may forget the general setup and consider the following example for concreteness' sake. Let φ be defined as

$$\varphi(x) = e^{-\pi\|x\|^2}.$$

This is radial, non-negative, radially decreasing and we have $\|\varphi\|_1 = 1$. The above is just the kernel h_1 .

For the remainder of the discussion, one may forget the general setup and consider the following example for concreteness' sake. Let φ be defined as

$$\varphi(x) = e^{-\pi\|x\|^2}.$$

This is radial, non-negative, radially decreasing and we have $\|\varphi\|_1 = 1$. The above is just the kernel h_1 .

For $t > 0$, we have

$$\varphi_t(x) = t^{-n} \exp\left(-\frac{\pi}{t^2}\|x\|^2\right).$$

Less abstract, more concrete

For the remainder of the discussion, one may forget the general setup and consider the following example for concreteness' sake. Let φ be defined as

$$\varphi(x) = e^{-\pi\|x\|^2}.$$

This is radial, non-negative, radially decreasing and we have $\|\varphi\|_1 = 1$. The above is just the kernel h_1 .

For $t > 0$, we have

$$\varphi_t(x) = t^{-n} \exp\left(-\frac{\pi}{t^2}\|x\|^2\right).$$

The above isn't exactly the heat kernel h_t .

Less abstract, more concrete

For the remainder of the discussion, one may forget the general setup and consider the following example for concreteness' sake. Let φ be defined as

$$\varphi(x) = e^{-\pi\|x\|^2}.$$

This is radial, non-negative, radially decreasing and we have $\|\varphi\|_1 = 1$. The above is just the kernel h_1 .

For $t > 0$, we have

$$\varphi_t(x) = t^{-n} \exp\left(-\frac{\pi}{t^2}\|x\|^2\right).$$

The above isn't exactly the heat kernel h_t . Rather, it is h_{t^2} .

Less abstract, more concrete

For the remainder of the discussion, one may forget the general setup and consider the following example for concreteness' sake. Let φ be defined as

$$\varphi(x) = e^{-\pi\|x\|^2}.$$

This is radial, non-negative, radially decreasing and we have $\|\varphi\|_1 = 1$. The above is just the kernel h_1 .

For $t > 0$, we have

$$\varphi_t(x) = t^{-n} \exp\left(-\frac{\pi}{t^2}\|x\|^2\right).$$

The above isn't exactly the heat kernel h_t . Rather, it is h_{t^2} .

It is now clear that proving the Main Theorem will show that (\star) holds for $x \in \text{Leb}(f)$.

Some final notation

S^{n-1} will denote the $n - 1$ sphere in \mathbb{R}^n .

S^{n-1} will denote the $n - 1$ sphere in \mathbb{R}^n . More explicitly,

$$S^{n-1} = \{x \in \mathbb{R}^n : \|x\| = 1\}.$$

S^{n-1} will denote the $n - 1$ sphere in \mathbb{R}^n . More explicitly,

$$S^{n-1} = \{x \in \mathbb{R}^n : \|x\| = 1\}.$$

For $x \in \mathbb{R}^n$ and $r > 0$, $B(x, r) = \{y \in \mathbb{R}^n : \|y - x\| < r\}$.

S^{n-1} will denote the $n - 1$ sphere in \mathbb{R}^n . More explicitly,

$$S^{n-1} = \{x \in \mathbb{R}^n : \|x\| = 1\}.$$

For $x \in \mathbb{R}^n$ and $r > 0$, $B(x, r) = \{y \in \mathbb{R}^n : \|y - x\| < r\}$.

V_n is the volume of the unit ball $B(0, 1)$.

S^{n-1} will denote the $n - 1$ sphere in \mathbb{R}^n . More explicitly,

$$S^{n-1} = \{x \in \mathbb{R}^n : \|x\| = 1\}.$$

For $x \in \mathbb{R}^n$ and $r > 0$, $B(x, r) = \{y \in \mathbb{R}^n : \|y - x\| < r\}$.

V_n is the volume of the unit ball $B(0, 1)$. Thus, we have

$$\int_{B(x, r)} 1 = V_n r^n$$

for $x \in \mathbb{R}^n$ and $r > 0$.

Recap on Polar Coordinates

Recall that given $f \in L^1(\mathbb{R}^n)$, we may compute its integral as

Recap on Polar Coordinates

Recall that given $f \in L^1(\mathbb{R}^n)$, we may compute its integral as

$$\int_{\mathbb{R}^n} f(x) \, dx = \int_0^\infty \int_{S^{n-1}} r^{n-1} f(r\omega) \, d\omega \, dr.$$

Recall that given $f \in L^1(\mathbb{R}^n)$, we may compute its integral as

$$\int_{\mathbb{R}^n} f(x) \, dx = \int_0^\infty \int_{S^{n-1}} r^{n-1} f(r\omega) \, d\omega \, dr.$$

In particular, if f is a radial function

Recall that given $f \in L^1(\mathbb{R}^n)$, we may compute its integral as

$$\int_{\mathbb{R}^n} f(x) \, dx = \int_0^\infty \int_{S^{n-1}} r^{n-1} f(r\omega) \, d\omega \, dr.$$

In particular, if f is a radial function and g is such that $f(x) = g(\|x\|)$, then

Recall that given $f \in L^1(\mathbb{R}^n)$, we may compute its integral as

$$\int_{\mathbb{R}^n} f(x) \, dx = \int_0^\infty \int_{S^{n-1}} r^{n-1} f(r\omega) \, d\omega \, dr.$$

In particular, if f is a radial function and g is such that $f(x) = g(\|x\|)$, then

$$\int_{\mathbb{R}^n} f(x) \, dx = \omega(S^{n-1}) \int_0^\infty r^{n-1} g(r) \, dr.$$

Proof of the Main Theorem

- 1 Recap
- 2 Notations and Setup
- 3 Proof of the Main Theorem
- 4 The Stronger Theorem

Proof of the Main Theorem

Fix a point $x \in \text{Leb}(f)$ and let $\epsilon > 0$ be arbitrary.

Proof of the Main Theorem

Fix a point $x \in \text{Leb}(f)$ and let $\epsilon > 0$ be arbitrary.
By definition of Lebesgue set, there exists $\delta > 0$ such that

Proof of the Main Theorem

Fix a point $x \in \text{Leb}(f)$ and let $\epsilon > 0$ be arbitrary.
By definition of Lebesgue set, there exists $\delta > 0$ such that

$$\frac{1}{r^n} \int_{\|u\| < r} |f(x-u) - f(x)| \, du < \epsilon \quad (L)$$

Proof of the Main Theorem

Fix a point $x \in \text{Leb}(f)$ and let $\epsilon > 0$ be arbitrary.

By definition of Lebesgue set, there exists $\delta > 0$ such that

$$\frac{1}{r^n} \int_{\|u\| < r} |f(x-u) - f(x)| \, du < \epsilon \quad (L)$$

for all $0 < r \leq \delta$.

Proof of the Main Theorem

Fix a point $x \in \text{Leb}(f)$ and let $\epsilon > 0$ be arbitrary.

By definition of Lebesgue set, there exists $\delta > 0$ such that

$$\int_{\|u\| < r} |f(x-u) - f(x)| \, du < \epsilon r^n \quad (L)$$

for all $0 < r \leq \delta$.

Proof of the Main Theorem

Fix a point $x \in \text{Leb}(f)$ and let $\epsilon > 0$ be arbitrary.

By definition of Lebesgue set, there exists $\delta > 0$ such that

$$\int_{\|u\| < r} |f(x-u) - f(x)| \, du < \epsilon r^n \quad (L)$$

for all $0 < r \leq \delta$.

Note that for all $t > 0$, we have $\int_{\mathbb{R}^n} \varphi_t = \int_{\mathbb{R}^n} \varphi = \int_{\mathbb{R}^n} |\varphi| = 1$.

Still proving the Main Theorem

Still proving the Main Theorem

Thus, for all $t > 0$, we have

Still proving the Main Theorem

Thus, for all $t > 0$, we have

$$|(f * \varphi_t)(x) - f(x)| = \left| \int_{\mathbb{R}^n} f(x - u) \varphi_t(u) \, du - f(x) \right|$$

Still proving the Main Theorem

Thus, for all $t > 0$, we have

$$\begin{aligned}|(f * \varphi_t)(x) - f(x)| &= \left| \int_{\mathbb{R}^n} f(x-u)\varphi_t(u) \, du - f(x) \right| \\ &= \left| \int_{\mathbb{R}^n} f(x-u)\varphi_t(u) \, du - f(x) \cdot 1 \right|\end{aligned}$$

Still proving the Main Theorem

Thus, for all $t > 0$, we have

$$\begin{aligned}|(f * \varphi_t)(x) - f(x)| &= \left| \int_{\mathbb{R}^n} f(x - u) \varphi_t(u) \, du - f(x) \right| \\ &= \left| \int_{\mathbb{R}^n} f(x - u) \varphi_t(u) \, du - f(x) \int_{\mathbb{R}^n} \varphi_t(u) \, du \right|\end{aligned}$$

Thus, for all $t > 0$, we have

$$\begin{aligned}|(f * \varphi_t)(x) - f(x)| &= \left| \int_{\mathbb{R}^n} f(x - u) \varphi_t(u) \, du - f(x) \right| \\&= \left| \int_{\mathbb{R}^n} f(x - u) \varphi_t(u) \, du - f(x) \int_{\mathbb{R}^n} \varphi_t(u) \, du \right| \\&= \left| \int_{\mathbb{R}^n} [f(x - u) - f(x)] \varphi_t(u) \, du \right|\end{aligned}$$

Still proving the Main Theorem

Thus, for all $t > 0$, we have

$$\begin{aligned} |(f * \varphi_t)(x) - f(x)| &= \left| \int_{\mathbb{R}^n} f(x-u) \varphi_t(u) \, du - f(x) \right| \\ &= \left| \int_{\mathbb{R}^n} f(x-u) \varphi_t(u) \, du - f(x) \int_{\mathbb{R}^n} \varphi_t(u) \, du \right| \\ &= \left| \int_{\mathbb{R}^n} [f(x-u) - f(x)] \varphi_t(u) \, du \right| \\ &\leq \left| \int_{\|u\| < \delta} [f(x-u) - f(x)] \varphi_t(u) \, du \right| \\ &\quad + \left| \int_{\|u\| \geq \delta} [f(x-u) - f(x)] \varphi_t(u) \, du \right| \end{aligned}$$

Still proving the Main Theorem

Thus, for all $t > 0$, we have

$$\begin{aligned} |(f * \varphi_t)(x) - f(x)| &= \left| \int_{\mathbb{R}^n} f(x-u) \varphi_t(u) \, du - f(x) \right| \\ &= \left| \int_{\mathbb{R}^n} f(x-u) \varphi_t(u) \, du - f(x) \int_{\mathbb{R}^n} \varphi_t(u) \, du \right| \\ &= \left| \int_{\mathbb{R}^n} [f(x-u) - f(x)] \varphi_t(u) \, du \right| \\ &\leq \left| \int_{\|u\| < \delta} [f(x-u) - f(x)] \varphi_t(u) \, du \right| \rightsquigarrow I_1 \\ &\quad + \left| \int_{\|u\| \geq \delta} [f(x-u) - f(x)] \varphi_t(u) \, du \right| \rightsquigarrow I_2 \end{aligned}$$

Still proving the Main Theorem

Thus, for all $t > 0$, we have

$$\begin{aligned} |(f * \varphi_t)(x) - f(x)| &= \left| \int_{\mathbb{R}^n} f(x-u) \varphi_t(u) \, du - f(x) \right| \\ &= \left| \int_{\mathbb{R}^n} f(x-u) \varphi_t(u) \, du - f(x) \int_{\mathbb{R}^n} \varphi_t(u) \, du \right| \\ &= \left| \int_{\mathbb{R}^n} [f(x-u) - f(x)] \varphi_t(u) \, du \right| \\ &\leq \left| \int_{\|u\| < \delta} [f(x-u) - f(x)] \varphi_t(u) \, du \right| \rightsquigarrow l_1(t) \\ &\quad + \left| \int_{\|u\| \geq \delta} [f(x-u) - f(x)] \varphi_t(u) \, du \right| \rightsquigarrow l_2(t) \end{aligned}$$

First, we note that

First, we note that

$$\int_{r/2 \leq \|u\| \leq r} \varphi(u) \, du$$

First, we note that

$$\int_{r/2 \leq \|u\| \leq r} \psi_0(r) \, du \leq \int_{r/2 \leq \|u\| \leq r} \varphi(u) \, du$$

First, we note that

$$\left(\int_{r/2 \leq \|u\| \leq r} 1 \right) \psi_0(r) \leq \int_{r/2 \leq \|u\| \leq r} \varphi(u) \, du$$

First, we note that

$$V_n \left(1 - \frac{1}{2^n}\right) r^n \psi_0(r) \leq \int_{r/2 \leq \|u\| \leq r} \varphi(u) \, du$$

First, we note that

$$V_n \left(1 - \frac{1}{2^n}\right) r^n \psi_0(r) \leq \int_{r/2 \leq \|u\| \leq r} \varphi(u) \, du \rightarrow 0,$$

First, we note that

$$V_n \left(1 - \frac{1}{2^n}\right) r^n \psi_0(r) \leq \int_{r/2 \leq \|u\| \leq r} \varphi(u) \, du \rightarrow 0,$$

as $r \rightarrow 0$

First, we note that

$$V_n \left(1 - \frac{1}{2^n}\right) r^n \psi_0(r) \leq \int_{r/2 \leq \|u\| \leq r} \varphi(u) \, du \rightarrow 0,$$

as $r \rightarrow 0$ or as $r \rightarrow \infty$.

First, we note that

$$V_n \left(1 - \frac{1}{2^n}\right) r^n \psi_0(r) \leq \int_{r/2 \leq \|u\| \leq r} \varphi(u) \, du \rightarrow 0,$$

as $r \rightarrow 0$ or as $r \rightarrow \infty$.

Thus, $r^n \psi_0(r) \rightarrow 0$

First, we note that

$$V_n \left(1 - \frac{1}{2^n} \right) r^n \psi_0(r) \leq \int_{r/2 \leq \|u\| \leq r} \varphi(u) \, du \rightarrow 0,$$

as $r \rightarrow 0$ or as $r \rightarrow \infty$.

Thus, $r^n \psi_0(r) \rightarrow 0$ as r tends to 0 or ∞ .

First, we note that

$$V_n \left(1 - \frac{1}{2^n}\right) r^n \psi_0(r) \leq \int_{r/2 \leq \|u\| \leq r} \varphi(u) \, du \rightarrow 0,$$

as $r \rightarrow 0$ or as $r \rightarrow \infty$.

Thus, $r^n \psi_0(r) \rightarrow 0$ as r tends to 0 or ∞ .

Hence, there exists $A > 0$ such that

First, we note that

$$V_n \left(1 - \frac{1}{2^n}\right) r^n \psi_0(r) \leq \int_{r/2 \leq \|u\| \leq r} \varphi(u) \, du \rightarrow 0,$$

as $r \rightarrow 0$ or as $r \rightarrow \infty$.

Thus, $r^n \psi_0(r) \rightarrow 0$ as r tends to 0 or ∞ .

Hence, there exists $A > 0$ such that $r^n \psi_0(r) \leq A$

First, we note that

$$V_n \left(1 - \frac{1}{2^n}\right) r^n \psi_0(r) \leq \int_{r/2 \leq \|u\| \leq r} \varphi(u) \, du \rightarrow 0,$$

as $r \rightarrow 0$ or as $r \rightarrow \infty$.

Thus, $r^n \psi_0(r) \rightarrow 0$ as r tends to 0 or ∞ .

Hence, there exists $A > 0$ such that $r^n \psi_0(r) \leq A$ for $r \in (0, \infty)$.

First, we note that

$$V_n \left(1 - \frac{1}{2^n}\right) r^n \psi_0(r) \leq \int_{r/2 \leq \|u\| \leq r} \varphi(u) \, du \rightarrow 0,$$

as $r \rightarrow 0$ or as $r \rightarrow \infty$.

Thus, $r^n \psi_0(r) \rightarrow 0$ as r tends to 0 or ∞ .

Hence, there exists $A > 0$ such that $r^n \psi_0(r) \leq A$ for $r \in (0, \infty)$.

Using this, we first show that $I_2(t) \xrightarrow{t \rightarrow 0} 0$.

$$l_2(t)$$

$$I_2(t) = \left| \int_{\|u\| \geq \delta} [f(x-u) - f(x)] \varphi_t(u) \, du \right|$$

$$I_2(t) \leq \int_{\|u\| \geq \delta} |f(x-u)| \varphi_t(u) \, du + |f(x)| \int_{\|u\| \geq \delta} \varphi_t(u) \, du.$$

$$l_2(t) \leq \int_{\|u\| \geq \delta} |f(x-u)| \varphi_t(u) \, du + |f(x)| \int_{\|u\| \geq \delta} \varphi_t(u) \, du.$$

The second term goes to 0 as $t \rightarrow 0$

$$I_2(t) \leq \int_{\|u\| \geq \delta} |f(x-u)| \varphi_t(u) \, du + |f(x)| \int_{\|u\| \geq \delta} \varphi_t(u) \, du.$$

The second term goes to 0 as $t \rightarrow 0$ since $\{\varphi_t\}_{t>0}$ is an approximate identity.

$$I_2(t) \leq \int_{\|u\| \geq \delta} |f(x-u)| \varphi_t(u) \, du + |f(x)| \int_{\|u\| \geq \delta} \varphi_t(u) \, du.$$

The second term goes to 0 as $t \rightarrow 0$ since $\{\varphi_t\}_{t>0}$ is an approximate identity.

Now, let χ_δ denote the characteristic function of $\{u \in \mathbb{R}^n : \|u\| \geq \delta\}$.

$$I_2(t) \leq \int_{\|u\| \geq \delta} |f(x-u)| \varphi_t(u) \, du + |f(x)| \int_{\|u\| \geq \delta} \varphi_t(u) \, du.$$

The second term goes to 0 as $t \rightarrow 0$ since $\{\varphi_t\}_{t>0}$ is an approximate identity.

Now, let χ_δ denote the characteristic function of $\{u \in \mathbb{R}^n : \|u\| \geq \delta\}$.

We see that the first term is at most $\|f\|_1 \|\chi_\delta \varphi_t\|_\infty$.

$$I_2(t) \leq \int_{\|u\| \geq \delta} |f(x-u)| \varphi_t(u) \, du + |f(x)| \int_{\|u\| \geq \delta} \varphi_t(u) \, du.$$

The second term goes to 0 as $t \rightarrow 0$ since $\{\varphi_t\}_{t>0}$ is an approximate identity.

Now, let χ_δ denote the characteristic function of $\{u \in \mathbb{R}^n : \|u\| \geq \delta\}$.

We see that the first term is at most $\|f\|_1 \|\chi_\delta \varphi_t\|_\infty$. Since φ is radially decreasing, we see that

$$I_2(t) \leq \int_{\|u\| \geq \delta} |f(x-u)| \varphi_t(u) \, du + |f(x)| \int_{\|u\| \geq \delta} \varphi_t(u) \, du.$$

The second term goes to 0 as $t \rightarrow 0$ since $\{\varphi_t\}_{t>0}$ is an approximate identity.

Now, let χ_δ denote the characteristic function of $\{u \in \mathbb{R}^n : \|u\| \geq \delta\}$.

We see that the first term is at most $\|f\|_1 \|\chi_\delta \varphi_t\|_\infty$. Since φ is radially decreasing, we see that

$$\|\chi_\delta \varphi_t\|_\infty = \sup_{\|u\| \geq \delta} \varphi_t(u)$$

$$I_2(t) \leq \int_{\|u\| \geq \delta} |f(x-u)| \varphi_t(u) \, du + |f(x)| \int_{\|u\| \geq \delta} \varphi_t(u) \, du.$$

The second term goes to 0 as $t \rightarrow 0$ since $\{\varphi_t\}_{t>0}$ is an approximate identity.

Now, let χ_δ denote the characteristic function of $\{u \in \mathbb{R}^n : \|u\| \geq \delta\}$.

We see that the first term is at most $\|f\|_1 \|\chi_\delta \varphi_t\|_\infty$. Since φ is radially decreasing, we see that

$$\|\chi_\delta \varphi_t\|_\infty = \sup_{\|u\| \geq \delta} t^{-n} \varphi(u/t)$$

$$I_2(t) \leq \int_{\|u\| \geq \delta} |f(x-u)| \varphi_t(u) \, du + |f(x)| \int_{\|u\| \geq \delta} \varphi_t(u) \, du.$$

The second term goes to 0 as $t \rightarrow 0$ since $\{\varphi_t\}_{t>0}$ is an approximate identity.

Now, let χ_δ denote the characteristic function of $\{u \in \mathbb{R}^n : \|u\| \geq \delta\}$.

We see that the first term is at most $\|f\|_1 \|\chi_\delta \varphi_t\|_\infty$. Since φ is radially decreasing, we see that

$$\|\chi_\delta \varphi_t\|_\infty = \sup_{\|u\| \geq \delta} t^{-n} \varphi(u/t) = t^{-n} \psi_0(\delta/t)$$

$$I_2(t) \leq \int_{\|u\| \geq \delta} |f(x-u)| \varphi_t(u) \, du + |f(x)| \int_{\|u\| \geq \delta} \varphi_t(u) \, du.$$

The second term goes to 0 as $t \rightarrow 0$ since $\{\varphi_t\}_{t>0}$ is an approximate identity.

Now, let χ_δ denote the characteristic function of $\{u \in \mathbb{R}^n : \|u\| \geq \delta\}$.

We see that the first term is at most $\|f\|_1 \|\chi_\delta \varphi_t\|_\infty$. Since φ is radially decreasing, we see that

$$\|\chi_\delta \varphi_t\|_\infty = \sup_{\|u\| \geq \delta} t^{-n} \varphi(u/t) = \delta^{-n} (\delta/t)^n \psi_0(\delta/t)$$

$$I_2(t) \leq \int_{\|u\| \geq \delta} |f(x-u)| \varphi_t(u) \, du + |f(x)| \int_{\|u\| \geq \delta} \varphi_t(u) \, du.$$

The second term goes to 0 as $t \rightarrow 0$ since $\{\varphi_t\}_{t>0}$ is an approximate identity.

Now, let χ_δ denote the characteristic function of $\{u \in \mathbb{R}^n : \|u\| \geq \delta\}$.

We see that the first term is at most $\|f\|_1 \|\chi_\delta \varphi_t\|_\infty$. Since φ is radially decreasing, we see that

$$\|\chi_\delta \varphi_t\|_\infty = \sup_{\|u\| \geq \delta} t^{-n} \varphi(u/t) = \delta^{-n} (\delta/t)^n \psi_0(\delta/t) \rightarrow 0,$$

$$l_2(t) \leq \int_{\|u\| \geq \delta} |f(x-u)| \varphi_t(u) \, du + |f(x)| \int_{\|u\| \geq \delta} \varphi_t(u) \, du.$$

The second term goes to 0 as $t \rightarrow 0$ since $\{\varphi_t\}_{t>0}$ is an approximate identity.

Now, let χ_δ denote the characteristic function of $\{u \in \mathbb{R}^n : \|u\| \geq \delta\}$.

We see that the first term is at most $\|f\|_1 \|\chi_\delta \varphi_t\|_\infty$. Since φ is radially decreasing, we see that

$$\|\chi_\delta \varphi_t\|_\infty = \sup_{\|u\| \geq \delta} t^{-n} \varphi(u/t) = \delta^{-n} (\delta/t)^n \psi_0(\delta/t) \rightarrow 0,$$

as $t \rightarrow 0$.

$$I_2(t) \leq \int_{\|u\| \geq \delta} |f(x-u)| \varphi_t(u) \, du + |f(x)| \int_{\|u\| \geq \delta} \varphi_t(u) \, du.$$

The second term goes to 0 as $t \rightarrow 0$ since $\{\varphi_t\}_{t>0}$ is an approximate identity.

Now, let χ_δ denote the characteristic function of $\{u \in \mathbb{R}^n : \|u\| \geq \delta\}$.

We see that the first term is at most $\|f\|_1 \|\chi_\delta \varphi_t\|_\infty$. Since φ is radially decreasing, we see that

$$\|\chi_\delta \varphi_t\|_\infty = \sup_{\|u\| \geq \delta} t^{-n} \varphi(u/t) = \delta^{-n} (\delta/t)^n \psi_0(\delta/t) \rightarrow 0,$$

as $t \rightarrow 0$. Thus, $I_2(t) \xrightarrow{t \rightarrow 0} 0$.

l_2 down, l_1 to go

Let us now define

$$g(r) = \int_{S^{n-1}} |f(x - r\omega) - f(x)| \, d\omega$$

Let us now define

$$g(r) = \int_{S^{n-1}} |f(x - r\omega) - f(x)| \, d\omega$$

and

$$G(r) = \int_0^r s^{n-1} g(s) \, ds.$$

Let us now define

$$g(r) = \int_{S^{n-1}} |f(x - r\omega) - f(x)| \, d\omega$$

and

$$G(r) = \int_0^r s^{n-1} g(s) \, ds.$$

Thus, the Lebesgue set condition (L) from earlier translates to

Let us now define

$$g(r) = \int_{S^{n-1}} |f(x - r\omega) - f(x)| \, d\omega$$

and

$$G(r) = \int_0^r s^{n-1} g(s) \, ds.$$

Thus, the Lebesgue set condition (L) from earlier translates to

$$G(r) \leq \epsilon r^n$$

Let us now define

$$g(r) = \int_{S^{n-1}} |f(x - r\omega) - f(x)| \, d\omega$$

and

$$G(r) = \int_0^r s^{n-1} g(s) \, ds.$$

Thus, the Lebesgue set condition (L) from earlier translates to

$$G(r) \leq \epsilon r^n \quad \text{for } r \leq \delta.$$

Note that $G(0) = 0$.

Let us now define

$$g(r) = \int_{S^{n-1}} |f(x - r\omega) - f(x)| \, d\omega$$

and

$$G(r) = \int_0^r s^{n-1} g(s) \, ds.$$

Thus, the Lebesgue set condition (L) from earlier translates to

$$G(r) \leq \epsilon r^n \quad \text{for } r \leq \delta.$$

Note that $G(0) = 0$.

With these notations, we do some more calculations.

Some more calculations

We have

Some more calculations

We have

$$I_1(t) = \left| \int_{\|u\| < \delta} [f(x-u) - f(x)] \varphi_t(u) \, du \right|$$

Some more calculations

We have

$$I_1(t) \leq \int_{\|u\| < \delta} |f(x-u) - f(x)| \varphi_t(u) \, du$$

Some more calculations

We have

$$I_1(t) \leq \int_{\|u\| < \delta} |f(x-u) - f(x)| t^{-n} \varphi(u/t) \, du$$

Some more calculations

We have

$$\begin{aligned} I_1(t) &\leq \int_{\|u\| < \delta} |f(x-u) - f(x)| t^{-n} \varphi(u/t) \, du \\ &= \int_0^\delta r^{n-1} g(r) t^{-n} \psi_0(r/t) \, dr \end{aligned}$$

Some more calculations

We have

$$\begin{aligned} I_1(t) &\leq \int_{\|u\| < \delta} |f(x-u) - f(x)| t^{-n} \varphi(u/t) \, du \\ &= \int_0^\delta r^{n-1} g(r) t^{-n} \psi_0(r/t) \, dr \end{aligned}$$

Integrate by parts

Some more calculations

We have

$$\begin{aligned} I_1(t) &\leq \int_{\|u\| < \delta} |f(x-u) - f(x)| t^{-n} \varphi(u/t) \, du \\ &= \int_0^\delta r^{n-1} g(r) t^{-n} \psi_0(r/t) \, dr \\ &= G(r) t^{-n} \psi_0(r/t) \Big|_0^\delta - \int_0^\delta G(r) \, d(t^{-n} \psi_0(r/t)) \end{aligned}$$

Some more calculations

We have

$$\begin{aligned} I_1(t) &\leq \int_{\|u\| < \delta} |f(x-u) - f(x)| t^{-n} \varphi(u/t) \, du \\ &= \int_0^\delta r^{n-1} g(r) t^{-n} \psi_0(r/t) \, dr \\ &= G(\delta) t^{-n} \psi_0(\delta/t) - \int_0^\delta G(r) \, d(t^{-n} \psi_0(r/t)) \end{aligned}$$

Some more calculations

We have

$$\begin{aligned} I_1(t) &\leq \int_{\|u\| < \delta} |f(x-u) - f(x)| t^{-n} \varphi(u/t) \, du \\ &= \int_0^\delta r^{n-1} g(r) t^{-n} \psi_0(r/t) \, dr \\ &= G(\delta) t^{-n} \psi_0(\delta/t) - \int_0^\delta G(r) \, d(t^{-n} \psi_0(r/t)) \\ &\leq \epsilon \delta^n t^{-n} \psi_0(\delta/t) - \int_0^{\delta/t} G(ts) t^{-n} \, d(\psi_0(s)) \end{aligned}$$

Some more calculations

We have

$$\begin{aligned} I_1(t) &\leq \int_{\|u\| < \delta} |f(x-u) - f(x)| t^{-n} \varphi(u/t) \, du \\ &= \int_0^\delta r^{n-1} g(r) t^{-n} \psi_0(r/t) \, dr \\ &= G(\delta) t^{-n} \psi_0(\delta/t) - \int_0^\delta G(r) \, d(t^{-n} \psi_0(r/t)) \\ &\leq \epsilon(\delta/t)^n \psi_0(\delta/t) - \int_0^{\delta/t} G(ts) t^{-n} \, d(\psi_0(s)) \end{aligned}$$

Some more calculations

We have

$$\begin{aligned} I_1(t) &\leq \int_{\|u\| < \delta} |f(x-u) - f(x)| t^{-n} \varphi(u/t) \, du \\ &= \int_0^\delta r^{n-1} g(r) t^{-n} \psi_0(r/t) \, dr \\ &= G(\delta) t^{-n} \psi_0(\delta/t) - \int_0^\delta G(r) \, d(t^{-n} \psi_0(r/t)) \\ &\leq \epsilon A - \int_0^{\delta/t} G(ts) t^{-n} \, d(\psi_0(s)) \end{aligned}$$

Some more calculations

We have

$$\begin{aligned} I_1(t) &\leq \int_{\|u\| < \delta} |f(x-u) - f(x)| t^{-n} \varphi(u/t) \, du \\ &= \int_0^\delta r^{n-1} g(r) t^{-n} \psi_0(r/t) \, dr \\ &= G(\delta) t^{-n} \psi_0(\delta/t) - \int_0^\delta G(r) \, d(t^{-n} \psi_0(r/t)) \\ &\leq \epsilon A - \int_0^{\delta/t} G(ts) t^{-n} \, d(\psi_0(s)) \\ &\leq \epsilon A - \int_0^{\delta/t} \epsilon (ts)^n t^{-n} \, d(\psi_0(s)) \end{aligned}$$

Note that $d\psi_0 \leq 0$.

Some more calculations

We have

$$\begin{aligned} I_1(t) &\leq \int_{\|u\| < \delta} |f(x-u) - f(x)| t^{-n} \varphi(u/t) \, du \\ &= \int_0^\delta r^{n-1} g(r) t^{-n} \psi_0(r/t) \, dr \\ &= G(\delta) t^{-n} \psi_0(\delta/t) - \int_0^\delta G(r) \, d(t^{-n} \psi_0(r/t)) \\ &\leq \epsilon A - \int_0^{\delta/t} G(ts) t^{-n} \, d(\psi_0(s)) \\ &\leq \epsilon A - \int_0^{\delta/t} \epsilon s^n \, d(\psi_0(s)) \end{aligned}$$

Note that $d\psi_0 \leq 0$.

Some more calculations

We have

$$\begin{aligned} I_1(t) &\leq \int_{\|u\| < \delta} |f(x-u) - f(x)| t^{-n} \varphi(u/t) \, du \\ &= \int_0^\delta r^{n-1} g(r) t^{-n} \psi_0(r/t) \, dr \\ &= G(\delta) t^{-n} \psi_0(\delta/t) - \int_0^\delta G(r) \, d(t^{-n} \psi_0(r/t)) \\ &\leq \epsilon A - \int_0^{\delta/t} G(ts) t^{-n} \, d(\psi_0(s)) \\ &\leq \epsilon A - \int_0^{\delta/t} \epsilon s^n \, d(\psi_0(s)) \leq \epsilon \left(A - \int_0^\infty s^n \, d(\psi_0(s)) \right) \end{aligned}$$

Note that $d\psi_0 \leq 0$.

Some more calculations

We have

$$\begin{aligned} I_1(t) &\leq \int_{\|u\| < \delta} |f(x-u) - f(x)| t^{-n} \varphi(u/t) \, du \\ &= \int_0^\delta r^{n-1} g(r) t^{-n} \psi_0(r/t) \, dr \\ &= G(\delta) t^{-n} \psi_0(\delta/t) - \int_0^\delta G(r) \, d(t^{-n} \psi_0(r/t)) \\ &\leq \epsilon A - \int_0^{\delta/t} G(ts) t^{-n} \, d(\psi_0(s)) \\ &\leq \epsilon A - \int_0^{\delta/t} \epsilon s^n \, d(\psi_0(s)) \leq \epsilon \left(A - \int_0^\infty s^n \, d(\psi_0(s)) \right) \end{aligned}$$

Note that $d\psi_0 \leq 0$.

The Green Integral

The green integral can be calculated exactly quite simply.

The green integral can be calculated exactly quite simply.

$$- \int_0^\infty s^n d(\psi_0(s))$$

The green integral can be calculated exactly quite simply.

$$-\int_0^\infty s^n d(\psi_0(s)) = -s^n \psi_0(s) \Big|_0^\infty + n \int_0^\infty s^{n-1} \psi_0(s) ds$$

The green integral can be calculated exactly quite simply.

$$\begin{aligned} - \int_0^\infty s^n d(\psi_0(s)) &= -s^n \psi_0(s) \Big|_0^\infty + n \int_0^\infty s^{n-1} \psi_0(s) ds \\ &= 0 + \frac{n}{\omega(S^{n-1})} \int_{\mathbb{R}^n} \varphi \end{aligned}$$

The green integral can be calculated exactly quite simply.

$$\begin{aligned} - \int_0^\infty s^n d(\psi_0(s)) &= -s^n \psi_0(s) \Big|_0^\infty + n \int_0^\infty s^{n-1} \psi_0(s) ds \\ &= \frac{n}{\omega(S^{n-1})}. \end{aligned}$$

The green integral can be calculated exactly quite simply.

$$\begin{aligned} - \int_0^\infty s^n d(\psi_0(s)) &= -s^n \psi_0(s) \Big|_0^\infty + n \int_0^\infty s^{n-1} \psi_0(s) ds \\ &= \frac{n}{\omega(S^{n-1})}. \end{aligned}$$

Putting this back, we get

The green integral can be calculated exactly quite simply.

$$\begin{aligned} - \int_0^\infty s^n d(\psi_0(s)) &= -s^n \psi_0(s) \Big|_0^\infty + n \int_0^\infty s^{n-1} \psi_0(s) ds \\ &= \frac{n}{\omega(S^{n-1})}. \end{aligned}$$

Putting this back, we get

$$I_1(t) \leq \epsilon \left(A + \frac{n}{\omega(S^{n-1})} \right)$$

The green integral can be calculated exactly quite simply.

$$\begin{aligned} - \int_0^\infty s^n d(\psi_0(s)) &= -s^n \psi_0(s) \Big|_0^\infty + n \int_0^\infty s^{n-1} \psi_0(s) ds \\ &= \frac{n}{\omega(S^{n-1})}. \end{aligned}$$

Putting this back, we get

$$I_1(t) \leq \epsilon \left(A + \frac{n}{\omega(S^{n-1})} \right) = \epsilon B.$$

The green integral can be calculated exactly quite simply.

$$\begin{aligned} - \int_0^\infty s^n d(\psi_0(s)) &= -s^n \psi_0(s) \Big|_0^\infty + n \int_0^\infty s^{n-1} \psi_0(s) ds \\ &= \frac{n}{\omega(S^{n-1})}. \end{aligned}$$

Putting this back, we get

$$I_1(t) \leq \epsilon \left(A + \frac{n}{\omega(S^{n-1})} \right) = \epsilon B.$$

Thus, we have bounded I_1

The green integral can be calculated exactly quite simply.

$$\begin{aligned} - \int_0^\infty s^n d(\psi_0(s)) &= -s^n \psi_0(s) \Big|_0^\infty + n \int_0^\infty s^{n-1} \psi_0(s) ds \\ &= \frac{n}{\omega(S^{n-1})}. \end{aligned}$$

Putting this back, we get

$$I_1(t) \leq \epsilon \left(A + \frac{n}{\omega(S^{n-1})} \right) = \epsilon B.$$

Thus, we have bounded I_1 *independent of t*

The green integral can be calculated exactly quite simply.

$$\begin{aligned} - \int_0^\infty s^n d(\psi_0(s)) &= -s^n \psi_0(s) \Big|_0^\infty + n \int_0^\infty s^{n-1} \psi_0(s) ds \\ &= \frac{n}{\omega(S^{n-1})}. \end{aligned}$$

Putting this back, we get

$$I_1(t) \leq \epsilon \left(A + \frac{n}{\omega(S^{n-1})} \right) = \epsilon B.$$

Thus, we have bounded I_1 *independent of t and of f* .

We had

We had

$$|(f * \varphi_t)(x) - f(x)| \leq I_1(t) + I_2(t).$$

We had

$$|(f * \varphi_t)(x) - f(x)| \leq I_1(t) + I_2(t).$$

We showed that I_1 is bounded independent of t .

We had

$$|(f * \varphi_t)(x) - f(x)| \leq I_1(t) + I_2(t) \leq \epsilon B + I_2(t).$$

We showed that I_1 is bounded independent of t .

We had

$$|(f * \varphi_t)(x) - f(x)| \leq I_1(t) + I_2(t) \leq \epsilon B + I_2(t).$$

We showed that I_1 is bounded independent of t . We also showed that $I_2(t) \xrightarrow{t \rightarrow 0} 0$.

We had

$$|(f * \varphi_t)(x) - f(x)| \leq I_1(t) + I_2(t) \leq \epsilon B + I_2(t).$$

We showed that I_1 is bounded independent of t . We also showed that $I_2(t) \xrightarrow{t \rightarrow 0} 0$. Thus, for t sufficiently small, we have

We had

$$|(f * \varphi_t)(x) - f(x)| \leq I_1(t) + I_2(t) \leq \epsilon B + I_2(t).$$

We showed that I_1 is bounded independent of t . We also showed that $I_2(t) \xrightarrow{t \rightarrow 0} 0$. Thus, for t sufficiently small, we have

$$|(f * \varphi_t)(x) - f(x)| < \epsilon(B + 1).$$

We had

$$|(f * \varphi_t)(x) - f(x)| \leq I_1(t) + I_2(t) \leq \epsilon B + I_2(t).$$

We showed that I_1 is bounded independent of t . We also showed that $I_2(t) \xrightarrow{t \rightarrow 0} 0$. Thus, for t sufficiently small, we have

$$|(f * \varphi_t)(x) - f(x)| < \epsilon(B + 1).$$

This completes the proof.

The Stronger Theorem

- 1 Recap
- 2 Notations and Setup
- 3 Proof of the Main Theorem
- 4 The Stronger Theorem

Concluding Remark

The theorem which I have proven is actually a weaker version of something more general.

Concluding Remark

The theorem which I have proven is actually a weaker version of something more general. Forget all the notation and hypothesis that we had so far.

Concluding Remark

The theorem which I have proven is actually a weaker version of something more general. Forget all the notation and hypothesis that we had so far.

Theorem (General Theorem)

Suppose $\varphi \in L^1(\mathbb{R}^n)$.

Concluding Remark

The theorem which I have proven is actually a weaker version of something more general. Forget all the notation and hypothesis that we had so far.

Theorem (General Theorem)

Suppose $\varphi \in L^1(\mathbb{R}^n)$. Let $\psi(y) = \operatorname{ess\,sup}_{\|z\| \geq \|y\|} |\varphi(z)|$

Concluding Remark

The theorem which I have proven is actually a weaker version of something more general. Forget all the notation and hypothesis that we had so far.

Theorem (General Theorem)

Suppose $\varphi \in L^1(\mathbb{R}^n)$. Let $\psi(y) = \operatorname{ess\,sup}_{\|z\| \geq \|y\|} |\varphi(z)|$ and for $t > 0$, let $\varphi_t(y) = t^{-n} \varphi(y/t)$.

Concluding Remark

The theorem which I have proven is actually a weaker version of something more general. Forget all the notation and hypothesis that we had so far.

Theorem (General Theorem)

Suppose $\varphi \in L^1(\mathbb{R}^n)$. Let $\psi(y) = \operatorname{ess\,sup}_{\|z\| \geq \|y\|} |\varphi(z)|$ and for $t > 0$, let $\varphi_t(y) = t^{-n}\varphi(y/t)$. If $\psi \in L^1(\mathbb{R}^n)$ and $f \in L^p(\mathbb{R}^n)$, $1 \leq p \leq \infty$,

Concluding Remark

The theorem which I have proven is actually a weaker version of something more general. Forget all the notation and hypothesis that we had so far.

Theorem (General Theorem)

Suppose $\varphi \in L^1(\mathbb{R}^n)$. Let $\psi(y) = \operatorname{ess\,sup}_{\|z\| \geq \|y\|} |\varphi(z)|$ and for $t > 0$, let $\varphi_t(y) = t^{-n} \varphi(y/t)$. If $\psi \in L^1(\mathbb{R}^n)$ and $f \in L^p(\mathbb{R}^n)$, $1 \leq p \leq \infty$, then $\lim_{t \rightarrow 0} (f * \varphi_t)(x) = f(x) \int_{\mathbb{R}^n} \varphi(t) \, dt$

Concluding Remark

The theorem which I have proven is actually a weaker version of something more general. Forget all the notation and hypothesis that we had so far.

Theorem (General Theorem)

Suppose $\varphi \in L^1(\mathbb{R}^n)$. Let $\psi(y) = \operatorname{ess\,sup}_{\|z\| \geq \|y\|} |\varphi(z)|$ and for $t > 0$, let $\varphi_t(y) = t^{-n} \varphi(y/t)$. If $\psi \in L^1(\mathbb{R}^n)$ and $f \in L^p(\mathbb{R}^n)$, $1 \leq p \leq \infty$, then $\lim_{t \rightarrow 0} (f * \varphi_t)(x) = f(x) \int_{\mathbb{R}^n} \varphi(t) \, dt$ whenever $x \in \operatorname{Leb}(f)$.

Concluding Remark

The theorem which I have proven is actually a weaker version of something more general. Forget all the notation and hypothesis that we had so far.

Theorem (General Theorem)

Suppose $\varphi \in L^1(\mathbb{R}^n)$. Let $\psi(y) = \operatorname{ess\,sup}_{\|z\| \geq \|y\|} |\varphi(z)|$ and for $t > 0$, let $\varphi_t(y) = t^{-n} \varphi(y/t)$. If $\psi \in L^1(\mathbb{R}^n)$ and $f \in L^p(\mathbb{R}^n)$, $1 \leq p \leq \infty$, then $\lim_{t \rightarrow 0} (f * \varphi_t)(x) = f(x) \int_{\mathbb{R}^n} \varphi(t) \, dt$ whenever $x \in \operatorname{Leb}(f)$.

Reference: *Introduction to Fourier Analysis on Euclidean Spaces* by Stein and Weiss