

## Section 7: Vector Fields

---

(1) Our focus this week is two-fold: on **arc lengths** and **vector-valued functions**. First, we spend some time talking about a quick and dirty **derivation of the arc-length formula** (this is just meant as a heuristic!).

(2) Now, let's talk about **vector-valued functions** i.e. functions  $\mathbf{F} : \mathbb{R}^n \rightarrow \mathbb{R}^n, n \geq 2$ , that take on vector values. So much can be said about these functions, but on the auspicious occasion of  $\pi$ -day, we focus on one idea:

### An inversion problem

When can we conclude that  $\mathbf{F}$  is the gradient of some real-valued function  $f : \mathbb{R}^n \rightarrow \mathbb{R}$ ?

Why care about this? Well, consider the differentiable functions  $f : \mathbb{R}^3 \rightarrow \mathbb{R}$ . These have gradients  $\nabla f(x)$  defined at each point. Therefore, we can define the vector-valued function  $\nabla f : \mathbb{R}^3 \rightarrow \mathbb{R}^3$ , taking  $x \mapsto \nabla f(x)$ . This means that, for every differentiable function, there is a corresponding vector-valued function.

A natural question to ask, then, is the 'inversion' - if I gave you a vector-valued function  $\mathbf{F} : \mathbb{R}^3 \rightarrow \mathbb{R}^3$ , can you give me a function whose gradient is  $\mathbf{F}$ ?

**The answer is no!**

**(Def. 1)** For  $n$ -dimensions, let  $\nabla = [\frac{\partial f}{\partial x_1}, \frac{\partial f}{\partial x_2} \dots \frac{\partial f}{\partial x_n}]$ . We call this the 'del' operator. For example, in  $\mathbb{R}^3, \nabla = [\frac{\partial f}{\partial x}, \frac{\partial f}{\partial y}, \frac{\partial f}{\partial z}]$ . We'll use it in this form herein.

**(Def. 2)** Thinking of  $\nabla$  as a vector, we can dot it and cross it with other vectors in  $\mathbb{R}^3$ . Let's do this with a vector-valued function,  $\mathbf{F}$ , to derive the **divergence** and **curl** of a vector-valued function.

$$\operatorname{div} \mathbf{F}_{(x,y,z)} = \nabla \cdot \mathbf{F}_{(x,y,z)}$$

$$\operatorname{curl} \mathbf{F}_{(x,y,z)} = \nabla \times \mathbf{F}_{(x,y,z)}$$

Given these ideas<sup>1</sup>, we have the following theorem.

**(Th. 1 (Gradients are curl-free))** If  $\nabla \times \mathbf{F} \neq 0$ , then  $\mathbf{F}$  is not the gradient of any function.

An easy proof involving brute computation can be found in Dr. Wang's lecture notes on divergence and curl. .

---

<sup>1</sup>We won't consider deeply into what these notions signify, because we'll do so later in the course. For the curious, however, I suggest 3Blue1Brown's video.

There is a more intuitive condition to check. Suppose that  $\mathbf{F}$  is the gradient of some function  $f$ , i.e.  $\nabla f = [\frac{\partial f}{\partial x}, \frac{\partial f}{\partial y}, \frac{\partial f}{\partial z}] = [F_1, F_2, F_3]$ . Then, since mixed partials are equal, we'd have that  $\frac{\partial f}{\partial x \partial y} = \frac{\partial F_1}{\partial y} = \frac{\partial f}{\partial y \partial x} = \frac{\partial F_2}{\partial x}$ .

Therefore, if mixed partials are unequal, i.e.  $\frac{\partial F_1}{\partial y} \neq \frac{\partial F_2}{\partial x}$ , then  $F$  can't be a gradient. Thus, we get the following...

**(Th. 2 (Identity of mixed partials))** If  $\frac{\partial F_1}{\partial y} \neq \frac{\partial F_2}{\partial x}$ , then  $\mathbf{F}$  is not the gradient of any function.

(3) Unfortunately, most vector fields are not gradients. However, there is a weaker sense in which this is possible - specifically, it may be that vector fields are gradients of a function along some path...

**(Def. 3)** Given  $\mathbf{F} : \mathbb{R}^n \rightarrow \mathbb{R}^n$ , if there is a path  $\mathbf{c} : \mathbb{R} \rightarrow \mathbb{R}^n$  such that  $\mathbf{c}'(t) = \mathbf{F}(\mathbf{c}(t))$ , then  $\mathbf{c}$  is called a **flow line** for  $\mathbf{F}$ .

**Q1.** The **arc-length** function,  $s(t)$ , for a path  $c(t)$  is given by  $s(x) = \int_{t_0}^x \|c'(t)\| dt$ . It represents the distance travelled until time  $x$ . Suppose you are walking on a helix staircase, so that your path is  $c(t) = (\cos(t), \sin(t), t)$ . Compute  $s(x)$  as you walk along this path.

**Q2.** Consider  $f(x, y) = x^2 + y^2$ . Sketch  $f$ , the gradient vector field of  $f$ , and some additional flow lines. What do you observe?

**Q3.** Is the curl of  $\mathbf{F}$  perpendicular to  $\mathbf{F}$ ? Consider  $\mathbf{F} = (-y, x, 1)$ .