

Logic of the Code

Aryana Haghjoo

March 14, 2023

1 Derivation of the Covariance Matrix

To find the analytic form of the covariance matrix, we need to find the second derivative of the chi-square with respect to the parameters of the model. Thus, we start with the matrix definition of the chi-square:

$$\chi^2 = (d - A(m))^T N^{-1} (d - A(m)) \quad (1)$$

Where d is an array containing the data, A is the theoretical model, which is dependent on the values of parameters (the dependency can be non-linear), and N is the noise matrix. If the noise is uncorrelated, the diagonal values of the noise matrix are just the square of the error bar on each data point.

$$\frac{d\chi^2}{dm} = -\left(\frac{dA(m)}{dm}\right)^T N^{-1} (d - A(m)) - (d - A(m))^T N^{-1} \frac{dA(m)}{dm} \quad (2)$$

Since we know that:

$$(N^{-1})^T = N^{-1} \quad (3)$$

$$\left[\frac{dA(m)}{dm} N^{-1} (d - A(m))\right]^T = (d - A(m))^T N^{-1} \frac{dA(m)}{dm} \quad (4)$$

Substituting in 2, we get:

$$\frac{d\chi^2}{dm} = -2\left(\frac{dA(m)}{dm}\right)^T N^{-1} (d - A(m)) \quad (5)$$

$$(6)$$

Thus, we can calculate the second derivative:

$$\frac{d^2\chi^2}{dm^2} = -2\left(\frac{d^2A(m)}{dm^2}\right)^T N^{-1} (d - A(m)) - 2\left(\frac{dA(m)}{dm}\right)^T N^{-1} \left(-\frac{d\chi^2}{dm}\right) \quad (7)$$

$$(8)$$

We can neglect the first term because of following two reasons:

1. First: The $(d - A(m))$ component, which is the residual, can take both negative and positive values; Thus on average, it will be close to zero.
2. Second: The fitted model should be very close to the data. Therefore, we expect that $(d - A(m)) \approx 0$

Finally, we are left with:

$$\frac{d^2\chi^2}{dm^2} = 2\left(\frac{dA(m)}{dm}\right)^T N^{-1} \left(\frac{dA(m)}{dm}\right) \quad (9)$$

Which is the definition of the **Covariance Matrix**

2 The Levenberg-Marquardt Algorithm

The Levenberg-Marquardt (LM), also known as the damped least-squares (DLS) method, is a fitting algorithm used for non-linear least-squares problems. This algorithm is iterative, and on each step, the parameters m will be replaced by $m + \delta m$. We aim to find δm by minimizing the chi-square calculated based on the perturbed parameters.

We knew that:

$$\chi^2(m) = (d - A(m))^T N^{-1} (d - A(m)) \quad (10)$$

$$\chi^2(m + \delta m) = \chi^2(m) + \frac{d\chi^2}{dm} \delta m \quad (11)$$

$$\frac{d\chi^2(m + \delta m)}{dm} = \frac{d}{dm}(\chi^2) + \frac{d}{dm}\left(\frac{d\chi^2}{dm} \delta m\right) \quad (12)$$

Since we calculated the first order derivative of chi-square in 2, we have:

$$\frac{d\chi^2(m + \delta m)}{dm} = -2\left(\frac{dA(m)}{dm}\right)^T N^{-1} (d - A(m)) + \frac{d^2\chi^2}{dm^2} \delta m + \frac{d\chi^2}{dm} \frac{d}{dm}(\delta m) \quad (13)$$

Where the last term is zero since δm does not have any dependencies on m , and we have already calculated the second derivative of chi-square in 9. Thus, we are left with:

$$\frac{d\chi^2(m + \delta m)}{dm} = -2\left(\frac{dA(m)}{dm}\right)^T N^{-1} (d - A(m)) + 2\left(\frac{dA(m)}{dm}\right)^T N^{-1} \left(\frac{dA(m)}{dm}\right) \delta m \quad (14)$$

We define $d - A(m) \equiv r$, and $\frac{dA(m)}{dm} \equiv A'$, we get:

$$A'^T N^{-1} A' \delta m = A'^T N^{-1} r \quad (15)$$

$$\delta m = (A'^T N^{-1} A')^{-1} A'^T N^{-1} r \quad (16)$$

Which is the basis for **Newton's method**. However, this method does not always converge, especially on complicated likelihood surfaces. To solve this issue, we add a new term to the

left-hand side of the above equation, which involves a control parameter named Λ that is updated on each iteration.

$$(A'^T N^{-1} A' + \Lambda I) \delta m = A'^T N^{-1} r \quad (17)$$

$$\delta m = (A'^T N^{-1} A' + \Lambda I)^{-1} A'^T N^{-1} r \quad (18)$$

On each iteration, we compare the chi-square to its value in the last step. If we encounter a higher value, the Λ will be multiplied to a constant arbitrary number (> 1). Otherwise, it will be divided with another constant value (> 1). For practical purposes, if Λ takes a value lower than a constant small arbitrary number, we set it equal to zero. If the Λ is zero, and the chi-square is less than an arbitrary threshold value, we declare the convergence of the algorithm.

3 Combining the Levenberg-Marquardt and Markov Chain Monte Carlo

There are various ways to force a MCMC to converge faster. One of which, involves using a posterior distribution that accounts for correlations between different parameters. To generate this posterior distribution, we first use a Levenberg-Marquardt fitter to obtain a set of best-fit parameters and the covariance matrix of parameters based on the local derivatives of the likelihood at the best-fit point. We then draw samples from this covariance matrix to generate the posterior distribution.

The logic of the above-mentioned algorithm can be summarized in the following chart.

