

A working doctor workflow with separate IT ownership

The local practice deployment is operational: a separate IT server manages access and native configuration; the doctor client opens matching records and full supplied images, starts a real Sol/Astra case handoff, and leaves report saving to the physician-role interface.

Outcome	Observed evidence
Separate IT guest	kauvery-it: 2 vCPU, 2 GiB RAM, 10 GiB virtual disk; persistent and autostart enabled
Hospital archive	22 studies / 1358 instances after five new practice cases
Native models	gpt-6.1-sol: coordination, preparation and drafting. gpt-6-astra: image interpretation and independent checking.
Doctor boundary	20 live HTTPS access checks passed; configuration and permanent approvals remain IT controlled.
Persistence	Native physician-role note save/reload passed. IT guest reboot, telemetry persistence and encrypted backup restore passed.

Decision: suitable for supervised local workflow practice

Keep clinician review mandatory. This evidence does not qualify the system for live patient deployment, autonomous diagnosis, electronic signing/release, individual SSO or patient-isolated history.

Doctor: <https://doctor.radiology.demo/worklist/>
IT: <https://it.radiology.demo/operations/>
Private logins: .private/HOSPITAL-DEMO-ACCESS.txt

Five modalities, explicit limits

Case / accession	Native supplied study	Assistant display packet
PRACTICE001 / ORD-22 DX	1 AP chest image	1 rendered frontal view
PRACTICE002 / ORD-23 CT	241 source instances	16 axial positions, lung and mediastinal windows
PRACTICE003 / ORD-24 MRI	95 source instances, 3 series	16/19 T2, 16/19 ADC, 16/57 DWI instances
PRACTICE004 / ORD-25 US	1 object, 227 spatial frames	16 spatial frames; not temporal cine
PRACTICE005 / ORD-26 MG	4 supplied source views	4 resized views; fine-detail limits disclosed

Actual native handoff rehearsal

The attached-image chest case completed through Sol Clinical Assistant, Astra Image Assistant and Astra Report Check. Identifiers, context and coverage were supplied from read-only native integrations; both Astra roles inspected the assigned pixels. The assistant returned a provisional report and did not write the hospital record.

Interaction measurement	GUI navigation rehearsal	Attached-image rehearsal
Observer one-shot reviews	67; early manual reviews excluded	2 teammate handoffs; no navigation reviews
Observed elapsed interval	44.4 minutes	5.1 minutes

Historical single-case engineering comparison, with different access paths and initial GUI trust/navigation friction. These are observation intervals and approval counts, not a controlled physician-efficiency study or a diagnostic accuracy comparison. Neither workflow was professionally clinically adjudicated.

Current permission policy: all six roles use native automatic tool review. A fresh Sol / Astra image-read / Astra report-check rehearsal completed in 302.0 seconds with 0 human tool approvals. Provider connection retries recovered during the run. Clinical record writes remain the physician's actions; the provider sandbox and IT configuration restrictions remain active. This is one integration observation, not a clinical validation.

All five full supplied studies rendered visible pixels in OHIF. All 11 packet assets matched SHA256 manifests. Missing packets were refused before native submission. Sampling, source-date shifts, missing history and display limitations remain visible to the doctor.

Verified operation; bounded claims

What passed

Authenticated doctor and IT origins; doctor configuration/SOUL/execution/permanent-permission denial; native IT management; actual gateway session renewal; tunnel-loss alert and recovery; IT guest reboot with preserved native state and telemetry; encrypted IT configuration backup and exact-bundle/SQLite restore; physician-role provisional note save and matching displayed content after reload.

The hospital stayed available during the IT transport failure. TLS checks used the trusted hospital CA without certificate bypass. The IT backup restored 300 telemetry samples with SQLite integrity verified. The on-machine backup has no external replica.

Try the deployment

1. Open Radiology Assistant or the doctor worklist and use the private login handout.
2. Choose PRACTICE002 / ORD-23 for a fresh CT case; verify the chart and full supplied images.
3. Inspect packet coverage and choose Prepare this case. Routine permissions are reviewed automatically.
4. Review and edit the proposal. Expand the Bahmni order, enter Radiology Notes, Save and reload.

PRACTICE001 / ORD-22 contains the clearly labelled practice note used for save/reload verification. Four other practice cases remain unsaved. No professional clinical sign-off is implied.

Clinical evidence remains separate

The existing Sol/Astra evaluation PDF reports the 106-study primary comparison, physician-report assertion measures, published-case diagnosis recognition and supplementary challenges. Workflow availability does not improve or replace those diagnostic findings. Consult Radiology-Frontier-Model-Evaluation.pdf before choosing any clinical use.

Source evidence

docs/evidence/hospital-deployment/: access-checks.json; doctor-packet-rehearsal.json; doctor-modalities-ui.json; doctor-save-reopen.json; it-recovery.json; it-backup-restore.json; packet-provenance.json. Operational guide: docs/HOSPITAL-DEPLOYMENT.md. Public TCIA image attribution: config/doctor/packets/README.md. Failed attempts are retained in IMPLEMENTATION-LOG.md.