

Lecture 33

Last Time

Def : A non-zero vec $\vec{v} \in \mathbb{R}^n$ is an eigenvector for $A \in M_{n \times n}$ with eigenvalue $\lambda \in \mathbb{R}$ if $A\vec{v} = \lambda\vec{v}$

• A is diagonalizable i.f.f. \exists basis for \mathbb{R}^n consisting e-vecs $\vec{v}_1, \dots, \vec{v}_n$ for A .

On Exam: No defs on eigenvectors/eigenvalues

↳ e-vecs/e-vals pretty only computational.

• $\lambda \in \mathbb{R}$ is an e-val of A i.f.f. $\det(A - \lambda I) = 0$

• then $\vec{v} \in \mathbb{R}^n$ is an e-vec with e-val λ i.f.f. $\vec{v} \in \ker(A - \lambda I)$.

Ex

$$A = \begin{bmatrix} 3 & 1 & 0 \\ 0 & 2 & 1 \\ 0 & 0 & -1 \end{bmatrix}$$

Find evals/e-vecs of A

$$\det(A - \lambda I) = 0 \quad : \quad \begin{bmatrix} 3-\lambda & 1 & 0 \\ 0 & 2-\lambda & 1 \\ 0 & 0 & -1-\lambda \end{bmatrix} \rightarrow (3-\lambda)(2-\lambda)(-1-\lambda) = 0$$
$$\lambda = 3, 2, -1$$

Moral: If A is upper triangular, the e-vals of A are diagonal entries of A.

e-vals	e-vecs
3	$\begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}$
2	$\begin{bmatrix} -1 \\ 1 \\ 0 \end{bmatrix}$
-1	$\begin{bmatrix} -4 \\ 12 \\ 1 \end{bmatrix}$

$$\lambda = 3$$

$$[A - 3I | 0] \rightarrow \left[\begin{array}{ccc|c} 0 & 1 & 0 & 0 \\ 0 & -1 & 1 & 0 \\ 0 & 0 & -4 & 0 \end{array} \right] \rightarrow \left[\begin{array}{ccc|c} 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 \end{array} \right] \quad \vec{v} = x_1 \begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix}$$

$$\lambda = 2$$

$$[A - 2I | 0] \rightarrow \left[\begin{array}{ccc|c} 1 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & -3 & 0 \end{array} \right] \rightarrow \left[\begin{array}{ccc|c} 1 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 \end{array} \right] \quad \begin{array}{l} x_2 \text{ is free} \\ \text{so} \end{array}$$
$$\vec{v} = x_2 \begin{pmatrix} -1 \\ 1 \\ 0 \end{pmatrix}$$

$$\lambda = -1$$

$$[A - (-1)I | 0] \rightarrow \left[\begin{array}{ccc|c} 4 & 1 & 0 & 0 \\ 0 & 3 & 1 & 0 \\ 0 & 0 & 0 & 0 \end{array} \right] \rightarrow \left[\begin{array}{ccc|c} 12 & 0 & -1 & 0 \\ 0 & 3 & 1 & 0 \\ 0 & 0 & 0 & 0 \end{array} \right]$$

x_3 is free

$$\text{so } x_3 \begin{bmatrix} 1/12 \\ -1/3 \\ 1 \end{bmatrix}$$

you can scale by 12

$$\text{to get } \begin{bmatrix} 1 \\ -4 \\ 12 \end{bmatrix}$$

$$A = S \Lambda S^{-1} \quad S = \begin{bmatrix} 1 & -1 & 1 \\ 0 & 1 & -4 \\ 0 & 0 & 12 \end{bmatrix} \quad \leftarrow \begin{array}{l} \text{each column} \\ \text{is an} \\ \text{e-vec} \end{array}$$

$$\Lambda = \begin{bmatrix} 3 & 0 & 0 \\ 0 & 2 & 0 \\ 0 & 0 & -1 \end{bmatrix} \quad \leftarrow \begin{array}{l} \text{notice each} \\ \text{entry is} \\ \text{the e-val} \\ \text{corresponding to} \\ \text{the col in } S. \end{array}$$

Proposition: A matrix $A \in M_{n \times n}$ is diagonalizable if it has eigenvectors $\vec{v}_1, \dots, \vec{v}_n$ with distinct e-vals $\lambda_1, \dots, \lambda_n \in \mathbb{R}$

Follows from

Lem: If $\vec{v}_1, \dots, \vec{v}_k \in \mathbb{R}^n$ are e-vecs for A with distinct e-vals then $\vec{v}_1, \dots, \vec{v}_k$ are LI.

Pf of Lem: Suppose not, i.e. \exists non-trivial relation

$$\vec{0} = c_1 \vec{v}_1 + \dots + c_k \vec{v}_k$$

$$\rightarrow \begin{pmatrix} \vec{0} = A \vec{0} = c_1 \lambda_1 \vec{v}_1 + \dots + c_k \lambda_k \vec{v}_k \\ \vec{0} = \lambda_k \vec{0} = c_1 \lambda_k \vec{v}_1 + \dots + c_k \lambda_k \vec{v}_k \end{pmatrix}$$

$$\vec{0} = c_1 (\lambda_1 - \lambda_k) \vec{v}_1 + \dots + c_{k-1} (\lambda_{k-1} - \lambda_k) \vec{v}_{k-1}$$

\rightarrow non-trivial relation among $\vec{v}_1, \dots, \vec{v}_k$

\hookrightarrow because $\vec{v}_k \neq \vec{0}$ and $\lambda_1 \neq \lambda_k$ when $i \neq k$.

$$\rightarrow \dots \rightarrow \vec{0} = \tilde{c}_1 \vec{v}_1$$

but this is impossible because $\tilde{c}_1 \neq 0$ and $\vec{v}_1 \neq \vec{0}$

$\therefore \vec{v}_1, \dots, \vec{v}_k$ are independent \square


Ex (not covered by Prop) : $A = [\text{proj}_W]$ where $W \subset \mathbb{R}^3$ is a plane through $\vec{0}$. Then

A is similar to $\begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 0 \end{bmatrix} = \lambda$

$$Av_1 = v_1$$

$$Av_2 = v_2$$

$$Av_3 = 0$$



$\lambda = \text{e-val of } A \in M_{n \times n} \iff \det(A - \lambda I) = 0$

$P(\lambda)$

//characteristic
polynomial
of A //

• Obs : $P(\lambda) = \text{poly of degree } n$

Def : Let $\lambda \in \mathbb{R}$ be an eigenvalue of $A \in M_{n \times n}$

• The algebraic multiplicity of λ is its multiplicity as a root of

$$\det(A - \lambda I) = (\lambda - \lambda_1)^n \cdot \text{stuff}$$

• The geometric multiplicity of λ_1 is $\dim \ker(A - \lambda_1 I)$