

Lecture 17

Types of recurrences:

1. Non-linear : Hard

2. Linear : Easy

Linear Recurrences:

• Just follow the steps.

Linear Homogeneous Recurrences

Ex

$$a_n = 2a_{n-1} + a_{n-2} - 2a_{n-3}, \quad a_0 = 7, \quad a_1 = 0, \quad a_2 = 10$$

Step 1

Move all terms to one side:

$$\rightarrow a_n - 2a_{n-1} - a_{n-2} + 2a_{n-3} = 0$$

$$\rightarrow x^3 - 2x^2 - x + 2 = 0$$

Side Example of characteristic eq:

$$a_n - a_{n-3} = 0$$

$$\rightarrow x^3 - 1 = 0$$

Step 2

Factorization

$$(x-2)(x-1)(x+1) = 0$$

$$x_1 = 2, \quad x_2 = 1, \quad x_3 = -1$$

Step 3

Get general solution by raising all roots to the power by n , and multiply each by different constants.

$$a_n = A(2)^n + B(1)^n + C(-1)^n$$

Step 4 Use initial conditions to set up system.

$$\begin{aligned} a_0 &= A(2)^0 + B(1)^0 + C(-1)^0 \\ \left. \begin{aligned} a_0 &= A + B + C = 7 \\ a_1 &= 2A + B - C = 0 \\ a_2 &= 4A + B + C = 10 \end{aligned} \right\} \text{System} \end{aligned}$$

Step 5 Solve system:

$$A = 1$$

$$B = 2$$

$$C = 4$$

Step 6 Write general solution with constants plugged in to get actual solution.

Another Example

$$a_n = 4a_{n-1} - 4a_{n-2}$$

$$a_n - 4a_{n-1} + 4a_{n-2} = 0$$

$$x^2 - 4x + 4 = 0$$

$$(x-2)^2 = 0$$

$$x_1 = x_2 = 2$$

$$\text{So then } a_n = A2^n + Bn2^n$$

Another Example

If you get $(x-2)^4$ as your roots to characteristic equation:

$$a_n = A2^n + Bn2^n + Cn^22^n + Dn^32^n$$

Linear Non-homogeneous Recurrences

[Ex]

$$f_n = \underbrace{4f_{n-1} - 3f_{n-2}}_{f_n^{(h)}} + \underbrace{4^n}_{f_n^{(p)}}$$

$$f_n = f_n^{(h)} + f_n^{(p)}$$

1. Solve for $f_n^{(h)}$

$$f_n = 4f_{n-1} - 3f_{n-2}$$

$$f_n - 4f_{n-1} - 3f_{n-2} = 0$$

$$x^2 - 4x + 3 = 0$$

$$(x-1)(x-3) = 0$$

$$x_1 = 1, x_2 = 3$$

$$f_n^{(h)} = A(1)^n + B(3)^n$$

2. Solve for $f_n^{(p)}$

↳ Since we have 4^n , we assume a root of the characteristic equation is $x=4$

↳ So you just assume $f_n^{(p)} = A4^n$

↳ then plug in $f_n^{(p)}$ into f_n and set it equal to $f_n^{(p)}$

$$f_n^{(p)} = 4f_{n-1}^{(p)} - 3f_{n-2}^{(p)} + 4^n$$

$$A4^n = 4(A4^{n-1}) - 3(A4^{n-2}) + 4^n$$

Solve
for
A

$$A = \frac{16}{3}$$

$$\underline{\underline{\text{So } f_n^{(p)} = \frac{16}{3} 4^n}}$$

So $f_n = f_n^{(h)} + f_n^{(p)}$

$$f_n = A(1)^n + B(3)^n + \frac{16}{3} 4^n$$

Ex $f_n = \underbrace{6f_{n-1} - 9f_{n-2}}_{f_n^{(h)}} + \underbrace{2 \cdot 3^n}_{f_n^{(p)}}$

1. Solve $f_n^{(h)}$

$$(x^2 - 6x + 9) = 0$$

$$(x-3)^2 = 0$$

$$x_1 = 3, x_2 = 3$$

$$f_n^{(h)} = A3^n + Bn3^n$$

2. Solve for $f_n^{(p)}$

1. Notice the root is $x=3$

2. Also notice $f_n^{(h)}$ has 2 roots equal to 3

↳ from $f_n^{(p)}$ we found another root equal to 3

↳ So now you know there are three roots equal to 3

$$\text{↳ so } f_n^{(p)} = Cn^2 3^n$$

$$\text{Now do } f_n^{(p)} = 6f_{n-1}^{(p)} - 9f_{n-2}^{(p)} + 2 \cdot 3^n$$

↳ from this $C=1$

So: $f_n = A3^n + Bn(3)^n + n^2 \cdot 3^n$