

ARYAN SINGH SIKARWAR

24BCE0403

RESEARCH PAPER



EFFECT OF NANO-FORMULATED AGROCHEMICALS ON RHIZOSPHERIC COMMUNITIES IN MILLET

• **Abstract**

Nanotechnology is an emerging technology that combines materials science and engineering fundamentals, altogether synthesizing materials at the nanoscale to solve various current problems. Agrochemicals are often used in modern intensive agriculture to protect crops from biotic stress, provide vital nutrients, and boost millet growth and yield. Although they are beneficial in the short term, their long-term and persistent applications damage soil fertility and negatively affect the rhizospheric microbiome. Nanotechnology in the form of nano-based agro-formulations is an innovative, environmentally acceptable, and practical solution for substitute synthetic fertilizers. Nano-technological formulations and nanoparticles in the form of nano-pesticides, nano-herbicides, nano-gels, nano fertilizer, and nanofungicides are reported to have efficacy in continuous release of nutrients regulated distribution to plant nutrients very effectively that helps in fighting phytopathological diseases, promotes plant and microbiome

productivity. In addition, they are effective in alleviating biotic and abiotic stress in crop plants. However, despite their extraordinary effectiveness, they also have several drawbacks, such as a tedious manufacturing process, shaky delivery, and dosage-sensitive effectiveness. Learning and acknowledging the influence of nano-based agrochemicals on a highly nutritious crops like millets is important for further commercialization and wide utilization in agriculture. Hence, this chapter focuses on the usage of nanoformulations on millet crops, their effects on soil rhizospheric communities, and soil fertility. Keywords: Nanoformulations, rhizosphere, millets, soil biota, millet-grown soils.

Keywords: Nanoformulations, rhizosphere, millets, soil biota, millet-grown soils.

1. Introduction

The sustainable increase in food production and eradication of hunger and poverty are the main goals of agricultural sustainability and food safety in world countries. The inorganic synthetic fertilizers provide macronutrients and micronutrients for plant growth, but many agrochemical applications in agricultural fields have shown several disadvantages in the long term. Nutrient Use Efficiency (NUE) of soil reduces with continuous crop growth and application of various chemicals in the soil, leading to imbalanced nutrients and low nutrient uptake by plants. More than 90% of the applied chemicals are lost in the environment, while others are unable to reach the target locations required for efficient use (Nizzetto et al. 2016). For many years, fertilizers and insecticides have been crucial in ensuring food production demands are met (Guo et al. 2018). However, excessive usage of these chemicals also impacts sustainable agriculture development. Nanotechnology has revolutionized its application in agriculture, medicine, and pharmaceuticals (Raliya et al. 2017; Singh et al. 2021). Nano-based agrochemical formulations are introduced in the agroindustry to reduce the excessive utilization of chemical fertilizers and improve soil nutrition with minor alternations to soil properties and microbial communities (Arora et al. 2022). The engineered nanoparticles such as nano fertilizers, nano pesticides, nano insecticides, nano rodenticides, and other nano formulations are utilized in agriculture for their broader scope, and distinct advantages such as controlled release of loaded substances, loss of nutrients, bioavailability, solubility, and desirable synthesis methods and are found as alternatives for conventional chemical fertilizers. A very limited number of highly tiny particles are used in the nanoformulation of agricultural chemicals, which act on the plant rhizosphere as active ingredients in fertilizers or other agricultural chemicals (Ul Haq and Ijaz 2019). They have been extensively tested for their positive and negative impacts on soil health for the past two decades since they are considered

as future enhancers of food production (Kumari and Singh 2020; An et al. 2022). Millets are the most popular crops grown in arid and semi-arid regions of Asia and southern Europe. The agrochemicals such as fertilizers and pesticides have been reported to influence the yield, soil fertility, microbial communities, and enzymatic activities in agriculture. The rhizosphere communities of millets are less focused than rice, wheat, and Cite this chapter Karthika, S., G K Dinesh., et al. (2023). Effect of Nano-Formulated Agrochemicals on Rhizospheric Communities in Millets. In: Pudake, R.N., Kumari, M., Sapkal, D.R., Sharma, A.K. (eds) Millet Rhizosphere. Rhizosphere Biology. Springer, Singapore. https://doi.org/10.1007/978-981-99-2166-9_15 For more updates: <https://www.researchgate.net/profile/G-K-Dinesh> 4 maize, and these microbial populations participate in all biogeochemical cycles and enhance the nutrient availability to millets (Xu et al. 2018). The first part of this chapter focuses on the influence of agrochemicals and nanoformulations on soil microbiota (rhizosphere communities, soil properties, soil fauna). The latter focuses on existing nano-based agroformulations and possible impacts on millet rhizosphere communities.

2. Current trends in Nano-agrochemical formulations

In recent years, nanotechnology interventions in agriculture have extended substantially, including nanoscale carriers, nano sensors, clay nanoparticles, nano polymeric composites, nano emulsions, and nano food packaging materials. These novel technologies are also utilized to combat plant diseases, identify pathogens, boost plant resistance, and reduce post-harvest losses. Nano-based agrochemical formulations provide a wide range of benefits (more effectiveness and durability, better distribution and permeability, ability to disintegrate in the soil and environment, lack of toxicity, and photo generative nature) as well as a reduced amount of agro-inputs with convenient pesticide properties to protect crops against insect pests and diseases effectively (Fig. 1). Nano carriers are utilized for slow and sustained delivery of fertilizers, fungicides, herbicides, pesticides, growth regulators, and other target active compounds (Vega-Vásquez et al. 2020). Nano polymeric compounds from plants, animals, and microbe-based byproducts create opportunities for exploring innovative and value-added nano materials and products. For example, clay-based nanomaterials can serve as versatile carriers of nutrients and pesticides with the advantage of extended release and could reduce 70-80% of pesticide use (Merino et al. 2020).



Fig. 1. Application of nanotechnology in agriculture

2.1. Nano carriers for Agrochemical formulations

Potential applications of nanotechnology for integrated pest management include the controlled administration of Artificial Intelligence (AIs) with increased activity at lessening drug concentrations. Nano carriers significantly improve the chemical stability of agrochemicals while protecting them from harsh environmental factors (such as radiation and high temperatures) (Kumar et al. 2019). Nano encapsulation, also known as an encapsulation of active ingredients with nanoparticles, is a technique for the sustained but effective release of chemicals to a specific host plant. The biphasic dispersion of two or more immiscible liquids (water in oil or oil in water) to form droplets stabilized using any amphiphilic surfactant to create an emulsion droplet less than 100 nm is referred to as nanoemulsions. Nanoemulsions are transparent or translucent oil-water emulsions used in agriculture to promote physio-chemical stability, improve the properties of pesticides (bioavailability), and sustainable release of chemicals (Bhattacharyya et al. 2016). The chemical or physical confinement of gels in nano size, which serves as nanocarriers for delivery of agrochemicals or drugs, is referred to as nanogels. These nanogels are swellable cross-linked polymer networks with a high ability to hold water without dissolving it into an aqueous medium. These polymers are either natural or synthetic and combination of both with properties such as amphiphilicity, softness, porosity, and Cite this chapter Karthika, S., G K Dinesh., et al. (2023). Effect of Nano-Formulated Agrochemicals on Rhizospheric Communities in Millets. In: Pudake, R.N., Kumari, M., Sapkal, D.R., Sharma, A.K. (eds) Millet Rhizosphere. Rhizosphere Biology. Springer, Singapore. https://doi.org/10.1007/978-981-99-2166-9_15 For more updates: <https://www.researchgate.net/profile/G-K-Dinesh> 6 degradability according to the variability in composition. The materials (i.e.,

nanoparticles) loaded into the nanogels protect the loaded material from degradation and elimination, and additionally, they actively participate in the delivery at the target site. The stimuli-responsive behavior of nanogels with stability makes them suitable for the carrier of desired nanoparticles. Polymeric nanogels are a new hybrid to overcome poor colloidal stability, elimination, and low aqueous solubility. A wide variety of molecules can be incorporated to increase their utility (Soni et al. 2016). For instance, the core-shell type lignin nanogels formed using lignin methacrylate, emulsified and graft copolymerized along with acrylic acid, were stimuli-responsive (pH-responsive swelling). Therefore, lignin bio-based nanogels could be used as a nanocarrier for the controlled release of pesticides or fungicides (Yiamsawas et al. 2021). The different types of nano-agrochemical formulations are used in the manufacturing of nano-fertilizer, nano-pesticides, nano-herbicides, nano-fungicides, etc.




Fig. 3. Commercially used nano-based agrochemicals

3. Soil rhizosphere communities:

Role, interactions, and importance in crop production Nanotechnology provides a novel agrochemical tool that helps in high crop productivity, hazardous chemical reduction, plant-pathogen management, and crop protection Cite this chapter Karthika, S., G K Dinesh., et al. (2023). Effect of Nano-Formulated Agrochemicals on Rhizospheric Communities in Millets. In: Pudake, R.N., Kumari, M., Sapkal, D.R., Sharma, A.K. (eds) Millet Rhizosphere. Rhizosphere Biology. Springer, Singapore.

https://doi.org/10.1007/978-981-99-2166-9_15 For more updates:

<https://www.researchgate.net/profile/G-K-Dinesh> 7 through plant adaptation (Shang et

al. 2019). Nanoparticles may have an impact on the soil microorganisms in the environment via (1) physical direct contact, (2) alteration of available nutrients and toxins, and (3) indirect effects due to their interaction with soil organic materials, therefore, leading to alterations in cell surface morphology and growth behavior, disruption of biofilm and core microorganisms of the millet rhizosphere including many organisms (Xie et al. 2017). Microbial communities in the rhizosphere are well known to be closely linked to plant roots. In addition, rhizosphere bacteria significantly impact nutrient uptake by increasing or lowering nutrient availability. Microbial communities in the rhizosphere are a subset of the soil microbial community but are frequently distinct from those in the bulk soil (Dotaniya and Meena 2015). Furthermore, rhizosphere microorganisms are recognized to have a crucial role in plant health, and plant health influences rhizosphere microbes in millets (Babu et al. 2019). It has been found that the rhizosphere microbial communities associated with healthy and diseased plants differ, although rhizospheres associated with infected and dead plants have comparable microbial compositions (Liu et al. 2020). In millets, the rhizosphere is the most active micro-domain for plant-microbe interactions, and it contains microorganisms vital for plant growth and health maintenance. Rhizosphere bacteria, for example, can provide nutrients to plants by speeding up nutrient cycling or nitrogen fixation and generating hormones or antibiotics that improve plant drought, salt, and disease tolerance. The microbiota can either hurt or help the host plant. Beneficial microbes in the soil rhizosphere can aid plant adaptation to be natural and artificial environmental challenges and climate change (Zhou et al. 2009). Microorganisms in the rhizosphere have positive and negative interactions with plants and roots because soil habitats vary due to various circumstances, defining the microbiome for millets becomes critical (Elhady et al. 2018). Due to their quick sensitivity to environmental changes, microorganisms serve as early warning biological indices of soil and root health. Microbial diversity in the rhizosphere fluctuates as a result of such variables, as well as farming patterns and climate (Rosen and Allan 2007). The microbial diversity in the rhizosphere of finger millet is unknown at this time; thus, the millet microbiome is essential for future nutritional and food security. Furthermore, the finger millet plant indicates its potential involvement in reducing pathogenic strains and offering medicinal and nutritional benefits to the linked plants. In Cite this chapter Karthika, S., G K Dinesh., et al. (2023). Effect of Nano-Formulated Agrochemicals on Rhizospheric Communities in Millets. In: Pudake, R.N., Kumari, M., Sapkal, D.R., Sharma, A.K. (eds) Millet Rhizosphere. Rhizosphere Biology. Springer, Singapore. https://doi.org/10.1007/978-981-99-2166-9_15 For more updates: <https://www.researchgate.net/profile/G-K-Dinesh> 8 environmentally problematic settings such as high salinity and locations with severe water shortages, it would be worthwhile to include beneficial species of this genus as bioinoculants to boost stress tolerance and produce finger millet (Kour et al. 2020). Intraspecific variability of millet rhizodeposition

and, more broadly, phenomena leading to root-driven soil aggregation and microbiological diversity are significant guides to long-term studies on climatic stress and early warning cues of aphids and other pathogen attacks. To measure the diversity of the ratio between the mass of root-adhering soil and the mass of root tissue, researchers looked at the relationships between soil adherence to roots, aggregation, microbiological activity, and root exudation (Amellal et al. 1998). Furthermore, the nutritional quality of the sediment and the vegetation mix of the patches may influence plant species' ability to shape their rhizosphere microbial community (Zhang et al. 2020). Microbes in the rhizosphere are essential for plant health and ecosystem functioning in terrestrial ecosystems. Yet, our knowledge of microbial co-occurrences and assembly processes in the millet rhizosphere is limited.

4. Influence of agrochemicals on soil microbiota

The unsustainable agricultural practices in many developing countries resulted in direct and indirect ecosystem contamination. The new technologies have further influenced the adverse impact of unsustainable practices, leading to land, water, and atmospheric degradation. The various agrochemicals such as herbicides, pesticides, insecticides, molluscicides, nematicides, rodenticides, fungicides, and many other chemical or inorganic compounds have impacted soil microbiota (Meena et al. 2020). A large quantity of agrochemicals used in agriculture has affected soil microbiota directly, which are considered a biological indicator for soil fertility. The nutrient status of soil, microbial carbon biomass, and non-target biota are disturbed by the impact of various pesticides. They also affect the soil enzymes such as urease, dehydrogenase, nitrate reductase, oxidoreductase, and hydrolases (Wang et al. 2006; Hussain et al. 2009).

4.1 Effects of fungicides on soil microbiota

Cite this chapter Karthika, S., G K Dinesh., et al. (2023). Effect of Nano-Formulated Agrochemicals on Rhizospheric Communities in Millets. In: Pudake, R.N., Kumari, M., Sapkal, D.R., Sharma, A.K. (eds) Millet Rhizosphere. Rhizosphere Biology. Springer, Singapore. https://doi.org/10.1007/978-981-99-2166-9_15 For more updates: <https://www.researchgate.net/profile/G-K-Dinesh> 9 The harmful effects on microbial growth, functions, metabolism, and survival alter soil fertility. Some of the negative impacts of fungicides on microbiota are: (i). mancozeb and chlorothalonil could reduce the nitrification and denitrification process; (ii). Carbendazim is highly toxic to a potential biocontrol, *Trichoderma harzianum* (anti-fungal against *Fusarium*, *Rhizoctonia*, and *Pythium*); (iii). Copper-based fungicides affect the N-fixing bacterial population; (iv). Dimethomorph inhibits ammonification and nitrification in sandy soil; (v). Fludioxonil is toxic to algae; (vi). Captan decreases nitrogenase activity and reduces viable counts of nitrifying and denitrifying bacteria, *Rhizobium ciceri*; (vi). Dinocap affects the

ammonifying bacteria (Kyei-Boahen et al. 2001; Virág et al. 2007; Cycoń et al. 2010). In addition, my non-target fungicides cause inhibitory effects by reducing the beneficial microbiota in soil .

4.2. Impact of herbicides on soil microbiota

Based upon the type of molecule present in the applied herbicide, the microbial population decreases within 7 to 30 days. Microbes' biosynthetic mechanisms or physiology are affected indirectly, affecting soil enzymes, protein biosynthesis, and plant growth regulators (Ethylene, gibberellins, and Indoleacetic Acid) (Kremer and Means 2009; Meena et al. 2020). The effects of herbicides on soil microbiota depend on concentration, degradability, adsorption, desorption, persistence, bioactivity, and toxic dosage level. Some microbes which can withstand a single application (Butachlor) are more sensitive to combination (Butachlor with cadmium) (Hussain et al. 2009). The inorganic fertilizers combined with heavy metals suppress soil microbial activities, and secondary metabolites produced by the transformation of herbicides are more severe than direct effects. For instance, metabolites produced by 2,4-D affected a group of gram-negative bacteria, *Burkholderia cepacia*. Even after the N fertilization, prosulfuron in soil affects the nitrification and denitrification process. Atrazine and simazine (10 mg/L) were toxic for the growth and biological activities of *Xanthobacter autotrophic*. Acetochlor and its derivatives negatively impact bacterial communities such as *florescens*, *Mycobacterium phlei*, and *Bacillus subtilis*. The enzymatic activities such as dehydrogenase and invertase are affected by atrazine and metolachlor (Sáez et al. 2006).

4.3. Effects of insecticides on soil microbiota Cite this chapter Karthika, S., G K Dinesh., et al. (2023). Effect of Nano-Formulated Agrochemicals on Rhizospheric Communities in Millets. In: Pudake, R.N., Kumari, M., Sapkal, D.R., Sharma, A.K. (eds) Millet Rhizosphere. Rhizosphere Biology. Springer, Singapore.
https://doi.org/10.1007/978-981-99-2166-9_15 For more updates:
<https://www.researchgate.net/profile/G-K-Dinesh>

10 The growth and population of bacterial species such as *Azotobacter*, *Pseudomonas*, *Bacillus*, *Penicillium*, and many others are affected by long-term insecticides such as fenthion parathion and malathion. The soil polluted with 15 organochlorine pesticides exerted severe ecological risk, among which 76% of ecotoxicity was exhibited by endosulfan. The bioindicators, including 61005 bacterial species and 33397 fungal species were not found in organochlorine pesticide-affected soil, whereas they were present in uncontaminated soil (Egbe et al. 2021). In addition, the short-term application of thiamethoxam reduced the microbial diversity, bacterial abundance, and structure during short-term application (*Actinobacteria*) (Wu et al. 2021). Like other pesticides, insecticides inhibit the soil enzyme activities (urease, phosphatase, and nitrogenase), microbial biomass, and beneficial species population reduction.

5. Application of Nano-formulated agrochemicals for various crops

5.1. Enhancing crop quality and growth using nano-formulated fertilizers

Nano-fertilizers are a noteworthy recent development in decreasing the loss of applied fertilizers to the environment. By combining plant nutrients with nanomaterials, coating nutrient molecules with a thin layer of nanomaterials, and creating nano-sized emulsions, nano-fertilizers are created that intelligently increase soil fertility and bioavailability (Sidorowicz et al. 2019). Additionally, studies of nutrients on slow/controlled release or control loss of nano-fertilizers conducted in soil and water have confirmed that the long-term availability of all doped nutrients to the plant over the whole crop period of cultivation is essential for promoting germination, growth, flowering, and fruiting (Lateef et al. 2016). Different nanomaterials have been used to encapsulate macronutrients like carbon (C), nitrogen (N), potassium (K), phosphorus (P), calcium (Ca), sulphur (S), and magnesium (Mg) to increase crop fertilizer uptake and reduce fertilizer outflow (Singh et al. 2021). For instance, urea-hydroxyapatite (HA) Nano formulations have the potential to increase release time and decrease nitrogen fertilizer consumption. Recent research has demonstrated that NPs, including CNTs, silicon dioxide (SiO₂) NPs, zinc oxide (ZnO) NPs, titanium dioxide (TiO₂) NPs, and even gold (Au) NPs, have favorable impacts on seed germination in agricultural plants, including tomato, wheat, rice, pearl millet, soybean, barley, and maize (Singh et al. 2016). Compared to untreated plants, the germination rate of pearl millet was dramatically improved by Au Nano formulations (Parveen et al. 2016). Numerous studies Cite this chapter Karthika, S., G K Dinesh., et al. (2023). Effect of Nano-Formulated Agrochemicals on Rhizospheric Communities in Millets. In: Pudake, R.N., Kumari, M., Sapkal, D.R., Sharma, A.K. (eds) Millet Rhizosphere. Rhizosphere Biology. Springer, Singapore. https://doi.org/10.1007/978-981-99-2166-9_15 For more updates: <https://www.researchgate.net/profile/G-K-Dinesh> 11 have demonstrated the benefits of using nanoparticles on germination, plant growth, and development (Table 1). Manufacturing proteins, carbohydrates, and hormones (such as auxins) are all regulated by micronutrients, which shield plants from infections and pests. Fe/SiO₂ nanoparticles have a great potential to enhance seed germination in barley and maize (Najafi Disfani et al. 2017). In addition to germination, nanomaterials such as ZnO, TiO₂, MWCNTs, FeO, CuZnFe₄O₄-oxide, and hydroxyl fullerenes are said to promote crop growth and development with quality improvement in a variety of crop species, including mustard, onion, spinach, tomato, potato, and wheat (Shalaby et al. 2016). For

example, Zn nanoparticles were used to boost grain output in a field crop called pearl millet (*Pennisetum americanum* L.) by roughly 38% (Moghaddasi et al. 2017). In several horticultural crop plants, including garden pea, cucumber, spinach, tomato, potato, eggplants, chilli, coriander, and onion, the usefulness of zinc-based nano fertilizers is documented (Ahmed et al. 2018; Pullagurala et al. 2018) (Table 1).

5.2. Nano-formulated chemicals for crop development and protection

Insecticide, herbicide, fungicide, and bactericide nano-formulation or encapsulation using nanomaterials holds immense potential for reducing agrochemical doses, boosting crop productivity, and fostering sustainable development (Liu and Lal 2015). Various nanoformulations of agricultural chemicals are summarized in Table 2. Additionally, using fewer pesticides lowers the cost of growing crops. In these situations, the application of NPs is reported as an effective alternative to directly decrease pest and disease infection and activity, resulting in higher crop development and output (Servin et al. 2015). Polymeric NPs (like chitosan and solid lipids), inorganic nonmetallic NPs (like silica NPs and nanoclays), and metallic NPs (like Cu NPs and ZnO NPs) are examples of agrochemical nanocarriers (Shukla et al. 2020). Furthermore, numerous studies have demonstrated that nano-formulated pesticides are more effective at eliminating pests without harming people or the environment. For instance, when applied to *Drosophila melanogaster*, spinosad- and permethrin-loaded chitosan NPs showed enhanced bioavailability even at lower dosages than free spinosad and free permethrin, and the nanocomposites caused less harm to people and the environment (Sharma et al. 2019). The nano-insecticide particles shrink and become more concentrated after being Cite this chapter Karthika, S., G K Dinesh., et al. (2023). Effect of Nano-Formulated Agrochemicals on Rhizospheric Communities in Millets. In: Pudake, R.N., Kumari, M., Sapkal, D.R., Sharma, A.K. (eds) Millet Rhizosphere. Rhizosphere Biology. Springer, Singapore. https://doi.org/10.1007/978-981-99-2166-9_15 For more updates: <https://www.researchgate.net/profile/G-K-Dinesh> 12 encapsulated by NPs, giving them stability and a gradual release capability. Furthermore, pesticides can exert systemic effects to suppress chewing or sucking-type insects like aphids by increasing their ability to penetrate through plants and reach the cell sap (Li et al. 2007). For instance, several nano-insecticides benefit from the poisonous properties of metallic NPs. Compared to bulk Al₂O₃ treatment, aluminum oxide (Al₂O₃) NPs showed excellent capability for eradicating *Sitophilus oryzae* on stored rice (Das et al. 2019). It has been established that using pheromones to manage insect populations is a promising and successful strategy (Dweck et al. 2015). The foliar application of polysaccharide NPs loaded with metsulfuron-methyl to wheat-growing weeds dramatically reduced weed biomass compared to conventional pesticides.

Additionally, the cytotoxicity of conventional and nano-herbicides was determined by culturing cells, and the results revealed that the herbicide-loaded NPs were less hazardous than the conventional herbicides (Kumar et al. 2017). According to a different study, solid lipid NP-based nanoherbicides had improved release patterns and herbicidal action than conventional herbicides. *Raphanus raphanistrum* (a weed species) development was considerably slowed down when atrazine was encapsulated by solid lipid NPs as opposed to the untreated group. Additionally, *Zea mays* were unaffected by the tested nanoherbicide concentration (de Oliveira et al. 2015). In weeds treated to SiO₂ nanoparticles, (Sharifi-Rad et al. 2018) found that germination, root and shoot lengths, fresh and dried weights, and photosynthetic pigments with total protein all fell dramatically. Similar to this, (Kumar et al. 2017) demonstrated that pectin (polysaccharide) nanoparticles loaded with the herbicide (metsulfuron methyl) are more cytotoxic to *Chenopodium album* plants both in the laboratory and in field conditions, and only a very low amount of AI is needed in comparison to the commercial herbicide. Commercial herbicides only control or eradicate the weeds' aboveground sections, leaving their underground components like rhizomes or tubers unaffected. As a result, weeds begin to grow again; however, nanoherbicides stop this from happening (Dwivedi et al. 2016).

6. Impact of nanoformulations on plant growth **(Root and plant endophytes)**

The interaction and uptake of nanoparticles are influenced by several factors like the nature of nanoparticles, plant physiology, the interaction of nanomaterials with the environment, etc. The mechanism of entry of nanoparticles, their mode of application, and their penetration through plant cells are detailed in Fig. 3. The common factors influencing nanoparticle uptake and transport in plants are, a) Nature of nanoparticle, its surface charge, and chemical properties; b) Method of application of nanoparticle; c) Nanoparticles interact with microorganism in soil, which might impede their absorption into the plant system via several tissues like epidermis, endodermis, casparian strips, and cuticle till vascular bundles; d) Mechanisms involved in nanoparticle internalization within the plant tissues, such as mediation of carrier proteins, pore formation, endocytosis, etc. (Pérez-de-Luque 2017). The entry of Nanoparticles occurs via. root system through cells using symplastic pathway or around the cells to reach xylem via apoplastic pathway (Lv et al. 2019). The nanoparticle uptake involves diffusion, gated channel confirmation, symporter, and antiporter active transport. Diffusion involves the movement of ion molecules from a region of higher concentration to a region of lower concentration. There are two types of gated ion channels, one, the Voltage-gated ion channels, and two, ligand-gated ion channels. These are the class of transmembrane proteins. In the case of voltage-gated ion channels, it forms the ion channel activated

near the channel by changes in the electrical membrane potential, which alters the conformation of the channel proteins, regulating their opening and closing. While in the case of ligand-gated ion channels, they provide a means of converting the biochemical signals into ion flux and electrical events near the membrane (Banerjee et al. 2019). In plants, the intracellular and extracellular signaling ligands are represented by cyclic nucleotides and glutamate, respectively. A symporter and the antiporters are integral membrane proteins and a type of co-transporter and exchanger, respectively. The symporter in the plasma membrane involves in exchange of two dissimilar molecules in the same direction through the cell membrane, whereas the antiporter exchange two or more different molecules or ions across a phospholipid membrane termed “secondary active transport” in opposite directions, one into the cell and one out of the cell (Lv et al. 2019).

9. Advantages and disadvantages

There is enormous pressure on the agriculture sector to find a solution for the continuous and bulky usage of chemicals in agriculture productivity to feed the growing population. This is because the nutrients applied to the soil are leached or lost than absorbed by plants. On the other hand, the soil rhizosphere communities of millets and other crops require fertilizers to retain their metabolisms and growth while increasing crop productivity. The development of nanoformulations with crucial properties is gaining attention worldwide; therefore, both sides of the coin need to be addressed for sustainable usage (Fig. 4).




Fig.4. Benefits and limitations of nano-based agroformulations

10. Research gaps and future perspective

In agriculture, more than 232 nano-enabled products are manufactured by 75 companies in 26 countries worldwide, most of which are in the fertilizer and pesticides category (Vishnu and Dhandapani 2021). The optimal NPK fertilizers ratio of 4:2:1 is ideal for crop productivity, while the current ratio is maintained at 10:2.7:1 in India, with more heavily subsidized urea (Sharma et al. 2021). A clear understanding of the interaction between nanoparticles and rhizosphere reactions is necessary to assess the impact of nanoformulated agrochemicals on crop production through improved disease resistance, enhanced nutrient use efficiency, and better crop yield, mainly in millets. The interaction of nanoformulated agrochemicals in millets crops should have a better understanding of regarding the important aspects such as (i) mechanism behind the improvement of growth, resistance, and yield of millets, (ii) impact on the rhizosphere environment, (iii) adverse effects on millets if any. However, extensive research in millet crops regarding the impact of nanoformulated agrochemicals is needed. It is also essential to consider the factors such as environmental safety, cost efficiency, sustainability, and availability to farmers for largescale applications. Cite this chapter Karthika, S., G K Dinesh., et al. (2023). Effect of Nano-Formulated Agrochemicals on Rhizospheric Communities in Millets. In: Pudake, R.N., Kumari, M., Sapkal, D.R., Sharma, A.K. (eds) Millet Rhizosphere. Rhizosphere Biology. Springer, Singapore. https://doi.org/10.1007/978-981-99-2166-9_15 For more updates: <https://www.researchgate.net/profile/G-K-Dinesh> 33 As a result, fresh ways are required to safeguard the food plants with minimum or no adverse environmental consequences. Developing several nano-formulated agrochemicals has demonstrated higher agrochemical activity, controlled release capacity, and targeted delivery qualities. They can also restore the oxidative stress symptoms induced by numerous stresses in roots and plants and stimulate plant adaptation to unfavorable conditions, making them an important agricultural tool. Though nanotechnology finds a vast relevance in agriculture, it is still in a very nascent stage, especially on nano biomaterials for agriculture use. The nano products developed so far have not yet penetrated the market and have not reached the common farmer. Concerted efforts are needed to promote the application of nanotechnology with multi-stakeholder participation, including the Government, Private Sector industries, educational institutions, and other relevant stakeholders engaged in agriculture. Formulating suitable policies and educating and encouraging the farmers for wide-scale use will sustainably assist in food and nutritional security, contribute to a safeguarding environment, and combat climate change.

11. Conclusion

The chapter discussed the state of nanoformulations in rhizospheric communities focusing on the current status and possible effects on rhizosphere communities of millets. This groundwork on millets will help formulate further research on the influence of nanoformulations of microbial communities in millet farming and how it may pose challenges to humans through the food chain. The alteration in soil microbiota such as microbial biomass, the microbial population including rhizosphere communities, enzyme activities, and function are positively explored during nano agrochemical application. But the uncertainty of nanomaterials in agriculture must be explored more, and a life cycle assessment of each nano product should be discovered before commercialization. The possible effects of millet rhizosphere communities may differ based on the dosage and concentration of nanoformulations and synthesis method. Hence, exploration of toxicity on the environment should be a research priority, especially in millets, since research findings on millet rhizosphere communities are meager comparatively. Biosynthesis of nano agrochemicals should be encouraged to improve yield and productivity sustainably.

References

Cite this chapter Karthika, S., G K Dinesh., et al. (2023). Effect of Nano-Formulated Agrochemicals on Rhizospheric Communities in Millets. In: Pudake, R.N., Kumari, M., Sapkal, D.R., Sharma, A.K. (eds) Millet Rhizosphere. Rhizosphere Biology. Springer, Singapore. https://doi.org/10.1007/978-981-99-2166-9_15 For more updates: <https://www.researchgate.net/profile/G-K-Dinesh>

34 Abdel-Aziz H, Hasaneen MN, Omar A (2018) Effect of foliar application of nano chitosan NPK fertilizer on the chemical composition of wheat grains. Egypt J Bot 58:87–95

Abdel-Hafez SII, Nafady NA, Abdel-Rahim IR, et al (2016) Assessment of protein silver nanoparticles toxicity against pathogenic *Alternaria solani*. 3 Biotech 6:1–12

Adams M, Xie J, Xie J, et al (2020) The effect of carrier addition on Anammox start-up and microbial community: a review. Rev Environ Sci Bio/Technology 19:355–368

Adisa IO, Pullagurala VLR, Peralta-Videa JR, et al (2019) Recent advances in nano-enabled fertilizers and pesticides: a critical review of mechanisms of action. Environ Sci Nano 6:2002–2030. <https://doi.org/10.1039/C9EN00265K>

Adisa IO, Reddy Pullagurala VL, Rawat S, et al (2018) Role of cerium compounds in Fusarium wilt suppression and growth enhancement in tomato (*Solanum lycopersicum*). J Agric Food Chem 66:5959–5970

Ahmed B, Khan MS, Musarrat J (2018a) Toxicity assessment of metal oxide nano-pollutants on tomato (*Solanum lycopersicon*): A study on growth dynamics and plant cell death. Environ Pollut 240:802–816

Ahmed B, Shahid M, Khan MS, Musarrat J (2018b) Chromosomal aberrations, cell suppression and oxidative stress generation induced by metal oxide nanoparticles in onion (*Allium cepa*) bulb. Metallomics 10:1315–1327

Ali M, Ahmed T, Wu W, et al (2020) Advancements in plant and microbe-based synthesis of

metallic nanoparticles and their antimicrobial activity against plant pathogens. *Nanomaterials* 10:1146 Almutairi ZM (2016) Effect of nano-silicon application on the expression of salt tolerance genes in germinating tomato ('*Solanum lycopersicum*'L.) seedlings under salt stress. *Plant Omics* 9:106–114 Amellal N, Burtin G, Bartoli F, Heulin T (1998) Colonization of wheat roots by an exopolysaccharide-producing *Pantoea agglomerans* strain and its effect on rhizosphere soil aggregation. *Appl Environ Microbiol* 64:3740–3747 An C, Sun C, Li N, et al (2022) Nanomaterials and nanotechnology for the delivery of agrochemicals: strategies towards sustainable agriculture. *J Nanobiotechnology* 20:1–19 Arora S, Murmu G, Mukherjee K, et al (2022) A Comprehensive Overview of Nanotechnology in Sustainable Agriculture. *J Biotechnol* Avestan S, Ghasemnezhad M, Esfahani M, Byrt CS (2019) Application of nano-silicon dioxide improves salt stress tolerance in strawberry plants. *Agronomy* 9:246 Ayoub HA, Khairy M, Elsaid S, et al (2018) Pesticidal activity of nanostructured metal oxides for generation of alternative pesticide formulations. *J Agric Food Chem* 66:5491– 5498