

Random Experiment: A **random experiment** is a process by which we observe something uncertain. After the experiment, the result of the random experiment is known. An **outcome** is a result of a random experiment.

Example: Throwing a die, a pack of cards.

Trial: Any particular performance of a random experiment is called a ***trial***

Sample space: The set of all possible outcomes of a statistical experiment is called the **sample space**. An **event** is a subset of a sample space.

Example: roll a die; sample space: $S = \{1, 2, 3, 4, 5, 6\}$. If we would like to know the probability of getting even number, then event is $\{2, 4, 6\}$.

Statistical Definition:

If a random experiment or trail results in ' n ' possible outcomes(or cases), out of which ' m ' are favourable to the occurrence of an event E , then the probability ' p ' of occurrence of E , denoted by $P(E)$, is given by:

$$p = P(E) = \frac{\text{Number of favourable cases}}{\text{Total number of exhaustive cases}} = \frac{m}{n}$$

(i) Since $m \geq 0, n \geq 0$ and $m \leq n$

$$P(E) \geq 0 \text{ and } P(E) \leq 1 \implies 0 \leq P(E) \leq 1$$

(ii) $P(E) + P(\bar{E}) = 1$

Algebra of Events:

For events A, B, C

(i) $A \cup B = \{\omega \in S: \omega \in A \text{ or } \omega \in B\}$

(ii) $A \cap B = \{\omega \in S: \omega \in A \text{ and } \omega \in B\}$

(iii) $\bar{A} = \{\omega \in S: \omega \notin A\}$

(iv) $A - B = \{\omega \in S: \omega \in A \text{ but } \omega \notin B\}$

(v) $A \subset B$ for every $\omega \in A, \omega \in B$

(vi) $A = B$ if and only if A and B have same elements, *i.e.*, $A \subset B$ and $B \subset A$

(vii) A and B disjoint events (mutually exclusive) $\Rightarrow A \cap B = \Phi$ (*null set*)

(viii) $A \cup B$ can be denoted by $A + B$ if A and B are disjoint.

(ix) $A \Delta B$ denotes those ω belonging to exactly one of A and B, *i.e.*,

$$A \Delta B = \bar{A} B \cup A \bar{B} = \bar{A} B + A \bar{B} \quad (\text{disjoint events})$$

Theorems of Probability

1) Addition Theorem: $P(A \cup B) = P(A) + P(B) - P(A \cap B)$

2) $P(\cup_{i=1}^n A_i) = \sum_{i=1}^n P(A_i) - \sum \sum P(A_i \cap A_j) + \dots + (-1)^{n-1} P(A_1 \cap A_2 \cap \dots \cap A_n)$

Conditional Probability: $P(A|B) = \frac{P(A \cap B)}{P(B)}$

If the events A and B are independent, $P(A|B) = P(A)$

$$P(B|A) = P(B)$$

Multiplication theorem of Probability for Independent events:

$$P(A \cap B) = P(A) \cdot P(B)$$

For n events, $P(A_1 \cap A_2 \cap \dots \cap A_n) = P(A_1) P(A_2|A_1) P(A_3|A_1 \cap A_2) \dots \times$

$$P(A_n|A_1 \cap A_2 \cap \dots \cap A_{n-1})$$

Bayes Theorem of Probability:

Theorem: If $E_1, E_2, \dots, E_n$ are mutually disjoint events with $P(E_i) \neq 0$, ($i = 1, 2, \dots, n$), then for any arbitrary event A which is a subset of $\cup_{i=1}^n (E_i)$ such that $P(A) > 0$, we have

$$P(E_i|A) = \frac{P(E_i) P(A|E_i)}{\sum_{i=1}^n P(E_i) P(A|E_i)} ; i = 1, 2, \dots, n$$

A real number X connected with the outcome of a random experiment E .

For example, if E consists of two tosses the random variable which is the number of **heads** (0,1 or 2).

<i>Outcome</i>	:	<i>HH</i>	<i>HT</i>	<i>TH</i>	<i>TT</i>
<i>Value of X</i>	:	2	1	1	0

Thus to each outcome ω , there corresponds a real number $X(\omega)$.

Random Variable

Let S be the sample space associated with a given random experiment $\mathbf{E}$.

A real-valued function defined on S and range is $\mathbf{R} (-\infty, \infty)$, called a *one-dimensional random variable*.

If the function values are ordered pairs of real numbers (*i.e.*, vectors in two-space), the function is said to be a *two-dimensional random variable*.

More generally, an n -dimensional random variable is simply a function whose domain is S and whose range is a collection of n -tuples of real numbers (vectors in n -space).

Definition: A random variable is a function that associates a real number with each element in the sample space.

Example: Two balls are drawn in succession without replacement from an urn containing 4 red balls and 3 black balls. The possible outcomes and the values y of the random variable: Y , where Y is the number of red balls, are

Sample Space	y
RR	2
RB	1
BR	1
BB	0

Note: One-dimensional *r.v.* will be denoted by capital letters, $X, Y, Z, \dots$ etc.

The values which $X, Y, Z, \dots$ etc, can assume are denoted by lower case letters., $x, y, z, \dots$ etc.

Discrete Random Variable

Let $\mathbf{X}$ be a finite random variable on a sample space $\mathbf{S}$, that is, $\mathbf{X}$ assigns only finite number or countably infinite number of values to $\mathbf{S}$.

$$\text{Say, } R_X = \{x_1, x_2, \dots, x_n, \dots, \infty\}$$

Example: marks obtained in a test, number of accidents per month, number of telephone calls per unit time, number of successes in n trials and so on.

Then, $\mathbf{X}$ induces a function $\mathbf{p}$ which assigns probabilities to the points in $\mathbf{R}_X$ as follows:

$$p(x_i) = p_X(x_i) = P(X = x_i) = P\{s \in S : X(s) = x_i\} \text{ for } i = 1, 2, \dots, n$$

Probability Mass function: If $\mathbf{X}$ is a discrete random variable with distinct values $x_1, x_2, \dots, x_n$ then the function $\mathbf{p}_X(\mathbf{x})$ is defined as : $\mathbf{p}_X(x_i) = P(X = x_i) = p_i$.

p_i is called the **probability mass function** of random variable $\mathbf{X}$.

The numbers $p(x_i); i = 1, 2, \dots$ must satisfy the following conditions:

$$(i) \quad p(x_i) \geq 0 \quad \forall \quad i$$

$$(ii) \quad \sum_{i=1}^n p(x_i) = 1$$

Probability distribution :

The collection of pairs $\{x_i, p_i\}$ is called the probability distribution of the R.V. X .

$X = x_i$	x_1		x_i
p_i	p_1		p_i

Example:

I have an unfair coin for which $P(H) = p$, where $0 < p < 1$. I toss the coin repeatedly until I observe a heads for the first time. Let Y be the total number of coin tosses. Find the distribution of Y .

Answer:

First, we note that the random variable Y can potentially take any positive integer, so we have $R_Y = \mathbb{N} = \{1, 2, 3, \dots\}$. To find the distribution of Y , we need to find $P_Y(k) = P(Y = k)$ for $k = 1, 2, 3, \dots$. We have

$$P_Y(1) = P(Y = 1) = P(H) = p,$$

$$P_Y(2) = P(Y = 2) = P(TH) = (1 - p)p,$$

$$P_Y(3) = P(Y = 3) = P(TTH) = (1 - p)^2 p,$$

$$\begin{array}{cccc} \cdot & \cdot & \cdot & \cdot \\ \cdot & \cdot & \cdot & \cdot \\ \cdot & \cdot & \cdot & \cdot \end{array}$$

$$P_Y(k) = P(Y = k) = P(TT \dots TH) = (1 - p)^{k-1} p.$$

Thus, we can write the PMF of Y in the following way

$$P_Y(y) = \begin{cases} (1 - p)^{y-1} p & \text{for } y = 1, 2, 3, \dots \\ 0 & \text{otherwise} \end{cases}$$

2. If $p = \frac{1}{2}$, find $P(2 \leq Y < 5)$.

Answer:

2. if $p = \frac{1}{2}$, to find $P(2 \leq Y < 5)$, we can write

$$\begin{aligned} P(2 \leq Y < 5) &= \sum_{k=2}^4 P_Y(k) \\ &= \sum_{k=2}^4 (1-p)^{k-1} p \\ &= \frac{1}{2} \left(\frac{1}{2} + \frac{1}{4} + \frac{1}{8} \right) \\ &= \frac{7}{16}. \end{aligned}$$

Continuous Random Variables

A random variable X is said to be a continuous random variable if it takes all possible values between certain limits or in an interval which may be finite or infinite.

Example: operating time between two failures of a computer

Probability density function (p.d.f):

The function $f(x)$ is a **probability density function** (pdf) for the continuous random variable X , defined over the set of real numbers, if

1. $f(x) \geq 0$, for all $x \in R$,
2. $\int_{-\infty}^{\infty} f(x) \, dx = 1$,
3. $P(a < X < b) = \int_a^b f(x) \, dx$.

Distribution Function (or) Cumulative Distribution Function (cdf)

If X is a R.V, discrete or continuous, then $F(x) = P(X \leq x)$ is called the cumulative distribution function of X .

i) If X is discrete then
$$F(x) = \sum_j p_j$$

ii) If X is continuous then
$$F(x) = P(-\infty \leq X \leq x) = \int_{-\infty}^x f(x)dx$$

Relationship between cumulative distribution function and probability distribution function

$$f(x) = \frac{dF(x)}{dx}$$

If $p(x)$ is the p.m.f of a random variable 'X' then

$$\text{Mean} = \sum_{n=0}^{\infty} xp(x)$$

$$\text{Variance} = \sum_{n=0}^{\infty} (x - \text{mean})^2 p(x)$$

$$\text{Moment about the mean} = \sum_{n=0}^{\infty} (x - \text{mean})^r p(x)$$

If $f(x)$ is the p.d.f of a random variable 'X' which defined in the interval (a, b) then

$$\text{(i) Mean} = \int_a^b xf(x)dx \quad \text{(ii) Variance} = \int_a^b (x - \text{mean})^2 f(x)dx$$

$$\text{(iii) Moment about mean} = \int_a^b (x - \text{mean})^r f(x)dx$$

Problem 1:

If the random variable takes the values 1,2,3 and 4 such that

$2P(X = 1) = 3P(X = 2) = P(X = 3) = 5P(X = 4)$, find the probability distribution and the cumulative distribution function of X.

Solution:

Let $2P(X = 1) = 3P(X = 2) = P(X = 3) = 5P(X = 4) = k$

That is, $P(X = 1) = \frac{k}{2}$, $P(X = 2) = \frac{k}{3}$, $P(X = 3) = k$, $P(X = 4) = \frac{k}{5}$

We know that, $\sum_i p_i = 1$

That is, $\frac{k}{2} + \frac{k}{3} + k + \frac{k}{5} = 1 \Rightarrow k = \frac{30}{61}$

The probability distribution of X is given by

x_i	1	2	3	4
$P(x_i)$	$\frac{15}{61}$	$\frac{10}{61}$	$\frac{30}{61}$	$\frac{6}{61}$

The c.d.f. $F(x)$ is defined as $F(x) = P(X \leq x)$.

When $x < 1$, $F(x) = 0$

When $1 \leq x < 2$, $F(x) = P(X = 1) = \frac{15}{61}$

When $2 \leq x < 3$, $F(x) = P(X = 1) + P(X = 2) = \frac{25}{61}$

When $3 \leq x < 4$, $F(x) = P(X = 1) + P(X = 2) + P(X = 3) = \frac{55}{61}$

When $x \geq 4$, $F(x) = P(X = 1) + P(X = 2) + P(X = 3) + P(X = 4) = 1$

Problem 2:

A random variable X has the following probability function.

Values of X	0	1	2	3	4	5	6	7	8
$P(X = x)$	a	3a	5a	7a	9a	11a	13a	15a	17a

(i) Find the value of a (ii) Find $P(X < 3)$, $P(0 < X < 3)$, $P(X \geq 3)$

(iii) Find cumulative distribution function of X.

Solution:

i) We know that , $\sum_i p_i = 1$

$$a + 3a + 5a + 7a + 9a + 11a + 13a + 15a + 17a = 1$$

$$81a = 1 \Rightarrow a = \frac{1}{81}$$

ii) $P(X < 3) = P(X = 0) + P(X = 1) + P(X = 2) = a + 3a + 5a$
 $= 9a = \frac{1}{9}$

$$P(0 < X < 3) = P(X = 1) + P(X = 2) = 3a + 5a = 8a = \frac{8}{81}$$

$$P(X \geq 3) = 1 - P(X < 3) = 1 - \frac{1}{9} = \frac{8}{9}$$

iii) Distribution function $F(x)$ of X

Values of X	0	1	2	3	4	5	6	7	8
$F(x) = P(X \leq x)$	a	$4a$	$9a$	$16a$	$25a$	$36a$	$49a$	$64a$	$81a$
	$\frac{1}{81}$	$\frac{4}{81}$	$\frac{9}{81}$	$\frac{16}{81}$	$\frac{25}{81}$	$\frac{36}{81}$	$\frac{49}{81}$	$\frac{64}{81}$	1

Problem 3: Let X be a random variable with PDF given by

$$f_X(x) = \begin{cases} cx^2 & |x| \leq 1 \\ 0 & \text{otherwise} \end{cases}$$

- Find the constant c .
- Find EX and $\text{Var}(X)$.
- Find $P(X \geq \frac{1}{2})$.

a. To find c , we can use $\int_{-\infty}^{\infty} f_X(u)du = 1$:

$$\begin{aligned} 1 &= \int_{-\infty}^{\infty} f_X(u)du \\ &= \int_{-1}^1 cu^2du \\ &= \frac{2}{3}c. \end{aligned}$$

Thus, we must have $c = \frac{3}{2}$.

b. To find EX , we can write

$$\begin{aligned} EX &= \int_{-1}^1 uf_X(u)du \\ &= \frac{3}{2} \int_{-1}^1 u^3du \\ &= 0. \end{aligned}$$

In fact, we could have guessed $EX = 0$ because the PDF is symmetric around $x = 0$. To find $\text{Var}(X)$, we have

$$\begin{aligned} \text{Var}(X) &= EX^2 - (EX)^2 = EX^2 \\ &= \int_{-1}^1 u^2 f_X(u)du \\ &= \frac{3}{2} \int_{-1}^1 u^4du \\ &= \frac{3}{5}. \end{aligned}$$

c. To find $P(X \geq \frac{1}{2})$, we can write

$$P(X \geq \frac{1}{2}) = \frac{3}{2} \int_{\frac{1}{2}}^1 x^2 dx = \frac{7}{16}.$$

Problem 4: Let X be a continuous random variable with PDF given by

$$f_X(x) = \frac{1}{2}e^{-|x|}, \quad \text{for all } x \in \mathbb{R}.$$

If $Y = X^2$, find the CDF of Y .

Answer: First, we note that $R_Y = [0, \infty)$. For $y \in [0, \infty)$, we have

$$\begin{aligned} F_Y(y) &= P(Y \leq y) \\ &= P(X^2 \leq y) \\ &= P(-\sqrt{y} \leq X \leq \sqrt{y}) \\ &= \int_{-\sqrt{y}}^{\sqrt{y}} \frac{1}{2}e^{-|x|} dx \\ &= \int_0^{\sqrt{y}} e^{-x} dx \\ &= 1 - e^{-\sqrt{y}}. \end{aligned}$$

Thus,

$$F_Y(y) = \begin{cases} 1 - e^{-\sqrt{y}} & y \geq 0 \\ 0 & \text{otherwise} \end{cases}$$

Problem 5: Let X be a continuous random variable with PDF

$$f_X(x) = \begin{cases} 4x^3 & 0 < x \leq 1 \\ 0 & \text{otherwise} \end{cases}$$

Find $P(X \leq \frac{2}{3} | X > \frac{1}{3})$.

Answer:

We have

$$\begin{aligned} P(X \leq \frac{2}{3} | X > \frac{1}{3}) &= \frac{P(\frac{1}{3} < X \leq \frac{2}{3})}{P(X > \frac{1}{3})} \\ &= \frac{\int_{\frac{1}{3}}^{\frac{2}{3}} 4x^3 dx}{\int_{\frac{1}{3}}^1 4x^3 dx} \\ &= \frac{3}{16}. \end{aligned}$$

Problem 6: **EXAMPLE:** Suppose that X is a continuous random variable whose probability density function is given by

$$f(x) = \begin{cases} C(4x - 2x^2) & 0 < x < 2 \\ 0 & \text{otherwise} \end{cases}$$

(a) What is the value of C ?

(b) Find $P\{X > 1\}$.

SOLUTION (a) Since f is a probability density function, we must have that $\int_{-\infty}^{\infty} f(x) dx = 1$, implying that

$$C \int_0^2 (4x - 2x^2) dx = 1$$

or

$$C \left[2x^2 - \frac{2x^3}{3} \right] \Big|_{x=0}^{x=2} = 1$$

or

$$C = \frac{3}{8}$$

(b) Hence

$$P\{X > 1\} = \int_1^{\infty} f(x) dx = \frac{3}{8} \int_1^2 (4x - 2x^2) dx = \frac{1}{2} \quad \blacksquare$$

Problem 7:

If a random variable 'X' has the p.d.f. $f(x) = \begin{cases} \frac{(x+1)}{2} & , -1 < x < 1 \\ 0 & , otherwise \end{cases}$ Then find the mean and variance.

Answer:

$$\text{We know that, Mean} = \int_{-1}^1 xf(x)dx = \int_{-1}^1 x \frac{(x+1)}{2} dx = \frac{1}{2} \int_{-1}^1 (x^2 + x) dx = \frac{1}{3}$$

$$\begin{aligned} \text{Variance} &= \int_{-1}^1 (x - \text{mean})^2 f(x) dx = \int_{-1}^1 \left(x - \frac{1}{3}\right)^2 \frac{(x+1)}{2} dx \\ &= \int_{-1}^1 (9x^2 + 1 - 6x)(x+1) dx = \frac{2}{9} \end{aligned}$$

Two-dimensional Random Variable

Our study of random variables and their probability distributions in the preceding sections is restricted to one-dimensional sample spaces.

In that we recorded outcomes of an experiment as values assumed by a single random variable.

There will be situations, however, where we may find it desirable to record the simultaneous outcomes of several random variables

Example:

For instance, blood pressure and cholesterol for each individual are measured simultaneously.

In a study to determine the likelihood of success in college based on high school data, we might use a three dimensional sample space and record for each individual his or her aptitude test score, high school class rank, and grade-point average at the end of freshman year in college.

Definition: Let S be a sample space associated with a random experiment E . Let X and Y be two random variables defined on S . then the pair (X, Y) is called a Two-dimensional random variable.

The value of (X, Y) at a point $s \in S$ is given by the ordered pair of real numbers $(X(s), Y(s)) = (x, y)$ where $X(s) = x$, $Y(s) = y$.

Two –Dimensional discrete random variable: If the possible values of (X, Y) are finite or countably infinite, then (X, Y) is called a two-dimensional discrete random variable.

When (X, Y) is a two-dimensional discrete random variable the possible values of (X, Y) may be represented as (x_i, y_j) , $i = 1, 2, 3, \dots n$, $j = 1, 2, 3, \dots m$.

Example: Consider the experiment of tossing a coin twice.

The sample space is $S = \{HH, HT, TH, TT\}$. Let X denote the number of heads obtained in the first toss and Y denote the number of heads in the second toss.

s	HH	HT	TH	TT
$X(s)$	1	1	0	0
$Y(s)$	1	0	1	0

(X, Y) is a two-dimensional random variable or bi-variate random variable. The range space of (X, Y) is $\{(1,1), (1,0), (0,1), (0,0)\}$ which is finite and so (X, Y) is a two-dimensional discrete random variables.

Joint probability distribution of Discrete R.V

The function $f(x, y)$ is a **joint probability distribution** or **probability mass function** of the discrete random variables X and Y if

1. $f(x, y) \geq 0$ for all (x, y) ,
2. $\sum_x \sum_y f(x, y) = 1$,
3. $P(X = x, Y = y) = f(x, y)$.

$X \backslash Y$	y_1	y_2	y_3	$\dots$	y_j	$\dots$	y_m	Total
x_1	p_{11}	p_{12}	p_{13}	$\dots$	p_{1j}	$\dots$	p_{1m}	$p_{1\cdot}$
x_2	p_{21}	p_{22}	p_{23}	$\dots$	p_{2j}	$\dots$	p_{2m}	$p_{2\cdot}$
x_3	p_{31}	p_{32}	p_{33}	$\dots$	p_{3j}	$\dots$	p_{3m}	$p_{3\cdot}$
$\vdots$	$\vdots$	$\vdots$	$\vdots$	$\vdots$	$\vdots$	$\vdots$	$\vdots$	$\vdots$
x_i	p_{i1}	p_{i2}	p_{i3}	$\dots$	p_{ij}	$\dots$	p_{im}	$p_{i\cdot}$
$\vdots$	$\vdots$	$\vdots$	$\vdots$	$\vdots$	$\vdots$	$\vdots$	$\vdots$	$\vdots$
x_n	p_{n1}	p_{n2}	p_{n3}	$\dots$	p_{nj}	$\dots$	p_{nm}	$p_{n\cdot}$
Total	$p_{\cdot 1}$	$p_{\cdot 2}$	$p_{\cdot 3}$	$\dots$	$p_{\cdot j}$	$\dots$	$p_{\cdot m}$	1

Roll two dice. Let X be the value on the first die and let T be the total on both dice. Here is the joint probability table:

$X \backslash T$	2	3	4	5	6	7	8	9	10	11	12
1	1/36	1/36	1/36	1/36	1/36	1/36	0	0	0	0	0
2	0	1/36	1/36	1/36	1/36	1/36	1/36	0	0	0	0
3	0	0	1/36	1/36	1/36	1/36	1/36	1/36	0	0	0
4	0	0	0	1/36	1/36	1/36	1/36	1/36	1/36	0	0
5	0	0	0	0	1/36	1/36	1/36	1/36	1/36	1/36	0
6	0	0	0	0	0	1/36	1/36	1/36	1/36	1/36	1/36

Exercise:

Suppose a car showroom has 10 cars of particular brand out of which 5 are good, 2 have defective transmission 3 have defective steering. If 2 cars are selected at random, find the joint probability distribution table.

Answer:

Let X denotes the number of cars with DT

Y denotes the number of cars with DS

$$P(X = 0, Y = 0) = \frac{\binom{5}{2}}{\binom{10}{2}} = \frac{2}{9}$$
$$P(X = 0, Y = 1) = \frac{\binom{5}{1}\binom{3}{1}}{\binom{10}{2}} = \frac{1}{3}$$

and so on....

$X \backslash Y$	0	1	2
0	$10/45$	$15/45$	$3/45$
1	$10/45$	$6/45$	0
2	$1/45$	0	0

Joint probability distribution of Continuous R.V

The function $f(x, y)$ is a **joint density function** of the continuous random variables X and Y if

1. $f(x, y) \geq 0$, for all (x, y) ,
2. $\int_{-\infty}^{\infty} \int_{-\infty}^{\infty} f(x, y) \, dx \, dy = 1$,
3. $P[(X, Y) \in A] = \int \int_A f(x, y) \, dx \, dy$, for any region A in the xy plane.

Exercise:

$$f(x, y) = \begin{cases} \frac{2}{5}(2x + 3y), & 0 \leq x \leq 1, 0 \leq y \leq 1, \\ 0, & \text{elsewhere.} \end{cases}$$

(a) Verify condition 2

(b) Find $P[(X, Y) \in A]$, where $A = \{(x, y) \mid 0 < x < \frac{1}{2}, \frac{1}{4} < y < \frac{1}{2}\}$.

Answer:

(a) The integration of $f(x, y)$ over the whole region is

$$\begin{aligned} \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} f(x, y) \, dx \, dy &= \int_0^1 \int_0^1 \frac{2}{5}(2x + 3y) \, dx \, dy \\ &= \int_0^1 \left(\frac{2x^2}{5} + \frac{6xy}{5} \right) \Big|_{x=0}^{x=1} dy \\ &= \int_0^1 \left(\frac{2}{5} + \frac{6y}{5} \right) dy = \left(\frac{2y}{5} + \frac{3y^2}{5} \right) \Big|_0^1 = \frac{2}{5} + \frac{3}{5} = 1. \end{aligned}$$

(b) To calculate the probability, we use

$$\begin{aligned} P[(X, Y) \in A] &= P\left(0 < X < \frac{1}{2}, \frac{1}{4} < Y < \frac{1}{2}\right) \\ &= \int_{1/4}^{1/2} \int_0^{1/2} \frac{2}{5}(2x + 3y) \, dx \, dy \\ &= \int_{1/4}^{1/2} \left(\frac{2x^2}{5} + \frac{6xy}{5}\right) \Big|_{x=0}^{x=1/2} dy = \int_{1/4}^{1/2} \left(\frac{1}{10} + \frac{3y}{5}\right) dy \\ &= \left(\frac{y}{10} + \frac{3y^2}{10}\right) \Big|_{1/4}^{1/2} \\ &= \frac{1}{10} \left[\left(\frac{1}{2} + \frac{3}{4}\right) - \left(\frac{1}{4} + \frac{3}{16}\right)\right] = \frac{13}{160}. \end{aligned}$$