

Patient Discharge Instructions

Synthetic healthcare document for RAG testing

Document Notice

This file is synthetic and created only for healthcare AI assistant RAG testing. It does not contain real patient data, PHI, confidential client data, or actual medical advice.

1. Purpose

This document provides general discharge instructions for patients leaving a healthcare facility. It is synthetic and should not replace clinician advice.

2. After Discharge

Patients should follow the discharge summary, take medications as prescribed, attend follow-up appointments, monitor symptoms, maintain hydration and nutrition as advised, and keep emergency contact information available.

- Patients should follow the discharge summary, take medications as prescribed, attend follow-up appointments, monitor symptoms, maintain hydration and nutrition as advised, and keep emergency contact information available.

3. Medication Safety

Patients should understand medication names, doses, timing, side effects, and refill instructions before leaving. They should not stop or change medication unless advised by a clinician.

4. Wound Care

If wound care is required, patients should keep the area clean and dry, follow dressing instructions, wash hands before care, and report redness, swelling, pus, fever, or worsening pain.

- If wound care is required, patients should keep the area clean and dry, follow dressing instructions, wash hands before care, and report redness, swelling, pus, fever, or worsening pain.

5. Follow-Up

Patients should schedule follow-up visits according to the discharge plan. Missing follow-up appointments may delay recovery and medication review.

6. Warning Signs

Patients should seek urgent medical attention for severe chest pain, difficulty breathing, uncontrolled bleeding, sudden weakness, confusion, fainting, high fever, or rapidly worsening symptoms.

- Patients should seek urgent medical attention for severe chest pain, difficulty breathing, uncontrolled bleeding, sudden weakness, confusion, fainting, high fever, or rapidly worsening symptoms.

7. Activity and Diet

Patients should follow activity restrictions and diet recommendations given by the clinician. Heavy lifting or driving may be restricted after certain procedures or medications.

8. AI Assistant Boundary

The AI assistant may summarize discharge instructions from documents but must not diagnose symptoms or replace emergency medical care.

Quick Reference Table

Question	Policy Answer
What should patients do after discharge?	Follow discharge summary, take medicines, attend follow-up, monitor symptoms, and keep emergency contacts available.
What wound warning signs should be reported?	Redness, swelling, pus, fever, or worsening pain.
When to seek urgent care?	Severe chest pain, breathing difficulty, uncontrolled bleeding, sudden weakness, confusion, fainting, high fever, or worsening symptoms.

Recommended RAG Test Prompts

- What should patients do after discharge?
- What wound warning signs should be reported?
- When to seek urgent care?
- What information is not available in this document?
- Summarize the main rules from this document.

Version: 1.0 | Data Type: Synthetic | Intended Use: Hackathon prototype and RAG demonstration.