

Ch - 1 → Units And Measurements

- SIGNIFICANT FIGURES :

(1) All non-zero digits are significant.

Eg. 23.459 → 5 significant figures

(2) Zeroes between 2 non-zero digits are significant.

Eg - 10.25  
 102.5  
 1.025 → 4 significant figures

(3) If the number is without decimal, trailing zeroes are not significant.

Eg - 100 → 1 significant figure  
 25000 → 2 significant figures  
 300000 → 1 sign. fig.

(4) If number is with decimal, the zeroes to the right of non-zero digit, but to the left of the decimal are significant.

Eg - 250.56 → 5 sig. fig.  
 3800.60 → 5 sig. fig.

(5) If number is with decimal, other zeroes to the right side of decimal point are significant.

Eg - 2500.00 → because they indicate precision level of  
 0.01 × instrument.

(6) If number is < 1, zeroes to the left but right of decimal point before non-zero are not significant.

Eg. 0.012 → 2 significant figures.

Date: 01/01/24

(7) Significant figures do not depend on system of unit.

$$\frac{25 \text{ km}}{25000 \text{ m}} \rightarrow 2 \text{ sig. fig}$$

Trailing zeros after decimal point are not significant (1)

\* Extra Q

• Counting no. of significant figures in rounded numbers (5)

$$\begin{aligned} ① 0.0009 &\rightarrow 1 \text{ sig. fig.} \\ ② 5.049 &\rightarrow 4 \\ ③ 3000 &\rightarrow 1 \end{aligned}$$

④ 0.020 m → 2 significant figures in product with IT (2)

$$⑤ 0.0200 \rightarrow 3 \text{ sig. fig.} = 0.01 \rightarrow 1$$

$$⑥ 201.0 \rightarrow 4 \text{ sig. figs.} = 0.0020$$

$$⑦ 71 \rightarrow 2 \text{ sig. figs.} = 0.00008$$

$$⑧ 4.50 \times 10^3 \rightarrow 2 \text{ sig. figs.}$$

⑨  $0.23 \times 10^{-6}$  → 2 significant figures in product with IT (2)

⑩  $2301.01 \times 10^2$  → 6 significant figures in product with IT (2)

• Round off the following digits 22.008 - p7

$$22.008 \rightarrow 22.008$$

$$① 18.35 \text{ (3 digits)} \rightarrow 18.4$$

② 18967 (3) → 19000, trailing digits in product with IT (2)

$$③ 18967 (2) \rightarrow 19000 \quad \text{trailing digits zero}$$

$$④ 248337 (3) \rightarrow 248000 \text{ (2.0000) } \rightarrow 6$$

$$⑤ 32.1135 \times 10^3 (4) \rightarrow 321.1 \times 10^4 \rightarrow 6$$

$$⑥ 101.55 \times 10^6 (3) \rightarrow 102 \times 10^6$$

$$⑦ 0.7995 (1) \rightarrow 0.8$$

Significant figures in product with IT (2)

• Trailing zeros are not significant (1)

• Trailing zeros are not significant (1)

M T W T F S S  
Page No.: 05  
Date: 5/4/24

Addition & Subtraction in terms of significant figures:

0.0008

0.002

0.003

0.004

0.005

0.006

0.007

0.008

0.009

0.010

0.011

0.012

0.013

0.014

0.015

0.016

0.017

0.018

0.019

0.020

0.021

0.022

0.023

0.024

0.025

0.026

0.027

0.028

0.029

0.030

0.031

0.032

0.033

0.034

0.035

0.036

0.037

0.038

0.039

0.040

0.041

0.042

0.043

0.044

0.045

0.046

0.047

0.048

0.049

0.050

0.051

0.052

0.053

0.054

0.055

0.056

0.057

0.058

0.059

0.060

0.061

0.062

0.063

0.064

0.065

0.066

0.067

0.068

0.069

0.070

0.071

0.072

0.073

0.074

0.075

0.076

0.077

0.078

0.079

0.080

0.081

0.082

0.083

0.084

0.085

0.086

0.087

0.088

0.089

0.090

0.091

0.092

0.093

0.094

0.095

0.096

0.097

0.098

0.099

0.100

0.101

0.102

0.103

0.104

0.105

0.106

0.107

0.108

0.109

0.110

0.111

0.112

0.113

0.114

0.115

0.116

0.117

0.118

0.119

0.120

0.121

0.122

0.123

0.124

0.125

0.126

0.127

0.128

0.129

0.130

0.131

0.132

0.133

0.134

0.135

0.136

0.137

0.138

0.139

0.140

0.141

0.142

0.143

0.144

0.145

0.146

0.147

0.148

0.149

0.150

0.151

0.152

0.153

0.154

0.155

0.156

0.157

0.158

0.159

0.160

0.161

0.162

0.163

0.164

0.165

0.166

0.167

0.168

0.169

0.170

0.171

0.172

0.173

0.174

0.175

0.176

0.177

0.178

0.179

0.180

0.181

0.182

0.183

0.184

0.185

0.186

0.187

0.188

0.189

0.190

0.191

0.192

0.193

0.194

0.195

0.196

0.197

0.198

0.199

0.200

0.201

0.202

0.203

0.204

0.205

0.206

0.207

0.208

0.209

0.210

0.211

0.212

0.213

0.214

0.215

0.216

0.217

0.218

0.219

0.220

0.221

0.222

0.223

0.224

0.225

0.226

0.227

0.228

0.229

0.230

0.231

0.232

0.233

0.234

0.235

0.236

0.237

0.238

0.239

0.240

0.241

0.242

0.243

0.244

Q - Radius of sphere is 1.41 cm. Express its volume in app. significant digit

$$\rightarrow \text{Volume} = \frac{4}{3} \pi r^3$$

$$\rightarrow \text{Vol} = \frac{4}{3} \times 22 \times \frac{1}{100} \times \frac{1.41}{100} \times \frac{1.41}{100}$$

$$[11.7 \text{ cm}^3]$$

$$Q - \frac{75.5 \times 125.2}{10 \times 16 \times 100} \approx 0.51$$

$$\rightarrow 4820.8260$$

$$\rightarrow 4800$$

$$[22.5]$$

$$Q - 6.2 + 4.33 + 17.456 \text{ Ans. Done}$$

$$Q - 187.20$$

$$- 63.54$$

$$- 123.66$$

$$(d) 123.5$$

$$0.85$$

$$Q - \frac{1.51 \times 10^4}{-3.9 \times 10^3}$$

$$(\cancel{1.51 \times 10^4})$$

$$3 \times 10^5$$

$$3.90 \times 10^4$$

$$- 2.5 \times 10^4$$

$$36.5 \times 10^4$$

$$\rightarrow 36.0 \times 10^4$$

$$390000 \\ - 25000 \\ \hline 365000$$

$$365000 \\ - 20000 \\ \hline 345000$$

$$345000 \\ - 10000 \\ \hline 335000$$

$$335000 \\ - 141000 \\ \hline 194000$$

$$194000 \\ - 56400 \\ \hline 137600$$

$$137600 \\ - 19100 \\ \hline 118500$$

$$118500 \\ - 19880 \\ \hline 98620$$

$$98620 \\ - 25964 \\ \hline 72956$$

$$72956 \\ - 95240 \\ \hline 7616$$

$$7616 \\ - 1203 \\ \hline 6412$$

$$6412 \\ - 751 \\ \hline 5661$$

$$5661 \\ - 1555 \\ \hline 4106$$

$$4106 \\ - 3775 \\ \hline 331$$

$$331 \\ - 38505 \\ \hline 4127$$

$$4127 \\ - 38505 \\ \hline 1252$$

$$1252 \\ - 2147010 \\ \hline 1925250$$

$$1925250 \\ - 1925250 \\ \hline 0$$

## # DIMENSIONAL FORMULA (also known as H.T. eqn)

### • Symbols

i) Length  $\rightarrow [L]$

ii) Mass  $\rightarrow [M]$

iii) Time  $\rightarrow [T]$

iv) Temperature  $\rightarrow [K]$

v) Current  $\rightarrow [A]$

vi) Amount of substance  $\rightarrow [\text{mol}]$

vii) Luminous intensity  $\rightarrow [cd]$

$F_g \rightarrow \text{Force} = \text{mass} \times \text{acceleration}$

$= \text{mass} \times \frac{\text{length}}{\text{time}^2} = \text{kg} \cdot \text{m}^2 \cdot \text{s}^{-2} \text{ N}$

$\text{Force} = [M^1 L^1 T^{-2}] \rightarrow \text{Dimensional formula for force.}$

Try!! (1) Power =  $Wt$   $[L^2 T^3 M]$  =  $\text{Joules} \text{ per } \text{sec}$  (J)

(2) Work  $\rightarrow Fd$

(3) Energy  $\rightarrow \frac{1}{2} mv^2$   $L^2 T^2 M$  =  $\text{Joules}$  (J)

(4) Momentum  $\rightarrow mv$   $L^1 T^1 M$  =  $\text{kg m/s}$  (N)

(5) Plane angle  $\rightarrow \frac{\text{arc}}{\text{radius}}$   $L^2 T^2 M$  =  $\text{radians}$  (rad)

$[A^2 T^2 M] = \frac{L^2 T^2 M}{L^2} =$   $\frac{T^2 M}{T^2} =$   $T^2 M$  =  $\text{radians}$  (rad)

$$\text{1) Power} = \cancel{F \times t} \rightarrow \text{Force} \times \text{displacement}$$

$$= \frac{\text{kgm}}{\text{s}^2} \times m$$

$$= \frac{\text{kgm}^2}{\text{s}^2}$$

$$= [M^1 L^2 T^{-2}]$$

$$(2) \text{Power} = \frac{\text{kgm}^2}{\text{s}^2} \times \frac{1}{\text{s}} \quad [?] \leftarrow \text{constant force}$$

$$= [M^1 L^2 T^{-3}] \quad [A] \leftarrow \text{time}$$

$$(3) \text{Energy} = \cancel{\frac{1}{2} \times \text{kg} \times (\frac{\text{m}}{\text{s}})^2} \quad [?] \leftarrow \text{constant force}$$

$$= [M^1 L^2 T^{-2}]$$

$$(4) \text{Momentum} = \text{mass} \times \text{velocity} = \cancel{\text{kgm} \times \text{ms}^{-1}}$$

$$= [M^1 L^1 T^{-1}]$$

$$(5) \text{Plane angle} = [M^0 L^0 T^0]$$

$$(6) \text{Electric charge} = It$$

$$= As = [M^1 L^0 T^1 A^1]$$

$$(7) \text{Potential difference} : W/Q$$

$$= \frac{\text{kgm}^2}{As^3} = [M^1 L^2 T^3 A^1]$$

$$(8) \text{Resistance} = \frac{W}{Q}$$

$$= \frac{\text{kgm}^2}{A^2 s^3} \rightarrow [M^1 L^2 T^{-3} A^{-2}]$$

$$(9) \text{Resistivity} = \cancel{\rho} \frac{RA}{l}$$

$$= \cancel{\frac{\rho}{l}} = [M^1 L^3 T^{-3} A^{-2}]$$

$$(10) \text{Gravitational constant} = \cancel{\frac{F r^2}{m_1 m_2}}$$

$$= \frac{\text{kgm}}{\text{s}^2} \times \frac{m^2}{r^2} \times \frac{1}{\rho \text{kgz}} = [M^1 L^2 T^2]$$

$$(11) \text{Heat Energy} = VT$$

$$= \frac{\text{kgm}^2}{As^3} \times A \times s = [M^1 L^2 T^{-2}]$$

Einheitsmaß Flussintensität ist  $\text{A/m}$ .

$$(12) \text{Solar Energy} = [M^1 L^2 T^{-2}]$$

$$(13) \text{Mechanical Energy} = [M^1 L^2 T^{-2}]$$

### ~~Applications of Dimensional Formulae~~

App-1) To check dimensional consistency of given eqn.

$$\text{Eq: } f = m^x n^y$$

$$RHS = \text{kg} \times \left(\frac{\text{m}}{\text{s}}\right)^2$$

$$RHS = [M^1 L^2 T^{-2}]$$

$$\frac{LHS}{T} = F \times v = Fd$$

$$= [M^1 L^2 T^{-2}]$$

$\therefore LHS = RHS$

$$\text{Eq. } \textcircled{1} F = m^2 \frac{v}{r}$$

$$\text{LHS} = F \cdot s \cdot t^{-1} \cdot r^{-1} \\ = [M^1 L^1 T^{-2}]$$

$$\text{RHS} = [M^2 L^0 T^{-1}] \quad \text{Ans. } \textcircled{1} \text{ is dimensionally consistent. (P)}$$

$$\therefore \text{LHS} \neq \text{RHS. } T^2 \cancel{[M]} - \cancel{M^2}$$

Hence, the given equation is dimensionally inconsistent.  $\textcircled{1}$

$$\textcircled{2} T = 2\pi \sqrt{\frac{l}{g}} \times \frac{m}{c}$$

$$\Rightarrow \text{LHS} = [M^0 L^0 T^2]$$

$$\Rightarrow \text{RHS} = [M^0 L^0 T^2] \times \pi, \text{ Ans. } \textcircled{2} \text{ is dimensionally consistent. (P)}$$

$$\therefore \text{LHS} = \text{RHS, and thus eqn is dimensionally consistent.} \quad [S \cdot T \cdot L \cdot M] \text{ - apparent ratio. (SI)}$$

$$\textcircled{3} V = \sqrt{\frac{P}{s}}$$

$$\Rightarrow \text{LHS} = [M^0 L^1 T^{-1}] \quad [S \cdot T \cdot L \cdot M] \text{ - apparent ratio. (SI)}$$

$$\Rightarrow \text{RHS} = \sqrt{\frac{M}{L^2 s^2}} \quad \cancel{\text{App. 1: Dimensional job with phys.}}$$

"If we put  $\frac{M}{L^2 s^2}$  retains dimensionality of (SI)"

$$\Rightarrow \text{RHS} = \sqrt{\frac{M}{L^2 s^2}} = [M^0 L^1 T^{-1}]$$

$$\therefore \text{LHS} = \text{RHS, it is consistent if - 2nd}$$

$$\textcircled{4} F = \frac{1}{2} mv^2 = \frac{1}{2} mu^2$$

$$\text{LHS} = [M^1 L^1 T^{-2}] [L^2] \Rightarrow [L^3] \\ = [M^1 L^2 T^{-2}]$$

$$\text{RHS} = \frac{1}{2} mu^2 \\ = \frac{1}{2} [M^1 L^1 T^{-2}] [L^2] = [M^1 L^2 T^{-2}]$$

$\therefore$  As LHS = RHS, the eqn is dimensionally consistent.

$$\textcircled{5} I = \frac{1}{2} \sqrt{\frac{L}{Mg}}$$

$$\Rightarrow \text{LHS} = I \\ = \frac{1}{2} = [M^0 L^0 T^{-1}]$$

$$\Rightarrow \text{RHS} = \frac{1}{2} \sqrt{\frac{M L^2}{M g}} = \frac{1}{2} \sqrt{L} = [M^0 L^0 T^{-1}]$$

$$\Rightarrow \text{RHS} = \frac{1}{2} \times \sqrt{L} = [M^0 L^0 T^{-1}]$$

As LHS = RHS,

$\therefore$  the equation is dimensionally consistent.  $\textcircled{5}$

# App. 2  $\rightarrow$  To obtain the formula of one physical quantity in terms of other physical quantities.

Eg.  $T \propto$  length of string

$\propto$  mass of bob

$\propto$  acc. due to gravity

$$T \propto \sqrt{L} \quad T \propto M \quad T \propto g$$

$$T \propto l^a m^b g^c$$

$$\Rightarrow T = K l^a m^b g^c \quad [l^{\frac{1}{2}} T^{\frac{1}{2}}] = 2^{1/2}$$

$$\Rightarrow [l^a m^b g^c] = K l^a m^b g^c$$

$$\Rightarrow [M^0 L^0 T^1] = K [M^0 L^1 T^0]^a [M^1 L^0 T^0]^b [M^0 L^1 T^{-2}]^c$$

$$\Rightarrow [M^0 L^0 T^1] = K [M^0 L^2 T^0] [M^b L^0 T^0] [M^0 L^c T^{-2c}]$$

$$\Rightarrow [M^0 L^0 T^1] = K [M^b L^{a+c} T^{-2c}]$$

$$\boxed{T} \quad \boxed{l} = f \quad (1)$$

So, on comparison, we get,

$$b = 0$$

$$a+c = 0 \quad [l^{\frac{1}{2}} T^{\frac{1}{2}}] = \frac{1}{T} \quad \therefore a = -c$$

$$-2c = 1$$

$$c = -\frac{1}{2}$$

$$\rightarrow \boxed{c = -\frac{1}{2}}, \boxed{a = \frac{1}{2}}, \boxed{b = 0}$$

∴ on putting in eqn (1) we get, with no time p. ∴

$$T = K l^a m^b g^c$$

with p. being, with p.

$$\Rightarrow T = 2\pi l^{1/2} g^{1/2} \quad (K = 2\pi)$$

$$\Rightarrow \boxed{T = 2\pi \sqrt{\frac{l}{g}}} \quad \text{factors of } l \text{ and } g \text{ cancel out}$$

Practice Q

Q-1 (1) The velocity of water waves depends on wavelength ( $\lambda$ ); density ( $\rho$ ), and ( $g$ ) acc. due to gravity.

$$\rightarrow v \propto \lambda^a \rho^b g^c$$

$$\Rightarrow v = K \lambda^a \rho^b g^c$$

$$\Rightarrow [v] = [M^0 L^1 T^0] [M^1 L^{-3} T^0] [M^2 L^0 T^2] \quad (\text{For } K=1)$$

$$\Rightarrow [M^0 L^1 T^1] = [M^0 L^0 T^0] [M^b L^{-3b} T^0] [M^0 L^c T^{-2c}]$$

On comparison, we get,

$$a = 0, b = 0$$

$$a - 3b = 1 \quad a - 3b + c = 1 \quad \text{impossible with } a$$

$$-2c = -1$$

$$\Rightarrow \boxed{c = \frac{1}{2}}, \boxed{a = \frac{1}{2}}, \boxed{b = 0}$$

Putting values in eqn,

$$\rightarrow v = \lambda^{\frac{1}{2}} \rho^0 g^{\frac{1}{2}}$$

$$(1 \text{ is red}) \Rightarrow \boxed{v = \sqrt{\lambda g} [l^{\frac{1}{2}}] [T^0] [M^0] [L^0] [T^0]} \quad [l^{\frac{1}{2}}] [T^0] [M^0] [L^0] [T^0]$$

(2) Frequency ( $\gamma$ ) is proportional to

$$\rightarrow \gamma \propto l^a \rho^b T^c$$

$$\Rightarrow \gamma = K l^a \rho^b T^c$$

$$\Rightarrow [M^1 L^{-1} T^0] = [M^0 L^1 T^0] [M^1 L^0 T^0] [M^2 L^0 T^2] \quad (\text{For } K=1)$$

$$\Rightarrow [M^0 L^0 T^0] = [M^0 L^0 T^0] [M^b T^b T^0] [M^c L^c T^{-2c}]$$

On comparison, we get:

$$b+c=0$$

$$a+b+c=0$$

$$-2c=-1$$

$$\therefore \boxed{c = \frac{1}{2}, b = -\frac{1}{2}, a = -\frac{1}{2}}$$

Putting values in eq<sup>n</sup>,

$$\gamma = L^b M^c T^{-b}$$

$$\boxed{\gamma = \frac{1}{2} \sqrt{\frac{T}{M}}}$$

(for circular motion)

$$(3) T = K v^a M^b G^c \quad \therefore a = 0$$

$$\Rightarrow [M^0 L^0 T^1] = [M^0 L^0 T^0] [M^1 L^0 T^0] [M^0 L^3 T^{-2c}] \quad (\text{for } K=1)$$

$$\Rightarrow [M^0 L^0 T^1] = [M^0 L^0 T^0] [M^b T^0 T^0] [M^c L^{3c} T^{-2c}]$$

On comparison, we get:

$$b-c=0$$

$$a+3c=0$$

$$\therefore \boxed{c = \frac{1}{2}, b = \frac{1}{2}, a = -\frac{3}{2}}$$

$$\therefore \boxed{c = \frac{1}{2}, b = \frac{1}{2}, a = -\frac{3}{2}}$$

Putting in eq<sup>n</sup>, we get,

$$T = r^{3/2} M^{1/2} G^{-1/2} \quad \text{or} \quad \boxed{T = r^{3/2} M^{1/2} G^{-1/2}}$$

# 3rd Appli → To convert units of physical quantities from one system to other.

$$\text{Formula} \rightarrow \boxed{\left[ \frac{n_2}{n_1} \right] = \left[ \frac{M_1}{M_2} \right]^a \left[ \frac{L_1}{L_2} \right]^b \left[ \frac{T_1}{T_2} \right]^c}$$

Eg. (Convert P.J to Joules)

$$\text{1 Joule energy} = [M^1 L^2 T^{-2}]$$

$$\rightarrow a=1, b=2, c=-2$$

$$\Rightarrow n_2 = 1 \left[ \frac{1 \text{ kg}}{1 \text{ g}} \right] \left[ \frac{1 \text{ m}}{1 \text{ cm}} \right] \left[ \frac{1 \text{ sec}}{1 \text{ sec}} \right]$$

$$\Rightarrow n_2 = 1 \left[ \frac{1000 \text{ g}}{1 \text{ g}} \right] \left[ \frac{100 \text{ cm}}{1 \text{ cm}} \right] \left[ \frac{1 \text{ m}}{1 \text{ m}} \right] \quad (\text{convert big to small units})$$

$$\Rightarrow n_2 = 1 \times 10^3 \times 10^4 \times 1 \times 10^2$$

$$\Rightarrow n_2 = 10^7$$

$$\therefore \boxed{1 \text{ Joule} = 10^7 \text{ erg}}$$

Q. The value of  $G$  in CGS is  $6.67 \times 10^{-8}$  dyne ( $\text{cm}^2/\text{g}^2$ ). Calculate its value in SI.

$$\Rightarrow n_2 = 6.67 \times 10^{-8} \quad [ \text{If we put } g \text{ in gilbert} ]$$

$$\Rightarrow G = [ \text{M}^1 \text{L}^{-3} \text{T}^{-2} ]$$

$$\Rightarrow n_2 = 6.67 \times 10^{-8} \left[ \frac{1\text{g}}{1\text{kg}} \right]^{-1} \left[ \frac{1\text{cm}}{1\text{m}} \right]^3 \left[ \frac{1\text{s}}{1\text{s}} \right]^{-2}$$

$$\Rightarrow n_2 = 6.67 \times 10^{-8} \left[ \frac{1}{1000} \right] \left[ \frac{1}{100} \right]^3 \left[ 10^8 \right]$$

$$\Rightarrow n_2 = 6.67 \times 10^{-8} \times 10^3 \times 10^6 \times 10^8$$

$$\Rightarrow n_2 = 6.67 \times 10 \Rightarrow n_2 = 6.67 \times 10^{-11} \text{ N m}^2/\text{kg}^2$$

$$\therefore G = 6.67 \text{ N}$$

$$\therefore G = 6.67 \times 10^{-11} \text{ N m}^2/\text{kg}^2$$

Q. Convert 1 N into dyne.

$$\Rightarrow F = [ \text{N}^1 \text{L}^1 \text{T}^{-2} ] \quad a=1, b=1, c=-2$$

$$\Rightarrow n_2 = 1 \times \left[ \frac{1\text{kg}}{1\text{g}} \right] \times \left[ \frac{1\text{m}}{1\text{cm}} \right]^1 \left[ \frac{1\text{sec}}{1\text{sec}} \right]^{-2}$$

$$\Rightarrow n_2 = 1 \times 10^3 \times 10^2$$

$$\therefore n_2 = 10^5$$

$$\text{pure FOI = decimal 1}$$

$$\therefore 1\text{N} = 10^5 \text{ dyne}$$

$$1\text{N} = 10^5 \text{ dyne} \quad 1\text{N} = 10^5 \text{ N}$$

Q. Find the value of  $60 \text{ J/m}$  on a system that has 100g, 100cm and 1min as fundamental units.

$$\Rightarrow 60 \text{ J/m} = \frac{60 \text{ J/s}}{60}$$

$$= [1\text{J}/\text{s}] = [\text{N}^1 \text{L}^2 \text{T}^{-3}]$$

$$n_2 = 1 \times \left[ \frac{100\text{g}}{100} \right] \left[ \frac{100\text{cm}}{100} \right]^2 \left[ \frac{1}{60} \right]$$

$$\Rightarrow n_2 = 1 \times 10 \times 1 \times 60^2$$

$$\Rightarrow n_2 = 216 \times 10^2 \times (10)$$

$$= 216 \times 10^6 \text{ nanotes}$$

tried to find between step 1 & step 2  
out come : true & false  
with help of 10^6 & 10^-3  
of help calculate with 10^-3

analogical quantity

unitless quantity or dimensionless quantity

area requires two unit  
length taken with 10^-3

Dimensionless unit = dimensionless unit

Dimensionless unit  
unitless quantity

u-2

## MOTION IN STRAIGHT LINE

Reference Point or Frame of Reference = A1 T 2d

1) 1D  $\rightarrow$   $x/y/z \rightarrow 1\text{ coord.}$

2) 2D  $\rightarrow$   $xy/yz/xz \rightarrow 2\text{ coord.}$

3) 3D  $\rightarrow$   $xyz \rightarrow 3\text{ coord.}$

### Parameters

$$\text{Distance} / \text{path length} = (\text{SI unit}) \times 1 \times 01 \times 1 = \text{m}$$

Amount of path covered by an object.

SI unit: metres

It is always positive.

It is scalar quantity.

### Displacement / change in position

Shortest path covered from initial to final position.

SI unit: metres

It may be positive or negative or zero.

It is a vector quantity.

### Speed

$$\frac{\text{Distance travelled}}{\text{time}} = \frac{\text{m}}{\text{s}} \text{ or } [\text{m/s}] \text{ SI unit}$$

It is a scalar quantity

It is always positive.

(4)

### Velocity

$$\frac{\text{Displacement}}{\text{Time}} = \frac{\text{m}}{\text{s}} \text{ or } [\text{m/s}] \text{ - unit}$$

It is a vector quantity.

It can be positive, negative or 0.

In straight line motion, distance = displacement  
speed = velocity

### Types of speed

$$\text{Instantaneous Speed} = V_I = \frac{dx}{dt} \quad (\text{at instant of time})$$

(at given interval) Avg. Speed

$$V_{\text{Avg}} = \frac{\text{Total Distance}}{\text{Total Time}}$$

$$V_{\text{Avg}} = \frac{\Delta x}{\Delta t}$$

Note  $\rightarrow V = \frac{dx}{dt}$   $\Rightarrow$  rate of change of position w.r.t. time

### Uniform and Non-Uniform Motion

Uniform Motion  $\rightarrow$  Position changes equally in equal intervals of time.

Non-Uniform Motion  $\rightarrow$  Position changes unequally in equal intervals of time.

Q- A body covers distance  $s = 5t^2 + 3t + 2$  in  $t$  seconds. Calculate the speed at  $t = 1$  second.

$$\Rightarrow v = \frac{ds}{dt} (5t^2 + 3t + 2) \quad [v = \frac{dx}{dt}]$$

$$\Rightarrow v = 10t + 3$$

Time  $\rightarrow$   $t$  in seconds with  $\omega$   
Distance  $\rightarrow$   $s$  in meters, velocity  $v$  in m/s

For  $t = 1$ ,

$$\Rightarrow v = 10 + 3 = 13 \text{ m/s}$$

Initial position and time given  $s$

Q- The position  $s = 10t^4 + 4t^2 + 2t$ , find speed at  $t = 2$  sec.

$$\Rightarrow v = \frac{ds}{dt} (10t^4 + 4t^2 + 2t) \quad [v = \frac{dx}{dt}]$$

Time  $\rightarrow$   $t$  in seconds with  $\omega$   
Distance  $\rightarrow$   $s$  in meters

$$\Rightarrow v = 40t^3 + 8t + 2$$

For  $t = 2$  sec,

$$\Rightarrow v = 40(2)^3 + 8(2) + 2$$

$$\Rightarrow v = 320 + 16 + 2$$

Initial position and time given  $s$

$$Q- s = \frac{4}{5}t^3 - \frac{2}{5}t^2 + 9 \text{ at } t = 1 \text{ sec.}$$

Initial position and time given  $s$

$$\Rightarrow v = \frac{ds}{dt} (\frac{4}{5}t^3 - \frac{2}{5}t^2 + 9) \quad [v = \frac{dx}{dt}]$$

Initial position and time given  $s$

$$\Rightarrow v = 12t^2 - 4 +$$

Initial position and time given  $s$

For  $t = 1$ ,

$$\Rightarrow v = \frac{12}{5} - 4$$

Initial position and time given  $s$

$$\Rightarrow v = \frac{8}{5} \text{ m/sec} = 1.6 \text{ m/s}$$

Q-  $s = 1.5t^2 + 2t$ . Find  $v$  at  $t = 3$  sec.

$$\Rightarrow v = \frac{ds}{dt} (1.5t^2 + 2t)$$

$$= 2 \cdot 3t + 2$$

$$= 9 + 2 = 11 \text{ m/s.} \quad (\text{for } t = 3)$$

Q- The relation between time  $t$  & position  $x$  is  $t = ax^2 + bx$  (abstact). Find out instantaneous velocity.

$$\Rightarrow \cancel{v = \frac{dt}{dx}} = \frac{d}{dx} (ax^2 + bx) \cancel{t = ax^2 + bx}$$

$$= 2ax + b$$

$$\Rightarrow v = \frac{dt}{dx}$$

~~$$\Rightarrow v = \frac{dt}{dx} = \frac{d}{dx} (t) = \frac{d}{dx} (ax^2 + bx)$$~~

~~$$\Rightarrow v = \frac{1}{b} - \frac{2ax}{b}$$~~

~~$$\Rightarrow v = \frac{dt}{dx} = \frac{d}{dx} (ax^2 + bx)$$~~

~~$$\Rightarrow \frac{dx}{dt} = \frac{1}{2ax + b}$$~~

$$\frac{dx}{dt} = \frac{1}{2ax + b}$$

$$\Rightarrow v = \frac{1}{2ax + b}$$

$$\Rightarrow v = (2ax + b)^{-1}$$

Q -  $x = 8.5 + 2.5t^2$ ,  $v$  between  $t=0$  sec to  $t=2$  sec

$$\rightarrow v = \frac{dx}{dt} (8.5 + 2.5t^2) \quad v = \frac{dx}{dt}$$

$$v = \frac{x_2 - x_1}{t_2 - t_1} \quad (1)$$

for  $t = 0$  sec,

$$v = 5 \text{ m/s}$$

$$\text{When } t = 1, x_1 = 8.5 + 8.5 = 17 \text{ m}$$

$$\text{When } t = 2, x_2 = 8.5 + 10 = 18.5 \text{ m}$$


for  $t = 2$  sec,

$$v = 10 \text{ m/s}$$

$$\therefore v = \frac{18.5 - 11}{2} = 3.75 \text{ m/s}$$


## AVERAGE SPEED & VELOCITY

Q - A body travels from A to B at 40 m/s and from B to A at 60 m/s. Calculate the avg. speed & avg. velocity.



$$\text{Avg. speed} = \frac{60 + 40}{2} = 50 \text{ m/s}$$

Avg. velocity = 0, as displacement is zero.



Q - Avg. speed =  $\frac{\text{total distance}}{\text{total time}} = \frac{2AB}{\frac{AB}{V_1} + \frac{AB}{V_2}}$

$$= \frac{2V_1 V_2}{V_1 + V_2} \rightarrow \text{Harmonic Average}$$

$$= \frac{2 \times 40 \times 60}{100} = \frac{4800}{100} = 48 \text{ m/s}$$

\* If the distances are same, then avg. speed is given by harmonic average:

$$v_{\text{avg}} = \frac{2V_1 V_2}{V_1 + V_2}$$

Q - Velocity =  $\frac{\text{displacement}}{\text{time}} = 0 \text{ m/s}$

Q - On a 60 km track, a train travels first 30 km with speed of 30 km/h. How fast must the train travel the next 30 km so avg. speed for the entire trip becomes 40 km/h.

$$\rightarrow \text{Average speed} = \frac{\text{Total distance}}{\text{Total time}}$$

$$\Rightarrow 40 \text{ km/h} = \frac{2V_1 V_2}{V_1 + V_2} \quad [\text{Harmonic Avg.}]$$

$$\Rightarrow 40(30 + x) = 60x \quad (\text{Let } V_2 \text{ be } x \text{ km/h})$$

$$\Rightarrow 1200 + 40x = 60x \quad \cancel{- 40x} \quad \cancel{+ 1200}$$

$$\Rightarrow 1200 = 20x \quad \cancel{- 1200} \quad \cancel{+ 20x}$$

$$\Rightarrow x = 60 \text{ km/h}$$

Q - A train moves with a speed of 80 km/h in the first 15 min, with another speed of 40 km/h in next 15 min. Find average speed.

Date: 19/4/24

$\text{Avg} \rightarrow V_{\text{avg}} = \frac{v_1 + v_2}{2} \dots V_A \text{ [Arithmetic Avg]}$

$\text{if intervals are not equal}$

$30 \text{ km/h} \quad 40 \text{ km/h}$   
 $15 \text{ min} \quad 15 \text{ min}$

$$V_{\text{avg}} = \frac{30+40}{2} = \boxed{35 \text{ km/h}}$$

\* If time intervals are same, we use arithmetic average.

$$V_{\text{avg}} = \frac{v_1 + v_2 + \dots + v_n}{n} \text{ [geometric]}$$

Q- A car travels along a straight line for the first half time with 50 km/h, second half time with 60 km/h. Calculate avg speed.

$$\rightarrow V_{\text{avg}} = \frac{2 \times \text{total distance}}{\text{total time}} = \frac{50+60}{2} = \boxed{55 \text{ km/h}}$$

$$= \frac{2 \times 50 \times 20}{50+60} = \boxed{60 \text{ km/h}}$$

~~50\*60/2~~ ~~total distance~~ ~~total time~~ ~~average speed~~

Q- A body covers  $\frac{1}{3}$  rd of journey with 20 km/h, next  $\frac{1}{3}$  rd with 40 km/h and last  $\frac{1}{3}$  rd of journey with 60 km/h. Avg. speed?

$$\rightarrow V_{\text{avg}} = \frac{3v_1 v_2 v_3}{v_1 + v_2 + v_3} \text{ [od = v0t + v0s]}$$

$$= \frac{3 \times 20 \times 40 \times 60}{20+40+60} = \boxed{32 \text{ km/h}}$$

$$20+40+60 = 20 \times 40 + 40 \times 60 + 60 \times 20$$

$$800 + 2400 + 1200 = \boxed{32 \times 10 \times 60}$$

Date: 19/4/24

$$\Rightarrow V_{\text{avg}} = \frac{360}{11} = \boxed{32.73 \text{ km/h}}$$

## # ACCELERATION

↳ Rate of change of velocity w.r.t. time.

$$a = \frac{\Delta v}{\Delta t}$$

$$\text{then } a = \lim_{\Delta t \rightarrow 0} \frac{\Delta v}{\Delta t}$$

$$\Rightarrow a = \frac{dv}{dt} \rightarrow a = \frac{d}{dt} \left( \frac{dx}{dt} \right)$$

$$= \frac{d^2x}{dt^2} \text{, 2nd order derivative}$$

Q-  $x = 7t^4 + 5t^3 - 3t^2 + 9$ . Find a at  $t = 2$ .

$$v = \frac{dx}{dt} = (28t^3 + 15t^2 - 6t)$$

$$dv = 28t^2 + 15t^2 - 6t$$

$$\frac{d^2x}{dt^2} = (28t^3 + 15t^2 - 6t)$$

$$\Rightarrow a = 84t^2 + 30t - 6$$

For  $t = 2$

$$a = 84 \times 4 + 30(2) - 6 = \boxed{336 \text{ m/s}^2}$$

$$a = 336 + 60 = \boxed{396 \text{ m/s}^2}$$

$$= \boxed{396 \text{ m/s}^2}$$

Integration

Acceleration  $\rightarrow \int dt$   $\rightarrow$  Velocity:  $\rightarrow \int dt$  Position

$$\text{Q- } v = \frac{3}{2}t^3 + 4t^2 + 3.$$

at time  $t=1s$  &  $t=2s$ .

$$\rightarrow \int \frac{3}{2}t^3 dt + \int 4t^2 dt + \int 3 dt$$

$$\Rightarrow dv = \left[ \frac{3}{2}t^4 \right]_0^1 + \left[ 4t^3 \right]_0^1 + \left[ 3t \right]_0^1$$

$$\text{For } t=2 \rightarrow 6 + 32 + 6 = 44$$

$$\text{For } t=1 \rightarrow \frac{3}{2} + 4 + 3 = 11$$

$$\frac{44 - 11}{2} = \frac{33}{2} = 16.5$$

~~$$\Rightarrow \frac{68}{3} - \frac{11}{6} = \frac{125}{6}$$~~
~~$$= \frac{136 - 11}{6} = \frac{125}{6}$$~~
~~$$= \frac{125}{6} \times \frac{1}{6} = \frac{125}{36}$$~~
~~$$= \frac{125}{36} \times 100 = 34.72$$~~

$$\text{Q- } v = 4.2t^2 + 2.1t$$

$$\text{Find } dv \text{ b/w } t=0 \text{ to } t=1$$

$$\Rightarrow dv = 4.2 \int t^2 dt + 2.1 \int t dt$$

$$\Rightarrow dv = \frac{4.2}{3} t^3 + 2.1 t^2$$

$$\text{For } t=0 \rightarrow dv = 0$$

$$\text{For } t=1 \rightarrow dv = 1.4 + 2.1$$

$$dv = \frac{2.8 + 2.1}{2} =$$

~~$$= \frac{3.9}{2} = 1.95$$~~

$$\therefore 16.5 = 2.45m$$

$$\therefore [16.5 = 2.45m]$$

$$\text{Q. } a = 5t^5 + 4t^4 + 3t^3 + 2t + 1$$

$$\Rightarrow dv = 5 \int t^5 dt + 4 \int t^4 dt + 3 \int t^3 dt + 2 \int t dt + \int 1 dt$$

$$\Rightarrow dv = \left[ \frac{5}{6}t^6 \right]_0^1 + \left[ \frac{4}{5}t^5 \right]_0^1 + \left[ \frac{3}{4}t^4 \right]_0^1 + \left[ t^2 \right]_0^1 + \left[ t \right]_0^1$$

$$\text{For } t=1, dv = \frac{5}{6} + \frac{4}{5} + \frac{3}{4} + 1 + 1$$

$$= \frac{50 + 48 + 45 + 120}{60} = \frac{263}{60}$$

~~$$\text{For } t=2, dv = \frac{5}{6} \times 64 + \frac{4}{5} \times 32 + \frac{3}{4} \times 16 + 4 + 2$$~~
~~$$= \frac{5}{6} \times 128 + \frac{4}{5} \times 128 + \frac{3}{4} \times 128 + 18 + 2$$~~
~~$$= \frac{1600 + 768 + 540}{30} = \frac{2908}{30} = \frac{5553}{60}$$~~


$$v = \frac{5553}{60} - \frac{263}{60} = \frac{5816}{60}$$

TS

POSITION - TIME GRAPH

$$\text{Distance} \leftarrow 0 = t \cdot 0$$

$$1 + 1 \cdot t \leftarrow 1 = t \cdot 0$$


Inference - (1) About motion

- At Rest

[App. C - 3c]

(2) Motion

→ 1. Velocity



The slope will give velocity.

- If slope = 0,  $v \rightarrow 0$
- If slope  $> 0$ ,  $v > 0$  (+ve)
- If slope  $< 0$ ,  $v < 0$  (-ve)






Eg. (1). Slope  $\rightarrow 0$ ,  $v \rightarrow 0$ 

$$1 + 1 \cdot t \leftarrow 1 = t \cdot 0$$

$$\text{Ex. } \frac{1}{2} \cdot 2t + 1 = t \cdot 0$$

(2) Slope  $\rightarrow +ve$ ,  $v \rightarrow +ve$ 

$$S + P + vt \times \frac{1}{2} + v \cdot t \times \frac{1}{2} + r \cdot t \times \frac{1}{2} = vt, S = t \cdot 0$$

(3) Slope  $\rightarrow -ve$ ,  $v \rightarrow -ve$ (4) Slope  $\rightarrow$  constant & +ve,  $v \rightarrow$  constant & +ve(5) Slope  $\rightarrow$  constant & -ve,  $v \rightarrow$  constant & -ve(6) Slope  $\rightarrow$  variable,  $v \rightarrow$  variablenon-uniform motion no constant time  $\Delta t$ 2. Slope  $\rightarrow$  initial  $\Rightarrow$ 1. distance - time  $\frac{\text{slope}}{\text{time}}$  speed2. Displacement - time  $\frac{\text{slope}}{\text{time}}$   $\rightarrow$  Velocity0 eqn,  $s = \frac{1}{2}at^2$ ,  $s = ut + \frac{1}{2}at^2$ 3. Velocity - time  $\frac{\text{slope}}{\text{time}}$  acceleration












(3) Types of motion no uniform motion


Uniform motion in straight line	TYPES	Non-Uniform motion
motion form - equal $\Delta t$ & $\Delta s$	1. If line is straight	2. If line is curved
and $a = 0$	→ Velocity - Constant → Slope - Constant → $a = 0$	→ Velocity - Variable → Slope - Variable → $a = \text{Variable}$ (depends on $x$ )

## (4) Acceleration

→ Acceleration occurs only in non-uniform motion.

- Slope - Variable, velocity - variable.





(1) B has a higher velocity as it has more displacement in less time, i.e., its slope is steeper than the slope of A.

$$(2) \frac{v_A}{v_B} = \frac{\tan 60^\circ}{\tan 30^\circ} = \frac{\sqrt{3}}{1}$$

$$\text{Distance} = \frac{1}{2} \times \text{length of hypotenuse} \times \text{parallel side} = \frac{1}{2} \times \sqrt{3} \times \sqrt{3} = \frac{3}{2}$$

$$\therefore v_A : v_B = 1 : 3$$

∴ motion is uniform.

∴ motion is uniform.

∴ motion is uniform.


Q. A person walking on a road takes 5 steps forward and 3 steps backward and plots the position-time graph.

∴ motion is uniform.


For uniform motion, distance  $\propto t$   
displacement  $\propto t$ ,  $a=0$ .  $\therefore$  motion is uniform.

∴ motion is uniform.

∴ Not possible as time can't be negative.



→ Does not exist as distance can't decrease.



→ (1) indicates uniform motion.

But  $s = ut + \frac{1}{2}at^2$ ,  $a \neq 0$ ,  $t = \text{constant}$  → Non-uniform motion.

∴ It is not possible to have two diff. positions for same time.  
∴ This doesn't exist.

∴ uniform motion,  $s = ut + \frac{1}{2}at^2$ ,  $a = 0$ ,  $t = \text{constant}$

∴ uniform motion,  $s = ut + \frac{1}{2}at^2$ ,  $a \neq 0$ ,  $t = \text{constant}$

∴ uniform motion,  $s = ut + \frac{1}{2}at^2$ ,  $a = 0$ ,  $t = \text{constant}$

∴ uniform motion,  $s = ut + \frac{1}{2}at^2$ ,  $a \neq 0$ ,  $t = \text{constant}$

∴ uniform motion,  $s = ut + \frac{1}{2}at^2$ ,  $a = 0$ ,  $t = \text{constant}$

∴ uniform motion,  $s = ut + \frac{1}{2}at^2$ ,  $a \neq 0$ ,  $t = \text{constant}$

∴ uniform motion,  $s = ut + \frac{1}{2}at^2$ ,  $a = 0$ ,  $t = \text{constant}$

∴ uniform motion,  $s = ut + \frac{1}{2}at^2$ ,  $a \neq 0$ ,  $t = \text{constant}$

∴ uniform motion,  $s = ut + \frac{1}{2}at^2$ ,  $a = 0$ ,  $t = \text{constant}$

∴ uniform motion,  $s = ut + \frac{1}{2}at^2$ ,  $a \neq 0$ ,  $t = \text{constant}$

∴ uniform motion,  $s = ut + \frac{1}{2}at^2$ ,  $a = 0$ ,  $t = \text{constant}$

∴ uniform motion,  $s = ut + \frac{1}{2}at^2$ ,  $a \neq 0$ ,  $t = \text{constant}$

∴ uniform motion,  $s = ut + \frac{1}{2}at^2$ ,  $a = 0$ ,  $t = \text{constant}$

∴ uniform motion,  $s = ut + \frac{1}{2}at^2$ ,  $a \neq 0$ ,  $t = \text{constant}$

∴ uniform motion,  $s = ut + \frac{1}{2}at^2$ ,  $a = 0$ ,  $t = \text{constant}$

∴ uniform motion,  $s = ut + \frac{1}{2}at^2$ ,  $a \neq 0$ ,  $t = \text{constant}$

∴ uniform motion,  $s = ut + \frac{1}{2}at^2$ ,  $a = 0$ ,  $t = \text{constant}$

∴ uniform motion,  $s = ut + \frac{1}{2}at^2$ ,  $a \neq 0$ ,  $t = \text{constant}$


→ Body is in motion

→  $v_{av} = 0$ , as displacement = 0

→  $a > 0$ ; then  $a = 0$

→ Non-uniform motion

Distance =  $t \times v + \text{const}$



### \* Velocity-time Graph

Distances  $\rightarrow$ \* Slope of  $v-t$  graph  $\rightarrow$  acceleration

$$\text{Slope} = 0, a = 0$$

$$\text{Slope} = +\text{ve}, a = +\text{ve}$$

$$\text{Slope} = -\text{ve}, a = -\text{ve}$$

$$\text{Slope} = \text{const.}, a = \text{const}$$

 $\hookrightarrow v \rightarrow \text{constant}$ 

$$\text{Slope} = 0, a = 0$$

uniform motion

(2)

 $\rightarrow v \rightarrow$  constant (increasing)  
slope - constant,  $a = \text{constant, +ve}$ 

Horizontal motion  $\rightarrow$  uniform motion. E.g. increasing speed  
with time of motorcycle.

(3)

 $\rightarrow v \rightarrow$  uniform (decreasing)

$\rightarrow$  slope  $\rightarrow$  negative,  $a = \text{negative const}$   
retarded uniform motion E.g. decreasing speed of motorcycle

motorcycle placed

$$a = \text{constant, } a = \text{const}$$

2. Area under  $v-t$  graph  $\rightarrow$  Displacement

motorcycle placed

$$\text{Area} \rightarrow v \times t = \text{displacement}$$

1)

✓

$\left. \begin{array}{l} \text{Slope} = -\text{ve}, v \rightarrow \downarrow, \text{constant} \\ a < 0 \& a \rightarrow \text{constant} \end{array} \right\} \rightarrow v \uparrow \text{in opposite direction.}$


$a > 0 \& a \rightarrow \text{constant}$  Eg. Throwing ball

$\left. \begin{array}{l} \text{Speed} \\ \text{of} \\ \text{ball} \end{array} \right\} \rightarrow \text{Speed of ball}$

$\rightarrow$  First decelerated ( $a < 0$ ) then accelerated ( $a > 0$ ) motion

$\left. \begin{array}{l} \text{Time} \\ \text{of} \\ \text{fall} \end{array} \right\} \rightarrow$

A ball is dropped from a certain height and rebounds from the floor with reduced speed. It goes on hitting the ground and ball comes to rest. Plot velocity-time graph




$\rightarrow$   $(0.5 \times 0.5 \times 1) + (0.5 \times 2.5 \times 0.5) + (0.5 \times 2.0 \times 1) = 2.5$

$\rightarrow$   $2 + 2 + 2 \times 2 = 6$

$\rightarrow$   $(0.5 \times 0.5 \times 1) + (0.5 \times 2.5 \times 0.5) + (0.5 \times 2.0 \times 1) = 2.5$


$\rightarrow$   $2 + 2 + 2 \times 2 = 6$



without graph  $v = \text{const}$

$$\text{Distance} = \text{Area under graph}$$


$$= \frac{1}{2} \times 10 \times 12 = 60 \text{ m}$$



$\Rightarrow$  Area =  $\frac{1}{2} \times (1 \times 5 \times 50) + (\frac{1}{2} \times 3 \times 50) + 50$

$$= 125 + 75 + 100 \leftarrow$$

$$= [300 \text{ m}]$$



$$\rightarrow \text{Area} = \left( \frac{1}{2} \times 30 \times 0.5 \right) + (0.25 \times 20) + \frac{1}{2} \left( 0.25 \times 15 \right)$$

$$+ (0.25 \times 20) + \frac{1}{2} \times 50 \times 2$$

$$= 50 + 5 + 3.75 + 5 + 50$$


$$= [137.5 \text{ m}]$$

$$\rightarrow \text{Area} = \left( \frac{1}{2} \times 0.5 \times 20 \right) + (0.25 \times 20) + \left( \frac{1}{2} \times 0.25 \times 30 \right)$$

$$+ (20 \times 0.25) + \frac{1}{2} \times 50$$

$$= 5 + 5 + 3.75 + 5 + 25 \leftarrow [48.75 \text{ m}]$$

## \* Kinematics Equation of Motion for Uniform Acceleration



$$(1) \text{ Slope of AB} = \tan \theta = \frac{BC}{AC}$$

$$a = \frac{v-u}{t} \quad (1)$$

$$\Rightarrow [v = u + at] = \text{Velocity time Equation}$$

(2) Area under OABC (Trapezium)

$\Rightarrow$  Area of  $\triangle ABC +$  Area of  $\square ACDO$ .

$$S = \frac{1}{2} \times AC \times BC + (AC \times CD)$$

$$S = \frac{1}{2} \times t \times (v-u) + (t \times u)$$

$$S = ut + \frac{1}{2} at^2 \quad (\text{From } (1), v-u = at)$$

(Position-time relation)

$\frac{D}{S} = \text{const}$  for uniform acc only

$$(3) S = ut + \frac{1}{2}(v-u)t + (1 - \frac{ut}{v-u}) \quad (\text{from } (2))$$

$$\Rightarrow S = \frac{1}{2} ut + \frac{1}{2} vt \quad (\text{const} = \frac{1}{2})$$

$$\Rightarrow S = \frac{1}{2} t(v+u) = \frac{1}{2} vt \quad (v = \text{avg. velocity})$$



$$\Rightarrow 25 \times 10^{-6} + \frac{1}{2} (10)^2 \times 25 \times 10^{18}$$

for point effect

$$\Rightarrow 25 \times 10^{-6} + \frac{1}{2} \times 25 \times 10^{-6}$$

$$\Rightarrow s = 25 \times 10^{-6} \left( 1 + \frac{1}{2} \right)$$

$$\boxed{s = 37.5 \times 10^{-6} \text{ m}}$$

Q- A driver takes 0.20 sec to apply brakes after he sees a need for it. The driving speed of car is 54 km/h, & the deceleration is 6 m/s<sup>2</sup> ( $a = -6$ ), find distance travelled by car.

$$\rightarrow t = 0.20$$

$$v = 54 \text{ km/h} = 15 \text{ m/s}$$

$$s = ?$$

$$a = -6 \text{ m/s}^2$$

$$s = ut + \frac{1}{2} at^2$$

$$\Rightarrow s = 15(0.20) + \frac{1}{2} (-6)(0.20)^2$$

$$\cancel{\Rightarrow s = 3 + \frac{1}{2}(-3)(0.04)}$$

$$\cancel{\Rightarrow s = 3 - 0.12}$$

$$\Rightarrow s = 2.88 \text{ m}$$

$$\rightarrow \text{Dist. travelled before breaking} = 0.20 \times 15$$

$$\boxed{13 \text{ m}}$$

$$\text{Dist. travelled after breaking} = BC$$

(Using  $s = ut + \frac{1}{2} at^2$ )

$$= 13 + \frac{1}{2} \times 6 \times 4^2 = 25.2 \text{ m}$$

$$\Rightarrow BC \rightarrow 0 - (15)^2 = 2(-6)s$$

$$\frac{-225}{-12} = s$$

$$\boxed{s = 18.75 \text{ m}}$$

$$\rightarrow \text{Total dist. travelled} = 18.75 + 3$$

$$= \boxed{21.75 \text{ m}}$$


Q- A body covers 12 m in 2nd second and 20 m in 4th second. How much distance will it cover in 4 seconds after 5th second.

$$\rightarrow s_2 = 12 \text{ m}$$

$$s_4 = 20 \text{ m}$$

$$s_5 \rightarrow s_2 \text{ m}$$

$$s_9 \rightarrow s_4 \text{ m}$$



$$s_2 = u + \frac{a}{2} (2n-1) \quad \left\{ s_4 = u + \frac{a}{2} (2n-1) \right.$$

$$\Rightarrow 12 = u + \frac{a}{2} (2(2)-1)$$

$$\Rightarrow 12 = u + \frac{3a}{2} \quad \left. \begin{array}{l} \cancel{12 = u + \frac{7a}{2}} \\ \cancel{12 = u + 7a} \end{array} \right\} \quad \textcircled{1}$$

$$\Rightarrow 20 = u + \frac{7a}{2} \quad \textcircled{2}$$

$$\frac{7a}{2} + u = 20$$

$$\frac{3a}{2} = -12 \quad \left( \begin{array}{l} \cancel{12 = u + 7a} \\ \cancel{12 = u + \frac{7a}{2}} \end{array} \right)$$

$$4a = 8$$

$$a = 4 \text{ m/s}^2$$

→ Substituting in  $\textcircled{1}$ ,

$$12 = u + \frac{3a}{2}$$

$$12 = u + \frac{3 \times 4}{2}$$

$$12 = u + 6$$

$$\boxed{u = 6 \text{ m/s}}$$

$$\therefore s_1 = ut + \frac{1}{2} at^2$$

$$\therefore s_1 = u(t) + \frac{1}{2} t^2 (a)$$

$$\therefore s_1 = 54 + 16t^2 = [216 \text{ m}]$$

$$\therefore s_2 = u(s) + \frac{1}{2} \times t^2 (s)^2 [216]$$

$$s_2 = 30t + 50t^2 \text{ m} \quad \text{bus is at 1 sec interval}$$

$$= 80 \text{ m}$$

$$\Rightarrow \Delta x = 216 - 80 = 2$$

$$= [136 \text{ m}]$$

Q - A bus starts from rest moving with constant acceleration  $5 \text{ m/s}^2$ . At some time, a car travels with constant velocity  $50 \text{ m/s}$  overtakes & passes the bus. (i) Find at what distance will the bus overtake the car.

(ii) How fast the bus will be travelling?

$$\rightarrow \begin{array}{l} 0 \text{ m/s} \\ 5 \text{ m/s}^2 \\ 0 \end{array} \quad \begin{array}{l} 50 \text{ m/s} \\ 0 \end{array} \quad \begin{array}{l} (1 \text{ sec}) \\ (1 \text{ sec}) \end{array} \quad \begin{array}{l} s_1 = ut + \frac{1}{2} at^2 \\ s_2 = ut \end{array}$$

$$\text{Dist. travelled by bus} = ut + \frac{1}{2} at^2$$

$$s_B = 0 + \frac{1}{2} (5) (t)^2$$

$$\boxed{s_B = \frac{5t^2}{2}}$$

$$\begin{aligned} \text{Dist. travelled by car} &= 50 \times t \\ &= \underline{\underline{50t}} \\ &= 1000 \text{ m} \end{aligned}$$

Now, the dist travelled by car & bus to overtaking is equal:

$$\therefore 50t = \frac{5t^2}{2}$$

$$\Rightarrow \frac{100}{5} = 5t$$

$$\Rightarrow \boxed{t = 20 \text{ s}}$$


- Q2 V vs t graph

$$v = u + at$$

$$v = 0 + 5 \times 20$$

$$\boxed{v = 100 \text{ m/s}}$$

Q - An object is moving with uniform acceleration. Its velocity after 5 sec is  $25 \text{ m/s}$  & after 8 sec is  $34 \text{ m/s}$ . Find out dist. travelled in 12 th second.



$$\Rightarrow v_1 = u + at$$

$$25 = u + a(5) \quad \text{--- (1)}$$

$$\Rightarrow u + 5a = 25$$

$$v_2 = u + at$$

$$34 = u + a(8) \quad \text{--- (2)}$$

$$\Rightarrow u + 8a = 34$$

$$\therefore u + 5a = 25 \quad \text{--- (1)}$$

$$\therefore u + 8a = 34 \quad \text{--- (2)}$$

$$\therefore 3a = 9$$

$$\therefore a = 3 \text{ m/s}^2$$

$$\therefore u = 10 \text{ m/s}$$

$$\therefore s_{12} = u + \frac{a}{2} (2(12) - 1)$$

$$\therefore s_{12} = 10 + \frac{3}{2} (23)$$

$$\therefore s_{12} = 10 + \frac{3}{2} (23)$$

## Motion Under Gravity

$$1) v = u + gt \quad (a = g)$$

$$2) s = ut + \frac{1}{2}gt^2$$

$$3) v^2 - u^2 = 2gs$$

Note → In upward motion,  $g \rightarrow$  negative,  $w = a$   
 In downward motion,  $g \rightarrow$  positive,  $w = a$

Q- A ball is thrown vertically upwards with velocity  $20\text{ m/s}$  from top of a building of height  $25\text{ m}$ . How high the ball will rise?

$$\begin{aligned} \Rightarrow v^2 - u^2 &= 2as \\ \Rightarrow 0 - 400 &= 2 \times 9.8 \times h \\ \Rightarrow h &= \frac{400}{2 \times 9.8} = 20.4 \text{ m} \end{aligned}$$

$$\Rightarrow h = 20\text{ m}$$

$$\begin{aligned} \text{Total dist. from ground} &= 20 + 25 = 45\text{ m} \\ \Rightarrow s &= 45\text{ m} \end{aligned}$$

→ How long will it take for ball to hit the ground.

$$\begin{aligned} \Rightarrow s &= ut + \frac{1}{2}at^2 \\ \Rightarrow 45 &= 0 + \frac{1}{2} \times 9.8 \times t^2 \\ \Rightarrow t^2 &= \frac{45}{9.8} = ? \\ \Rightarrow t &= \sqrt{3.57} = 1.88 \text{ s} \\ \Rightarrow t_1 &= 1.88 \text{ s} \\ \Rightarrow t_2 &= 1.88 + 1.88 = 3.76 \text{ s} \end{aligned}$$

$$(1 - 1.88) \approx 1.0 = 1.2 \text{ s}$$

Q- A ball thrown up, caught by the thrower after  $4\text{ s}$ . How high did it go? What is initial velocity?

→ Time for upward motion = Time for downward motion

$$\Rightarrow \text{Total time taken} = 4\text{ s}$$

$$\Rightarrow t_d = 2\text{ s} \quad 2 + t_d = 4\text{ s}$$

### Upward motion

$$u = 0 \quad v = 0$$

$$g = 10\text{ m/s}^2 \quad t = 2\text{ s}$$

$$v^2 - u^2 = 2as$$

$$\Rightarrow -400 = -20s$$

$$\Rightarrow s = 20\text{ m}$$

$$\Rightarrow v = u + at$$

$$\Rightarrow 0 = 0 + 10t$$

$$\Rightarrow t = 2\text{ s}$$

③ How far was it below the highest point at  $3\text{ sec}$  after it was thrown.

$$\Rightarrow s = ut + \frac{1}{2}gt^2$$

$$\Rightarrow s = 0 + \frac{1}{2} \times 10 \times 1$$

$$\Rightarrow s = 5\text{ m}$$

Q. A balloon is moving up at a rate of  $9.8 \text{ m/s}$ . At a height of  $39.2$  above the ground a food packet is thrown downwards. After how much time and with what velocity does it reach to the ground?

$$\rightarrow s = ut + \frac{1}{2}at^2$$

$$\rightarrow s = 9.8t + \frac{1}{2} \times 9.8t^2$$

$$\rightarrow 39.2 = 9.8t + 4.9t^2$$

$$\rightarrow t^2 + 2t - 8 = 0$$

$$\Rightarrow t^2 + 4t - 2t - 8 = 0 \Rightarrow t^2 - 2t + 8 = 0$$

$$\Rightarrow t(t+4) - (t+4) = 0 \Rightarrow t^2 - 4t$$

$$\Rightarrow (t-4)(t+4) = 0$$

$$\rightarrow -39.2 = 9.8t + 4.9t^2$$

$$\Rightarrow -8 = 2t - t^2$$

$$\Rightarrow t^2 - 2t - 8 = 0$$

$$\Rightarrow t^2 + 4t - 2t - 8 = 0 \text{ (cancel out common terms, ref next)} \quad (8)$$

$$\Rightarrow t(t-4) + 2(t-4) = 0$$

$$\Rightarrow (t-4)(t+2) = 0$$


As time can't be negative,  $t = 4 \text{ s}$ .

$$\rightarrow v = u + at$$

$$\Rightarrow v = +9.8 + 9.8(4)$$

$$\Rightarrow v = +9.8 + 39.2$$

$$\Rightarrow v = +29.4 \text{ m/s}$$



Q. A food packet is released from a helicopter which is moving upwards with  $2 \text{ m/s}$ . After  $2 \text{ s}$ , what is velocity of packet? How far is it below the helicopter?

$$t = 2 \text{ s}, u = 2 \text{ m/s}, g = -10 \text{ m/s}^2$$

$$s = ?$$

$$\rightarrow v = u + gt$$

$$\Rightarrow v = 2 + (-10) \cdot 2$$

$$\boxed{v = -18 \text{ m/s}}$$

$$\rightarrow v^2 - u^2 = 2as \quad s = ut + \frac{1}{2}gt^2$$

$$\Rightarrow s = 2(2) + \frac{1}{2}(-10) \times 2^2$$

$$2) s = 4 - 16 = \boxed{-12 \text{ m}}$$


(Neg. as direction is opposite)

→ Dist covered by helo in  $2 \text{ sec} \rightarrow 2 \times 2$

$$= \boxed{4 \text{ m}}$$

$$\therefore \text{Total distance} = 4 + 16 = \boxed{20 \text{ m}} \quad (\text{As direction is opposite})$$

Q. Two balls are thrown simultaneously. Ball A upward with  $20 \text{ m/s}$ , ball B downward  $20 \text{ m/s}$  at a height of  $40 \text{ m}$  with the same line. At what point they both collide?  $v = 20 \text{ m/s}$



Balla failed example because  
it's not a complete sentence.

$$\rightarrow s = ut + \frac{1}{2}gt^2$$

$$\Rightarrow x = 20t + \frac{1}{2}(-10)t^2 \quad |_{t=0} = 0 \quad |_{t=2} = 20 \times 2 - 5 \times 2^2 = 40 - 20 = 20$$

$$\Rightarrow u = 2at - 5t^2 \quad - (1)$$

Ball B

$$\rightarrow s = ut + \frac{1}{2}gt^2$$

$$\Rightarrow 40 - n = 20t + \frac{1}{2}(10)t^2$$

$$\Rightarrow y_0 - n = 20t + 5t^2$$

$$m = -5t^2 - 20t - 40$$

Subtracting ② from ①,

$$\Rightarrow y_0 - x_0 = 20 + 5t^2 \quad \text{From (1),}$$

$\rightarrow 40 \cdot (20 + 50 + 2) = 1200 + 200 + 40 = 1440$

$$\Rightarrow 46 = 20t + 3t^2 + 5t^3 \quad \text{Solving for } t \text{ gives } t = 1$$

$$\Rightarrow t = 15 \text{ min}$$

Sub. value of  $t$  in (1)

$$n = 26 - 5$$
$$= \boxed{15m}$$

- d - A rocket is moving upward with  $a = 10 \text{ m/s}^2$ . The fuel finish in 1 minute and it continues to up. What is the max height?

$$\rightarrow \underline{\text{Dist OA}} \quad a = 10 \text{ m/s}^2 \quad v = 0 \\ t = 6 \text{ s}$$

$$\Rightarrow s = ut + \frac{1}{2}gt^2$$

$$25 \neq 1 \times \frac{5}{2} \times 3600$$

$$\Rightarrow s = 18000 \text{ m} \approx \underline{\underline{18 \text{ km}}}$$

$$\rightarrow v = u + at$$

$$\Rightarrow v = 10 \times 60 = \underline{\underline{600 \text{ m/s}}}$$

$$\text{Dist AB} \quad a = -1 \text{ m/s}^2 \quad u = 600 \text{ m/s}$$

$$\rightarrow v^2 - u^2 = 2as$$

$$\Rightarrow 360000 = 2x - 10x + 18000$$


$$\Rightarrow -18000 = s$$

$$\text{Total dist.} \rightarrow 18000 + 18000 = \underline{\underline{36000 \text{ km}}}$$


(Ch-3)  $\rightarrow$  MOTION IN A PLANE (minimum two dimensions)

Position Vector ( $\vec{r}$ )

→ Denoted by  $\vec{r}$



Displacement Vector



Vector symbol ( $\rightarrow$ )

A or  $\vec{A} \rightarrow$  bold


$$\rightarrow \vec{r} = \vec{r}_2 - \vec{r}_1$$

### # Key Terms

(1) Equal vectors: Vectors having same magnitude and same direction are known as equal vectors.

$$\vec{A} \quad \vec{B} \quad \rightarrow \vec{A} = \vec{B}$$

(2) Parallel Vectors: Vectors having same direction but magnitude need not be same are called parallel vectors.



(3) Negative of vectors: Opposite vector of a given vector.

$$\vec{A} \quad \vec{B} \quad \rightarrow \vec{B} = -\vec{A}$$

(4) Modulus of vector:  $|\vec{A}|$  or  $|A|$ . → A real quantity To obtain magnitude of vector.

(5) Unit vector: Vector which has magnitude = 1 (unity)

$$\text{Cap } \hat{A} = \frac{\vec{A}}{|\vec{A}|}$$

'x' unit vector -  $\hat{i}$

'y' unit vector -  $\hat{j}$

'z' unit vector -  $\hat{k}$

} To indicate direction of vector.

(6) Null vector:  $\vec{0}$ , vector having zero magnitude.

Ex (1) At rest,  $\vec{v} = \vec{0}$ .

(2) At origin,  $\vec{r} = \vec{0}$

(3) Uniform velocity,  $\vec{a} = \vec{0}$

(4)  $\vec{A} \& \vec{B} = -\vec{A}$

$$\vec{A} + \vec{B} = \vec{0}$$

(7) Multiplication of vector by real number:

$$\vec{A} \times 2 = \vec{2A}$$


~~Non-commutative |  $\vec{A}$  will give result for  $2\vec{A}$  and  $\vec{A}$  for  $\vec{A} \times 2$~~

~~(magnitude & direction), (sign) 2 is 1 with only sign~~

~~(vector & direction)~~


## \* Triangle Law of Vector Addition

### 1) Diagram / Pictorial form




### 2) Subtraction form


$$\vec{A} - \vec{B} = \vec{A} + (-\vec{B})$$



## \* Parallelogram Law of Vector Addition



\* Vector Addition of Two Vectors using triangle / parallelogram law of Vector Addition (Sine & Cosine), (Magnitude & Direction of resultant vector).



$$\text{For } \triangle PQR, \rightarrow \sin \theta = \frac{PQ}{PR}$$

$$\Rightarrow PQ \sin \theta = PR$$

$$\Rightarrow PR = \underline{PQ \sin \theta} \quad \text{--- (1)}$$

$$\Rightarrow \cos \theta = \frac{SR}{PR}$$

$$\Rightarrow SR = \underline{PQ \cos \theta} \quad \text{--- (2)}$$

$$\begin{aligned} \text{In } \triangle PQR, \quad R^2 &= PR^2 + (PQ)^2 \\ &= PR^2 + (PS+SR)^2 \\ \Rightarrow R^2 &= \underline{PQ^2 + (B \sin \theta)^2 + (A + B \cos \theta)^2}. \end{aligned}$$

$$\Rightarrow R^2 = B^2 \sin^2 \theta + A^2 + B^2 \cos^2 \theta + 2AB \cos \theta$$

~~$$\text{magnitude} = \sqrt{R^2} = \sqrt{A^2 + B^2 + 2AB \cos \theta}$$~~

\*  $\theta$  is angle b/w  $\vec{A}$  &  $\vec{B}$

\* ~~Angle direction~~  $\rightarrow \tan \theta = \frac{|B| \sin \theta}{|A| + |B| \cos \theta}$

- Q. Two forces  $5N$  &  $7N$  act at an angle of  $60^\circ$  b/w them.  
Find resultant force.

$$\vec{A} = 5N \quad \vec{B} = 7N$$

$$\begin{aligned}\vec{R}^2 &= \sqrt{25+49+70\cos60^\circ} \\ &= \sqrt{74 + 70 \times \frac{1}{2}} \\ &= \sqrt{109} \quad N = 10.4N \text{ approx}\end{aligned}$$

$$\begin{aligned}\text{Direction} &= \tan^{-1} \frac{7}{5} \\ &= \frac{7 \times \sqrt{3}}{5} \\ &= \frac{7 \times \frac{\sqrt{3}}{2}}{5} \\ &= \frac{7\sqrt{3}}{10} \times \frac{10}{17} = \frac{7\sqrt{3}}{17} = \frac{120.84}{17}\end{aligned}$$

$$= \frac{7\sqrt{3}}{17} = \tan^{-1} \left( \frac{7\sqrt{3}}{17} \right)$$


- Q. A body has two velocities  $30\text{ m/s}$  due East &  $40\text{ m/s}$  due North. Find resultant velocity.

$$\vec{R}^2 = \sqrt{1600+900}$$

$$\vec{R} = 50\text{ m/s}$$

$$\tan \beta = \frac{v_2 \sin 90^\circ}{v_1 + v_2 \cos 90^\circ}$$

$$\beta = \tan^{-1} \frac{40}{30} = 53.8^\circ$$



- Q. Two equal forces have their resultant equal to either. What is the angle between them?

$$\rightarrow \vec{R} = \sqrt{\vec{P}^2 + \vec{Q}^2 + 2PQ \cos 0^\circ}$$

Imp

$$\cos^{-1} \left( \frac{1}{2} \right) = 120^\circ$$

As all three forces are equal,

$$\vec{R}^2 = \sqrt{2\vec{R}^2 + 2\vec{R}^2 \cos 0^\circ}$$

$$\rightarrow \vec{R}^2 = 2\vec{R}^2 + 2\vec{R}^2 \cos 0^\circ$$

$$\rightarrow -\vec{R}^2 = 2\vec{R}^2 \cos 0^\circ$$

$$\rightarrow \cos 0^\circ = -\frac{1}{2}$$

$$\rightarrow \theta = \cos^{-1} \left( -\frac{1}{2} \right)$$

$$= 120^\circ$$

- Q. A boat moving towards north with  $25\text{ km/hr}$  & water flowing with  $10\text{ km/hr}$  in direction of  $60^\circ$  east of south. Find resultant velocity.

$$\rightarrow \vec{R} = \sqrt{\vec{v}_1^2 + \vec{v}_2^2 + 2v_1 v_2 \cos 0^\circ} \quad (\theta = 120^\circ)$$

$$\rightarrow \vec{R}^2 = \sqrt{625 + 100 + 500 \cos 120^\circ}$$


$$\rightarrow \vec{R} = \sqrt{725 + 500 \times -\frac{1}{2}}$$

$$\rightarrow \vec{R} = \sqrt{475} = [21.8 \text{ km/hr}]$$

$$\tan \beta = \frac{v_2 \sin \theta}{v_1 + v_2 \cos \theta}$$

$$\tan \beta = \frac{(\tan 1)}{\frac{10\sqrt{3}}{25+10\sqrt{3}} \times \frac{1}{2}} = \frac{\tan^{-1}(\sqrt{3})}{4}$$

Q - Rain was falling vertically with 35 m/s and wind is blowing in east to west with 12 m/s. In which direction should a boy hold his umbrella.



$$\rightarrow \vec{R} = \sqrt{\vec{v}_1^2 + \vec{v}_2^2 + 2v_1 v_2 \cos \theta}$$

$$= \sqrt{144 + 1225 + 840 \times 0}$$

$$= \sqrt{144 + 1225} = \sqrt{1369}$$

$$= [37.1 \text{ m}]$$

$\tan \beta = \frac{v_2 \sin 90^\circ}{v_1 + v_2 \cos 90^\circ}$ , then convert primum toad A - ?

$v_1 + v_2 \cos 90^\circ$  for wait with in all and all the

$$= \frac{35}{12} = \left| \tan^{-1} \left( \frac{35}{12} \right) \right|$$


~~$$35^2 + 12^2 = 50^2$$~~

~~$$35^2 + 12^2 = 50^2$$~~

~~$$35^2 + 12^2 = 50^2$$~~

Q - A force is at  $45^\circ$  to the horizontal component. Its component in horizontal direction is 50 N.

Find component in vertical direction.



$$\rightarrow f_x = f \cos \theta$$

$$\Rightarrow f = \frac{f_x}{\cos \theta}$$

$$\Rightarrow f = \frac{50}{\frac{1}{\sqrt{2}}} = [50\sqrt{2}]$$

$$f_y = f \sin \theta$$

~~$$\Rightarrow f_y = 50\sqrt{2} \times \frac{1}{\sqrt{2}} = [50 \text{ N}]$$~~

x — X — x — x — x

$$\rightarrow \vec{A} = 3\hat{i} + 4\hat{j} + 2\hat{k} \rightarrow \text{magnitude of } A$$

$$\vec{B} = \hat{i} + 2\hat{j} + 5\hat{k}$$

$$\vec{A} + \vec{B} = 4\hat{i} + 6\hat{j} + 7\hat{k}$$

$$\text{Magnitude of } \vec{A} = \sqrt{3^2 + 4^2 + 2^2}$$

$$= \sqrt{9+16+4} = \sqrt{29}$$

$$|\vec{B}| = \sqrt{1^2 + 2^2 + 5^2}$$

$$= \sqrt{30}$$

$$|\vec{A} + \vec{B}| = \sqrt{4^2 + 6^2 + 7^2} = \sqrt{101}$$

in rectangular direction  
in one direction

$$\hat{A} = \frac{\vec{A}}{|\vec{A}|}$$

$$= \frac{3\hat{i} + 4\hat{j} + 2\hat{k}}{\sqrt{29}}$$

$$\hat{B} = \frac{\vec{B}}{|\vec{B}|} = \frac{\hat{i} + 2\hat{j} + 5\hat{k}}{\sqrt{30}}$$

$$\hat{A} + \hat{B} = \frac{4\hat{i} + 6\hat{j} + 7\hat{k}}{\sqrt{101}}$$

$\vec{A} \cdot \vec{B} = |\vec{A}| |\vec{B}| \cos \theta$

$\rightarrow \theta = \text{angle b/w } \vec{A} \text{ & } \vec{B}$

$$\hat{i} \cdot \hat{i} = |\hat{i}| |\hat{i}| \cos 0^\circ = 1$$

$$\hat{j} \cdot \hat{j} = 1$$

$$\hat{k} \cdot \hat{k} = 1$$

$$\hat{i} \cdot \hat{j} = (\hat{i} \cdot 1) \hat{j} = 0 \quad \checkmark$$

$$\hat{i} \cdot \hat{k} = 0$$

$$\hat{j} \cdot \hat{k} = 0$$

Ex.  $\vec{A} = 2\hat{i} + 3\hat{j} + \hat{k}$

$$\vec{B} = \hat{i} - \hat{j} - \hat{k}$$

$$= \sqrt{2^2 + (-1)^2 + (-1)^2} = \sqrt{3}$$

$$\vec{A} \cdot \vec{B} = (2\hat{i} + 3\hat{j} + \hat{k}) \cdot (\hat{i} - \hat{j} - \hat{k})$$

$$= 2 - 3 - 1 = \boxed{-2}$$

$$\rightarrow \cos \theta = \frac{\vec{A} \cdot \vec{B}}{|\vec{A}| |\vec{B}|} = \frac{-2}{\sqrt{101} \times \sqrt{3}}$$

$$= \frac{-2}{\sqrt{42}} = \frac{-2}{\sqrt{42}}$$

$$\therefore \theta = \cos^{-1} \left( \frac{-2}{\sqrt{42}} \right)$$

$$\vec{A} = \hat{i} + 2\hat{j} + 3\hat{k}$$

$$\vec{B} = 2\hat{i} - \hat{j} + \hat{k}$$

$$\vec{A} \cdot \vec{B} = (\hat{i} + 2\hat{j} + 3\hat{k}) \cdot (2\hat{i} - \hat{j} + \hat{k})$$

$$= 2 - 2 + 0 = \boxed{0}$$

$$\vec{A} = 5\hat{i} + 7\hat{j} - 3\hat{k}$$

$$\vec{B} = 2\hat{i} + 2\hat{j} - \hat{k}$$

$$\rightarrow \vec{A} \cdot \vec{B} = (5\hat{i} + 7\hat{j} - 3\hat{k}) \cdot (2\hat{i} + 2\hat{j} - \hat{k}) = 0.$$

$$(5)(1) + (7)(2) + (-3)(-1) = 0 + 14 + 3 = 17$$

$$17 - 24 = -7$$

$$(5)(1) + (7)(2) + (-3)(-1) = 0 + 14 + 3 = 17$$

$$Q. \quad |\vec{A} + \vec{B}| = |\vec{A} - \vec{B}|$$

Find angle between A & B.

$$\rightarrow |\vec{A} + \vec{B}|^2 = |\vec{A} - \vec{B}|^2$$

$$\Rightarrow A^2 + 2\vec{A} \cdot \vec{B} + B^2 = A^2 - 2\vec{A} \cdot \vec{B} + B^2$$

$$\Rightarrow 4\vec{A} \cdot \vec{B} = 0$$

$$\Rightarrow 4AB \cos\theta = 0$$

$$\Rightarrow \cos\theta = 0$$

$$\therefore \theta = 90^\circ$$

Q.  $\vec{P}, \vec{Q}$  &  $\vec{R}$  have magnitudes 5, 12 & 13 resp. If  $P+Q=R$ . Find out angle between Q & R.

$$\rightarrow \vec{P} = \vec{R} - \vec{Q}$$

Squaring both sides,

$$\rightarrow P^2 = R^2 + Q^2 + 2\vec{R} \cdot \vec{Q}$$

$$\rightarrow 25 = 144 + 169 - 2\vec{R} \cdot \vec{Q}$$

$$\rightarrow 25 = -125 + 2\vec{R} \cdot \vec{Q} \Rightarrow 25 = 313 - 2\vec{R} \cdot \vec{Q}$$

$$\rightarrow \vec{R} \cdot \vec{Q} = 25 \Rightarrow \vec{R} \cdot \vec{Q} = 144$$

$$\therefore RQ \cos\theta = 25 \Rightarrow RQ \cos\theta = 144$$

$$\therefore \cos\theta =$$

$$\therefore \cos\theta = \frac{12}{144} = \frac{1}{12}$$

Q.  $\vec{A} + \vec{B} = \vec{C}$  and  $A^2 + B^2 = C^2$ . Prove that A is perpendicular to B.

$$\rightarrow \vec{A} + \vec{B} = \vec{C}$$

(i) Squaring;

$$\rightarrow A^2 + B^2 + 2\vec{A} \cdot \vec{B} = C^2$$

$$\rightarrow C^2 + 2\vec{A} \cdot \vec{B} = C^2$$

$$\rightarrow 2AB \cos\theta = 0$$

$$\rightarrow \cos\theta = 0$$

$$\rightarrow \theta = 90^\circ$$

X — X — X — X — X



\* Vector Product

$$\rightarrow \vec{A} \times \vec{B} = |\vec{A}| |\vec{B}| \sin\theta \hat{n}$$

$\hat{n}$  = normal vector (perpendicular vector)

→ normal vector to vectors  $\vec{A}$  &  $\vec{B}$  both

$$\begin{aligned} \# \quad \hat{i} \times \hat{j} &= \hat{k} \quad \# \quad \hat{j} \times \hat{i} = -\hat{k} \\ \hat{j} \times \hat{k} &= \hat{i} \quad \hat{k} \times \hat{j} = -\hat{i} \\ \hat{k} \times \hat{i} &= \hat{j} \quad \hat{i} \times \hat{k} = -\hat{j} \end{aligned}$$



$$\# \quad \hat{i} \times \hat{i} = 0 \quad \hat{j} \times \hat{j} = 0 \quad \hat{k} \times \hat{k} = 0$$

$$\hat{i} \times \hat{j} = 0$$

$$\hat{j} \times \hat{k} = 0 \quad (\text{Normal to } \vec{A} \text{ and } \vec{B})$$

$$\text{Q- } \vec{A} = \hat{i} - 2\hat{j} + \hat{k}$$

$$\vec{B} = 2\hat{i} + 4\hat{j} - \hat{k}$$

$$\text{1. } \vec{A} \times \vec{B} = \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ 1 & -2 & 1 \\ 2 & 4 & -1 \end{vmatrix} = \hat{i}(2)(-1) - \hat{j}(-1-2) + \hat{k}(4-1) = -2\hat{i} + 3\hat{j} + 3\hat{k}$$

Always left to right

Cross multiplication.  $\theta = \text{Btw } \vec{A} \text{ & } \vec{B}$

$O = \text{Btw } \vec{A}$

$$\text{2. Angle b/w } \vec{A} \text{ & } \vec{B} \rightarrow \sin \theta = \frac{|\vec{A} \times \vec{B}|}{|\vec{A}| |\vec{B}|} \quad [O \text{ P} = \theta]$$

$$\Rightarrow \sin \theta = \frac{\sqrt{77}}{\sqrt{21} \sqrt{6}} \quad \text{[if both rgt. angled]}$$

$$\text{2) } \theta = \sin^{-1} \left( \frac{\sqrt{77}}{\sqrt{21} \sqrt{6}} \right)$$

$$\text{3. Find normal vector } \hat{n} = \frac{\vec{A} \times \vec{B}}{|\vec{A} \times \vec{B}|}$$

$$= \frac{-2\hat{i} + 3\hat{j} + 3\hat{k}}{\sqrt{77}}$$

$$\text{Q. } \vec{A} = 2\hat{i} - 3\hat{j} - \hat{k}$$

$$\vec{B} = -6\hat{i} + 9\hat{j} + 3\hat{k} \quad [O = \vec{A} \times \vec{B}]$$

Prove  $\vec{A} \text{ & } \vec{B}$  are parallel:  $\vec{A} \times \vec{B}$

$$\Rightarrow \vec{A} \times \vec{B} = \hat{i}(-9 - (-9)) - \hat{j}(6 - 6) + \hat{k}(18 - 18) = 0\hat{i} - 0\hat{j} + 0\hat{k} = 0$$

parallel to

$\vec{A} \parallel \vec{B}$ , hence perpend.

$$\text{Q- } \vec{A} = 3\hat{i} + 4\hat{j} + 0\hat{k}$$

$$\vec{B} = -3\hat{i} + 7\hat{j} + 0\hat{k}$$

$$|\vec{A} \times \vec{B}|$$

are the sides of parallelogram. find area of lgm.

$$\Rightarrow \vec{A} \times \vec{B} = \hat{i}(0+0) - \hat{j}(0+0) + \hat{k}(21 - (-12))$$

$$\Rightarrow \vec{A} \times \vec{B} = 33\hat{k}$$

$$\Rightarrow |\vec{A} \times \vec{B}| = \sqrt{33^2}$$

$$= \sqrt{1089} = 33 \text{ sq. unit}$$

Q-  $\vec{A} \text{ & } \vec{B}$  are sides of lgm. Its area is  $\frac{|\vec{A}| |\vec{B}|}{2}$ . Find angle between  $\vec{A} \text{ & } \vec{B}$ .

$$\Rightarrow |\vec{A} \times \vec{B}| = \frac{|\vec{A}| |\vec{B}|}{2}$$

$$\Rightarrow |\vec{A}| |\vec{B}| \sin \theta = \frac{|\vec{A}| |\vec{B}|}{2}$$

$$\Rightarrow \sin \theta = \frac{1}{2}$$

$$\Rightarrow \theta = 30^\circ$$

$$\text{Q. } \vec{A} = 2\hat{i} - 3\hat{j} + 6\hat{k}, ((\vec{A}) - P) \vec{s} = 5\hat{x} \vec{A}$$

$$\vec{B} = \hat{i} + \hat{j} - \hat{k}$$

Find a vector whose length is 7 & perpendicular to A & B.

$$\Rightarrow \vec{A} \times \vec{B} = \hat{i}(3+6) - \hat{j}(-2+6) + \hat{k}(2+3)$$

$$\Rightarrow \vec{A} \times \vec{B} = -3\hat{i} - 8\hat{j} + 5\hat{k}$$

$$|\vec{A} \times \vec{B}| = \sqrt{9+64+25} = 10$$

$$((\vec{A}) - P) \vec{s} + (\vec{A} \times \vec{B}) \vec{s} = 5\hat{x} \vec{A}$$

$$\hat{n} = \frac{\vec{A} \times \vec{B}}{|\vec{A} \times \vec{B}|}$$

$$= \frac{-3\hat{i} - 8\hat{j} + 5\hat{k}}{\sqrt{98}}$$

$$= \frac{-3\hat{i} - 8\hat{j} + 5\hat{k}}{7\sqrt{2}}$$


$$\Rightarrow \hat{n} = \frac{-3\hat{i} - 8\hat{j} + 5\hat{k}}{7\sqrt{2}}$$

$$\therefore \hat{n} = \boxed{\frac{-3\hat{i} - 8\hat{j} + 5\hat{k}}{\sqrt{2}}}$$

$$\frac{1}{s} = B \sin \theta$$

$$\boxed{\theta = 90^\circ}$$

## # Projectile Motion



$\Rightarrow$  Horizontal Direction  $\rightarrow v_x \rightarrow$  constant (always)

Velocity  $\rightarrow$  axis is  $\alpha$

$\therefore$  direction never changes

$\Rightarrow$  Vertical Direction  $\rightarrow$  v changes with time

Direction of velocity changes

$\ddot{x} \rightarrow$	Variable uniform
$\ddot{a} \rightarrow$	variable

accelerated motion

Imp   
  $\star \star$

① Equation of Trajectory

(Displacement)

Horizontal Distance  $\rightarrow$   $x = v t$

$$x = v_{x0} t$$

$$\boxed{t = \frac{x}{v_{x0}}} \quad \text{--- (1)}$$

Vertical Displacement  $\rightarrow v_y = u \sin \theta + at = -g t$

$$y = ut + \frac{1}{2} at^2$$

$$\Rightarrow y = u \sin \theta x + \frac{1}{2} (-g) + \left( \frac{u \cos \theta}{g} \right)^2$$

$$\Rightarrow y = x \tan \theta + -\left( \frac{g}{2u^2 \cos^2 \theta} \right) x^2$$

Let  $\tan \theta = p$   $\Rightarrow [y = px - \frac{g}{2} x^2] = \text{eqn of parabola}$

$$\frac{g}{2u^2 \cos^2 \theta} = q$$

## ② Time of maximum height ( $t_m$ )

For vertical,  $v = u + at$

$$\Rightarrow 0 = u \sin \theta - gt_m$$

$$t_m = \frac{u \sin \theta}{g}$$

equation for wait time  $\textcircled{1}$

(time up)

$t_w = 2t_m$  (time of descent)

$$t_w = 2t_m$$

$$\textcircled{1} \rightarrow \frac{t_w}{t_w} = 1$$

## Ch 8 $\star$ Laws of Motion

\* Inertia  $\rightarrow$  Resistance to change in state of motion.

↳ of rest - Tendency to stay at rest.

↳ of motion - Tendency to stay in motion

↳ of direction - Tendency to retain motion in one direction.

### # 1st Law Of Motion - "Law of Inertia"

↳ Every body continues in the state of rest or state of uniform motion unless compelled by external force.

## Projectile Motion

### ③ Time of flight

$\rightarrow$  Displacement in vertical direction = 0

~~$y = u_y t + \frac{1}{2} at^2$~~

$$0 = u \sin \theta t - \frac{1}{2} gt^2$$

$$\Rightarrow t = \frac{2u \sin \theta}{g} \quad \text{— Time of flight}$$

Also, time of max height =  $\frac{u \sin \theta}{g}$

$\therefore$  Time of flight =  $2 \times$  Time of max height.