

# CLINICAL TRIAL PROTOCOL: INVESTIGATIONAL COMPOUND VX-709

Protocol ID: NCT05829104 | Sponsor: BioGenex Therapeutics | Phase: IIb

## 1. Study Synopsis & Objectives

This Phase IIb multicenter, randomized, double-blind, placebo-controlled study evaluates the therapeutic efficacy, safety, and pharmacokinetic profile of oral VX-709 in adult patients with moderate-to-severe ulcerative colitis.

## **2. Patient Inclusion Criteria**

- Age 18 to 70 years at the time of signing informed consent.
- Confirmed diagnosis of ulcerative colitis for  $\geq 6$  months established by clinical and endoscopic evidence.
- Total Mayo score of 6 to 12 points, including an endoscopic subscore of  $\geq 2$ .

### **3. Patient Exclusion Criteria**

- Prior exposure to JAK inhibitors or anti-TNF biologics within 8 weeks of randomization.
- History of severe cardiovascular events, deep vein thrombosis, or pulmonary embolism within 12 months.
- Absolute neutrophil count (ANC) < 1,500 cells/uL or serum creatinine > 1.8 mg/dL.

## 4. Dosage Regimens & Administration Schedule

Subjects are randomized 1:1:1 into three treatment arms: (Arm A) VX-709 15mg once daily; (Arm B) VX-709 30mg once daily; (Arm C) Matched Placebo once daily. Study medication must be taken in the morning with 240mL of water with or without food.

## 5. Primary & Secondary Endpoints

Primary Endpoint: Proportion of patients achieving clinical remission (Mayo score  $\leq 2$  with no individual subscore  $> 1$ ) at Week 12.

Secondary Endpoints: Endoscopic mucosal healing at Week 12 and Week 24; change from baseline in fecal calprotectin levels at Week 4, 8, and 12.

## **6. Adverse Event Reporting & Data Safety Monitoring**

All Serious Adverse Events (SAEs) must be transmitted to the Medical Monitor within 24 hours of site awareness. An independent Data Safety Monitoring Board (DSMB) reviews unblinded safety data every 6 weeks.