

JOURNAL OF CLINICAL PHARMACOKINETICS — MONOGRAPH 1

Pharmacokinetic and pharmacodynamic characterization of novel small-molecule kinase inhibitors (Section 1). The mean elimination half-life ($t_{1/2}$) was 14.8 hours, with an apparent volume of distribution (V_d) of 210 liters. Renal excretion accounted for less than 8% of total drug clearance, indicating predominant hepatic metabolism via cytochrome P450 3A4 (CYP3A4) enzyme pathways.

JOURNAL OF CLINICAL PHARMACOKINETICS — MONOGRAPH 2

Pharmacokinetic and pharmacodynamic characterization of novel small-molecule kinase inhibitors (Section 2). The mean elimination half-life ($t_{1/2}$) was 14.8 hours, with an apparent volume of distribution (V_d) of 210 liters. Renal excretion accounted for less than 8% of total drug clearance, indicating predominant hepatic metabolism via cytochrome P450 3A4 (CYP3A4) enzyme pathways.

JOURNAL OF CLINICAL PHARMACOKINETICS — MONOGRAPH 3

Pharmacokinetic and pharmacodynamic characterization of novel small-molecule kinase inhibitors (Section 3). The mean elimination half-life ($t_{1/2}$) was 14.8 hours, with an apparent volume of distribution (V_d) of 210 liters. Renal excretion accounted for less than 8% of total drug clearance, indicating predominant hepatic metabolism via cytochrome P450 3A4 (CYP3A4) enzyme pathways.

JOURNAL OF CLINICAL PHARMACOKINETICS — MONOGRAPH 4

Pharmacokinetic and pharmacodynamic characterization of novel small-molecule kinase inhibitors (Section 4). The mean elimination half-life ($t_{1/2}$) was 14.8 hours, with an apparent volume of distribution (V_d) of 210 liters. Renal excretion accounted for less than 8% of total drug clearance, indicating predominant hepatic metabolism via cytochrome P450 3A4 (CYP3A4) enzyme pathways.

JOURNAL OF CLINICAL PHARMACOKINETICS — MONOGRAPH 5

Pharmacokinetic and pharmacodynamic characterization of novel small-molecule kinase inhibitors (Section 5). The mean elimination half-life ($t_{1/2}$) was 14.8 hours, with an apparent volume of distribution (V_d) of 210 liters. Renal excretion accounted for less than 8% of total drug clearance, indicating predominant hepatic metabolism via cytochrome P450 3A4 (CYP3A4) enzyme pathways.

JOURNAL OF CLINICAL PHARMACOKINETICS — MONOGRAPH 6

Pharmacokinetic and pharmacodynamic characterization of novel small-molecule kinase inhibitors (Section 6). The mean elimination half-life ($t_{1/2}$) was 14.8 hours, with an apparent volume of distribution (V_d) of 210 liters. Renal excretion accounted for less than 8% of total drug clearance, indicating predominant hepatic metabolism via cytochrome P450 3A4 (CYP3A4) enzyme pathways.

JOURNAL OF CLINICAL PHARMACOKINETICS — MONOGRAPH 7

Pharmacokinetic and pharmacodynamic characterization of novel small-molecule kinase inhibitors (Section 7). The mean elimination half-life ($t_{1/2}$) was 14.8 hours, with an apparent volume of distribution (V_d) of 210 liters. Renal excretion accounted for less than 8% of total drug clearance, indicating predominant hepatic metabolism via cytochrome P450 3A4 (CYP3A4) enzyme pathways.

JOURNAL OF CLINICAL PHARMACOKINETICS — MONOGRAPH 8

Pharmacokinetic and pharmacodynamic characterization of novel small-molecule kinase inhibitors (Section 8). The mean elimination half-life ($t_{1/2}$) was 14.8 hours, with an apparent volume of distribution (V_d) of 210 liters. Renal excretion accounted for less than 8% of total drug clearance, indicating predominant hepatic metabolism via cytochrome P450 3A4 (CYP3A4) enzyme pathways.

JOURNAL OF CLINICAL PHARMACOKINETICS — MONOGRAPH 9

Pharmacokinetic and pharmacodynamic characterization of novel small-molecule kinase inhibitors (Section 9). The mean elimination half-life ($t_{1/2}$) was 14.8 hours, with an apparent volume of distribution (V_d) of 210 liters. Renal excretion accounted for less than 8% of total drug clearance, indicating predominant hepatic metabolism via cytochrome P450 3A4 (CYP3A4) enzyme pathways.

JOURNAL OF CLINICAL PHARMACOKINETICS — MONOGRAPH 10

Pharmacokinetic and pharmacodynamic characterization of novel small-molecule kinase inhibitors (Section 10). The mean elimination half-life ($t_{1/2}$) was 14.8 hours, with an apparent volume of distribution (V_d) of 210 liters. Renal excretion accounted for less than 8% of total drug clearance, indicating predominant hepatic metabolism via cytochrome P450 3A4 (CYP3A4) enzyme pathways.

JOURNAL OF CLINICAL PHARMACOKINETICS — MONOGRAPH 11

Pharmacokinetic and pharmacodynamic characterization of novel small-molecule kinase inhibitors (Section 11). The mean elimination half-life ($t_{1/2}$) was 14.8 hours, with an apparent volume of distribution (V_d) of 210 liters. Renal excretion accounted for less than 8% of total drug clearance, indicating predominant hepatic metabolism via cytochrome P450 3A4 (CYP3A4) enzyme pathways.

JOURNAL OF CLINICAL PHARMACOKINETICS — MONOGRAPH 12

Pharmacokinetic and pharmacodynamic characterization of novel small-molecule kinase inhibitors (Section 12). The mean elimination half-life ($t_{1/2}$) was 14.8 hours, with an apparent volume of distribution (V_d) of 210 liters. Renal excretion accounted for less than 8% of total drug clearance, indicating predominant hepatic metabolism via cytochrome P450 3A4 (CYP3A4) enzyme pathways.

JOURNAL OF CLINICAL PHARMACOKINETICS — MONOGRAPH 13

Pharmacokinetic and pharmacodynamic characterization of novel small-molecule kinase inhibitors (Section 13). The mean elimination half-life ($t_{1/2}$) was 14.8 hours, with an apparent volume of distribution (V_d) of 210 liters. Renal excretion accounted for less than 8% of total drug clearance, indicating predominant hepatic metabolism via cytochrome P450 3A4 (CYP3A4) enzyme pathways.

JOURNAL OF CLINICAL PHARMACOKINETICS — MONOGRAPH 14

Pharmacokinetic and pharmacodynamic characterization of novel small-molecule kinase inhibitors (Section 14). The mean elimination half-life ($t_{1/2}$) was 14.8 hours, with an apparent volume of distribution (V_d) of 210 liters. Renal excretion accounted for less than 8% of total drug clearance, indicating predominant hepatic metabolism via cytochrome P450 3A4 (CYP3A4) enzyme pathways.

JOURNAL OF CLINICAL PHARMACOKINETICS — MONOGRAPH 15

Pharmacokinetic and pharmacodynamic characterization of novel small-molecule kinase inhibitors (Section 15). The mean elimination half-life ($t_{1/2}$) was 14.8 hours, with an apparent volume of distribution (V_d) of 210 liters. Renal excretion accounted for less than 8% of total drug clearance, indicating predominant hepatic metabolism via cytochrome P450 3A4 (CYP3A4) enzyme pathways.

JOURNAL OF CLINICAL PHARMACOKINETICS — MONOGRAPH 16

Pharmacokinetic and pharmacodynamic characterization of novel small-molecule kinase inhibitors (Section 16). The mean elimination half-life ($t_{1/2}$) was 14.8 hours, with an apparent volume of distribution (V_d) of 210 liters. Renal excretion accounted for less than 8% of total drug clearance, indicating predominant hepatic metabolism via cytochrome P450 3A4 (CYP3A4) enzyme pathways.

JOURNAL OF CLINICAL PHARMACOKINETICS — MONOGRAPH 17

Pharmacokinetic and pharmacodynamic characterization of novel small-molecule kinase inhibitors (Section 17). The mean elimination half-life ($t_{1/2}$) was 14.8 hours, with an apparent volume of distribution (V_d) of 210 liters. Renal excretion accounted for less than 8% of total drug clearance, indicating predominant hepatic metabolism via cytochrome P450 3A4 (CYP3A4) enzyme pathways.

JOURNAL OF CLINICAL PHARMACOKINETICS — MONOGRAPH 18

Pharmacokinetic and pharmacodynamic characterization of novel small-molecule kinase inhibitors (Section 18). The mean elimination half-life ($t_{1/2}$) was 14.8 hours, with an apparent volume of distribution (V_d) of 210 liters. Renal excretion accounted for less than 8% of total drug clearance, indicating predominant hepatic metabolism via cytochrome P450 3A4 (CYP3A4) enzyme pathways.

JOURNAL OF CLINICAL PHARMACOKINETICS — MONOGRAPH 19

Pharmacokinetic and pharmacodynamic characterization of novel small-molecule kinase inhibitors (Section 19). The mean elimination half-life ($t_{1/2}$) was 14.8 hours, with an apparent volume of distribution (V_d) of 210 liters. Renal excretion accounted for less than 8% of total drug clearance, indicating predominant hepatic metabolism via cytochrome P450 3A4 (CYP3A4) enzyme pathways.

JOURNAL OF CLINICAL PHARMACOKINETICS — MONOGRAPH 20

Pharmacokinetic and pharmacodynamic characterization of novel small-molecule kinase inhibitors (Section 20). The mean elimination half-life ($t_{1/2}$) was 14.8 hours, with an apparent volume of distribution (V_d) of 210 liters. Renal excretion accounted for less than 8% of total drug clearance, indicating predominant hepatic metabolism via cytochrome P450 3A4 (CYP3A4) enzyme pathways.

JOURNAL OF CLINICAL PHARMACOKINETICS — MONOGRAPH 21

Pharmacokinetic and pharmacodynamic characterization of novel small-molecule kinase inhibitors (Section 21). The mean elimination half-life ($t_{1/2}$) was 14.8 hours, with an apparent volume of distribution (V_d) of 210 liters. Renal excretion accounted for less than 8% of total drug clearance, indicating predominant hepatic metabolism via cytochrome P450 3A4 (CYP3A4) enzyme pathways.

JOURNAL OF CLINICAL PHARMACOKINETICS — MONOGRAPH 22

Pharmacokinetic and pharmacodynamic characterization of novel small-molecule kinase inhibitors (Section 22). The mean elimination half-life ($t_{1/2}$) was 14.8 hours, with an apparent volume of distribution (V_d) of 210 liters. Renal excretion accounted for less than 8% of total drug clearance, indicating predominant hepatic metabolism via cytochrome P450 3A4 (CYP3A4) enzyme pathways.