CODA: Theory Notes

Guarantees for host-calibrated, shift-corrected device thresholds

Abdelaziz Saud
University of Sharjah

Abstract—We collect the finite-sample guarantees behind CODA: (i) the marginal union-bound guarantee with exact conformal ranks, (ii) its regression-adjusted variant and the data-reuse conditions it needs, (iii) a relative guarantee (device false-alarm rate at most host rate plus δ) that survives arbitrary calibration/test drift, with a training-conditional (PAC) version, (iv) a block version for dependent windows whose resolution is set by the number of blocks, and (v) tighter corrections. The union bound is sharp when only the two marginals are known (Makarov's bound), so tightening requires information about the joint law of score and shift: paired in-domain probes (CODA-J) or a structural assumption (CODA-conv). Every result is checked numerically in experiments/sim_theory.py.

I. Setting and the conformal quantile

A window x has host score $s(x)$ and device score $s_d(x) = s(x) + \Delta(x)$; $\Delta = \Delta_d$ is the device shift. An alarm is raised when the score exceeds a threshold. We have n host calibration windows $X_1, \dots, X_n$ with scores $S_i = s(X_i)$, m probe windows $Z_1, \dots, Z_m$ with shifts $D_j = \Delta(Z_j)$, and a normal test window X with $S = s(X)$, $D = \Delta(X)$. The device false-alarm rate of a (data-dependent) threshold t is $\mathrm{FAR}_d(t) = \mathbb{P}(s_d(X) > t)$, where $\mathbb{P}$ is over all randomness unless stated; $\mathrm{FAR}_{\mathrm{host}}(t) = \mathbb{P}(s(X) > t)$.

Definition 1 (Conformal quantile). For $v_1, \dots, v_N$ and $\beta \in (0,1)$ let $k_N(\beta) = \lceil (N+1)\beta \rceil$ and $\widehat{Q}_\beta(v_{1:N}) = v_{(k_N(\beta))}$, the $k_N(\beta)$-th smallest value, with $\widehat{Q}_\beta = +\infty$ if $k_N(\beta) > N$. This is conformal_quantile in coda/conformal.py. It is finite iff $N \geq (\beta)/(1-\beta)$, e.g. $N \geq 99$ for $\beta = 0.99$.

Lemma 1 (Split conformal [1], [2], [3]). If $(V_1, \dots, V_N, V_{N+1})$ is exchangeable, then $\mathbb{P}(V_{N+1} > \widehat{Q}_\beta(V_{1:N})) \leq 1 - \beta$. If moreover ties occur with probability zero and $k_N(\beta) \leq N$, then $\mathbb{P}(V_{N+1} > \widehat{Q}_\beta(V_{1:N})) \geq 1 - \beta - \frac{1}{N+1}$.

Proof. Write $k = k_N(\beta)$. If $k > N$ the bound is trivial. Otherwise $V_{N+1} > V_{(k)}$ among $V_{1:N}$ iff the rank of V_{N+1} among all $N+1$ values is $> k$ (counting ties in its favour). By exchangeability the rank is uniform on $\{1, \dots, N+1\}$ without ties and stochastically larger than a "favourable" tie-broken uniform rank otherwise, so the probability is $\leq (N+1-k)/(N+1) \leq 1 - \beta$; without ties it equals $(N+1-k)/(N+1) \geq 1 - \beta - 1/(N+1)$ because $k < (N+1)\beta + 1$. □

II. Marginal guarantee of CODA

CODA sets $\tau = \widehat{Q}_{1-\alpha}(S_{1:n})$, $\hat{q} = \widehat{Q}_{1-\delta}(D_{1:m})$ and $\tau_d = \tau + \hat{q}$.

Assumption 1. (A1) $(S_1, \dots, S_n, S)$ is exchangeable. (A2) $(D_1, \dots, D_m, D)$ is exchangeable.

No independence between calibration and probes is required: (A1) and (A2) are separate marginal statements, and both hold, e.g., when calibration, probe and test windows are i.i.d. draws from the same normal law, even if the probes are a subset of the calibration windows.

Theorem 1 (Marginal validity). Under Assumption 1,

$$\mathrm{FAR}_d(\tau_d) \leq \underbrace{\mathbb{P}(S > \tau)}_{\leq \alpha} + \underbrace{\mathbb{P}(D > \hat{q})}_{\leq \delta} \leq \alpha + \delta.$$

If host scores have no ties a.s. and $n \geq (1-\alpha)/\alpha$, the host part is tight: $\alpha - \frac{1}{n+1} \leq \mathrm{FAR}_{\mathrm{host}}(\tau) \leq \alpha$; likewise $\delta - \frac{1}{m+1} \leq \mathbb{P}(D > \hat{q}) \leq \delta$ for continuous shifts and $m \geq (1-\delta)/\delta$.

Proof. On $\{S \leq \tau\} \cap \{D \leq \hat{q}\}$ we have $s_d(X) = S + D \leq \tau + \hat{q}$. Hence $\{s_d(X) > \tau_d\} \subseteq \{S > \tau\} \cup \{D > \hat{q}\}$; apply the union bound and Lemma 1 to each term with (A1), (A2). □

Remark 1 (No matching lower bound for the device). Unlike the host, $\mathrm{FAR}_d(\tau_d)$ has no useful lower bound: the gap to $\alpha + \delta$ contains $\mathbb{P}(S > \tau,\ S + D \leq \tau + \hat{q})$, the host alarms removed by raising the threshold. With a heavy-tailed shift, $\hat{q}$ is large compared with the score spread near τ, so the slack is close to α; this is the observed FAR ≈ 0.02 at $\alpha = 0.05$ (W4A8, ESP-DL). Simulation T1: FAR 0.026 vs. bound 0.06.

Proposition 1 (The union bound is sharp given only marginals). Let F_S, F_Δ be fixed marginal laws. For every t,

$$\sup_{\text{couplings}} \mathbb{P}(S + D > t) = \min\Big\{1,\ \inf_u \big[\mathbb{P}(S > u) + \mathbb{P}(D \geq t - u)\big]\Big\}$$

up to the usual left-limit conventions (exact for continuous marginals) [4]. In particular, Theorem 1 cannot be improved without assumptions on the joint law.

Example attaining the bound. Let $U \sim \mathrm{Unif}(0,1)$, $S = U$ (so $\tau^* = 1-\alpha$), and $D = \eta U$ if $U \leq 1-\alpha-\delta$, $D = 1+U$ if $U \in (1-\alpha-\delta, 1-\alpha]$, $D = 1 - \frac{1}{2}(U - 1 + \alpha)$ if $U > 1-\alpha$, with $0 < \eta$ small and $\alpha + 2\delta \leq 1$. Then D is continuous with $\mathbb{P}(D > 1) = \delta$, so $q^* = 1$. On $\{U > 1-\alpha\}$, $S + D = 1 +$

$\frac{1}{2}(U+1-\alpha) > 2-\alpha$; on the band, $S+D = 1+2U > 2-\alpha$; on the bulk, $S+D \le (1-\alpha-\delta)(1+\eta) < 2-\alpha$. Hence $\mathbb{P}(S+D > \tau^*+q^*) = \alpha+\delta$; as $n, m \to \infty$, $\tau \to \tau^*$, $\hat{q} \to q^*$ and the CODA FAR tends to $\alpha+\delta$. The general identity is Makarov's theorem. □

Remark 2. Proposition 1 says that when the calibration data only reveal the law of S and the probes only reveal the law of Δ (out-of-domain probes), no rule can be uniformly tighter than the union bound with the best split (α', δ') of the budget. All tighter constructions in Sec. VI therefore use extra information about the joint law of (s, Δ) on test windows.

III. Regression-adjusted CODA

coda_linear fits $g(v) = av + b$ by least squares to the probe pairs $(s(Z_j), s_d(Z_j))$, takes $\hat{r} = \widehat{Q}_{1-\delta}(R_{1:m})$ of residuals $R_j = s_d(Z_j) - g(s(Z_j))$, and returns $\tau_d^g = \widehat{Q}_{1-\alpha}(g(S_{1:n})) + \hat{r}$.

Theorem 2 (Validity of regression-adjusted CODA). Let g be any measurable function computed from data $\mathcal{D}_{\text{fit}}$. Assume that conditionally on $\mathcal{D}_{\text{fit}}$: (B1) $(S_1, \dots, S_n, S)$ is exchangeable, and (B2) $(R_1, \dots, R_{m'}, R)$ is exchangeable, where R_j are residuals on m' residual probes and $R = s_d(X) - g(S)$. Then $\text{FAR}_d(\tau_d^g) \le \alpha + \delta$. If g is increasing (e.g. $a > 0$), then $\widehat{Q}_{1-\alpha}(g(S_{1:n})) = g(\tau)$ and the host event is exactly $\{S > \tau\}$.

Proof. $s_d(X) = g(S) + R$, so $\{s_d(X) > \tau_d^g\} \subseteq \{g(S) > \widehat{Q}_{1-\alpha}(g(S_{1:n}))\} \cup \{R > \hat{r}\}$. Given $\mathcal{D}_{\text{fit}}$, g is fixed, $(g(S_1), \dots, g(S_n), g(S))$ is exchangeable by (B1), and Lemma 1 applies to both terms; integrate over $\mathcal{D}_{\text{fit}}$. Monotonicity of g is not needed for validity because the quantile is taken over $g(S_i)$, not $g(\tau)$. □

Remark 3 (Data reuse). (i) Calibration part. If probes are drawn from the calibration windows, g depends on some S_i and (B1) fails conditionally on $\mathcal{D}_{\text{fit}}$. The probes must be removed from the host calibration set, as experiments/e3_correction.py does (np.delete(fp_cal, idx)). For plain CODA (Theorem 1) this removal is not needed, but it is harmless. (ii) Residual part. The current implementation fits g and computes $\hat{r}$ on the same probes; in-sample residuals are not exchangeable with the test residual, so (B2) holds only approximately (the effect is of order p/m for $p = 2$ parameters). In simulation T2 the residual exceedance is 0.0095–0.0100 for $m \in \{100, 200, 500\}$ (target 0.01), i.e. not detectable. A provably valid version fits g on a separate fifth of the probes (or on any independent data, e.g. synthetic windows, which need not be exchangeable with the test windows at all) and computes $\hat{r}$ on the rest; this needs $m' \ge 99$ residual probes for $\delta = 0.01$. Jackknife+ residuals [5] give a no-split alternative at level 2δ.

IV. Relative guarantee under drift

The main metric is the excess false-alarm rate $\text{FAR}_d - \text{FAR}_{\text{host}}$. It needs no assumption on the calibration data.

Theorem 3 (Device matches host). Let τ be any threshold that is independent of the probes and the test window (for example a conformal quantile of calibration data from a different, drifted distribution). Let the test window have arbitrary law P_{test}, and let $Z_1, \dots, Z_m$ be i.i.d. from P_{probe}, independent of (τ, X). Assume shift dominance:

$$P_{\text{test}}(\Delta(X) > c) \le P_{\text{probe}}(\Delta(Z) > c) \quad \forall c, \tag{1}$$

which holds with equality if $P_{\text{probe}} = P_{\text{test}}$. Write $\text{FAR}_\bullet(t) = P_{\text{test}}(\cdot > t)$ conditional on the thresholds. Then, for every realization,

$$\text{FAR}_d(\tau + \hat{q}) - \text{FAR}_{\text{host}}(\tau) \le \pi(\hat{q}) := P_{\text{test}}\big(\Delta(X) > \hat{q},\ s(X) \le \tau\big),$$

and $\mathbb{E}[\pi(\hat{q})] \le \delta$. Moreover, if $\hat{q} = D_{(k)}$ with

$$k = k_{\text{PAC}}(m, \delta, \gamma) := \min\{k : \mathbb{P}(\text{Bin}(m, 1-\delta) \ge k) \le \gamma\},$$

then $\mathbb{P}\big(\text{FAR}_d(\tau + \hat{q}) - \text{FAR}_{\text{host}}(\tau) > \delta\big) \le \gamma$.

Proof. Pointwise, $\{s_d > \tau + \hat{q}\} \subseteq \{s > \tau\} \cup \{s \le \tau,\ \Delta > \hat{q}\}$, which gives the first inequality. By (1), $\pi(\hat{q}) \le P_{\text{probe}}(\Delta > \hat{q})$. Let D' be a fresh draw from P_{probe}; $(D_1, \dots, D_m, D')$ is i.i.d., so $\mathbb{E}[P_{\text{probe}}(\Delta > \hat{q})] = \mathbb{P}(D' > \hat{q}) \le \delta$ by Lemma 1. For the PAC part, let $\xi = \inf\{x : F(x) \ge 1-\delta\}$, F the c.d.f. of D'. If $D_{(k)} \ge \xi$ then $P_{\text{probe}}(\Delta > D_{(k)}) \le 1 - F(\xi) \le \delta$. Otherwise at least k of the D_j are $< \xi$, and $\mathbb{P}(D_j < \xi) \le 1-\delta$, so this has probability $\le \mathbb{P}(\text{Bin}(m, 1-\delta) \ge k) \le \gamma$. No continuity is needed. This is the training-conditional guarantee of [6] (see also [7]). □

Remark 4 (What the experiments measure). (i) The bound is one-sided and per-τ: it holds for the drifted "time" protocol as long as (1) holds. Probes taken from the training period do not guarantee (1) if drift moves inputs into regions with larger quantization error; this is an assumption on the shift only, which is much weaker than exchangeability of scores. (ii) The "viol" column (share of runs with excess $> \delta$) is a training-conditional quantity. With the marginal rank $k = \lceil (m+1)(1-\delta) \rceil$ the PAC level is only $\gamma = 0.44$ ($m = 500$), 0.40 ($m = 200$), 0.37 ($m = 100$); the observed $\approx 1\%$ comes from the union-bound slack, not from theory. For a guaranteed $\gamma = 0.05$ use $k_{\text{PAC}}(500, 0.01, 0.05) = 499$. Simulation T3: cal/test drift with host FAR 0.149 at $\alpha = 0.05$; mean excess -0.057, no run above δ.

V. Block-dependent windows

SMD windows overlap and are strongly dependent; the data are cut into shuffled 128-step blocks, and windows never cross blocks. The exchangeable units are blocks, not windows.

Assumption 2. Calibration blocks $B_1, \dots, B_K$ and the test block B_{K+1} are exchangeable as random multisets of window scores; block b holds $L_b \geq 1$ scores with empirical c.d.f. F_b. The test window is uniform within B_{K+1} (block-averaged FAR).

Definition 2 (Block conformal quantile). $\widehat{Q}^{\mathrm{blk}}_\beta = \inf\{t : \sum_{b=1}^K F_b(t) \geq \beta(K+1)\}$ ($+\infty$ if $\beta(K+1) > K$). With equal block sizes L it is the $\lceil \beta(K+1)L \rceil$-th smallest of the KL pooled scores.

Theorem 4 (Block split conformal). Under Assumption 2, $\tau^{\mathrm{blk}} = \widehat{Q}^{\mathrm{blk}}_{1-\alpha}$ satisfies $\mathbb{E}[1 - F_{K+1}(\tau^{\mathrm{blk}})] \leq \alpha$, i.e. the window-level FAR is at most α. Applying the same construction to probe blocks (K_p blocks of shifts) and combining as in Theorem 1 gives $\mathrm{FAR}_d \leq \alpha + \delta$, with $\hat{q}^{\mathrm{blk}}$ finite iff $K_p \geq (1-\delta)/\delta$.

Proof. Let $T = \inf\{t : \sum_{b=1}^{K+1} F_b(t) \geq (1-\alpha)(K+1)\}$. Adding $F_{K+1} \geq 0$ can only reach the target earlier, so $T \leq \tau^{\mathrm{blk}}$ and $1 - F_{K+1}(\tau^{\mathrm{blk}}) \leq 1 - F_{K+1}(T)$. T is a symmetric function of the $K+1$ blocks, so by exchangeability $\mathbb{E}[1 - F_{K+1}(T)] = \mathbb{E}\big[\frac{1}{K+1}\sum_{b=1}^{K+1}(1 - F_b(T))\big]$. By right-continuity $\sum_b F_b(T) \geq (1-\alpha)(K+1)$, so the bracket is $\leq \alpha$. □

Corollary 1 (Pooled window-level quantile). With equal block size L, the usual window-level rank $\lceil (KL+1)(1-\alpha) \rceil$ only guarantees FAR $\leq \alpha + (1-\alpha)\frac{L-1}{(K+1)L} < \alpha + \frac{1-\alpha}{K+1}$.

Proof. Same argument with T the k-th smallest of all $(K+1)L$ scores, $k = \lceil (KL+1)(1-\alpha) \rceil$: the fraction above T is $\leq 1 - k/((K+1)L) \leq \alpha + (1-\alpha)(L-1)/((K+1)L)$. □

Remark 5 (Resolution, and what it explains). The guarantee is in units of blocks: $1/(K+1)$ plays the role of $1/(n+1)$, and the bound cannot be improved without within-block assumptions (if all windows of a block share one score, the problem is conformal prediction with K exchangeable scalars). For $\delta = 0.01$ one needs $K_p \geq 99$ probe blocks ($\approx$ 12.7k steps at 128 steps per block). In E1 the probes come from the calibration blocks, roughly 30 blocks (15% of $\approx$200); Corollary 1 then allows a shift exceedance up to $\delta + 0.03$, consistent with the observed 1.5–2.3% exceedances at $\delta = 0.01$. Simulation T4 ($K = 33$, block-level heavy shift): window-level $\hat{q}$ is exceeded on 2.65% of test windows (target 1%) and host FAR is $0.056 > \alpha$; the block quantile is valid (0.028) but $\hat{q}^{\mathrm{blk}} = +\infty$ at $\delta = 0.01$. With $K = 150$: window-level 0.0137, block-level 0.0073.

Remark 6 (Approximate block exchangeability). Shuffled consecutive blocks of a stationary series are only approximately exchangeable (adjacent blocks are dependent at their boundaries). Barber et al. [8] bound the coverage loss of (weighted) split conformal by $\sum_i \tilde{w}_i \, d_{\mathrm{TV}}(Z, Z^i)$, where Z^i swaps unit i with the test unit and $\tilde{w}_i = w_i/(1+\sum_j w_j)$ for fixed weights $w_i \in [0,1]$ ($\tilde{w}_i = 1/(n+1)$ for unweighted split conformal on n units); applied with blocks as units, the loss is small when the series mixes on a scale shorter than the block length. Block-permutation conformal inference for dependent data is developed in [9]. A gap of a few windows between blocks reduces boundary dependence further.

VI. Tighter corrections

Proposition 1 limits what any marginal-only method can do. We list the options and their status.

A. Optimal split of the budget (valid, small gain)

Proposition 2. For any split $\alpha' + \delta' = \alpha + \delta$ fixed before seeing the data (or chosen on independent pilot data), $\tau(\alpha') + \hat{q}(\delta')$ is valid at $\alpha + \delta$. Choosing the split that minimizes the threshold on the same data over a grid G is only guaranteed at $\max_G \alpha' + \max_G \delta'$.

Proof. The first claim is Theorem 1. For the second, with g^* the minimizer, $\{s_d > \tau(\alpha'_{g^*}) + \hat{q}(\delta'_{g^*})\} \subseteq \bigcup_g \{S > \tau(\alpha'_g)\} \cup \bigcup_g \{D > \hat{q}(\delta'_g)\} = \{S > \tau(\max \alpha')\} \cup \{D > \hat{q}(\max \delta')\}$, since the thresholds are monotone in the level. □

In the population the optimal split equates the densities $f_S(\tau_{\alpha'}) = f_\Delta(q_{\delta'})$ (first-order condition of $\min_{\alpha'} F_S^{-1}(1-\alpha') + F_\Delta^{-1}(1-\alpha-\delta+\alpha')$); with a heavy-tailed shift it moves budget to δ. Simulation T5: $(0.04, 0.02)$ instead of $(0.05, 0.01)$ lowers the threshold by 14% of the gap to the oracle and raises FAR from 0.026 to 0.028 only. The split is not the main source of conservativeness.

B. CODA-J: paired in-domain probes (valid, dominates CODA)

Theorem 5 (CODA-J). Let the probes be disjoint from the calibration windows, and let the pairs $\big(s(Z_j), s_d(Z_j)\big)_{j \leq m}$ and $\big(s(X), s_d(X)\big)$ be exchangeable conditionally on the calibration data, with (A1) for the calibration data. Let $\tau = \widehat{Q}_{1-\alpha}(S_{1:n})$,

$$U_j = \begin{cases} s_d(Z_j) & s(Z_j) \leq \tau \\ -\infty & \text{otherwise} \end{cases}, \qquad t_J = \widehat{Q}_{1-\delta}(U_{1:m}).$$

Then (i) $\mathrm{FAR}_d(t_J) \leq \alpha + \delta$; (ii) $t_J \leq \tau + \hat{q}$ for the same probes (CODA-J never alarms less); (iii) under drift between calibration and test, $\mathbb{E}[\mathrm{FAR}_d(t_J) - \mathrm{FAR}_{\mathrm{host}}(\tau)] \leq \delta$ if the probe pairs and test pair are exchangeable.

Proof. (i) $\{s_d(X) > t_J\} \subseteq \{S > \tau\} \cup \{S \leq \tau,\ s_d(X) > t_J\} = \{S > \tau\} \cup \{U > t_J\}$ with U the test analogue of U_j. Given the calibration data, τ is fixed and $(U_1, \dots, U_m, U)$ is exchangeable; Lemma 1. (ii) $U_j \leq \tau + D_j$ for every j (equality of the first case uses $s(Z_j) \leq \tau$; the second is $-\infty$), and order statistics are monotone. (iii) As in Theorem 3. □

Remark 7. Disjointness matters here: if probes are calibration windows, τ depends on them and (U_j) is no longer exchangeable with U. As $\alpha' \to 0$, CODA-J becomes recalibration on the probes (B2m) at level δ'; at matched

budget, CODA-J($\alpha - \delta, \delta$) and B2m(α) give the same FAR (0.050) and recall in simulation T5, versus 0.026 for CODA. So with in-domain paired probes the union bound should not be used; CODA-J is the way to phrase "CODA with in-domain probes" so that it is never worse than either CODA or B2m. CODA-J is not valid for out-of-domain probes: if their host scores are lower than in-domain, U_j is too small.

C. CODA-conv: convolution of scores and shifts (valid under an extra assumption)

Assumption 3 (Conditional shift dominance). Probe shifts $D_1, \ldots, D_m$ are i.i.d. from a law P_Δ and independent of the calibration and test windows, and conditionally on $(S_1, \ldots, S_n, S)$ the test shift is dominated: $\mathbb{P}(D > c \mid S_{1:n}, S) \le P_\Delta((c, \infty))$ for all c. A sufficient condition: in the test domain $\Delta(X)$ is independent of $s(X)$ and has the probe shift law.

Theorem 6 (CODA-conv). Under (A1) and Assumption 3, let t_C be the k-th smallest of the nm sums $\{S_i + D_j\}$ with $k = \lceil (1-\beta)(n+1)(m+1) \rceil$ ($+\infty$ if $k > nm$). Then $\mathrm{FAR}_d(t_C) \le \beta$. No separate δ budget is spent.

Proof. Coupling: let $V \sim \mathrm{Unif}(0,1)$ independent of everything, $D = F^{-1}_{D \mid S_{1:n}, S}(V)$ (equal in law to the test shift) and $D' = F^{-1}_\Delta(V)$. Dominance gives $D \le D'$, and D' is independent of the S's with law P_Δ, so $(D_1, \ldots, D_m, D')$ is i.i.d. and independent of $(S_1, \ldots, S_n, S)$. Hence $\mathrm{FAR}_d(t_C) \le \mathbb{P}(S + D' > t_C)$. Put $S_{n+1} = S$, $D_{m+1} = D'$, and let T be the k-th smallest of all $(n+1)(m+1)$ sums. Since the calibration sums are a subset, $T \le t_C$. The joint law is invariant under separate permutations of the S-indices and of the D-indices, and T is invariant under both, so $\mathbb{P}(S_{n+1} + D_{m+1} > T)$ equals the expected fraction of the $(n+1)(m+1)$ cells above T, which is $\le 1 - k/((n+1)(m+1)) \le \beta$. □

Remark 8 (Strengths and the honest caveat). t_C is finite for small m: with $n = 2000$ it needs $m \ge 20$ at $\beta = 0.05$ and $m \ge 105$ at $\beta = 0.01$ (CODA at $\delta = 0.01$ needs $m \ge 99$ for any α). In simulation T5 with an independent heavy-tailed shift, CODA-conv(α) has FAR 0.048 (CODA 0.026), recall 0.45 vs. 0.27, and the threshold gap to the oracle shrinks from +0.86 to +0.07 (92% smaller), with in-domain or out-of-domain probes alike; at $m = 50$ it gives FAR 0.031 where CODA is $+\infty$. But Assumption 3 is not implied by any exchangeability and cannot be checked on the test domain with out-of-domain probes. When the shift grows with the score (positive dependence), CODA-conv is anti-conservative: FAR 0.055 with in-domain and 0.063 with out-of-domain probes at $\beta = \alpha = 0.05$. A partial diagnostic on the probes themselves is to test whether D_j depends on $s(Z_j)$ (e.g. Kendall's τ, or comparing upper shift quantiles in the top and bottom score halves); with paired in-domain probes the independence of s and Δ is directly testable.

Remark 9 (Monotone variant, valid but impractical). If $\Delta \mid s = v$ has the same law in both domains and is stochastically increasing in v, then shifts of probes with $s(Z) \ge v^*$ dominate $\Delta \mid s \le v^*$. With $v^* = \widehat{Q}_{1-\alpha_1}$ from one half of the calibration data, CODA-conv on the other half (scores above v^* masked to $-\infty$) with those probe shifts at level $1 - \alpha_2$ gives $\mathrm{FAR}_d \le \alpha_1 + \alpha_2$ by the same coupling. It needs (synthetic) probes whose host score exceeds v^* and is extremely conservative with heavy tails (T5: FAR 0.0005).

D. Per-run violations are not evidence of invalidity

All guarantees above except the PAC part of Theorem 3 are marginal (averaged over calibration and probe draws). Any method whose expected FAR is exactly α has a per-run FAR that fluctuates around α with standard deviation of order $\sqrt{\alpha(1-\alpha)/m_{\mathrm{eff}}}$, where m_{eff} is the number of exchangeable units used for its threshold (windows, or blocks in Sec. V). The share of runs with excess $> \delta$ is therefore large for every tight method. In simulation T5 (i.i.d. windows, $m = 500$), recalibration on the probes, B2m(α), which is exactly valid, has excess $> \delta$ in 18.8% of runs; the oracle threshold has 2.2%; CODA has 0% only because of its slack. The heuristic coda_convolution in coda/conformal.py shows excess $> \delta$ in 10–23% of SMD runs; this is in the range of B2m and is not by itself a sign of invalidity. The marginal check is the mean excess ($\le \delta$; CODA-conv(α) targets ≤ 0), and a per-run guarantee requires a PAC rank (Theorem 3) for every method, including B2m. With block-dependent data (≈ 30 blocks) the per-run spread is larger still, since $m_{\mathrm{eff}} \approx K$.

Remark 10 (Implementation detail of CODA-conv). coda_convolution returns np.quantile(sums, 1-alpha, method="higher"), i.e. rank $\lceil (1-\alpha)nm \rceil$ of the nm sums. Theorem 6 needs rank $\lceil (1-\alpha)(n+1)(m+1) \rceil$, which is larger by about $(1-\alpha)(n+m+1)$ positions (a quantile-level change of $\approx (1-\alpha)(1/n + 1/m)$, i.e. 0.0024 for $n = 2000$, $m = 500$). Using the theorem's rank makes the heuristic exactly valid under Assumption 3; it does not repair failures of that assumption.

E. Relation to conformal calibration transfer

Doula [10] (TCC) transfers labeled source calibration to a target space through unlabeled paired observations (x_s, x_t): a map f is learned on pairs, source calibration inputs are transported, and the remaining mismatch is corrected by a one-sided KS gap $\hat{\delta}$ between transported and real target inputs on a label-free surrogate T, inflated by two DKW terms, $\delta^+ = \hat{\delta} + \sqrt{\log(4/\eta)/(2m_{\mathrm{tgt}})} + \sqrt{\log(4/\eta)/(2m_{\mathrm{trans}})}$; calibration then uses level $\alpha^\star = \max\{0, \alpha - \delta^+\}$, with miscoverage $\le \alpha^\star + \delta^+ + \varepsilon$ with probability $\ge 1 - \eta$ over the unlabeled data, where ε bounds how much the surrogate gap understates the true score gap (Assumption A2 there). A weighted variant is valid with oracle density ratios.

Mapping to CODA. In CODA the "source" and "target" are the same window scored by two models, so every probe is a pair in TCC's sense, and the transport f corresponds to the score map g of coda_linear. Two differences matter. (i) In anomaly detection on normal data the target conformity score is label-free: the device score of an unlabeled normal window is the score whose quantile we need, so TCC's surrogate is exact ($\varepsilon = 0$) only when probes are normal windows from the test domain; but then the probes can be used directly (B2m, CODA-J), which is tighter. (ii) The KS/DKW correction costs $\sqrt{\log(4/\eta)/(2m)}$ per sample, ≈ 0.08 for $m = 500$, $\eta = 0.05$; this exceeds $\alpha = 0.05$, so $\alpha^\star = 0$ and TCC-KS is vacuous at our levels, whereas CODA's shift quantile costs $\delta = 0.01$ in expectation (or a binomial PAC rank, Theorem 3). Hence TCC does not give a tighter valid correction than the union bound in either probe regime: with in-domain paired probes, CODA-J/B2m dominate it, and with out-of-domain probes its target-sample term needs unlabeled target-domain inputs, which are exactly what CODA avoids putting on devices. Conceptually, CODA differs in what is transferred and what is assumed: TCC transports inputs and bounds a distribution gap (uniform over thresholds, via KS), while CODA transports a single threshold and bounds only the shift tail at that threshold (one quantile, no uniform bound), and it requires no learned map and no target-domain inputs, only that probe shifts dominate test shifts (Eq. (1)).

F. Other ideas (heuristics)

Weighted conformal on the host score. If $\Delta \mid s$ has the same law in both domains, the probe-to-test mismatch is a covariate shift in s; weighted conformal prediction [11] with the likelihood ratio $w(v) = f_{\text{test}}(v)/f_{\text{probe}}(v)$ (estimable from calibration and probe host scores) applied to U_j of CODA-J would be valid for a known ratio; with an estimated ratio and probes that rarely reach the score region near τ, it is a heuristic. Band decompositions $\{S > \tau\} \cup \{S \in (\tau - \epsilon, \tau], D > q_\epsilon\} \cup \{S \le \tau - \epsilon, D > q_\epsilon + \epsilon\}$ are a discretized CODA-J; their conformal estimates need paired in-domain data, and with out-of-domain probes they need the same invariance assumption as the weighted method.

VII. Summary of what is provable

TABLE I
Guarantees and assumptions ($\alpha = 0.05, \delta = 0.01$; simulation T5, independent heavy-tailed shift, $m = 500$)

Method	Bound	Needs	FAR
CODA	$\alpha + \delta$	(A1),(A2) or (1)	0.026
CODA, split (.04, .02)	$\alpha + \delta$	same, split fixed	0.028
CODA-lin (split fit)	$\alpha + \delta$	fit indep. of rest	–
CODA-J($\alpha - \delta, \delta$)	α	paired in-domain probes	0.050
B2m(α)	α	in-domain probes	0.050
CODA-conv(α)	α	Assumption 3	0.048
Block CODA	$\alpha + \delta$	block exch., $K_p \ge 99$	–

AI assistance

These notes and the simulation script were drafted with the help of an AI assistant (Claude) and checked by the author.

References

[1] V. Vovk, A. Gammerman, and G. Shafer, Algorithmic Learning in a Random World. New York, NY, USA: Springer, 2005, doi: 10.1007/b106715.

[2] J. Lei, M. G'Sell, A. Rinaldo, R. J. Tibshirani, and L. Wasserman, "Distribution-free predictive inference for regression," J. Amer. Statist. Assoc., vol. 113, no. 523, pp. 1094–1111, 2018, doi: 10.1080/01621459.2017.1307116.

[3] A. N. Angelopoulos and S. Bates, "Conformal prediction: A gentle introduction," Found. Trends Mach. Learn., vol. 16, no. 4, pp. 494–591, 2023, doi: 10.1561/2200000101.

[4] G. D. Makarov, "Estimates for the distribution function of a sum of two random variables when the marginal distributions are fixed," Theory Probab. Appl., vol. 26, no. 4, pp. 803–806, 1982 (Russian original 1981), doi: 10.1137/1126086.

[5] R. Foygel Barber, E. J. Candès, A. Ramdas, and R. J. Tibshirani, "Predictive inference with the jackknife+," Ann. Statist., vol. 49, no. 1, pp. 486–507, 2021, doi: 10.1214/20-AOS1965.

[6] V. Vovk, "Conditional validity of inductive conformal predictors," in Proc. Asian Conf. Mach. Learn. (ACML), PMLR vol. 25, pp. 475–490, 2012.

[7] S. Bates, A. Angelopoulos, L. Lei, J. Malik, and M. I. Jordan, "Distribution-free, risk-controlling prediction sets," J. ACM, vol. 68, no. 6, pp. 1–34, 2021, doi: 10.1145/3478535.

[8] R. Foygel Barber, E. J. Candès, A. Ramdas, and R. J. Tibshirani, "Conformal prediction beyond exchangeability," Ann. Statist., vol. 51, no. 2, pp. 816–845, 2023, doi: 10.1214/23-AOS2276.

[9] V. Chernozhukov, K. Wüthrich, and Y. Zhu, "Exact and robust conformal inference methods for predictive machine learning with dependent data," in Proc. Conf. Learn. Theory (COLT), PMLR vol. 75, pp. 732–749, 2018.

[10] A. Doula, "Conformal calibration transfer," in Proc. 43rd Int. Conf. Mach. Learn. (ICML), PMLR vol. 306, pp. 26291–26318, 2026; arXiv:2609.10737.

[11] R. J. Tibshirani, R. Foygel Barber, E. J. Candès, and A. Ramdas, "Conformal prediction under covariate shift," in Adv. Neural Inf. Process. Syst. (NeurIPS), vol. 32, 2019.