

# Elementary Calculus Manual

Santiago Angulo Díaz-Parreño (sangulo@ceu.es)  
Pablo Ares Gastesi (pablo.aresgastesi@ceu.es)  
José Rojo Montijano (jrojo.eps@ceu.es)  
Anselmo Romero Limón (arlimon@ceu.es)  
Alfredo Sánchez Alberca (asalber@ceu.es)

Sep 2016

Department of Applied Math and Statistics  
CEU San Pablo



CEU  
*Universidad  
San Pablo*




## License terms

This work is licensed under an Attribution-NonCommercial-ShareAlike 4.0 International Creative Commons License. <http://creativecommons.org/licenses/by-nc-sa/4.0/>

You are free to:

- Share – copy and redistribute the material in any medium or format
- Adapt – remix, transform, and build upon the material

Under the following terms:

-  **Attribution.** You must give appropriate credit, provide a link to the license, and indicate if changes were made. You may do so in any reasonable manner, but not in any way that suggests the licensor endorses you or your use.
-  **NonCommercial.** You may not use the material for commercial purposes.
-  **ShareAlike.** If you remix, transform, or build upon the material, you must distribute your contributions under the same license as the original.

No additional restrictions — You may not apply legal terms or technological measures that legally restrict others from doing anything the license permits.

## Contents

<b>1</b>	<b>Analytic geometry</b>	<b>3</b>
1.1	Vectors . . . . .	3
1.2	Lines . . . . .	7
1.3	Planes . . . . .	8
<b>2</b>	<b>Differential calculus with one real variable</b>	<b>10</b>
2.1	Concept of derivative . . . . .	10
2.2	Algebra of derivatives . . . . .	13
2.3	Analysis of functions . . . . .	14
2.4	Function approximation . . . . .	17
<b>3</b>	<b>Integrals</b>	<b>25</b>
3.1	Antiderivative of a function . . . . .	25
3.2	Elementary integrals . . . . .	26
3.3	Techniques of integration . . . . .	27
3.4	Definite integral . . . . .	31
3.5	Area calculation . . . . .	32
<b>4</b>	<b>Ordinary Differential Equations</b>	<b>34</b>
4.1	Ordinary Differential Equations . . . . .	34
4.2	Separable differential equations . . . . .	36
4.3	Homogeneous differential equations . . . . .	37
4.4	Linear differential equations . . . . .	38
<b>5</b>	<b>Several Variables Differentiable Calculus</b>	<b>41</b>
5.1	Vector functions of a single real variable . . . . .	41
5.2	Tangent line to a trajectory . . . . .	43
5.3	Functions of several variables . . . . .	47
5.4	Partial derivative notion . . . . .	50
5.5	Gradient . . . . .	52
5.6	Second order partial derivatives . . . . .	53
5.7	Matriz hessiana . . . . .	53

# 1 Analytic geometry

## Analytic geometry

## Contents

### 1.1 Vectors

#### Scalars

Some phenomena of Nature can be described by a number and a unit of measurement.

**Definition 1** (Scalar). A *scalar* is a number that expresses a magnitude without direction.

**Examples** The height or weight of a person, the temperature of a gas or the time it takes a vehicle to travel a distance.


However, there are other phenomena that cannot be described adequately by a scalar. If, for instance, a sailor wants to head for seaport and only knows the intensity of wind, he will not know what direction to take. The description of wind requires two elements: intensity and direction.

#### Vectors

**Definition 2** (Vector). A *vector* is a number that expresses a magnitude and has associated an orientation and a sense.

**Examples** The velocity of a vehicle or the force applied to an object.


Geometrically, a vector is represented by an directed line segment, that is, an arrow.



#### Vector representation

An oriented segment can be located in different places in a Cartesian space. However, regardless of where it is located, if the length and the direction of the segment does not change, the segment represents always the same vector.

This allows to represent all vectors with the same origin, the origin of the Cartesian coordinate system. Thus, a vector can be represented by the Cartesian *coordinates* of its final end in any Euclidean space.




### Vector from two points

Given two points  $P$  and  $Q$  of a Cartesian space, the vector that starts at  $P$  and ends at  $Q$  has coordinates  $\vec{PQ} = Q - P$ .

**Example** Given the points  $P = (1, 1)$  and  $Q = (3, 4)$  in the real plane  $\mathbb{R}^2$ , the coordinates of the vector that start at  $P$  and ends at  $Q$  are

$$\vec{PQ} = Q - P = (3, 4) - (1, 1) = (3 - 1, 4 - 1) = (2, 3).$$



### Module of a vector

**Definition 3** (Module of a vector). Given a vector  $\mathbf{v} = (v_1, \dots, v_n)$  in  $\mathbb{R}^n$ , the *module* of  $\mathbf{v}$  is

$$|\mathbf{v}| = \sqrt{v_1^2 + \dots + v_n^2}.$$

The module of a vector coincides with the length of the segment that represents the vector.

**Examples** Let  $\mathbf{u} = (3, 4)$  be a vector in  $\mathbb{R}^2$ , then its module is

$$|\mathbf{u}| = \sqrt{3^2 + 4^2} = \sqrt{25} = 5$$

Let  $\mathbf{v} = (4, 7, 4)$  be a vector in  $\mathbb{R}^3$ , then its module is

$$|\mathbf{v}| = \sqrt{4^2 + 7^2 + 4^2} = \sqrt{81} = 9$$


### Unit vectors

**Definition 4** (Unit vector). A vector  $\mathbf{v}$  in  $\mathbb{R}^n$  is a *unit vector* if its module is one, that is,  $|\mathbf{v}| = 1$ .

The unit vectors with the direction of the coordinate axes are of special importance and they form the *standard basis*.


In  $\mathbb{R}^2$  the standard basis is formed by two vectors

$$\mathbf{i} = (1, 0) \text{ and } \mathbf{j} = (0, 1)$$



In  $\mathbb{R}^3$  the standard basis is formed by three vectors

$$\mathbf{i} = (1, 0, 0), \mathbf{j} = (0, 1, 0) \text{ and } \mathbf{k} = (0, 0, 1)$$




### Sum of two vectors

**Definition 5** (Sum of two vectors). Given two vectors  $\mathbf{u} = (u_1, \dots, u_n)$  y  $\mathbf{v} = (v_1, \dots, v_n)$  de  $\mathbb{R}^n$ , the *sum* of  $\mathbf{u}$  and  $\mathbf{v}$  is

$$\mathbf{u} + \mathbf{v} = (u_1 + v_1, \dots, u_n + v_n).$$

**Example** Let  $\mathbf{u} = (3, 1)$  and  $\mathbf{v} = (2, 3)$  two vectors in  $\mathbb{R}^2$ , then the sum of them is

$$\mathbf{u} + \mathbf{v} = (3 + 2, 1 + 3) = (5, 4).$$




### Product of a vector by a scalar

**Definition 6** (Product of a vector by a scalar). Given a vector  $\mathbf{v} = (v_1, \dots, v_n)$  in  $\mathbb{R}^n$ , and a scalar  $a \in \mathbb{R}$ , the *product* of  $\mathbf{v}$  by  $a$  is

$$a\mathbf{v} = (av_1, \dots, av_n).$$

**Example** Let  $\mathbf{v} = (2, 1)$  a vector in  $\mathbb{R}^2$  and  $a = 2$  a scalar, then the product of  $a$  by  $\mathbf{v}$  is

$$a\mathbf{v} = 2(2, 1) = (4, 2).$$




### Expressing a vector as a linear combination of the standard basis

The sum of vectors and the product of vector by a scalar allow us to express any vector as a linear combination of the standard basis.

In  $\mathbb{R}^3$ , for instance, a vector with coordinates  $\mathbf{v} = (v_1, v_2, v_3)$  can be expressed as the linear combination

$$\mathbf{v} = (v_1, v_2, v_3) = v_1\mathbf{i} + v_2\mathbf{j} + v_3\mathbf{k}.$$



### Dot product of two vectors

**Definition 7** (Dot product of two vectors). Given the vectors  $\mathbf{u} = (u_1, \dots, u_n)$  and  $\mathbf{v} = (v_1, \dots, v_n)$  in  $\mathbb{R}^n$ , the *dot product* of  $\mathbf{u}$  and  $\mathbf{v}$  is

$$\mathbf{u} \cdot \mathbf{v} = u_1v_1 + \dots + u_nv_n.$$

**Example** Let  $\mathbf{u} = (3, 1)$  and  $\mathbf{v} = (2, 3)$  two vectors in  $\mathbb{R}^2$ , then the dot product of them is

$$\mathbf{u} \cdot \mathbf{v} = 3 \cdot 2 + 1 \cdot 3 = 9.$$

It holds that

$$\mathbf{u} \cdot \mathbf{v} = |\mathbf{u}||\mathbf{v}| \cos \alpha$$

where  $\alpha$  is the angle between the vectors.

### Parallel vectors

**Definition 8** (Parallel vectors). Two vectors  $\mathbf{u}$  and  $\mathbf{v}$  are *parallel* if there is a scalar  $a \in \mathbb{R}$  such that

$$\mathbf{u} = a\mathbf{v}.$$

**Example** The vectors  $\mathbf{u} = (-4, 2)$  and  $\mathbf{v} = (2, -1)$  in  $\mathbb{R}^2$  are parallel, as there is a scalar  $-2$  such that

$$\mathbf{u} = (-4, 2) = -2(2, -1) = -2\mathbf{v}.$$

### Orthogonal and orthonormal vectors

**Definition 9** (Orthogonal and orthonormal vectors). Two vectors  $\mathbf{u}$  and  $\mathbf{v}$  are *orthogonal* if their dot product is zero,

$$\mathbf{u} \cdot \mathbf{v} = 0.$$

If in addition both vectors are unit vectors,  $|\mathbf{u}| = |\mathbf{v}| = 1$ , then the vectors are *orthonormal*.

Orthogonal vectors are perpendicular, that is the angle between them is right.

**Example** The vectors  $\mathbf{u} = (2, 1)$  and  $\mathbf{v} = (-2, 4)$  in  $\mathbb{R}^2$  are orthogonal, as

$$\mathbf{u} \cdot \mathbf{v} = 2 \cdot (-2) + 1 \cdot 4 = 0,$$

but they are not orthonormal since  $|\mathbf{u}| = \sqrt{2^2 + 1^2} \neq 1$  and  $|\mathbf{v}| = \sqrt{(-2)^2 + 4^2} \neq 1$ .

The vectors  $\mathbf{i} = (1, 0)$  and  $\mathbf{j} = (0, 1)$  in  $\mathbb{R}^2$  are orthonormal, as

$$\mathbf{i} \cdot \mathbf{j} = 1 \cdot 0 + 0 \cdot 1 = 0, \quad |\mathbf{i}| = \sqrt{1^2 + 0^2} = 1, \quad |\mathbf{j}| = \sqrt{0^2 + 1^2} = 1.$$

## 1.2 Lines


### Vectorial equation of a straight line

**Definition 10** (Vectorial equation of a straight line). Given a point  $P = (p_1, \dots, p_n)$  and a vector  $\mathbf{v} = (v_1, \dots, v_n)$  of  $\mathbb{R}^n$ , the *vectorial equation of the line*  $l$  that passes through the point  $P$  with the direction of  $\mathbf{v}$  is

$$l : X = P + t\mathbf{v} = (p_1, \dots, p_n) + t(v_1, \dots, v_n) = (p_1 + tv_1, \dots, p_n + tv_n), \quad t \in \mathbb{R}.$$

**Example** Let  $l$  the line of  $\mathbb{R}^3$  that goes through  $P = (1, 1, 2)$  with the direction of  $\mathbf{v} = (3, 1, 2)$ , then the vectorial equation of  $l$  is

$$\begin{aligned} l : X = P + t\mathbf{v} &= (1, 1, 2) + t(3, 1, 2) = \\ &= (1 + 3t, 1 + t, 2 + 2t) \quad t \in \mathbb{R}. \end{aligned}$$



### Parametric and Cartesian equations of a line

From the vectorial equation of a line  $l : X = P + t\mathbf{v} = (p_1 + tv_1, \dots, p_n + tv_n)$  is easy to obtain the coordinates of the the points of the line with  $n$  *parametric equations*

$$x_1(t) = p_1 + tv_1, \dots, x_n(t) = p_n + tv_n$$

from where, if  $\mathbf{v}$  is a vector with non-null coordinates ( $v_i \neq 0 \forall i$ ), we can solve for  $t$  and equal the equations getting the *Cartesian equations*

$$\frac{x_1 - p_1}{v_1} = \dots = \frac{x_n - p_n}{v_n}$$



**Example** Given a line with vectorial equation  $l : X = (1, 1, 2) + t(3, 1, 2) = (1 + 3t, 1 + t, 2 + 2t)$  in  $\mathbb{R}^3$ , its parametric equations are

$$x(t) = 1 + 3t, \quad y(t) = 1 + t, \quad z(t) = 2 + 2t,$$

and the Cartesian equations are

$$\frac{x-1}{3} = \frac{y-1}{1} = \frac{z-2}{2}$$

### Point-slope equation of a line in the plane

In the particular case of the real plane  $\mathbb{R}^2$ , if we have a line with vectorial equation  $l : X = P + t\mathbf{v} = (x_0, y_0) + t(a, b) = (x_0 + ta, y_0 + tb)$ , its parametric equations are

$$x(t) = x_0 + ta, \quad y(t) = y_0 + tb$$

and its Cartesian equation is

$$\frac{x - x_0}{a} = \frac{y - y_0}{b}.$$

From this, moving  $b$  to the other side of the equation, we get

$$y - y_0 = \frac{b}{a}(x - x_0),$$

or renaming  $m = b/a$ ,


$$y - y_0 = m(x - x_0).$$

This equation is known as the *point-slope equation* of the line.

### Slope of a line in the plane

**Definition 11** (Slope of a line in the plane). Given a line  $l : X = P + t\mathbf{v}$  in the real plane  $\mathbb{R}^2$ , with direction vector  $\mathbf{v} = (a, b)$ , the *slope* of  $l$  is  $b/a$ .

Recall that given two points  $P = (x_1, y_1)$  and  $Q = (x_2, y_2)$  on the line  $l$ , we can take as a direction vector the vector from  $P$  to  $Q$ , with coordinates  $\vec{PQ} = Q - P = (x_2 - x_1, y_2 - y_1)$ . Thus, the slope of  $l$  is  $\frac{y_2 - y_1}{x_2 - x_1}$ , that is, the ratio between the changes in the vertical and horizontal axes.




## 1.3 Planes

### Vector equation of a plane in space

To get the equation of a plane in the real space  $\mathbb{R}^3$  we can take a point of the plane  $P = (x_0, y_0, z_0)$  and an orthogonal vector to the plane  $\mathbf{v} = (a, b, c)$ . Then, any point  $Q = (x, y, z)$  of the plane satisfies that the vector  $\vec{PQ} = (x - x_0, y - y_0, z - z_0)$  is orthogonal to  $\mathbf{v}$ , and therefore their dot product is zero.

$$\vec{PQ} \cdot \mathbf{v} = (x - x_0, y - y_0, z - z_0)(a, b, c) = a(x - x_0) + b(y - y_0) + c(z - z_0) = 0.$$

This equation is known as the *vector equation of the plane*.



### Scalar equation of a plane in space

From the vector equation of the plane we can get

$$a(x - x_0) + b(y - y_0) + c(z - z_0) = 0 \Leftrightarrow ax + by + cz = ax_0 + by_0 + cz_0,$$

that, renaming  $d = ax_0 + by_0 + cz_0$ , can be written as

$$ax + by + cz = d,$$

and is known as the *scalar equation of the plane*.

**Example** Given the point  $P = (2, 1, 1)$  and the vector  $\mathbf{v} = (2, 1, 2)$ , the vector equation of the plane that passes through  $P$  and is orthogonal to  $\mathbf{v}$  is

$$(x - 2, y - 1, z - 1)(2, 1, 2) = 2(x - 2) + (y - 1) + 2(z - 1) = 0,$$

and its scalar equation is

$$2x + y + 2z = 7.$$

## 2 Differential calculus with one real variable

### 2.1 Concept of derivative

#### Increment

**Definition 12** (Increment of a variable). An *increment* of a variable  $x$  is a change in the value of the variable; it is denoted  $\Delta x$ . The increment of a variable  $x$  along an interval  $[a, b]$  is given by

$$\Delta x = b - a.$$

**Definition 13** (Increment of a function). The *increment* of a function  $y = f(x)$  along an interval  $[a, b] \subseteq \text{Dom}(f)$  is given by

$$\Delta y = f(b) - f(a).$$

**Example** The increment of  $x$  along the interval  $[2, 5]$  is  $\Delta x = 5 - 2 = 3$  and the increment of the function  $y = x^2$  along the same interval is  $\Delta y = 5^2 - 2^2 = 21$ .

#### Average rate of change

The study of a function  $y = f(x)$  requires to understand how the function changes, that is, how the dependent variable  $y$  changes when we change the independent variable  $x$ .

**Definition 14** (Average rate of change). The *average rate of change* of a function  $y = f(x)$  in an interval  $[a, a + \Delta x] \subseteq \text{Dom}(f)$ , is the quotient between the increment of  $y$  and the increment of  $x$  in that interval, and is denoted by

$$\text{ARC } f[a, a + \Delta x] = \frac{\Delta y}{\Delta x} = \frac{f(a + \Delta x) - f(a)}{\Delta x}.$$


#### Average rate of change

*Example of the area of a square*

Let  $y = x^2$  be the function that measures the area of a metallic square of side length  $x$ .

If at any given time the side of the square is  $a$ , and we heat the square uniformly increasing the side by dilatation a quantity  $\Delta x$ , how much will the area of the square increase?

$$\begin{aligned} \Delta y &= f(a + \Delta x) - f(a) = (a + \Delta x)^2 - a^2 = \\ &= a^2 + 2a\Delta x + \Delta x^2 - a^2 = 2a\Delta x + \Delta x^2. \end{aligned}$$




What is the average rate of change in the interval  $[a, a + \Delta x]$ ?

$$\text{ARC } f[a, a + \Delta x] = \frac{\Delta y}{\Delta x} = \frac{2a\Delta x + \Delta x^2}{\Delta x} = 2a + \Delta x.$$

### Geometric interpretation of the average rate of change

The average rate of change of a function  $y = f(x)$  in an interval  $[a, a + \Delta x]$  is the slope of the *secant* line to the graph of  $f$  through the points  $(a, f(a))$  and  $(a + \Delta x, f(a + \Delta x))$ .



### Instantaneous rate of change

Often it is interesting to study the rate of change of a function, not in an interval, but in a point.

Knowing the tendency of change of a function in an instant can be used to predict the value of the function in nearby instants.

**Definition 15** (Instantaneous rate of change and derivative). The *instantaneous rate of change* of a function  $f(x)$  at a point  $x = a$ , is the limit of the average rate of change of  $f$  in the interval  $[a, a + \Delta x]$ , when  $\Delta x$  tends to 0, and is denoted by

$$\text{IRC } f(a) = \lim_{\Delta x \rightarrow 0} \text{ARC } f[a, a + \Delta x] = \lim_{\Delta x \rightarrow 0} \frac{\Delta y}{\Delta x} = \lim_{\Delta x \rightarrow 0} \frac{f(a + \Delta x) - f(a)}{\Delta x}$$

When this limit exists, the function  $f$  is said to be *differentiable* at the point  $a$ , and its value is called the *derivative* of  $f$  at  $a$ , and it is denoted  $f'(a)$  (Lagrange's notation) or  $\frac{df}{dx}(a)$  (Leibniz's notation).

### Instantaneous rate of change

*Example of the area of a square*

Let's take again the function  $y = x^2$  that measures the area of a metallic square of side length  $x$ .

If at any given time the side of the square is  $a$ , and we heat the square uniformly increasing the side, what is the tendency of change of the area in that moment?

$$\begin{aligned} \text{IRC } f(a) &= \lim_{\Delta x \rightarrow 0} \frac{\Delta y}{\Delta x} = \lim_{\Delta x \rightarrow 0} \frac{f(a + \Delta x) - f(a)}{\Delta x} = \\ &= \lim_{\Delta x \rightarrow 0} \frac{2a\Delta x + \Delta x^2}{\Delta x} = \lim_{\Delta x \rightarrow 0} 2a + \Delta x = 2a. \end{aligned}$$

Thus,

$$f'(a) = \frac{df}{dx}(a) = 2a,$$

indicating that the area of the square tends to increase the double of the side.

### Interpretation of the derivative

The derivative of a function  $f'(a)$  shows the growth rate of  $f$  at point  $a$ :

- $f'(a) > 0$  indicates an increasing tendency ( $y$  increases as  $x$  increases).
- $f'(a) < 0$  indicates a decreasing tendency ( $y$  decreases as  $x$  increases).


**Example** A derivative  $f'(a) = 3$  indicates that  $y$  tends to increase triple of  $x$  at point  $a$ . A derivative  $f'(a) = -0.5$  indicates that  $y$  tends to decrease half of  $x$  at point  $a$ .

### Geometric interpretation of the derivative

We have seen that the average rate of change of a function  $y = f(x)$  in an interval  $[a, a + \Delta x]$  is the slope of the *secant* line, but when  $\Delta x$  tends to 0, the secant line becomes the tangent line.


The instantaneous rate of change or derivative of a function  $y = f(x)$  at  $x = a$  is the slope of the *tangent line* to the graph of  $f$  at point  $(a, f(a))$ . Thus, the equation of the tangent line to the graph of  $f$  at the point  $(a, f(a))$  is

$$y - f(a) = f'(a)(x - a) \Leftrightarrow y = f(a) + f'(a)(x - a)$$



### Kinematic applications: Linear motion

Assume that the function  $y = f(t)$  describes the position of an object moving in the real line at time  $t$ . Taking as reference the coordinates origin  $O$  and the unitary vector  $\mathbf{i} = (1)$ , we can represent the position of the moving object  $P$  at every moment  $t$  with a vector  $\vec{OP} = x\mathbf{i}$  where  $x = f(t)$ .



**Remark** It also makes sense when  $f$  measures other magnitudes as the temperature of a body, the concentration of a gas, or the quantity of substance in a chemical reaction at every moment  $t$ .

**Kinematic interpretation of the average rate of change**

In this context, if we take the instants  $t = a$  and  $t = a + \Delta t$ , both in  $\text{Dom}(f)$ , the vector

$$\mathbf{v}_m = \frac{f(a + \Delta t) - f(a)}{\Delta t}$$

is known as the *average velocity* of the trajectory  $f$  in the interval  $[a, a + \Delta t]$ .

**Example** A vehicle makes a trip from Madrid to Barcelona. Let  $f(t)$  be the function that determine the position of the vehicle at every moment  $t$ . If the vehicle departs from Madrid (km 0) at 8:00 and arrives at Barcelona (km 600) at 14:00, then the average velocity of the vehicle in the path is

$$\mathbf{v}_m = \frac{f(14) - f(8)}{14 - 8} = \frac{600 - 0}{6} = 100 \text{ km/h.}$$

**Kinematic interpretation of the derivative**

In the same context of the linear motion, the derivative of the function  $f(t)$  at the moment  $t_0$  is the vector

$$\mathbf{v} = f'(a) = \lim_{\Delta t \rightarrow 0} \frac{f(a + \Delta t) - f(a)}{\Delta t},$$

that is known, as long as the limit exists, as the *instantaneous velocity* or simply *velocity* of the trajectory  $f$  at moment  $a$ .

That is, the derivative of the object position with respect to time is a vector field that is called *velocity along the trajectory*  $f$ .

**Example** Following with the previous example, what indicates the speedometer at any instant is the modulus of the instantaneous velocity vector at that moment.

**2.2 Algebra of derivatives****Properties of the derivative**

If  $y = c$ , is a constant function, then  $y' = 0$  at any point.

If  $y = x$ , is the identity function, then  $y' = 1$  at any point.

If  $u = f(x)$  and  $v = g(x)$  are two differentiable functions, then

- $(u + v)' = u' + v'$
- $(u - v)' = u' - v'$
- $(u \cdot v)' = u' \cdot v + u \cdot v'$
- $\left(\frac{u}{v}\right)' = \frac{u' \cdot v - u \cdot v'}{v^2}$

**Derivative of a composite function**

*The chain rule*

**Theorem 16** (Chain rule). *If the function  $y = f \circ g$  is the composition of two functions  $y = f(z)$  and  $z = g(x)$ , then*

$$(f \circ g)'(x) = f'(g(x))g'(x).$$

It is easy to prove this fact using the Leibniz notation

$$\frac{dy}{dx} = \frac{dy}{dz} \frac{dz}{dx} = f'(z)g'(x) = f'(g(x))g'(x).$$

**Example** If  $f(z) = \sin z$  and  $g(x) = x^2$ , then  $f \circ g(x) = \sin(x^2)$ . Applying the chain rule the derivative of the composite function is

$$(f \circ g)'(x) = f'(g(x))g'(x) = \cos(g(x))2x = \cos(x^2)2x.$$

On the other hand,  $g \circ f(z) = (\sin z)^2$ , and applying the chain rule again, its derivative is

$$(g \circ f)'(z) = g'(f(z))f'(z) = 2f(z) \cos z = 2 \sin z \cos z.$$

### Derivative of the inverse of a function

**Theorem 17** (Derivative of the inverse function). *Given a function  $y = f(x)$  with inverse  $x = f^{-1}(y)$ , then*

$$(f^{-1})'(y) = \frac{1}{f'(x)} = \frac{1}{f'(f^{-1}(y))},$$

*provided that  $f$  is differentiable at  $f^{-1}(y)$  and  $f'(f^{-1}(y)) \neq 0$ .*

Again, it is easy to prove this equality using the Leibniz notation

$$\frac{dx}{dy} = \frac{1}{dy/dx} = \frac{1}{f'(x)} = \frac{1}{f'(f^{-1}(y))}$$

### Derivative of the inverse of a function

*Example*

The inverse of the exponential function  $y = f(x) = e^x$  is the natural logarithm  $x = f^{-1}(y) = \ln y$ , so we can compute the derivative of the natural logarithm using the previous theorem and we get

$$(f^{-1})'(y) = \frac{1}{f'(x)} = \frac{1}{e^x} = \frac{1}{e^{\ln y}} = \frac{1}{y}.$$

**Example** Sometimes it is easier to apply the chain rule to compute the derivative of the inverse of a function. In this example, as  $\ln x$  is the inverse of  $e^x$ , we know that  $e^{\ln x} = x$ , so differentiating both sides and applying the chain rule to the left side we get

$$(e^{\ln x})' = x' \Leftrightarrow e^{\ln x}(\ln(x))' = 1 \Leftrightarrow (\ln(x))' = \frac{1}{e^{\ln x}} = \frac{1}{x}.$$

## 2.3 Analysis of functions

### Analysis of functions: increase and decrease

The main application of derivatives is to determine the variation (increase or decrease) of functions. For that we use the sign of the first derivative.

**Theorem 18.** *Let  $f(x)$  be a function with first derivative in an interval  $I \subseteq \mathbb{R}$ .*

- If  $\forall x \in I f'(x) > 0$  then  $f$  is increasing on  $I$ .
- If  $\forall x \in I f'(x) < 0$  then  $f$  is decreasing on  $I$ .

If  $f'(a) = 0$  then  $a$  is known as a *critical point* or *stationary point*. At this point the function can be increasing, decreasing or neither increasing nor decreasing.

**Example** The function  $f(x) = x^2$  has derivative  $f'(x) = 2x$ ; it is decreasing on  $\mathbb{R}^-$  as  $f'(x) < 0 \forall x \in \mathbb{R}^-$  and increasing on  $\mathbb{R}^+$  as  $f'(x) > 0 \forall x \in \mathbb{R}^+$ .


It has a critical point at  $x = 0$ , as  $f'(0) = 0$ ; at this point the function is neither increasing nor decreasing.

**Remark** A function can be increasing or decreasing on an interval and not have first derivative.

### Analysis of functions: increase and decrease

*Example*

Let us analyze the increase and decrease of the function  $f(x) = x^4 - 2x^2 + 1$ . Its first derivative is  $f'(x) = 4x^3 - 4x$ .



### Analysis of functions: relative extrema

As a consequence of the previous result we can also use the first derivative to determine the relative extrema of a function.

**Theorem 19** (First derivative test). Let  $f(x)$  be a function with first derivative in an interval  $I \subseteq \mathbb{R}$  and let  $a \in I$  be a critical point of  $f$  ( $f'(a) = 0$ ).

- If  $f'(x) > 0$  on an open interval extending left from  $a$  and  $f'(x) < 0$  on an open interval extending right from  $a$ , then  $f$  has a relative maximum at  $a$ .
- If  $f'(x) < 0$  on an open interval extending left from  $a$  and  $f'(x) > 0$  on an open interval extending right from  $a$ , then  $f$  has a relative minimum at  $a$ .
- If  $f'(x)$  has the same sign on both an open interval extending left from  $a$  and an open interval extending right from  $a$ , then  $f$  has an inflection point at  $a$ .


**Remark** A vanishing derivative is a necessary but not sufficient condition for the function to have a relative extrema at a point.

**Example** The function  $f(x) = x^3$  has derivative  $f'(x) = 3x^2$ ; it has a critical point at  $x = 0$ . However it does not have a relative extrema at that point, but an inflection point.



**Analysis of functions: relative extrema***Example*

Consider again the function  $f(x) = x^4 - 2x^2 + 1$  and let's analyze its relative extrema now. Its first derivative is  $f'(x) = 4x^3 - 4x$ .

**Analysis of functions: concavity**

The concavity of a function can be determined by the second derivative.

**Theorem 20.** Let  $f(x)$  be a function with second derivative in an interval  $I \subseteq \mathbb{R}$ .


- If  $\forall x \in I f''(x) > 0$  then  $f$  is concave up (convex) on  $I$ .
- If  $\forall x \in I f''(x) < 0$  then  $f$  is concave down (concave) on  $I$ .

**Example** The function  $f(x) = x^2$  has second derivative  $f''(x) = 2 > 0 \forall x \in \mathbb{R}$ , so it is concave up in all  $\mathbb{R}$ .

**Remark** A function can be concave up or down and not have second derivative.

**Analysis of functions: concavity***Example*


Let us analyze the concavity of the same function of previous examples  $f(x) = x^4 - 2x^2 + 1$ . Its second derivative is  $f''(x) = 12x^2 - 4$ .



## 2.4 Function approximation

### Approximating a function with the derivative

The tangent line to the graph of a function  $f(x)$  at  $x = a$  can be used to approximate  $f$  in a neighbourhood of  $a$ .



Thus, the increment of a function  $f(x)$  in an interval  $[a, a + \Delta x]$  can be approximated multiplying the derivative of  $f$  at  $a$  by the increment of  $x$

$$\Delta y \approx f'(a)\Delta x$$


### Approximating a function with the derivative

*Example of the area of a square*

In the previous example of the function  $y = x^2$  that measures the area of a metallic square of side  $x$ , if the side of the square is  $a$  and we increment it by a quantity  $\Delta x$ , then the increment on the area will be approximately

$$\Delta y \approx f'(a)\Delta x = 2a\Delta x.$$

In the figure below we can see that the error of this approximation is  $\Delta x^2$ , which is smaller than  $\Delta x$  when  $\Delta x$  tends to 0.



### Approximating a function by a polynomial

Another useful application of the derivative is the approximation of functions by polynomials. Polynomials are functions easy to calculate (sums and products) with very good properties:

- Defined in all the real numbers.
- Continuous.
- Differentiable of all orders with continuous derivatives.

#### Goal

Approximate a function  $f(x)$  by a polynomial  $p(x)$  near a point  $x = a$ .

### Approximating a function by a polynomial of order 0

A polynomial of degree 0 has equation

$$p(x) = c_0,$$

where  $c_0$  is a constant.


As the polynomial should coincide with the function  $f$  at  $a$ , it must satisfy

$$p(a) = c_0 = f(a).$$

Therefore, the polynomial of degree 0 that best approximates  $f$  near  $a$  is

$$p(x) = f(a).$$

### Approximating a function by a polynomial of order 0



### Approximating a function by a polynomial of order 1

A polynomial of degree 1 has equation

$$p(x) = c_0 + c_1x,$$

but it can also be written as

$$p(x) = c_0 + c_1(x - a).$$

Among all the polynomials of degree 1, the one that best approximates  $f$  near  $a$  is that which meets the following conditions

1.  $p$  and  $f$  coincide at  $a$ :  $p(a) = f(a)$ ,
2.  $p$  and  $f$  have the same rate of change at  $a$ :  $p'(a) = f'(a)$ .

The last condition guarantees that  $p$  and  $f$  have approximately the same tendency, but it requires the function  $f$  to be differentiable at  $a$ .

### The tangent line: Best approximating polynomial of order 1

Imposing the previous conditions we have


1.  $p(x) = c_0 + c_1(x - a) \Rightarrow p(a) = c_0 + c_1(a - a) = c_0 = f(a)$ ,
2.  $p'(x) = c_1 \Rightarrow p'(a) = c_1 = f'(a)$ .

Therefore, the polynomial of degree 1 that best approximates  $f$  near  $a$  is

$$p(x) = f(a) + f'(a)(x - a),$$

which turns out to be the tangent line to  $f$  at  $(a, f(a))$ .

### Approximating a function by a polynomial of order 1



### Approximating a function by a polynomial of order 2

A polynomial of degree 2 is a parabola with equation

$$p(x) = c_0 + c_1x + c_2x^2,$$

but it can also be written as

$$p(x) = c_0 + c_1(x - a) + c_2(x - a)^2.$$

Among all the polynomials of degree 2, the one that best approximates  $f(x)$  near  $a$  is that which meets the following conditions

1.  $p$  and  $f$  coincide at  $a$ :  $p(a) = f(a)$ ,
2.  $p$  and  $f$  have the same rate of change at  $a$ :  $p'(a) = f'(a)$ .
3.  $p$  and  $f$  have the same concavity at  $a$ :  $p''(a) = f''(a)$ .

The last condition requires the function  $f$  to be differentiable twice at  $a$ .

### Best approximating polynomial of order 2


Imposing the previous conditions we have

1.  $p(x) = c_0 + c_1(x - a) \Rightarrow p(a) = c_0 + c_1(a - a) = c_0 = f(a)$ ,
2.  $p'(x) = c_1 \Rightarrow p'(a) = c_1 = f'(a)$ .
3.  $p''(x) = 2c_2 \Rightarrow p''(a) = 2c_2 = f''(a) \Rightarrow c_2 = \frac{f''(a)}{2}$ .

Therefore, the polynomial of degree 2 that best approximates  $f$  near  $a$  is

$$p(x) = f(a) + f'(a)(x - a) + \frac{f''(a)}{2}(x - a)^2.$$

### Approximating a function by a polynomial of order 2



### Approximating a function by a polynomial of order $n$

A polynomial of degree  $n$  has equation

$$p(x) = c_0 + c_1x + c_2x^2 + \cdots + c_nx^n,$$

but it can also be written as

$$p(x) = c_0 + c_1(x-a) + c_2(x-a)^2 + \cdots + c_n(x-a)^n.$$

Among all the polynomials of degree  $n$ , the one that best approximate  $f(x)$  near  $a$  is that which meets the following  $n+1$  conditions

1.  $p(a) = f(a),$
2.  $p'(a) = f'(a),$
3.  $p''(a) = f''(a),$
- ...
- $n+1.$   $p^{(n)}(a) = f^{(n)}(a).$

Observe that these conditions require the function  $f$  to be differentiable  $n$  times at  $a$ .

### Coefficients calculation for the best approximating polynomial of order $n$

The successive derivatives of  $p$  are

$$\begin{aligned} p(x) &= c_0 + c_1(x-a) + c_2(x-a)^2 + \cdots + c_n(x-a)^n, \\ p'(x) &= c_1 + 2c_2(x-a) + \cdots + nc_n(x-a)^{n-1}, \\ p''(x) &= 2c_2 + \cdots + n(n-1)c_n(x-a)^{n-2}, \\ &\vdots \\ p^{(n)}(x) &= n(n-1)(n-2) \cdots 1c_n = n!c_n. \end{aligned}$$

Imposing the previous conditions we have

1.  $p(a) = c_0 + c_1(a-a) + c_2(a-a)^2 + \cdots + c_n(a-a)^n = c_0 = f(a),$
2.  $p'(a) = c_1 + 2c_2(a-a) + \cdots + nc_n(a-a)^{n-1} = c_1 = f'(a),$
3.  $p''(a) = 2c_2 + \cdots + n(n-1)c_n(a-a)^{n-2} = 2c_2 = f''(a) \Rightarrow c_2 = f''(a)/2,$
- ...
- $n+1.$   $p^{(n)}(a) = n!c_n = f^{(n)}(a) \Rightarrow c_n = \frac{f^{(n)}(a)}{n!}.$

**Taylor polynomial of order  $n$** 

**Definition 21** (Taylor polynomial). Given a function  $f(x)$  differentiable  $n$  times at  $a$ , the *Taylor polynomial* of order  $n$  of  $f$  at  $a$  is the polynomial with equation

$$\begin{aligned} p_{f,a}^n(x) &= f(a) + f'(a)(x-a) + \frac{f''(a)}{2}(x-a)^2 + \cdots + \frac{f^{(n)}(a)}{n!}(x-a)^n = \\ &= \sum_{i=0}^n \frac{f^{(i)}(a)}{i!}(x-a)^i. \end{aligned}$$

The Taylor polynomial of order  $n$  of  $f$  at  $a$  is the  $n$ th degree polynomial that best approximates  $f$  near  $a$ , as is the only one that meets the previous conditions.

**Taylor polynomial calculation**

*Example*

Let us approximate the function  $f(x) = \log x$  near the value 1 by a polynomial of order 3.

The equation of the Taylor polynomial of order 3 of  $f$  at  $a = 1$  is


$$p_{f,1}^3(x) = f(1) + f'(1)(x-1) + \frac{f''(1)}{2}(x-1)^2 + \frac{f'''(1)}{3!}(x-1)^3.$$

The derivatives of  $f$  at 1 up to order 3 are

$$\begin{array}{ll} f(x) = \log x & f(1) = \log 1 = 0, \\ f'(x) = 1/x & f'(1) = 1/1 = 1, \\ f''(x) = -1/x^2 & f''(1) = -1/1^2 = -1, \\ f'''(x) = 2/x^3 & f'''(1) = 2/1^3 = 2. \end{array}$$

And substituting into the polynomial equation we get

$$p_{f,1}^3(x) = 0 + 1(x-1) + \frac{-1}{2}(x-1)^2 + \frac{2}{3!}(x-1)^3 = \frac{2}{3}x^3 - \frac{3}{2}x^2 + 3x - \frac{11}{6}.$$

**Taylor polynomials of the logarithmic function**

**Maclaurin polynomial of order  $n$** 

The Taylor polynomial equation has a simpler form when the polynomial is calculated at 0. This special case of Taylor polynomial at 0 is known as the *Maclaurin polynomial*.

**Definition 22** (Maclaurin polynomial). Given a function  $f(x)$  differentiable  $n$  times at 0, the *Maclaurin polynomial* of order  $n$  of  $f$  is the polynomial with equation

$$\begin{aligned} p_{f,0}^n(x) &= f(0) + f'(0)x + \frac{f''(0)}{2}x^2 + \dots + \frac{f^{(n)}(0)}{n!}x^n = \\ &= \sum_{i=0}^n \frac{f^{(i)}(0)}{i!}x^i. \end{aligned}$$

**Maclaurin polynomial calculation**

*Example*

Let us approximate the function  $f(x) = \sin x$  near the value 0 by a polynomial of order 3.

The Maclaurin polynomial equation of order 3 of  $f$  is


$$p_{f,0}^3(x) = f(0) + f'(0)x + \frac{f''(0)}{2}x^2 + \frac{f'''(0)}{3!}x^3.$$

The derivatives of  $f$  at 0 up to order 3 are

$$\begin{array}{ll} f(x) = \sin x & f(0) = \sin 0 = 0, \\ f'(x) = \cos x & f'(0) = \cos 0 = 1, \\ f''(x) = -\sin x & f''(0) = -\sin 0 = 0, \\ f'''(x) = -\cos x & f'''(0) = -\cos 0 = -1. \end{array}$$

And substituting into the polynomial equation we get

$$p_{f,0}^3(x) = 0 + 1 \cdot x + \frac{0}{2}x^2 + \frac{-1}{3!}x^3 = x - \frac{x^3}{6}.$$

**Maclaurin polynomial of the sine function****Maclaurin polynomials of elementary functions**



$f(x)$	$p_{f,0}^n(x)$
$\sin x$	$x - \frac{x^3}{3!} + \frac{x^5}{5!} - \cdots + (-1)^k \frac{x^{2k-1}}{(2k-1)!}$ if $n = 2k$ or $n = 2k - 1$
$\cos x$	$1 - \frac{x^2}{2!} + \frac{x^4}{4!} - \cdots + (-1)^k \frac{x^{2k}}{(2k)!}$ if $n = 2k$ or $n = 2k + 1$
$\arctan x$	$x - \frac{x^3}{3} + \frac{x^5}{5} - \cdots + (-1)^k \frac{x^{2k-1}}{(2k-1)}$ if $n = 2k$ or $n = 2k - 1$
$e^x$	$1 + x + \frac{x^2}{2!} + \frac{x^3}{3!} + \cdots + \frac{x^n}{n!}$
$\log(1+x)$	$x - \frac{x^2}{2} + \frac{x^3}{3} - \cdots + (-1)^{n-1} \frac{x^n}{n}$

### Taylor remainder and Taylor formula

Taylor polynomials allow to approximate a function in a neighborhood of a value  $a$ , but most of the times there is an error in the approximation.

**Definition 23** (Taylor remainder). Given a function  $f(x)$  and its Taylor polynomial of order  $n$  at  $a$ ,  $p_{f,a}^n(x)$ , the *Taylor remainder* of order  $n$  of  $f$  at  $a$  is the difference between the function and the polynomial,

$$r_{f,a}^n(x) = f(x) - p_{f,a}^n(x).$$

The Taylor remainder measures the error in the approximation of  $f(x)$  by the Taylor polynomial and allow us to express the function as the Taylor polynomial plus the Taylor remainder

$$f(x) = p_{f,a}^n(x) + r_{f,a}^n(x).$$

This expression is known as the *Taylor formula* of order  $n$  or  $f$  at  $a$ .

It can be proved that

$$\lim_{h \rightarrow 0} \frac{r_{f,a}^n(a+h)}{h^n} = 0,$$

which means that the remainder  $r_{f,a}^n(a+h)$  is much smaller than  $h^n$ .

## 3 Integrals

### 3.1 Antiderivative of a function

#### Antiderivative of a function

**Definition 24** (Antiderivative of a function). Given a function  $f(x)$ , the function  $F(x)$  is an *antiderivative* or *primitive function* of  $f$  if it satisfies that  $F'(x) = f(x) \forall x \in \text{Dom}(f)$ .

**Example** The function  $F(x) = x^2$  is an antiderivative of the function  $f(x) = 2x$  as  $F'(x) = 2x$  on  $\mathbb{R}$ .

Roughly speaking, the calculus of antiderivatives is the reverse process of differentiation, and that is the reason for the name of antiderivative.

#### Indefinite integral of a function

As two functions that differs in a constant term have the same derivative, if  $F(x)$  is an antiderivative of  $f(x)$ , so will be any function of the form  $F(x) + k \forall k \in \mathbb{R}$ . This means that, when a function has an antiderivative, it has an infinite number of antiderivatives.

**Definition 25** (Indefinite integral). The *indefinite integral* of a function  $f(x)$  is the set of all its antiderivatives; it is denoted by

$$\int f(x) dx = F(x) + C$$

where  $F(x)$  is an antiderivative of  $f(x)$  and  $C$  is a constant.

**Example** The indefinite integral of the function  $f(x) = 2x$  is

$$\int 2x dx = x^2 + C.$$

#### Interpretation of the integral

We have seen in a previous chapter that the derivative of a function is the instantaneous rate of change of the function. Thus, if we know the instantaneous rate of change of the function at any point, we can compute the change of the function.

**Example** What is the space covered by an free falling object?

Assume that the only force acting upon an object drop is gravity, with an acceleration of  $9.8 \text{ m/s}^2$ . As acceleration is the the rate of change of the speed, that is constant at any moment, the antiderivative is the speed of the object,

$$v(t) = 9.8t \text{ m/s}$$

And as the speed is the rate of change of the space covered by object during the fall, the antiderivative of the speed is the space covered by the object,

$$s(t) = \int 9.8t dt = 9.8 \frac{t^2}{2}.$$

Thus, for instance, after 2 seconds, the covered space is  $s(2) = 9.8 \frac{2^2}{2} = 19.6 \text{ m}$ .

#### Linearity of integration

Given two integrable functions  $f(x)$  and  $g(x)$  and a constant  $k \in \mathbb{R}$ , it is satisfied that

$$1. \int (f(x) + g(x)) dx = \int f(x) dx + \int g(x) dx,$$

$$2. \int kf(x) dx = k \int f(x) dx.$$

This means that the integral of any linear combination of functions equals the same linear combination of the integrals of the functions.

## 3.2 Elementary integrals

### Elementary integrals

- $\int a dx = ax + C$ , with  $a$  constant.
- $\int x^n dx = \frac{x^{n+1}}{n+1} + C$  if  $n \neq -1$ .
- $\int \frac{1}{x} dx = \ln|x| + C$ .
- $\int e^x dx = e^x + C$ .
- $\int a^x dx = \frac{a^x}{\ln a} + C$ .
- $\int \sin x dx = -\cos x + C$ .
- $\int \cos x dx = \sin x + C$ .
- $\int \tan x dx = \ln|\sec x| + C$ .
- $\int \sec x dx = \ln|\sec x + \tan x| + C$ .
- $\int \csc x dx = \ln|\csc x - \cot x| + C$ .
- $\int \cot x dx = \ln|\sin x| + C$ .
- $\int \sec^2 x dx = \tan x + C$ .
- $\int \csc^2 x dx = -\cot x + C$ .
- $\int \sec x \tan x dx = \sec x + C$ .
- $\int \csc x \cot x dx = -\csc x + C$ .
- $\int \frac{dx}{\sqrt{a^2 - x^2}} = \arcsin \frac{x}{a} + C$ .
- $\int \frac{dx}{a^2 + x^2} = \frac{1}{a} \arctan \frac{x}{a} + C$ .
- $\int \frac{dx}{x\sqrt{x^2 - a^2}} = \frac{1}{a} \sec^{-1} \frac{x}{a} + C$ .
- $\int \frac{dx}{a^2 - x^2} = \frac{1}{2a} \ln \left| \frac{x+a}{x-a} \right| + C$ .

### 3.3 Techniques of integration

#### Techniques of integration

Unfortunately, unlike differential calculus, there is not a foolproof procedure to compute the antiderivative of a function. However, there are some techniques that allow to integrate some types of functions. The most common methods of integration are

- Integration by parts
- Integration by reduction
- Integration by substitution
- Integration of rational functions
- Integration of trigonometric functions

#### Integration by parts

Given two differentiable functions  $u(x)$  and  $v(x)$ , from the rule for differentiating a product we can get

$$\int u(x)v'(x) dx = u(x)v(x) - \int u'(x)v(x) dx,$$

or, writing  $u'(x)dx = du$  and  $v'(x)dx = dv$ ,

$$\int u dv = uv - \int v du.$$

To apply this method we have to choose the functions  $u$  and  $dv$  in a way so that the final integral is easier to compute than the original one.

**Example** To integrate  $\int x \sin x dx$  we have to choose  $u = x$  and  $dv = \sin x dx$ , so  $du = dx$  and  $v = -\cos x$ , getting

$$\int x \sin x dx = -x \cos x - \int (-\cos x) dx = -x \cos x + \sin x.$$

If we had chosen  $u = \sin x$  and  $dv = x dx$ , we would have got a more difficult integral.

#### Integration by reduction

The reduction technique is used when we have to apply the integration by parts several times.

If we want to compute the antiderivative  $I_n$  that depends on a natural number  $n$ , the reduction formulas allow us to write  $I_n$  as a function of  $I_{n-1}$ , that is, we have a recurrent relation

$$I_n = f(I_{n-1}, x, n)$$

so by computing the first antiderivative  $I_0$  we should be able to compute the others.

**Example** To compute  $I_n = \int x^n e^x dx$  applying integration by parts, we have to choose  $u = x^n$  y  $dv = e^x dx$ , so  $du = nx^{n-1} dx$  and  $v = e^x$ , getting

$$I_n = \int x^n e^x dx = x^n e^x - n \int x^{n-1} e^x dx = x^n e^x - n I_{n-1}.$$

Thus, for instance, for  $n = 3$  we have

$$\begin{aligned} \int x^3 e^x dx &= I_3 = x^3 e^x - 3I_2 = x^3 e^x - 3(x^2 e^x - 2I_1) = x^3 e^x - 3(x^2 e^x - (x e^x - I_0)) = \\ &= x^3 e^x - 3(x^2 e^x - (x e^x - e^x)) = e^x(x^3 - 3x^2 + 6x - 6). \end{aligned}$$

**Integration by substitution**

From the chain rule for differentiating the composition of two functions

$$f(g(x))' = f'(g(x))g'(x),$$

we can make a variable change  $u = g(x)$ , so  $du = g'(x)dx$ , and get

$$\int f'(g(x))g'(x) dx = \int f'(u) du = f(u) + C = f(g(x)) + C.$$

**Example** To compute the integral of  $\int \frac{1}{x \log x} dx$  we can make the substitution  $u = \log x$ , so  $du = \frac{1}{x} dx$ , and we have

$$\int \frac{dx}{x \log x} = \int \frac{1}{\log x} \frac{1}{x} dx = \int \frac{1}{u} du = \log |u| + C.$$

Finally, undoing the substitution we get

$$\int \frac{1}{x \log x} dx = \log |\log x| + C.$$

**Integration of rational functions***Partial fractions decomposition*

A rational function can be written as the sum of a polynomial (with an immediate antiderivative) plus a proper rational function, that is, a rational function in which the degree of the numerator is less than the degree of the denominator.

On the other hand, depending of the factorization of the denominator, a proper rational function can be expressed as a sum of simpler fractions of the following types

- Denominator with a single linear factor:  $\frac{A}{(x-a)}$
- Denominator with a linear factor repeated  $n$  times :  $\frac{A}{(x-a)^n}$
- Denominator with a single quadratic factor:  $\frac{Ax+B}{x^2+cx+d}$
- Denominator with a quadratic factor repeated  $n$  times:  $\frac{Ax+B}{(x^2+cx+d)^n}$

**Integration of rational functions***Antiderivatives of partial fractions*

Using the linearity of integration, we can compute the antiderivative of a rational function from the antiderivative of these partial fractions

$$\begin{aligned} \int \frac{A}{x-a} dx &= A \log |x-a| + C, \\ \int \frac{A}{(x-a)^n} dx &= \frac{-A}{(n-1)(x-a)^{n-1}} + C \text{ si } n \neq 1. \\ \int \frac{Ax+B}{x^2+cx+d} &= \frac{A}{2} \log |x^2+cx+d| + \frac{2B-Ac}{\sqrt{4d-c^2}} \arctan \frac{2x+c}{\sqrt{4d-c^2}} + C. \end{aligned}$$

**Integration of rational functions***Example of denominator with linear factors*

Consider the function  $f(x) = \frac{x^2 + 3x - 5}{x^3 - 3x + 2}$ .

The factorization of the denominator is  $x^3 - 3x + 2 = (x - 1)^2(x + 2)$ ; it has a single linear factor  $(x + 2)$  and a linear factor  $(x - 1)$ , repeated two times. In this case the decomposition in partial fractions is:

$$\begin{aligned}\frac{x^2 + 3x - 5}{x^3 - 3x + 2} &= \frac{A}{x - 1} + \frac{B}{(x - 1)^2} + \frac{C}{x + 2} = \\ &= \frac{A(x - 1)(x + 2) + B(x + 2) + C(x - 1)^2}{(x - 1)^2(x + 2)} = \\ &= \frac{(A + C)x^2 + (A + B - 2C)x + (-2A + 2B + C)}{(x - 1)^2(x + 2)}\end{aligned}$$

and equating the numerators we get  $A = 16/9$ ,  $B = -1/3$  and  $C = -7/9$ , so

$$\frac{x^2 + 3x - 5}{x^3 - 3x + 2} = \frac{16/9}{x - 1} + \frac{-1/3}{(x - 1)^2} + \frac{-7/9}{x + 2}.$$

Finally, integrating each partial fraction we have

$$\begin{aligned}\int \frac{x^2 + 3x - 5}{x^3 - 3x + 2} dx &= \int \frac{16/9}{x - 1} dx + \int \frac{-1/3}{(x - 1)^2} dx + \int \frac{-7/9}{x + 2} dx = \\ &= \frac{16}{9} \int \frac{1}{x - 1} dx - \frac{1}{3} \int (x - 1)^{-2} dx - \frac{7}{9} \int \frac{1}{x + 2} dx = \\ &= \frac{16}{9} \ln|x - 1| + \frac{1}{3(x - 1)} - \frac{7}{9} \ln|x + 2| + C.\end{aligned}$$

**Integration of rational functions***Example of denominator with simple quadratic factors*

Consider the function  $f(x) = \frac{x + 1}{x^2 - 4x + 8}$ .

In this case the denominator cannot be factorised as a product of linear factors, but we can write

$$x^2 - 4x + 8 = (x - 2)^2 + 4,$$

so

$$\begin{aligned}\int \frac{x + 1}{x^2 - 4x + 8} dx &= \int \frac{x - 2 + 3}{(x - 2)^2 + 4} dx = \\ &= \int \frac{x - 2}{(x - 2)^2 + 4} dx + \int \frac{3}{(x - 2)^2 + 4} dx = \\ &= \frac{1}{2} \ln|(x - 2)^2 + 4| + \frac{3}{2} \arctan\left(\frac{x - 2}{2}\right) + C.\end{aligned}$$

**Integration of trigonometric functions***Integration of  $\sin^n x \cos^m x$  with  $n$  or  $m$  odd*

If  $f(x) = \sin^n x \cos^m x$  with  $n$  or  $m$  odd, then we can make the substitution  $t = \sin x$  or  $t = \cos x$ , to convert the function into a polynomial.

**Example**

$$\int \sin^2 x \cos^3 x \, dx = \int \sin^2 x \cos^2 x \cos x \, dx = \int \sin^2 x (1 - \sin^2 x) \cos x \, dx,$$

and making the substitution  $t = \sin x$ , so  $dt = \cos x \, dx$ , we have

$$\int \sin^2 x (1 - \sin^2 x) \cos x \, dx = \int t^2 (1 - t^2) \, dt = \int t^2 - t^4 \, dt = \frac{t^3}{3} - \frac{t^5}{5} + C.$$

Finally, undoing the substitution we have

$$\int \sin^2 x \cos^3 x \, dx = \frac{\sin^3 x}{3} - \frac{\sin^5 x}{5} + C.$$

**Integration of trigonometric functions**

*Integration of  $\sin^n x \cos^m x$  with  $n$  and  $m$  even*

If  $f(x) = \sin^n x \cos^m x$  with  $n$  and  $m$  even, then we can make the following substitutions to simplify the integration

$$\begin{aligned}\sin^2 x &= \frac{1}{2}(1 - \cos(2x)) \\ \cos^2 x &= \frac{1}{2}(1 + \cos(2x)) \\ \sin x \cos x &= \frac{1}{2} \sin(2x)\end{aligned}$$

**Example**

$$\begin{aligned}\int \sin^2 x \cos^4 x \, dx &= \int (\sin x \cos x)^2 \cos^2 x \, dx = \int \left(\frac{1}{2} \sin(2x)\right)^2 \frac{1}{2} (1 + \cos(2x)) \, dx = \\ &= \frac{1}{8} \int \sin^2(2x) \, dx + \frac{1}{8} \int \sin^2(2x) \cos(2x) \, dx,\end{aligned}$$

the first integral is of the same type and the second one of the previous type, so

$$\int \sin^2 x \cos^4 x \, dx = \frac{1}{32}x - \frac{1}{32}\sin(2x) + \frac{1}{24}\sin^3(2x).$$

**Integration of trigonometric functions**

*Products of sines and cosines*

The equalities

$$\begin{aligned}\sin x \cos y &= \frac{1}{2}(\sin(x - y) + \sin(x + y)) \\ \sin x \sin y &= \frac{1}{2}(\cos(x - y) - \cos(x + y)) \\ \cos x \cos y &= \frac{1}{2}(\cos(x - y) + \cos(x + y))\end{aligned}$$

transform products in sums, simplifying the integration

**Example**

$$\begin{aligned}
\int \sin x \cos 2x \, dx &= \int \frac{1}{2} (\sin(x - 2x) + \sin(x + 2x)) \, dx = \\
&= \frac{1}{2} \int \sin(-x) \, dx + \frac{1}{2} \int \sin 3x \, dx = \\
&= \frac{1}{2} \cos(-x) - \frac{1}{6} \cos 3x + C.
\end{aligned}$$

**Integration of trigonometric functions**

*Rational functions of sines and cosines*

If  $f(x, y)$  is a rational function then the function  $f(\sin x, \cos x)$  can be transformed in an rational function of  $t$  with the following substitutions

$$\tan \frac{x}{2} = t \quad \sin x = \frac{2t}{1+t^2} \quad \cos x = \frac{1-t^2}{1+t^2} \quad dx = \frac{2}{1+t^2} dt.$$


**Example**

$$\int \frac{1}{\sin x} \, dx = \int \frac{1}{\frac{2t}{1+t^2}} \frac{2}{1+t^2} \, dt = \int \frac{1}{t} \, dt = \log |t| + C = \log \left| \tan \frac{x}{2} \right| + C.$$

**3.4 Definite integral****Definite integral**

**Definition 26** (Definite integral). Let  $f(x)$  be a function which is continuous on an interval  $[a, b]$ . Divide this interval into  $n$  subintervals of equal width  $\Delta x$  and choose an arbitrary point  $x_i$  from each interval. The *definite integral* of  $f$  from  $a$  to  $b$  is defined to be the limit

$$\int_a^b f(x) \, dx = \lim_{n \rightarrow \infty} \sum_{i=1}^n f(x_i) \Delta x.$$

**Definite integral**

**Theorem 27** (First fundamental theorem of Calculus). If  $f(x)$  is continuous on the interval  $[a, b]$  and  $F(x)$  is an antiderivative of  $f$  on  $[a, b]$ , then

$$\int_a^b f(x) \, dx = F(b) - F(a)$$



**Example.** Given the function  $f(x) = x^2$ , we have

$$\int_1^2 x^2 dx = \left[ \frac{x^3}{3} \right]_1^2 = \frac{2^3}{3} - \frac{1^3}{3} = \frac{7}{3}.$$

### Properties of the definite integral

Given two functions  $f(x)$  and  $g(x)$  integrable on  $[a, b]$  and  $k \in \mathbb{R}$  the following properties are satisfied:

- $\int_a^b (f(x) + g(x)) dx = \int_a^b f(x) dx + \int_a^b g(x) dx$  (linearity)
- $\int_a^b kf(x) dx = k \int_a^b f(x) dx$  (linearity)
- $\int_a^b f(x) dx \leq \int_a^b g(x) dx$  si  $f(x) \leq g(x) \forall x \in [a, b]$  (monotony)
- $\int_a^b f(x) dx = \int_a^c f(x) dx + \int_c^b f(x) dx$  for any  $c \in (a, b)$  (additivity)
- $\int_a^b f(x) dx = - \int_b^a f(x) dx$

## 3.5 Area calculation


### Area calculation

*Area between a positive function and the x axis*

If  $f(x)$  is an integrable function on the interval  $[a, b]$  and  $f(x) \geq 0 \forall x \in [a, b]$ , then the definite integral

$$\int_a^b f(x) dx$$

measures the area between the graph of  $f$  and the  $x$  axis on the interval  $[a, b]$ .




### Area calculation

*Area between a negative function and the x axis*

If  $f(x)$  is an integrable function on the interval  $[a, b]$  and  $f(x) \leq 0 \forall x \in [a, b]$ , then the area between the graph of  $f$  and the  $x$  axis on the interval  $[a, b]$  is

$$- \int_a^b f(x) dx.$$




### Area calculation

*Area between a function and the x axis*

In general, if  $f(x)$  is an integrable function on the interval  $[a, b]$ , no matter the sign of  $f$  on  $[a, b]$ , the area between the graph of  $f$  and the  $x$  axis on the interval  $[a, b]$  is

$$\int_a^b |f(x)| dx.$$




### Area calculation

*Area between two functions*

If  $f(x)$  and  $g(x)$  are two integrable functions on the interval  $[a, b]$ , then the area between the graph of  $f$  and  $g$  on the interval  $[a, b]$  is

$$\int_a^b |f(x) - g(x)| dx.$$



## 4 Ordinary Differential Equations

### 4.1 Ordinary Differential Equations

#### Ordinary Differential Equation

Often in Physics, Chemistry, Biology, Geometry, etc there arise equations that relate a function with its derivative, or successive derivatives.

**Definition 28** (Ordinary differential equation). An *ordinary differential equation* (O.D.E.) is a equation that relates an independent variable  $x$ , a function  $y(x)$  that depends on  $x$ , and the successive derivatives of  $y$ ,  $y', y'', \dots, y^{(n)}$ ; it can be written as

$$F(x, y, y', y'', \dots, y^{(n)}) = 0.$$

The *order* of a differential equation is the greatest order of the derivatives in the equation.

Thus, for instance, the equation  $y''' + \sin(x)y' = 2x$  is a differential equation of order 3.

#### Deducing a differential equation

To deduce a differential equation that explains a natural phenomenon is essential to understand what a derivative is and how to interpret it.

**Example** Newton's law of cooling states

*"The rate of change of the temperature of a body in a surrounding medium is proportional to the difference between the temperature of the body  $T$  and the temperature of the medium  $T_a$ ."*

The rate of change of the temperature is the derivative of temperature with respect to time  $dT/dt$ . Thus, Newton's law of cooling can be explained by the differential equation

$$\frac{dT}{dt} = k(T - T_a),$$

where  $k$  is a proportionality constant.

#### Solution of an ordinary differential equation

**Definition 29** (Solution of an ordinary differential equation). Given an ordinary differential equation  $F(x, y, y', y'', \dots, y^{(n)}) = 0$ , the function  $y = f(x)$  is a *solution of the ordinary differential equation* if it satisfies the equation, that is, if

$$F(x, f(x), f'(x), f''(x), \dots, f^{(n)}(x)) = 0.$$

The graph of a solution of the ordinary differential equation is known as *integral curve*.

Solving an ordinary differential equations consists on finding all its solutions in a given domain. For integral calculus is required.

The same manner than the indefinite integral is a family of antiderivatives, that differ in a constant term, after integrating an ordinary differential equation we get a family of solutions that differ in a constant. We can get particular solutions by giving values to this constant.

#### General solution of an ordinary differential equation

**Definition 30** (General solution of an ordinary differential equation). Given an ordinary differential equation  $F(x, y, y', y'', \dots, y^{(n)}) = 0$  of order  $n$ , the *general solution* of the differential equation is a family of functions

$$y = f(x, C_1, \dots, C_n),$$

depending on  $n$  constants, such that for any value of  $C_1, \dots, C_n$  we get a solution of the differential equation.

For every value of the constant we get *particular solution* of the differential equation. Thus, when a differential equation can be solved, it has infinite solutions.

Geometrically, the general solution of a differential equation corresponds to a family of integral curves of the differential equation.

Often, it is common to impose conditions to the solutions of a differential equation to reduce the number of solutions. In many cases, these conditions allow to determine the values of the constants in the general solution to get a particular solution.

### First order differential equations

In this chapter we are going to study first order differential equations

$$F(x, y, y') = 0.$$

The general solution of a first order differential equation is

$$y = f(x, C),$$

so to get a particular solution from the general one, it is enough to set the value of the constant  $C$ , and for that we only need to impose one initial condition.

**Definition 31** (Initial value problem). The group consisting of a first order differential equation and an initial condition is known as *initial value problem*:

$$\begin{cases} F(x, y, y') = 0, & \text{First order differential equation;} \\ y(x_0) = y_0, & \text{Initial condition.} \end{cases}$$

Solving an initial value problem consists in finding a solution of the first order differential equation that satisfies the initial condition.

### Solving an initial value problem

*Example*

Recall the first order differential equation of the Newton's law of cooling,

$$\frac{dT}{dt} = k(T - T_a),$$

where  $T$  is the temperature of the body and  $T_a$  is the temperature of the surrounding medium.

It is easy to check that the general solution of this equation is

$$T(t) = Ce^{kt} + T_a.$$

If we impose the initial condition that the temperature of the body at the initial instant is  $5^\circ\text{C}$ , that is,  $T(0) = 5$ , we have

$$T(0) = Ce^{k \cdot 0} + T_a = C + T_a = 5,$$

from where we get  $C = 5 - T_a$ , and this give us the particular solution


$$T(t) = (5 - T_a)e^{kt} + T_a.$$

**Integral curve of an initial value problem***Example*

If we assume in the previous example that the temperature of the surrounding medium is  $T_a = 0^\circ\text{C}$  and the cooling constant of the body is  $k = 1$ , the general solution of the differential equation is

$$T(t) = Ce^t,$$

that is a family of integral curves. Among all of them, only the one that passes through the point  $(0, 5)$  corresponds to the particular solution of the previous initial value problem.

**Existence and uniqueness of solutions**

**Theorem 32** (Existence and uniqueness of solutions of a first order ODE). *Given an initial value problem*

$$\begin{cases} y' = F(x, y); \\ y(x_0) = y_0; \end{cases}$$

*if  $F(x, y(x))$  is a function continuous on an open interval around the point  $(x_0, y_0)$ , then a solution of the initial value problem exists. If, in addition,  $\frac{\partial F}{\partial y}$  is continuous in an open interval around  $(x_0, y_0)$ , the solution is unique.*

Although this theorem guarantees the existence and uniqueness of a solution of a first order differential equation, it does not provide a method to compute it. In fact, there is not a general method to solve first order differential equations, but we will see how to solve some types:

- Separable differential equations
- Homogeneous differential equations
- Linear differential equations

**4.2 Separable differential equations****Separable differential equations**

**Definition 33** (Separable differential equation). *A separable differential equation is a first order differential equation that can be written as*

$$y'g(y) = f(x),$$

or what is the same,

$$g(y)dy = f(x)dx,$$

so the different variables are on different sides of the equality (the variables are separated).

The general solution for a separable differential equation comes after integrating both sides of the equation

$$\int g(y) dy = \int f(x) dx + C.$$

**Solving a separable differential equation***Example*

The differential equation of the Newton's law of cooling

$$\frac{dT}{dt} = k(T - T_a),$$

is a separable differential equation since it can be written as

$$\frac{1}{T - T_a} dT = k dt.$$

Integrating both sides of the equation we have

$$\int \frac{1}{T - T_a} dT = \int k dt \Leftrightarrow \log(T - T_a) = kt + C,$$

and solving for  $T$  we get the general solution of the equation

$$T(t) = e^{kt+C} + T_a = e^C e^{kt} + T_a = C e^{kt} + T_a,$$

rewriting  $C = e^C$  as an arbitrary constant.

**4.3 Homogeneous differential equations****Homogeneous functions**

**Definition 34** (Homogeneous function). A function  $f(x, y)$  is *homogeneous* of degree  $n$ , if it satisfies

$$f(kx, ky) = k^n f(x, y),$$

for any value  $k \in \mathbb{R}$ .

In particular, a homogeneous function of degree 0 always satisfies

$$f(kx, ky) = f(x, y).$$

Setting  $k = 1/x$  we have

$$f(x, y) = f\left(\frac{1}{x}x, \frac{1}{x}y\right) = f\left(1, \frac{y}{x}\right) = g\left(\frac{y}{x}\right).$$

This way, a homogeneous function of degree 0 always can be written as a function of  $u = y/x$ :

$$f(x, y) = g\left(\frac{y}{x}\right) = g(u).$$

**Homogeneous differential equations**

**Definition 35** (Homogeneous differential equation). A *homogeneous differential equation* is a first order differential equation that can be written as

$$y' = f(x, y),$$

where  $f(x, y)$  is a homogeneous function of degree 0.

We can solve a homogeneous differential equation by making the substitution

$$u = \frac{y}{x} \Leftrightarrow y = ux,$$

so the equation becomes

$$u'x + u = f(u),$$

that is a separable differential equation.

Once solved the separable differential equation, the substitution must be undone.

**Solving a homogeneous differential equation***Example*

Let us consider the following differential equation

$$4x - 3y + y'(2y - 3x) = 0.$$

Rewriting the equation in this way

$$y' = \frac{3y - 4x}{2y - 3x}$$

we can easily check that it is a homogeneous differential equation.

To solve this equation we have to do the substitution  $y = ux$ , and we get

$$u'x + u = \frac{3ux - 4x}{2ux - 3x} = \frac{3u - 4}{2u - 3}$$

that is a separable differential equation.

Separating the variables we have

$$u'x = \frac{3u - 4}{2u - 3} - u = \frac{-2u^2 + 6u - 4}{2u - 3} \Leftrightarrow \frac{2u - 3}{-2u^2 + 6u - 4} du = \frac{1}{x} dx.$$

**Solving a homogeneous differential equation***Example*

Now, integrating both sides of the equation we have

$$\begin{aligned} \int \frac{2u - 3}{-2u^2 + 6u - 4} du &= \int \frac{1}{x} dx \Leftrightarrow -\frac{1}{2} \log |u^2 - 3u + 2| = \log |x| + C \Leftrightarrow \\ &\Leftrightarrow \log |u^2 - 3u + 2| = -2 \log |x| - 2C, \end{aligned}$$

then, applying the exponential function to both sides and simplifying we get the general solution

$$u^2 - 3u + 2 = e^{-2 \log |x| - 2C} = \frac{e^{-2C}}{e^{\log |x|^2}} = \frac{C}{x^2},$$

rewriting the constant  $K = e^{-2C}$ .

Finally, undoing the initial substitution  $u = y/x$ , we arrive at the general solution of the homogeneous differential equation

$$\left(\frac{y}{x}\right)^2 - 3\frac{y}{x} + 2 = \frac{K}{x^2} \Leftrightarrow y^2 - 3xy + 2x^2 = K.$$

**4.4 Linear differential equations****Linear differential equations**

**Definition 36** (Linear differential equation). A linear differential equation is a first order differential equation that can be written as

$$y' + g(x)y = h(x).$$

To solve a linear differential equation we try to write the left side of the equation as the derivative of a product. For that we multiply both sides by a function  $f(x)$ , such that

$$f'(x) = g(x)f(x).$$

Thus, we get

$$\begin{aligned} y'f(x) + g(x)f(x)y &= h(x)f(x) \\ \Updownarrow \\ y'f(x) + f'(x)y &= h(x)f(x) \\ \Updownarrow \\ \frac{d}{dx}(yf(x)) &= h(x)f(x) \end{aligned}$$

### Solving a linear differential equation

Integrating both sides of the previous equation we get the solution

$$yf(x) = \int h(x)f(x) dx + C.$$

On the other hand, the unique function that satisfies  $f'(x) = g(x)f(x)$  is

$$f(x) = e^{\int g(x) dx},$$

so, substituting this function in the previous solution we arrive at the solution of the linear differential equation

$$ye^{\int g(x) dx} = \int h(x)e^{\int g(x) dx} dx + C,$$

or what is the same

$$y = e^{-\int g(x) dx} \left( \int h(x)e^{\int g(x) dx} dx + C \right).$$

### Solving a linear differential equation

*Example*

If in the differential equation of the Newton's law of cooling the temperature of the surrounding medium is a function of time  $T_a(t)$ , then the differential equation

$$\frac{dT}{dt} = k(T - T_a(t)),$$

is a linear differential equation since it can be written as

$$T' - kT = -kT_a(t),$$

where the independent term is  $-kT_a(t)$  and the coefficient of  $T$  is  $-k$ .

Substituting in the formula of the general solution of a linear differential equation we have

$$y = e^{-\int -k dt} \left( \int -kT_a(t)e^{\int -k dt} dt + C \right) = e^{kt} \left( -\int kT_a(t)e^{-kt} dt + C \right).$$



**Solving a linear differential equation***Example*

In the particular case that  $T_a(t) = t$ , and the proportionality constant  $k = 1$ , the general solution of the linear differential equation is

$$y = e^t \left( - \int t e^{-kt} dt + C \right) = e^t (e^{-t}(t + 1) + C) = Ce^t + t + 1.$$

If, in addition, we know that the temperature of the body at time  $t = 0$  is  $5^\circ\text{C}$ , that is, we have the initial condition  $T(0) = 5$ , then we can compute the value of the constant  $C$ ,

$$y(0) = Ce^0 + 0 + 1 = 5 \Leftrightarrow C + 1 = 5 \Leftrightarrow C = 4,$$

and we get the particular solution

$$y(t) = 4e^t + t + 1.$$

## 5 Several Variables Differentiable Calculus

### 5.1 Vector functions of a single real variable

#### Vector function of a single real variable

**Definition 37** (Vector function of a single real variable). A *vector function of a single real variable* or *vector field of a scalar variable* is a function that maps every scalar value  $t \in D \subseteq \mathbb{R}$  into a vector  $(x_1(t), \dots, x_n(t))$  in  $\mathbb{R}^n$ :

$$\begin{aligned} f: \mathbb{R} &\rightarrow \mathbb{R}^n \\ t &\rightarrow (x_1(t), \dots, x_n(t)) \end{aligned}$$

where  $x_i(t)$ ,  $i = 1, \dots, n$ , are real function of a single real variable known as *coordinate functions*.

The most common vector field of scalar variable are in the the real plane  $\mathbb{R}^2$ , where usually they are represented as


$$f(t) = x(t)\mathbf{i} + y(t)\mathbf{j},$$

and in the real space  $\mathbb{R}^3$ , where usually they are represented as


$$f(t) = x(t)\mathbf{i} + y(t)\mathbf{j} + z(t)\mathbf{k},$$

#### Graphic representation of vector fields

The graphic representation of a vector field in  $\mathbb{R}^2$  is a trajectory in the real plane.



The graphic representation of a vector field in  $\mathbb{R}^3$  is a trajectory in the real space.



#### Derivative of a vector field

The concept of derivative as the limit of the average rate of change of a function can be extended easily to vector fields.

**Definition 38** (Derivative of a vectorial field). A vectorial field  $f(t) = (x_1(t), \dots, x_n(t))$  is *differentiable* at a point  $t = a$  if the limit

$$\lim_{\Delta t \rightarrow 0} \frac{f(a + \Delta t) - f(a)}{\Delta t}.$$

exists. In such a case, the value of the limit is known as the *derivative* of the vector field at  $a$ , and it is written  $f'(a)$ .

### Derivative of a vector field

Many properties of real functions of a single real variable can be extended to vector fields through its component functions. Thus, for instance, the derivative of a vector field can be computed from the derivatives of its component functions.

**Theorem 39.** Given a vector field  $f(t) = (x_1(t), \dots, x_n(t))$ , if  $x_i(t)$  is differentiable at  $t = a$  for all  $i = 1, \dots, n$ , then  $f$  is differentiable at  $a$  and its derivative is

$$f'(a) = (x'_1(a), \dots, x'_n(a))$$

The proof for a vectorial field in  $\mathbb{R}^2$  is easy.

$$\begin{aligned} f'(a) &= \lim_{\Delta t \rightarrow 0} \frac{f(a + \Delta t) - f(a)}{\Delta t} = \lim_{\Delta t \rightarrow 0} \frac{(x(a + \Delta t), y(a + \Delta t)) - (x(a), y(a))}{\Delta t} = \\ &= \lim_{\Delta t \rightarrow 0} \left( \frac{x(a + \Delta t) - x(a)}{\Delta t}, \frac{y(a + \Delta t) - y(a)}{\Delta t} \right) = \\ &= \left( \lim_{\Delta t \rightarrow 0} \frac{x(a + \Delta t) - x(a)}{\Delta t}, \lim_{\Delta t \rightarrow 0} \frac{y(a + \Delta t) - y(a)}{\Delta t} \right) = (x'(a), y'(a)). \end{aligned}$$


### Kinematics: Curvilinear motion

The notion of derivative as a velocity along a trajectory in the real line can be generalized to a trajectory in any euclidean space  $\mathbb{R}^n$ .

In case of a two dimensional space  $\mathbb{R}^2$ , if  $f(t)$  describes the position of a moving object in the real plane at any time  $t$ , taking as reference the coordinates origin  $O$  and the unitary vectors  $\{\mathbf{i} = (1, 0), \mathbf{j} = (0, 1)\}$ , we can represent the position of the moving object  $P$  at every moment  $t$  with a vector  $\vec{OP} = x(t)\mathbf{i} + y(t)\mathbf{j}$ , where the coordinates

$$\begin{cases} x = x(t) \\ y = y(t) \end{cases} \quad t \in \text{Dom}(f)$$

are the *coordinate functions* of  $f$ .




### Velocity of a curvilinear motion in the plane

In this context the derivative of a trajectory  $f'(a) = (x'_1(a), \dots, x'_n(a))$  is the *velocity* vector of the trajectory  $f$  at moment  $t = a$ .

**Example** Given the trajectory  $f(t) = (\cos t, \sin t)$ ,  $t \in \mathbb{R}$ , whose image is the unit circumference centered in the coordinate origin, its coordinate functions are  $x(t) = \cos t$ ,  $y(t) = \sin t$ ,  $t \in \mathbb{R}$ , and its velocity is

$$\mathbf{v} = f'(t) = (x'(t), y'(t)) = (-\sin t, \cos t).$$

In the moment  $t = \pi/4$ , the object is in position  $f(\pi/4) = (\cos(\pi/4), \sin(\pi/4)) = (\sqrt{2}/2, \sqrt{2}/2)$  and it is moving with a velocity  $\mathbf{v} = f'(\pi/4) = (-\sin(\pi/4), \cos(\pi/4)) = (-\sqrt{2}/2, \sqrt{2}/2)$ .



Observe that the module of the velocity vector is always 1 as  $|\mathbf{v}| = \sqrt{(-\sin t)^2 + (\cos t)^2} = 1$ .

## 5.2 Tangent line to a trajectory

### Tangent line to a trajectory in the plane

#### Vectorial equation

Given a trajectory  $f(t)$  in the real plane, the vectors that are parallel to the velocity  $\mathbf{v}$  at a moment  $a$  are called *tangent vectors* to the trajectory  $f$  at the moment  $a$ , and the line passing through  $P = f(a)$  directed by  $\mathbf{v}$  is the tangent line to the graph of  $f$  at the moment  $a$ .

**Definition 40** (Tangent line to a trajectory). Given a trajectory  $f(t)$  in the real plane  $\mathbb{R}^2$ , the *tangent line* to the graph of  $f$  at  $a$  is the line with equation

$$\begin{aligned} l : (x, y) &= f(a) + tf'(a) = (x(a), y(a)) + t(x'(a), y'(a)) \\ &= (x(a) + tx'(a), y(a) + ty'(a)). \end{aligned}$$

### Tangent line to a trajectory in the plane

#### Example

We have seen that for the trajectory  $f(t) = (\cos t, \sin t)$ ,  $t \in \mathbb{R}$ , whose image is the unit circumference centered at the coordinate origin, the object position at the moment  $t = \pi/4$  is  $f(\pi/4) = (\sqrt{2}/2, \sqrt{2}/2)$  and its velocity  $\mathbf{v} = (-\sqrt{2}/2, \sqrt{2}/2)$ . Thus the equation of the tangent line to  $f$  at that moment is

$$\begin{aligned} l : (x, y) &= f(\pi/4) + t\mathbf{v} = \left( \frac{\sqrt{2}}{2}, \frac{\sqrt{2}}{2} \right) + t \left( \frac{-\sqrt{2}}{2}, \frac{\sqrt{2}}{2} \right) = \\ &= \left( \frac{\sqrt{2}}{2} - t\frac{\sqrt{2}}{2}, \frac{\sqrt{2}}{2} + t\frac{\sqrt{2}}{2} \right). \end{aligned}$$

### Tangent line to a trajectory in the plane

#### Cartesian and point-slope equations

From the vectorial equation of the tangent to a trajectory  $f(t)$  at the moment  $t = a$  we can get the coordinate functions

$$\begin{cases} x = x(a) + tx'(a) \\ y = y(a) + ty'(a) \end{cases} \quad t \in \mathbb{R},$$

and solving for  $t$  and equalling both equations we get the *Cartesian equation* of the tangent

$$\frac{x - x(a)}{x'(a)} = \frac{y - y(a)}{y'(a)},$$

if  $x'(a) \neq 0$  and  $y'(a) \neq 0$ .

From this equation it is easy to get the *point-slope equation* of the tangent

$$y - y(a) = \frac{y'(a)}{x'(a)}(x - x(a)).$$

### Tangent line to a trajectory in the plane

*Example of Cartesian and point-slope equations*

Using the vectorial equation of the tangent of the previous example

$$l : (x, y) = \left( \frac{\sqrt{2}}{2} - t \frac{\sqrt{2}}{2}, \frac{\sqrt{2}}{2} + t \frac{\sqrt{2}}{2} \right),$$

its Cartesian equation is

$$\frac{x - \sqrt{2}/2}{-\sqrt{2}/2} = \frac{y - \sqrt{2}/2}{\sqrt{2}/2}$$

and the point-slope equation is

$$y - \sqrt{2}/2 = \frac{-\sqrt{2}/2}{\sqrt{2}/2}(x - \sqrt{2}/2) \Rightarrow y = -x + \sqrt{2}.$$

### Normal line to a trajectory in the plane

We have seen that the tangent line to a trajectory  $f(t)$  at  $a$  is the line passing through the point  $P = f(a)$  directed by the velocity vector  $\mathbf{v} = f'(a) = (x'(a), y'(a))$ . If we take as direction vector a vector orthogonal to  $\mathbf{v}$ , we get another line that is known as *normal line* to the trajectory.

**Definition 41** (Normal line to a trajectory). Given a trajectory  $f(t)$  in the real plane  $\mathbb{R}^2$ , the *normal line* to the graph of  $f$  at moment  $t = a$  is the line with equation

$$l : (x, y) = (x(a), y(a)) + t(y'(a), -x'(a)) = (x(a) + ty'(a), y(a) - tx'(a)).$$

The Cartesian equation is

$$\frac{x - x(a)}{y'(a)} = \frac{y - y(a)}{-x'(a)},$$

and the point-slope equation is

$$y - y(a) = \frac{-x'(a)}{y'(a)}(x - x(a)).$$

The normal line is always perpendicular to the tangent line as their direction vectors are orthogonal.

### Normal line to a trajectory in the plane

*Example*

Considering again the trajectory of the unit circumference  $f(t) = (\cos t, \sin t)$ ,  $t \in \mathbb{R}$ , the normal line to the graph of  $f$  at moment  $t = \pi/4$  is

$$\begin{aligned} l : (x, y) &= (\cos(\pi/2), \sin(\pi/2)) + t(\cos(\pi/2), \sin(\pi/2)) = \\ &= \left( \frac{\sqrt{2}}{2}, \frac{\sqrt{2}}{2} \right) + t \left( \frac{\sqrt{2}}{2}, \frac{\sqrt{2}}{2} \right) = \left( \frac{\sqrt{2}}{2} + t \frac{\sqrt{2}}{2}, \frac{\sqrt{2}}{2} + t \frac{\sqrt{2}}{2} \right), \end{aligned}$$

the Cartesian equation is

$$\frac{x - \sqrt{2}/2}{\sqrt{2}/2} = \frac{y - \sqrt{2}/2}{\sqrt{2}/2},$$

and the point-slope equation is

$$y - \sqrt{2}/2 = \frac{\sqrt{2}/2}{\sqrt{2}/2}(x - \sqrt{2}/2) \Rightarrow y = x.$$

### Tangent and normal lines to a function

A particular case of tangent and normal lines to a trajectory are the tangent and normal lines to a function of one real variable. For every function  $y = f(x)$ , the trajectory that trace its graph is

$$g(x) = (x, f(x)) \quad x \in \mathbb{R},$$

and its velocity is

$$g'(x) = (1, f'(x)),$$

so that the tangent line to  $g$  at the moment  $a$  is

$$\frac{x - a}{1} = \frac{y - f(a)}{f'(a)} \Rightarrow y - f(a) = f'(a)(x - a),$$

and the normal line is

$$\frac{x - a}{f'(a)} = \frac{y - f(a)}{-1} \Rightarrow y - f(a) = \frac{-1}{f'(a)}(x - a),$$

### Tangent and normal lines to a function


#### Example

Given the function  $y = x^2$ , the trajectory that traces its graph is  $g(x) = (x, x^2)$  and its velocity is  $g'(x) = (1, 2x)$ . At the moment  $x = 1$  the trajectory passes through the point  $(1, 1)$  with a velocity  $(1, 2)$ . Thus, the tangent line at that moment is

$$\frac{x - 1}{1} = \frac{y - 1}{2} \Rightarrow y - 1 = 2(x - 1) \Rightarrow y = 2x - 1,$$

and the normal line is

$$\frac{x - 1}{2} = \frac{y - 1}{-1} \Rightarrow y - 1 = \frac{-1}{2}(x - 1) \Rightarrow y = \frac{-x}{2} + \frac{3}{2}.$$



### Tangent line to a trajectory in the space

The concept of tangent line to a trajectory can be easily extended from the real plane to the three-dimensional space  $\mathbb{R}^3$ .

If  $f(t) = (x(t), y(t), z(t))$ ,  $t \in \mathbb{R}$ , is a trajectory in the real space  $\mathbb{R}^3$ , then at the moment  $a$ , the moving object that follows this trajectory will be at the position  $P = (x(a), y(a), z(a))$  with a velocity  $\mathbf{v} = f'(t) = (x'(t), y'(t), z'(t))$ . Thus, the tangent line to  $f$  at this moment have the following vectorial equation

$$\begin{aligned} l : (x, y, z) &= (x(a), y(a), z(a)) + t(x'(a), y'(a), z'(a)) = \\ &= (x(a) + tx'(a), y(a) + ty'(a), z(a) + tz'(a)), \end{aligned}$$

and the Cartesian equations are

$$\frac{x - x(a)}{x'(a)} = \frac{y - y(a)}{y'(a)} = \frac{z - z(a)}{z'(a)},$$

provided that  $x'(a) \neq 0$ ,  $y'(a) \neq 0$  y  $z'(a) \neq 0$ .

### Tangent line to a trajectory in the space

*Example*

Given the trajectory  $f(t) = (\cos t, \sin t, t)$ ,  $t \in \mathbb{R}$  in the real space, at the moment  $t = \pi/2$  the trajectory passes through the point


$$f(\pi/2) = (\cos(\pi/2), \sin(\pi/2), \pi/2) = (0, 1, \pi/2),$$

with velocity

$$\mathbf{v} = f'(\pi/2) = (-\sin(\pi/2), \cos(\pi/2), 1) = (-1, 0, 1),$$

and the tangent line to the graph of  $f$  at that moment is

$$l : (x, y, z) = (0, 1, \pi/2) + t(-1, 0, 1) = (-t, 1, t + \pi/2).$$



### Normal plane to a trajectory in the space

In the three-dimensional space  $\mathbb{R}^3$ , the normal line to a trajectory is not unique. There are an infinite number of normal lines and all of them are in the normal plane.

If  $f(t) = (x(t), y(t), z(t))$ ,  $t \in \mathbb{R}$ , is a trajectory in the real space  $\mathbb{R}^3$ , then at the moment  $a$ , the moving object that follows this trajectory will be at the position  $P = (x(a), y(a), z(a))$  with a velocity  $\mathbf{v} = f'(t) = (x'(t), y'(t), z'(t))$ . Thus, using the velocity vector as normal vector the normal plane to  $f$  at this moment have the following vectorial equation

$$\begin{aligned} \Pi : (x - x(a), y - y(a), z - z(a))(x'(a), y'(a), z'(a)) &= 0 \\ &= x'(a)(x - x(a)) + y'(a)(y - y(a)) + z'(a)(z - z(a)) = 0. \end{aligned}$$

### Tangent line to a trajectory in the space

#### Example

For the trajectory of the previous example  $f(t) = (\cos t, \sin t, t)$ ,  $t \in \mathbb{R}$ , at the moment  $t = \pi/2$  the trajectory passes through the point


$$f(\pi/2) = (\cos(\pi/2), \sin(\pi/2), \pi/2) = (0, 1, \pi/2),$$

with velocity

$$\mathbf{v} = f'(\pi/2) = (-\sin(\pi/2), \cos(\pi/2), 1) = (-1, 0, 1),$$

and normal plane to the graph of  $f$  at that moment is

$$\Pi : \left(x - 0, y - 1, z - \frac{\pi}{2}\right) \cdot (-1, 0, 1) = 0 \Leftrightarrow -x + z - \frac{\pi}{2} = 0.$$



## 5.3 Functions of several variables

### Need for functions of several variables

A lot of problems in Geometry, Physics, Chemistry, Biology, etc. involve a variable that depend on two or more variables:

- The area of a triangle depends on two variables that are the base and height lengths.
- The volume of a perfect gas depends on two variables that are the pressure and the temperature.
- The way travelled by an object free falling depends on a lot of variables: the time, the area of the cross section of the object, the latitude and longitude of the object, the height above the sea level, the air pressure, the air temperature, the speed of wind, etc.

These dependencies are expressed with functions of several variables.

### Functions of several real variables

**Definition 42** (Functions of several real variables). A function of  $n$  real variables or a scalar field from a set  $A_1 \times \cdots \times A_n \subseteq \mathbb{R}^n$  in a set  $B \subseteq \mathbb{R}$ , is a relation that maps any tuple  $(a_1, \dots, a_n) \in A_1 \times \cdots \times A_n$  into a unique element of  $B$ , denoted by  $f(a_1, \dots, a_n)$ , that is known as the *image* of  $(a_1, \dots, a_n)$  by  $f$ .

$$\begin{aligned} f : A_1 \times \cdots \times A_n &\longrightarrow B \\ (a_1, \dots, a_n) &\longrightarrow f(a_1, \dots, a_n) \end{aligned}$$

#### Example

- The area of a triangle is a real function of two real variables

$$f(x, y) = \frac{xy}{2}.$$


- The volume of a perfect gas is a real function of two real variables

$$v = f(t, p) = \frac{nRt}{p}, \quad \text{with } n \text{ and } R \text{ constants.}$$




**Graph of a function of two variables**

The graph of a function of two variables  $f(x, y)$  is a surface in the real space  $\mathbb{R}^3$  where every point of the surface has coordinates  $(x, y, z)$ , with  $z = f(x, y)$ .

**Graph of a function of two variables**


*Area of a triangle*

The function  $f(x, y) = \frac{xy}{2}$  that measures the area of a triangle of base  $x$  and height  $y$  has the graph below.

**Graph of a function of two variables**

*Wave of a water drop*

The function  $f(x, y) = \frac{\sin(x^2 + y^2)}{\sqrt{x^2 + y^2}}$  has the peculiar graph below.



**Level set of a scalar field**

**Definition 43** (Level set). Given a scalar field  $f : \mathbb{R}^n \rightarrow \mathbb{R}$ , the *level set*  $c$  of  $f$  is the set

$$C_{f,c} = \{(x_1, \dots, x_n) : f(x_1, \dots, x_n) = c\},$$

that is, a set where the function takes on the constant value  $c$ .


**Level set of a scalar field**

*Example*


Given the scalar field  $f(x, y) = x^2 + y^2$  and the point  $P = (1, 1)$ , the level set of  $f$  that includes  $P$  is

$$C_{f,2} = \{(x, y) : f(x, y) = f(1, 1) = 2\} = \{(x, y) : x^2 + y^2 = 2\},$$

that is the circumference of radius  $\sqrt{2}$  centered at the origin.


**Level set of a scalar field**

Topographic maps



Level curves correspond to points with the same atmospheric pressure.

Weather maps (Isobars)



Level curves correspond to points with the same atmospheric pressure.

### Partial functions

**Definition 44** (Partial function). Given a scalar field  $f : \mathbb{R}^n \rightarrow \mathbb{R}$ , an  $i$ -th *partial function* of  $f$  is any function  $f_i : \mathbb{R} \rightarrow \mathbb{R}$  that results of substituting all the variables of  $f$  by constants, except the  $i$ -th variable, that is:

$$f_i(x) = f(c_1, \dots, c_{i-1}, x, c_{i+1}, \dots, c_n),$$

with  $c_j$  ( $j = 1, \dots, n, j \neq i$ ) constants.

**Example** If we take the function that measures the area of a triangle

$$f(x, y) = \frac{xy}{2},$$

and set the value of the base to  $x = c$ , then the area of the triangle depends only of the height, and  $f$  becomes a function of one variable, that is the partial function

$$f_1(y) = f(c, y) = \frac{cy}{2}, \quad \text{with } c \text{ constant.}$$

## 5.4 Partial derivative notion

### Variation of a function with respect to a variable

We can measure the variation of a scalar field with respect to each of its variables in the same way that we measured the variation of a one-variable function.

Let  $z = f(x, y)$  be a scalar field of  $\mathbb{R}^2$ . If we are at point  $(x_0, y_0)$  and we increase the value of  $x$  a quantity  $\Delta x$ , then we move in the direction of the  $x$ -axis from the point  $(x_0, y_0)$  to the point  $(x_0 + \Delta x, y_0)$ , and the variation of the function is

$$\Delta z = f(x_0 + \Delta x, y_0) - f(x_0, y_0).$$

Thus, the rate of change of the function with respect to  $x$  along the interval  $[x_0, x_0 + \Delta x]$  is given by the quotient

$$\frac{\Delta z}{\Delta x} = \frac{f(x_0 + \Delta x, y_0) - f(x_0, y_0)}{\Delta x}.$$

### Instantaneous rate of change of a scalar field with respect to a variable

If instead of measuring the rate of change in an interval, we measure the rate of change in a point, that is, when  $\Delta x$  approaches 0, then we get the instantaneous rate of change:

$$\lim_{\Delta x \rightarrow 0} \frac{\Delta z}{\Delta x} = \lim_{\Delta x \rightarrow 0} \frac{f(x_0 + \Delta x, y_0) - f(x_0, y_0)}{\Delta x}.$$


The value of this limit, if exists, it is known as the *partial derivative* of  $f$  with respect to the variable  $x$  at the point  $(x_0, y_0)$ ; it is written as

$$\frac{\partial f}{\partial x}(x_0, y_0).$$

This partial derivative measures the instantaneous rate of change of  $f$  at the point  $P = (x_0, y_0)$  when  $P$  when we move  $P$  in the  $x$ -axis direction.

### Geometric interpretation of partial derivatives

Geometrically, a two-variable function  $z = f(x, y)$  defines a surface. If we cut this surface with a plane of equation  $y = y_0$  (that is, the plane where  $y$  is the constant  $y_0$ ) the intersection is a curve, and the partial derivative of  $f$  with respect to  $x$  at  $(x_0, y_0)$  is the slope of the tangent line to that curve at  $x = x_0$ .



### Partial derivative

The concept of partial derivative can be extended easily from two-variable function to  $n$ -variables functions.

**Definition 45** (Partial derivative). Given a  $n$ -variables function  $f(x_1, \dots, x_n)$ ,  $f$  is *partially differentiable* with respect to the variable  $x_i$  at the point  $a = (a_1, \dots, a_n)$  if exists the limit

$$\lim_{\Delta x_i \rightarrow 0} \frac{f(a_1, \dots, a_{i-1}, a_i + \Delta x_i, a_{i+1}, \dots, a_n) - f(a_1, \dots, a_{i-1}, a_i, a_{i+1}, \dots, a_n)}{\Delta x_i}.$$

In such a case, the value of the limit is known as *partial derivative* of  $f$  with respect to  $x_i$  at  $a$ ; it is denoted

$$f'_{x_i}(a) = \frac{\partial f}{\partial x_i}(a).$$

**Remark** The definition of derivative for one-variable functions is a particular case of this definition for  $n = 1$ .

### Partial derivatives computation

When we measure the variation of  $f$  with respect to a variable  $x_i$  at the point  $a = (a_1, \dots, a_n)$ , the other variables remain constant. Thus, if we can consider the  $i$ -th partial function

$$f_i(x_i) = f(a_1, \dots, a_{i-1}, x_i, a_{i+1}, \dots, a_n),$$

the partial derivative of  $f$  with respect to  $x_i$  can be computed differentiating this function:

$$\frac{\partial f}{\partial x_i}(a) = f'_i(a_i).$$

### Rule

To differentiate partially  $f(x_1, \dots, x_n)$  with respect to the variable  $x_i$ , you have to differentiate  $f$  as a function of the variable  $x_i$ , considering the other variables as constants.

### Partial derivatives computation

*Example of the volume of a perfect gas*

Consider the function that measures the volume of a perfect gas

$$v(t, p) = \frac{nRt}{p},$$

where  $t$  is the temperature,  $p$  the pressure and  $n$  and  $R$  are constants.

The instantaneous rate of change of the volume with respect to the pressure is the partial derivative of  $v$  with respect to  $p$ . To compute this derivative we have to think in  $t$  as a constant and differentiate  $v$  as if the unique variable was  $p$ :

$$\frac{\partial v}{\partial p}(t, p) = \frac{d}{dp} \left( \frac{nRt}{p} \right)_{t=\text{cte}} = \frac{-nRt}{p^2}.$$

In the same way, the instantaneous rate of change of the volume with respect to the temperature is the partial derivative of  $v$  with respect to  $t$ :

$$\frac{\partial v}{\partial t}(t, p) = \frac{d}{dt} \left( \frac{nRt}{p} \right)_{p=\text{cte}} = \frac{nR}{p}.$$

## 5.5 Gradient

### Gradient

**Definition 46** (Gradient). Given a scalar field  $f(x_1, \dots, x_n)$ , the *gradiente* of  $f$ , denoted by  $\nabla f$ , is a function that maps every point  $a = (a_1, \dots, a_n)$  to a vector with coordinates the partial derivatives of  $f$  at  $a$ ,

$$\nabla f(a) = \left( \frac{\partial f}{\partial x_1}(a), \dots, \frac{\partial f}{\partial x_n}(a) \right).$$

Later we will show that the gradient in a point is a vector with the magnitude and direction of the maximum rate of change of the function in that point. Thus,  $\nabla f(a)$  shows the direction of maximum increase of  $f$  at  $a$ , while  $-\nabla f(a)$  show the direction of maximum decrease of  $f$  at  $a$ .

**Gradient computation**

*Example with a temperature function*

After heating a surface, the temperature  $t$  (in °C) at each point  $(x, y, z)$  (in m) of the surface is given by the function

$$t(x, y, z) = \frac{x}{y} + z^2.$$

In what direction will increase the temperature faster at point  $(2, 1, 1)$  of the surface? What magnitude will the maximum increase of temperature have?

The direction of maximum increase of the temperature is given by the gradient

$$\nabla t(x, y, z) = \left( \frac{\partial t}{\partial x}(x, y, z), \frac{\partial t}{\partial y}(x, y, z), \frac{\partial t}{\partial z}(x, y, z) \right) = \left( \frac{1}{y}, \frac{-x}{y^2}, 2z \right).$$

At point  $(2, 1, 1)$  the direction is given by the vector

$$\nabla t(2, 1, 1) = \left( \frac{1}{1}, \frac{-2}{1^2}, 2 \cdot 1 \right) = (1, -2, 2),$$

and its magnitude is

$$|\nabla f(2, 1, 1)| = |\sqrt{1^2 + (-2)^2 + 2^2}| = |\sqrt{9}| = 3 \text{ °C/m}.$$

**5.6 Second order partial derivatives****Second order partial derivatives**

As the partial derivatives of a function are also functions of several variables we can differentiate partially each of them.

If a function  $f(x_1, \dots, x_n)$  has a partial derivative  $f'_{x_i}(x_1, \dots, x_n)$  with respect to the variable  $x_i$  in a set  $A$ , then we can differentiate partially again  $f'_{x_i}$  with respect to the variable  $x_j$ . This second derivative, when exists, is known as *second order partial derivative* of  $f$  with respect to the variables  $x_i$  and  $x_j$ ; it is written as

$$\frac{\partial^2 f}{\partial x_j \partial x_i} = \frac{\partial}{\partial x_j} \left( \frac{\partial f}{\partial x_i} \right).$$

In the same way we can define higher order partial derivatives.

**Second order partial derivatives computation**

*Example*

The two-variables function

$$f(x, y) = x^y$$

has 4 second order partial derivatives:

$$\begin{aligned} \frac{\partial^2 f}{\partial x^2}(x, y) &= \frac{\partial}{\partial x} \left( \frac{\partial f}{\partial x}(x, y) \right) = \frac{\partial}{\partial x} (yx^{y-1}) = y(y-1)x^{y-2}, \\ \frac{\partial^2 f}{\partial y \partial x}(x, y) &= \frac{\partial}{\partial y} \left( \frac{\partial f}{\partial x}(x, y) \right) = \frac{\partial}{\partial y} (yx^{y-1}) = x^{y-1} + yx^{y-1} \log x, \\ \frac{\partial^2 f}{\partial x \partial y}(x, y) &= \frac{\partial}{\partial x} \left( \frac{\partial f}{\partial y}(x, y) \right) = \frac{\partial}{\partial x} (x^y \log x) = yx^{y-1} \log x + x^y \frac{1}{x}, \\ \frac{\partial^2 f}{\partial y^2}(x, y) &= \frac{\partial}{\partial y} \left( \frac{\partial f}{\partial y}(x, y) \right) = \frac{\partial}{\partial y} (x^y \log x) = x^y (\log x)^2. \end{aligned}$$

## 5.7 Matriz hessiana

### Hessian matrix and Hessian

**Definition 47** (Hessian matrix). Given a scalar field  $f(x_1, \dots, x_n)$ , with second order partial derivatives at the point  $a = (a_1, \dots, a_n)$ , the *Hessian matrix* of  $f$  at  $a$ , denoted by  $\nabla^2 f(a)$ , is the matrix

$$\nabla^2 f(a) = \begin{pmatrix} \frac{\partial^2 f}{\partial x_1^2}(a) & \frac{\partial^2 f}{\partial x_1 \partial x_2}(a) & \cdots & \frac{\partial^2 f}{\partial x_1 \partial x_n}(a) \\ \frac{\partial^2 f}{\partial x_2 \partial x_1}(a) & \frac{\partial^2 f}{\partial x_2^2}(a) & \cdots & \frac{\partial^2 f}{\partial x_2 \partial x_n}(a) \\ \vdots & \vdots & \ddots & \vdots \\ \frac{\partial^2 f}{\partial x_n \partial x_1}(a) & \frac{\partial^2 f}{\partial x_n \partial x_2}(a) & \cdots & \frac{\partial^2 f}{\partial x_n^2}(a) \end{pmatrix}$$

The determinant of this matrix is known as *Hessian* of  $f$  at  $a$ ; it is denoted  $Hf(a) = |\nabla^2 f(a)|$ .

### Hessian matrix computation

Consider again the two-variables function

$$f(x, y) = x^y.$$

Its Hessian matrix is

$$\nabla^2 f(x, y) = \begin{pmatrix} \frac{\partial^2 f}{\partial x^2} & \frac{\partial^2 f}{\partial x \partial y} \\ \frac{\partial^2 f}{\partial y \partial x} & \frac{\partial^2 f}{\partial y^2} \end{pmatrix} = \begin{pmatrix} y(y-1)x^{y-2} & x^{y-1}(y \log x + 1) \\ x^{y-1}(y \log x + 1) & x^y(\log x)^2 \end{pmatrix}.$$

At point  $(1, 2)$  is

$$\nabla^2 f(1, 2) = \begin{pmatrix} 2(2-1)1^{2-2} & 1^{2-1}(2 \log 1 + 1) \\ 1^{2-1}(2 \log 1 + 1) & 1^2(\log 1)^2 \end{pmatrix} = \begin{pmatrix} 2 & 1 \\ 1 & 0 \end{pmatrix}.$$

And its Hessian is

$$Hf(1, 2) = \begin{vmatrix} 2 & 1 \\ 1 & 0 \end{vmatrix} = 2 \cdot 0 - 1 \cdot 1 = -1.$$