

## EXAM OF STATISTICS (PROBABILITY AND RANDOM VARIABLES)

2nd Physiotherapy

Version A

May, 27 2019

Name:

DNI:

Group:

**Duration:** 1 hour and 15 minutes.

- (4 pts.) 1. It has been observed that the concentration of a metabolite in urine can be used as a diagnostic test for a disease. The concentration (in mg/dl) in healthy individuals follows a normal distribution with mean 90 and standard deviation 8, while in sick individuals follows a normal distribution with mean 120 and standard deviation 10.
- (a) If the cut-off point is set at 105 mg/dl (positive above and negative below), what is the sensitivity and the specificity of the test?
  - (b) If the cut-off point is set at 105 mg/dl and we assume a prevalence of 10%, what is the probability of a correct diagnostic?
  - (c) If we want a sensitivity of 95%, where must we set the cut-off point? What would the specificity of the test be ?
- (3 pts.) 2. Let  $A$  and  $B$  be two events of a random experiment, such that  $A$  is three times as likely as  $B$ ,  $P(A \cup B) = 0.8$  and  $P(A \cap B) = 0.2$ .
- (a) Compute  $P(A)$  and  $P(B)$ .
  - (b) Compute  $P(A - B)$  and  $P(B - A)$ .
  - (c) Compute  $P(\bar{A} \cup \bar{B})$  and  $P(\bar{A} \cap \bar{B})$ .
  - (d) Compute  $P(A|B)$  and  $P(B|A)$ .
  - (e) Are  $A$  and  $B$  independent?
- (3 pts.) 3. The employees of a courier company send an average of 246.2 messages in a period of 12 hours. It is also known that the mean of messages sent by males is 256.2 and by females is 237.4 in the same period.
- (a) Compute the probability that a random person of the company sends 5 messages in a period of half an hour.
  - (b) If we draw randomly 10 women of this company, what is the probability that at least 3 of them sends more than one message in a period of one hour?
  - (c) If we draw randomly 100 men of this company, what is the probability that none of them sends less than 2 messages in a period of a quarter of an hour?