EXAM OF STATISTICS
(PROBABILITY AND RANDOM VARIABLES)

2nd Physiotherapy **Version A** **May, 25 2020**

Name: DNI: Group:

Duration: 1 hour.

(2.5 pts.) 1. A hospital orders a DNA compatibility test to three labs A, B and C. Lab A performs 40 test a day, lab B 50, and lab C 60. It is known that the probability of a wrong diagnose is 20% in lab A, 18% in lab B and 22% in lab C. If we select a random test of the hospital,

(a) Compute the probability of wrong diagnose in that test.

(b) If the test is wrong, what is the probability that it has been performed by lab B?

(c) If the test is right, which lab is more likely to have performed the test?

(2 pts.) 2. An epidemiological study tries to determine the effectiveness of face masks to prevent the COVID19. In a sample 4000 persons without the virus and 1000 persons with it were selected. I was observed that in the group of infected people 120 had used face masks in the two previous weeks, while in the non-infected group, 1250 had used face masks in the two previous weeks.

(a) Compute the relative risk of been infected with face masks.

(b) Compute the odds ratio of been infected with face masks.

(c) Which association measure is more reliable?

(2.5 pts.) 3. During the COVID19 quarantine a telephone exchange with 4 telephone operators received an average of 12 calls per day. Assuming that the calls are equally distributed among the operators,

(a) Compute the probability that an operator received more than 3 calls a day.

(b) Compute the probability that all the the operators received some call a day.

(3 pts.) 4. In a course with 200 students the score of a test to measure the intelligence quotient follows a normal distribution. After applying the test to the students 10 of them got a score above 130 and 30 of them a score below 60.

(a) Compute the mean and the standard deviation of the score.

(b) How many students will have a score between 90 and 95?

(c) Compute the limits of the interval centered at the mean that accumulates 95% of the scores.