

alignaa

CLEAR ALIGNERS: A CLINICAL CASE

Dr. Tanvi Bore

BDS, MDS, Specialist in Orthodontics
MUMBAI, INDIA


Dr. Ganesh Shinde

BDS, Dental Surgeon, Specialist in Smile Designing
PUNE, INDIA

A 20-year-old female patient presents to the office accompanied by her Sister who reports that the patient "has anterior cross bite".

FACIAL ASSESSMENT

Patient came with a slightly mild vertical and sagittal discrepancy, potentially consistent with a borderline Class III skeletal tendency. In the frontal photo, a smile with anterior crossbite and peg laterals can be seen.



FIGS. 1, 2 ,3 - Initial extraoral photos

Radiographic analysis

The nasopharyngeal airway appears reasonably patent.No obvious gross cystic lesion or major bony abnormality is apparent.
Predominantly mandibular/chin retrusive appearance.
→ Some anterior irregularity/crowding
→ Vertical skeletal pattern
→ possible mild hyperdivergent tendency



FIG. 4,5 - Initial X-ray

EVALUATION OF THE DENTAL ARCHES

Some apparent crowding/space deficiency in the madibular dentition.
Skeletal Class III tendency / borderline Class III
Anterior cross bite with peg lateral incisors.





FIGS. 6-8 - Initial intraoral photos

TREATMENT

The treatment is performed using a Alignaa clear ALigners, with the aim of improving the anterior cross bite, and mesialise the upper arch. The appliance is positioned and used according to the protocol, remaining in the mouth for a period of about 12.5 months.

CLINICAL OUTCOME

Following the use of the alignaa aligners, an anterior cross-bite is expected to be resolved. The patient continues to undergo check-ups to monitor and assess stability.



FIGS. 9,10 - Comparison of pre- and post-upper arch treatment files

SCANS

INITIAL

END OF EXPANSION



FIGS. 11,12 - Comparison of pre- and post-treatment files in frontal, occlusal, and lateral views.