1. The probability of a leap year selected at random contain 53				
Sunday is:				
(a) 53/ 366	(b) 1/7	(c) 2/7	(d) 53/365	
2. A bag contains	3 red and 2 h	olue marbles. A	marble is drawn at	
random. The proba	ability of draw	wing a black ball	lis:	
(a) 3/5	(b) 2/5	(c) 0/5	(d) 1/5	
3. The probability	that it will ra	in tomorrow is ().85. What is the	
probability that it w				
(a) 0.25	(b) 0.145	(c) 3/20	(d) none of these	
4. What is the prob	pability that	a number select	ed from the numbers	
(1, 2, 3,,15) i	-			
		(c) 2/15		
5. What are the to	tal outcome	s when we throv	v three coins?	
• •	• •	(c) 8	• •	
6. The probability	that a prime	e number selecte	ed at random from the	
numbers (1,2,3,	•			
(a) 12/35	(b) 11/35	(c) 13/35	(d) none of these	
7. The sum of the	-			
) 0 (d) non		
		are given; choos	se the correct answer	
for that which is no				
			(d) none of these.	
		nultaneously, tha	an the probability of	
getting at least two	heads, is:			
(a) 1/4 (•		* *	
10. A letter is cho				
♦ ASSASSINATION				
(a) 6/13	(b) 7/13	(c) 1	(d) none of these.	
11. A dice is throw	-		ting an even number.	
(A) 2/3	(B) 1	(C) 5/6	(D) 1/2	
		e same time. Fin	d the probability of	
getting both heads		(D) 0		
(A) 3/4 (B) 1/4	(C) 1/2	(D) U		
13. Two dice are th	ırown simult	aneously. The p	robability of getting a	

1

sum of 9 is:

(4) 4 (4.0	(5) 0 (40	(0) 1 (0	(D) 4	10	
(A) 1/10	(B) 3/10	(C) 1/9	(D) 4	<i>'</i> 9	
14. 100 cards are numbered from 1 to 100. Find the probability of getting a prime number.					
(A) 3/4	(B) 27/50	(C) 1/4	(D)	29/100	
15. A bag contains 5 red balls and some blue balls .If the probability of drawing a blue ball is double that of a red ball, then the number of blue balls in a bag is:					
(A) 5	(B) 10 (C) 15	(D) 20		
16. A box of 600 bulbs contains 12 defective bulbs. One bulb is taken out at random from this box. Then the probability that it is non-defective bulb is:					
(A) 143/150) (B) 14//	150 (C)	1/25	(D) 1/50	
17. Cards marked with numbers 2 to 101 are placed in a box and mixed thoroughly. One card is drawn from this box randomly, then the probability that the number on card is a perfect square. (A) 9/100 (B) 1/10 (C) 3/10 (D) 19/100					
18. What is the probability of getting 53 Mondays in a leap year? (A) 1/7 (B) 53/366 (C) 2/7 (D) 7/366					
19. A card is drawn from a well shuffled deck of 52 cards. Find the probability of getting a king of red suit. (A) $1/26$ (B) $3/26$ (C) $7/52$ (D) $1/13$					
20. A game of chance consists of spinning an arrow which is equally likely to come to rest pointing to one of the number 1,2,312 ,then the probability that it will point to an odd number is: (A) 1/6 (B) 1/12 (C) 7/12 (D) 5/12					
21. A game consists of tossing a one rupee coin 3 times and noting its outcome each time. Aryan wins if all the tosses give the same result i.e. three heads or three tails and loses otherwise. Then the probability that Aryan will lose the game. (A) $3/4$ (B) $1/2$ (C) 1 (D) $1/4$					

22. Riya and Kajal are friends. Probability that both will have the same birthday is the same birthday is:						
•		-	(D) 1/133225			
23. A number x is chosen at random from the numbers -2, -1, 0, 1, 2. Then the probability that $x^2 < 2$ is? (A) $1/5$ (B) $2/5$ (C) $3/5$ (D) $4/5$						
24. A jar contains 24 marbles. Some are red and others are white. If a marble is drawn at random from the jar, the probability that it is red is 2/3, then the number of white marbles in the jar is: (A) 10 (B) 6 (C) 8 (D) 7						
Then the prob	25. A number is selected at random from first 50 natural numbers. Then the probability that it is a multiple of 3 and 4 is: (A) 7/50 (B) 4/25 (C) 1/25 (D) 2/25					
26. Consider a dice with the property that that probability of a face with n dots showing up is proportional to n. The probability of face showing 4 dots is?						
a) $\frac{1}{7}$	b) $\frac{5}{42}$	c) $\frac{1}{21}$	d) 4/21			
27. Runs scored by batsman in 5 one day matches are 50, 70, 82,						
	ne standard devia b) 25.49		d) 25.69			
28. Find median and mode of the messages received on 9 consecutive days 15, 11, 9, 5, 18, 4, 18, 13, 17.						
	b) 13, 18		d) 13, 16			
29. A coin is tossed up 4 times. The probability that tails turn up in 3 cases is						
a) $\frac{1}{2}$ 30. X is a variance	b) $^1\!/_3$ ate between 0 an	d 3. The value of	$\frac{d}{1}$ 6 d) $\frac{1}{6}$ 6 E(X²) is			
31. The random variables X and Y have variances 0.2 and 0.5 respectively. Let Z= 5X-2Y. The variance of Z is?						

probability?	?	•	n one is not poss	ible in		
a) $P(x) = 1$ c) $P(x) = 0.5$	b) ∑ x 5	x P(x) = 3 (x) = -0.5				
	2 and E(z) = 6 b) 6		•	sufficient data		
34.The cov	34. The covariance of two independent random variable is					
a) 1	b) 0	c) - 1	d) Ur	defined		
) = k² – 8 then b) 1	, the value o <mark>c) 3</mark>		sufficient data		
• •	0.5 and x = 4 , b) 0.5	• • •	? d) 2			
37.In a discrete probability distribution, the sum of all probabilities is always?						
a) 0	b) Infinite	c) 1	d) Und	defined		
•	obability of h	itting the tar	get is 0.4, find m	nean and		
variance. a) 0.4, 0.24	b) 0.6,	0.24	c) 0.4, 0.16	d) 0.6, 0.16		
•	% and if 10 bo		pped from a place opped, find mean 0.4, 0.16			
 40. Find the mean of tossing 8 coins. a) 2 b) 4 c) 8 d) 1 41. What is the mean and variance for standard normal distribution? 						


c) 5

d) 7

a) 3

b) 4

a) Mean is (c) Mean is (
42.Varianc a) E(X)						d) (E(X))2
43.Mean of a random variable X is given by a) E(X)					d) (E(X))2	
44.Mean of a constant 'a' is a) 0						
45.Variance of a constant 'a' is . a) 0						
46.Find the mean and variance of X?						
×	0	1	2	3	4	
f(x)	1/9	2/9	3/9	2/9	1/9	
a) 2, 4/3	b)	3, 4/3		c) 2, 2/3	}	d) 3, 2/3
47. Find the expectation of a random variable X?						



48. In a Binomial Distribution, if p, q and n are probability of success, failure and number of trials respectively then variance is given by

b) npq

c) np2q

d) npq2

49. If 'X' is a random variable, taking values 'x', probability of success and failure being 'p' and 'q' respectively and 'n' trials being conducted, then what is the probability that 'X' takes values 'x'? Use **Binomial Distribution.**

- a) P(X = x) = nCx px qx
- b) P(X = x) = nCx px q(n-x)
- c) P(X = x) = xCn qx p(n-x)
- d) P(x = x) = xCn pn qx

50. If 'p', 'q' and 'n' are probability pf success, failure and number of trials respectively in a Binomial Distribution, what is its Standard **Deviation?**

- a) \sqrt{np} b) \sqrt{pq} c) (np)2 d) \sqrt{npq}