

**SOLUTIONS FOR TERM TEST**  
**MARCH 23, 2018**  
**MATB43**

- (1) Let  $\sum_j a_j$  and  $\sum_j b_j$  be series of positive terms. Suppose there exists a constant  $C$  such that

$$1/C \leq a_j/b_j \leq C ,$$

for all  $j$ . Prove that then both series converge or both series diverge.

Rewrite the inequalities as

$$b_j \leq C a_j , \quad a_j \leq C b_j ,$$

for all  $j$ .

Applying the comparison test to the first inequality, we get: if  $\sum_j a_j$  converges, then  $\sum_j b_j$  converges, and if  $\sum_j b_j$  diverges, then  $\sum_j a_j$  diverges.

Using the second inequality, we see that: if  $\sum_j b_j$  converges, then  $\sum_j a_j$  converges, and if  $\sum_j a_j$  diverges, then  $\sum_j b_j$  diverges.

Combining these statements gives us: both series converge or both series diverge.

- (2) Let

$$\sinh x = \frac{e^x - e^{-x}}{2} , \quad \cosh x = \frac{e^x + e^{-x}}{2} .$$

- (a) Give power series expansions for  $\sinh x$  and  $\cosh x$ .

$$\begin{aligned}\sinh x &= \frac{e^x - e^{-x}}{2} \\ &= \frac{1}{2} \left( \sum_{n=0}^{\infty} \frac{x^n}{n!} - \sum_{n=0}^{\infty} \frac{(-x)^n}{n!} \right) = \sum_{n=0}^{\infty} \frac{x^{2n+1}}{(2n+1)!}\end{aligned}$$

$$\begin{aligned}\cosh x &= \frac{e^x + e^{-x}}{2} \\ &= \frac{1}{2} \left( \sum_{n=0}^{\infty} \frac{x^n}{n!} + \sum_{n=0}^{\infty} \frac{(-x)^n}{n!} \right) = \sum_{n=0}^{\infty} \frac{x^{2n}}{(2n)!}\end{aligned}$$

(b) Prove that

$$\cosh^2 x - \sinh^2 x = 1 .$$

We have

$$\cosh^2 x - \sinh^2 x = (\cosh x - \sinh x)(\cosh x + \sinh x) .$$

Now

$$\begin{aligned}\cosh x - \sinh x &= \frac{e^x + e^{-x}}{2} - \frac{e^x - e^{-x}}{2} = e^{-x} \\ \cosh x + \sinh x &= \frac{e^x + e^{-x}}{2} + \frac{e^x - e^{-x}}{2} = e^x\end{aligned}$$

Therefore

$$\cosh^2 x - \sinh^2 x = e^{-x} \cdot e^x = 1 .$$

- (3) Give an example of non-empty closed sets  $S_1 \subset S_2 \subset \cdots \subset S_j \subset \cdots$  such that  $S = \cup_{j=1}^{\infty} S_j$  is open.

Let  $S_j = [-1 + 1/j, 1 - 1/j]$ ,  $j = 1, 2, \dots$ . For any  $a \in (-1, 1)$ , there exists  $j > 0$  with

$$-1 + 1/j \leq a \leq 1 - 1/j .$$

Therefore  $S \supseteq (-1, 1)$ . On the other hand,

$$S_j \subset (-1, 1) ,$$

for all  $j$ . So  $S \subseteq (-1, 1)$ . Thus  $S = (-1, 1)$ , which is open.

- (4) Find all accumulation points of the following sets. Explain your answers.

(a)  $\{\frac{1}{n} \mid n \in \mathbb{N}\}$ .

Since  $\lim_{n \rightarrow \infty} \frac{1}{n} = 0$ , 0 is an accumulation point. For any other  $a \in \mathbb{R}$ , there exists a neighbourhood which does not contain any other point in the set. Therefore 0 is the only accumulation point.

(b)  $\{\frac{1}{m} + \frac{1}{n} \mid m, n \in \mathbb{N}\}$ .

Since  $\lim_{n \rightarrow \infty} (\frac{1}{m} + \frac{1}{n}) = \frac{1}{m}$ ,  $\frac{1}{m}$  is an accumulation point for all  $m \in \mathbb{N}$ . As well,  $\lim_{n \rightarrow \infty} (\frac{1}{n} + \frac{1}{n}) = \lim_{n \rightarrow \infty} \frac{1}{2n} = 0$  is an accumulation point. For any other  $a \in \mathbb{R}$ , there exists a neighbourhood which does not contain any other point in the set. Therefore  $\{\frac{1}{m} \mid m \in \mathbb{N}\} \cup \{0\}$  is the set of accumulation points.

(c)  $\{(-1)^n(1 + \frac{1}{n}) \mid n \in \mathbb{N}\}$ .

We have  $\{(-1)^n(1 + \frac{1}{n}) \mid n \in \mathbb{N}\} = \{(1 + \frac{1}{2n}) \mid n \in \mathbb{N}\} \cup \{(-1 - \frac{1}{2n+1}) \mid n \in \mathbb{N}\}$ . Since  $\lim_{n \rightarrow \infty} (1 + \frac{1}{2n}) = 1$  and  $\lim_{n \rightarrow \infty} (-1 - \frac{1}{2n+1}) = -1$ , 1 and -1 are accumulation points. For any other  $a \in \mathbb{R}$ , there exists a neighbourhood which does not contain any other point in the set. Therefore these are the only accumulation points.

Student Number 1002423727

Name POON KEEGAN  
LAST NAME FIRST NAME INITIAL  
(Please print in block letters)

BOOK NO. \_\_\_\_\_

TOTAL NUMBER  
OF BOOKS USED \_\_\_\_\_

**THIS BOOK MUST NOT BE TAKEN  
FROM THE EXAMINATION ROOM**

Course MATB43  
(e.g. ANTA01Y, PHL 202S)

Instructor JOHN SCHERK

Date of Examination \_\_\_\_\_

Room Number \_\_\_\_\_ Seat Number \_\_\_\_\_

Tut 03?  
Fri 11-12

**TERM**

**INSTRUCTIONS**

Write the information sought in the spaces above.

Write the answers on the RULED SIDE ONLY; all calculations or rough drafts of answers should be shown, preferably on the unruled side.

Clearly identify the question to which each answer applies; whenever the answer to a question is divided, note at the end of the first section "see also work on page \_\_\_\_\_".

If a page is left blank write on it "see work on page \_\_\_\_\_".

If more than one book is used, indicate the total number on the cover of each. At the conclusion of the examination, place all other books inside Book No. 1.

**DO NOT TEAR ANY PAPER OUT OF THIS BOOK.**

**EXAMINER'S REPORT**

1	25
2	25
3	25
4	25
5	
6	
7	
8	
9	
10	
11	
12	
TOTAL	

**University of Toronto  
Scarborough**

$$1) \quad 1/c \leq a_j/b_j \leq c$$

$$\Rightarrow a_j/c \leq a_j \leq b_j/c$$

$$0 \leq \frac{1}{c} \sum_{j=1}^{\infty} b_j \leq \sum_{j=1}^{\infty} a_j \leq c \sum_{j=1}^{\infty} b_j$$

So if  $b_j$  converges/diverges, we can immediately do a comparison test

Since if  $b_j$  diverges  $\Rightarrow \frac{1}{c} \sum b_j$  diverges  $\Rightarrow \sum a_j$  diverges & since  $\frac{1}{c} \sum b_j \leq \sum a_j \leq c \sum b_j$   $\Rightarrow a_j$  diverges as well since partials of  $b_j$  diverge.

Similarly if  $b_j$  converges,  $0 \leq a_j \leq c b_j$  &  $c \sum b_j$  converges.

So CT  $\Rightarrow a_j$  converges again.

Since all terms are positive,  $\Rightarrow c$  also positive &  $1/c$  also positive

$$\Rightarrow \text{Note that } (1/c)^+ \geq (a_j/b_j)^+ \geq (c)^+ \text{ as } x^+ \text{ decreasing function \& continuous on } (0, \infty)$$

$$\Rightarrow c \geq b_j/a_j \geq 1/c$$

$$\text{so } 1/c \leq b_j/a_j \leq c$$

Therefore,  $b_j$  also converges/diverges with  $a_j$ , symmetrically to how  $a_j$  was proven.



$$2a) e^x = \sum_{k=0}^{\infty} \frac{x^k}{k!} \text{ for all } x \in (-\infty, \infty) \text{ \& } \forall x, \left| \frac{e^x - e^{-x}}{2} \right| = \frac{\sum_{k=0}^{\infty} \frac{x^k}{k!} - \sum_{k=0}^{\infty} \frac{(-x)^k}{k!}}{2}$$

$$= \frac{1}{2} \sum_{k=0}^{\infty} \left[ \frac{x^k}{k!} - \frac{(-x)^k}{k!} \right] \rightarrow \text{on even } k = \frac{x^{2k}}{(2k)!} - \frac{(-x)^{2k}}{(2k)!} = \frac{x^{2k} - x^{2k}}{(2k)!} = 0$$

$$\Rightarrow \sinh x = \sum_{k=0}^{\infty} \frac{x^{2k+1}}{(2k+1)!} \text{ on odd } k = \frac{x^{2k+1}}{(2k+1)!} - \frac{(-x)^{2k+1}}{(2k+1)!} = 2 \frac{x^{2k+1}}{(2k+1)!}$$

with the same radius of convergence

$$\text{While for } \cosh x = \frac{e^x + e^{-x}}{2} = \frac{\sum_{k=0}^{\infty} \frac{x^k}{k!} + \sum_{k=0}^{\infty} \frac{(-x)^k}{k!}}{2} \quad 15$$

$$= \frac{1}{2} \sum_{k=0}^{\infty} \left( \frac{x^k}{k!} + \frac{(-x)^k}{k!} \right) \rightarrow \text{on even } k = \frac{x^{2k}}{(2k)!} + \frac{(-x)^{2k}}{(2k)!}$$

$$\Rightarrow \cosh x = \sum_{k=0}^{\infty} \frac{x^{2k}}{(2k)!} \text{ on odd } k = \frac{x^{2k+1}}{(2k+1)!} + \frac{(-x)^{2k+1}}{(2k+1)!} = 0$$

With the same radius of convergence.

$$b) \cosh^2 x - \sinh^2 x$$

$$= \left( \frac{e^x + e^{-x}}{2} \right)^2 - \left( \frac{e^x - e^{-x}}{2} \right)^2$$

$$= \frac{e^{2x} + 2e^{x-x} + e^{-2x}}{4} - \frac{e^{2x} - 2e^{x-x} + e^{-2x}}{4}$$

$$= \frac{4e^0}{4} = 1 //$$

$$3) S_n = [-1 + \frac{1}{n}, 1 - \frac{1}{n}] \quad -1 + \frac{1}{n+1} < -1 + \frac{1}{n} \quad \& \quad 1 - \frac{1}{n+1} > 1 - \frac{1}{n} \Rightarrow S_n \subset S_{n+1}$$

Then  $S = \bigcup_{i=1}^{\infty} S_i = (-1, 1)$  is open

Pf/  $0 \in S_n \forall n$

Let  $0 < |a| < 1$ , then  $1 - |a| = \varepsilon > 0$

$$\Rightarrow |a| = 1 - \varepsilon$$

But  $\forall \varepsilon > 0 \exists N$  s.t.  $n > N \Rightarrow x(\frac{1}{n}) < \varepsilon$  (since  $\lim_{n \rightarrow \infty} \frac{1}{n} = 0$ )

$$\Rightarrow |a| = 1 - \varepsilon < 1 - \frac{1}{n}$$

$\Rightarrow |a| < 1 - \frac{1}{n}$  so  $a \in (-1 + \frac{1}{n}, 1 - \frac{1}{n})$  for some  $n \in \mathbb{N}$

$\therefore a \in S$

But  $n > 0 \Rightarrow -1 + \frac{1}{n} > -1 \forall n \quad \& \quad 1 - \frac{1}{n} < 1 \forall n$

So  $S = (-1, 1)$

Now  $(-1, 1)$  is clearly open since for any  $x \in (-1, 1)$

$\varepsilon = |x| - 1 > 0$  so  $(x - \varepsilon, x + \varepsilon)$  is a neighborhood within  $S$  around  $x$ .



4a)  $0 \notin \{1/n \mid n \in \mathbb{N}\}$  is the only possible accumulation point. Since the sequence  $\{1/n\}$  converges to 0, there are infinitely many points within  $(0, 0+\epsilon)$ , that belong in the sequence.

Also know that  $\{1/n \mid n \in \mathbb{N}\} \subset (0, 1)$  so no accumulation points outside that range. Given any  $1/x$ ,  $x \in \mathbb{R}^+$ , there is at most 1 element of the set in  $(1/x - \delta, 1/x + \delta)$ , as the distance between  $1/n$  and  $1/(n+1)$  surrounding  $x > \delta$ , where  $\delta = \frac{1}{Lx+1} - \frac{1}{Lx+2}$   $Lx$  being the integer part of  $x$ .

b) Every point of the previous set  $\{1/n \mid n \in \mathbb{N}\}$  is an accumulation point in this set. Given the justification used earlier,  $1/n$  gets arbitrarily close to zero  $\Rightarrow$  infinitely many points near  $1/n$ , formally

Since  $\forall \epsilon > 0, \exists M > 0$  s.t.  $n > M \Rightarrow x/1/n < \epsilon$

$$\Rightarrow 1/n(1/1/n) + 1/n < \epsilon + 1/n \quad 1/n > 0$$

$$\Rightarrow 1/n < 1/1/n + 1/n < \epsilon + 1/n$$

since  $1/n + 1/n > 0$

$$1/n < 1/1/n + 1/n < \epsilon + 1/n \Rightarrow \frac{1}{m} + \frac{1}{n} \in (-\epsilon + 1/n - \epsilon, 1/n + \epsilon)$$

$\exists$  infinitely  $n > M$ , so for  $\epsilon$  precision to  $1/n$ ,  $\exists$  infinitely many points.

0 is also an accumulation point for similar reasons to the first,  $a/n = 0$  as  $n \rightarrow \infty$

c) The accumulation points are  $\{1, -1\}$

first examine  $\{1 + 1/n \mid n \in \mathbb{N}\}$   $\lim_{n \rightarrow \infty} 1/n \rightarrow 0$  so  $1 + 1/n \rightarrow 1$  as  $n \rightarrow \infty$ .

So again there must be accumulation points at 1 or -1

if we take the subsets  $E = \{1 + 1/2n \mid n \in \mathbb{N}\}$  and  $F = \{-1 - 1/2n \mid n \in \mathbb{N}\}$  we can that the even  $n$  generate a sequence whose limit is 1, so there is an infinite sequence of points converging to 1  $\Rightarrow$  infinitely many points around 1 exactly the same as done in b). Similarly for 0,  $-1/2n$  converges to 0, so the sequence generated by ordering each element by its value of  $n$ , converges to -1, giving another accumulation point.

Also know that for any  $x \neq 1, -1$ , Since  $\{1 + 1/n \mid n \in \mathbb{N}\}$  within  $\epsilon$  of 1 for any sufficiently large  $N$ ,  $\Rightarrow$  for sufficiently large  $n$ ,  $\exists$  neighbourhood around  $x$  such that no elements of  $\{1 + 1/n \mid n \in \mathbb{N}, n > N\}$  are in it  $\Rightarrow$  not accumulation point.