



ROOT CAUSE UNLOCKED · COMPLETE SET

Low Back and Hip, the *Complete* Set

A Whole-Person Approach to Healing, Recovery, and Resilience

Dr. Adam Seay, DC

Root Cause Unlocked, Garner, NC · rootcauseunlocked.com · (919) 662-0520

Low Back and Hip, the Complete Set

A Whole-Person Approach to Healing, Recovery, and Resilience

BEFORE YOU START

Disclaimer: *This ebook is for educational purposes only and does not constitute medical advice. Always consult with a qualified healthcare provider before starting any new treatment, supplement, exercise program, or dietary change. Do not start, stop, or change any prescribed medication without speaking to the prescriber. If you experience progressive weakness, loss of bowel or bladder control, fever with back pain, or symptoms that worsen rapidly, seek emergency medical attention immediately.*

A note on the evidence in this guide. *Every numbered citation in this edition points to a specific published study that was checked against the source paper. Where the research is weak, mixed, or negative, this guide says so rather than overstating it. Where a claim could not be traced to a real study, it was removed.*

Contents

1. Understanding Low Back Pain

2. Anatomy of the Lumbar Spine

3. Common Causes and Risk Factors

The Hip Connection: Why the Joint Below Often Drives the Pain

4. How Low Back Pain Is Diagnosed

5. The Healing Timeline: What to Expect

6. Supplements for Spinal Health and Pain Relief: What the Evidence Actually Says

7. Movement and Exercise for Low Back Recovery

8. Integrative and Physical Therapies

9. The Anti-Inflammatory Diet for Spinal Recovery

10. Lifestyle Modifications: Sleep, Stress, and Ergonomics

11. Mind-Body Approaches to Chronic Pain

12. The In-Clinic Pre-Treatment Protocol

13. Red Flags: When to Seek Immediate Care

14. Building Your Personal Recovery Plan

Final Thoughts

15. References

Working with Root Health

1. What Is Sciatica?

2. Understanding the Anatomy

3. Common Causes and Risk Factors

4. How Sciatica Is Diagnosed

5. The Healing Timeline: What to Expect

6. Evidence-Based Supplements for Nerve Health and Inflammation

7. Targeted Exercises for Sciatica Relief

8. Physical Therapies That Work

9. The Anti-Inflammatory Diet for Nerve Recovery

10. Lifestyle Modifications and Pain Management

11. The In-Clinic Pre-Treatment Protocol

12. Red Flags: When to Seek Immediate Care

13. Building Your Personal Recovery Plan

Final Thoughts

References

Working with Root Health

1. Understanding Hip Pain

2. Anatomy of the Hip: The Structures at Play

3. Common Causes of Hip Pain

4. The Real Story Behind Lateral Hip Pain: Gluteal Tendinopathy

5. How Hip Pain Is Diagnosed

6. Regional Interdependence: Why the Hip Rarely Works Alone

7. Movement and Exercise for Hip Recovery

8. Integrative and Physical Therapies

9. Nutrition for Joint and Tendon Health

10. Lifestyle Modifications: Sleep, Weight, and Footwear

11. The In-Clinic Pre-Treatment Protocol

12. Red Flags: When to Seek Immediate Care

13. Building Your Personal Recovery Plan

Final Thoughts

14. References

Read this part first

Why one guide covers the low back and the hip

If you were told you have... (your diagnosis, translated)

The one idea that makes this kit honest

The low back sequence

The hip sequence

The program

When to get evaluated instead of self-treating

Why nutrient evidence looks thinner than it is

Sort the tissue before you buy anything

The spine: tested and failed, said plainly

The arthritic hip: borrowed evidence, labeled as borrowed

Lateral hip pain: you have a tendon, and that changes everything

The floor under all three tissues

What I would actually run, and how to judge it

Interactions, and the ones people miss

Track it, so the program can be judged by data

Dr. Seay recommends: the exact products

Working with Root Health

References

1. Understanding Low Back Pain

Low back pain is one of the most common health complaints worldwide. It is now the leading cause of disability globally, and it occurs in high-income, middle-income, and low-income countries alike, across every age group from children to older adults.¹ If you are reading this, you are part of a very large community, and importantly, you are not alone.

The vast majority of low back pain is “nonspecific.” That means it is not caused by a serious underlying condition like cancer, infection, or fracture, and often no single structural cause can be definitively identified. For nearly all people with low back pain, it is not possible to identify a specific tissue source, and only a small proportion have a well understood pathological cause such as a vertebral fracture, malignancy, or infection.¹ This can be frustrating, but it is also encouraging: it means that your back is strong, adaptable, and capable of healing even without a precise diagnosis.

Types of Low Back Pain

Acute low back pain lasts less than 6 weeks. It is often triggered by a specific movement, lift, or strain, and it is highly likely to improve substantially on its own. Subacute low back pain lasts 6 to 12 weeks and may require more structured intervention. Chronic low back pain persists beyond 12 weeks, and this is where a biopsychosocial approach becomes essential.

The Biopsychosocial Model

Modern pain science recognizes that low back pain is not purely a physical problem. It is influenced by biological factors such as tissue health, inflammation, genetics, sleep, and nutrition; psychological factors such as stress, anxiety, depression, fear of movement, and catastrophizing; and social factors such as work demands, social support, financial stress, and cultural beliefs about pain.

Initial high pain intensity, psychological distress, and pain at multiple body sites all increase the risk that low back pain becomes persistent and disabling. There is increasing evidence that central pain-modulating mechanisms and pain beliefs play important roles in that process.¹

This ebook embraces that integrative perspective. Healing your back involves more than just treating your spine. It involves nourishing your whole self.

2. Anatomy of the Lumbar Spine

Your lower back is an engineering marvel. Understanding its basic anatomy helps demystify pain and gives you the confidence to move.

The Structures

Your lumbar vertebrae, L1 to L5, are the five largest vertebrae in your spine, designed to bear weight and allow movement. Between them sit the intervertebral discs, gel-filled cushions that act as shock absorbers and allow spinal motion. They are avascular (they have no direct blood supply) and rely on movement for nutrition. The facet joints are small paired joints at the back of each vertebra that guide movement, and spinal ligaments, tough bands including the anterior longitudinal ligament, posterior longitudinal ligament, ligamentum flavum, and interspinous and supraspinous ligaments, stabilize the spine. Your muscles do two different jobs: the deep stabilizers, multifidus, transversus abdominis, pelvic floor, and diaphragm, form your “core,” while the global movers, erector spinae, quadratus lumborum, psoas, gluteals, and hamstrings, handle larger movements. Finally, your nerves include the cauda equina (“horse’s tail”), a bundle of nerve roots that exit the spinal canal and travel down the legs.

Key Insight: Discs Need Movement

Your intervertebral discs do not have their own blood supply. They receive nutrients and expel waste through a process called imbibition, essentially the pumping action that occurs when you move. This is one reason prolonged sitting is unhelpful for spinal health, and why gentle, regular movement is foundational to recovery.

3. Common Causes and Risk Factors

Common Causes

Muscle strain or ligament sprain is the most common cause of acute low back pain, and it usually settles within days to weeks. Degenerative disc disease refers to age-related changes in disc hydration and height. This is a normal part of aging, not necessarily a disease, and it does not always cause pain. A herniated disc occurs when the nucleus pulposus pushes through the annulus fibrosus, potentially irritating a nerve root, while facet joint arthropathy is wear and tear of the small joints at the back of the spine. Spondylolisthesis is when one vertebra slips forward over another, and spinal stenosis is a narrowing of the spinal canal, often age-related. Sacroiliac (SI) joint dysfunction is pain originating from the joint connecting the spine to the pelvis, and myofascial pain syndrome involves trigger points in the muscles of the lower back, hips, and pelvis.

Risk Factors

People with physically demanding jobs, physical and mental health comorbidities, smokers, and people with obesity are at greatest risk of reporting low back pain.¹ Commonly discussed contributors include a sedentary lifestyle, since prolonged sitting reduces the movement that discs rely on for nutrition, and poor physical fitness, since weak core, glutes, and hip muscles increase spinal load. Excess body weight matters too: overweight and obesity are associated with a higher risk of low back pain, with the strongest associations for seeking care and for chronic low back pain.² Occupational factors such as heavy lifting, repetitive bending, twisting, and prolonged standing or sitting also contribute, as does smoking, which is associated with greater disc degeneration.³ Psychosocial stress, including anxiety, depression, job dissatisfaction, and catastrophizing, is linked to persistent, disabling low back pain.¹ A history of previous episodes is the factor that most consistently predicts a future episode,⁴ and age plays a role as well: disc degeneration increases with age, though pain does not always follow.⁵

The Pain-MRI Paradox

Research has consistently shown that MRI findings do not correlate well with pain. A landmark 2015 systematic review in the *American Journal of Neuroradiology* pooled imaging from 3,110 **asymptomatic** people, meaning people with no back pain at all, and reported that disc degeneration rose from **37%** of 20-year-olds to **96%** of 80-year-olds, disc bulge rose from **30%** of 20-year-olds to **84%** of 80-year-olds, disc protrusion rose from **29%** of 20-year-olds to **43%** of 80-year-olds, and annular fissure rose from **19%** of 20-year-olds to **29%** of 80-year-olds.⁵

None of these people had back pain. The authors concluded that many imaging-based degenerative features are likely part of normal aging and unassociated with pain, and must be interpreted alongside your clinical picture.⁵

This means that a “bad MRI” does not doom you to chronic pain, and a “good MRI” does not guarantee a pain-free life. Your back is more resilient than imaging sometimes suggests.

The Sacroiliac Joint: A Commonly Missed Driver of Chronic Low Back Pain

If you have lived with low back pain for months or years, and if you or your provider have also noticed that your glute muscles on one or both sides always feel tight, guarded, or overworked no matter how much you stretch them, it is worth asking a question that rarely gets asked: has anyone looked specifically at your sacroiliac joints?

The sacroiliac joints sit where your sacrum, the triangular bone at the base of your spine, meets the two wings of your pelvis, the ilia. Unlike the joints higher up in your lumbar spine, the sacroiliac joints are not built for large ranges of motion. They are built for load transfer, moving your body weight from your spine into your legs with every step, every time you stand up, and every time you shift your weight. That job is done almost entirely by ligaments, some of the largest and strongest in the human body, rather than by muscle control alone.

This matters because a sacroiliac sprain does not always announce itself the way an ankle sprain does. Many people never receive that diagnosis at all. They are simply told they have “low back pain,” given generic advice, and sent on their way, while the sacroiliac joint itself is never specifically examined. In one chart review of patients with recalcitrant chronic low back pain who went on to have their pain source formally identified through diagnostic injections at an academic spine center, the sacroiliac joint was confirmed as the source in about 18% of cases, a similar order of magnitude to facet joint pain, and its likelihood rose with age.⁴⁵ That is a meaningful share of chronic low back pain, and it is a joint that a standard exam does not always single out.

Part of the reason it gets missed is that reliably identifying sacroiliac joint pain on physical examination is genuinely harder than clinicians once assumed. Individual hands-on tests for how the joint is moving have poor agreement between examiners, and there is no simple gold-standard test that confirms the diagnosis on its own. What the research does support is that a cluster of provocation tests, positions and pressures applied to the joint that are designed to reproduce your familiar pain rather than assess how it moves, performs meaningfully better than any single test alone: three or more positive pain-reproduction tests out of a standard battery have been reported with roughly 91% sensitivity and 78% specificity for identifying the sacroiliac joint as a true pain source.⁴⁶ In practice, that means a careful, multi-test physical exam by someone specifically looking for it is what actually finds this, not a quick check of whether your back hurts when a spot is pressed.

There is also a real anatomical reason sacroiliac pain is so often felt in and around the glutes rather than in the low back itself. Research mapping where sacroiliac joint injections produce sensation in otherwise pain-free volunteers found that the affected area centers over the buttock, extending down from the bony landmark at the back of the pelvis, the posterior superior iliac spine, and outward toward the hip.⁴⁷ That is precisely the region where patients and providers alike tend to feel, and blame, tight glutes. A joint problem

can present as a muscle problem, which is one more reason it gets waved off as ordinary tightness rather than investigated as its own condition.

In my clinical experience, this is a pattern worth taking seriously: a patient with longstanding, low-grade chronic low back pain whose glutes stay persistently tight and guarded on manual exam, even after conventional stretching, soft tissue work, and general core strengthening, often turns out to have an underlying sacroiliac issue that was never separately assessed. My explanation for why this becomes chronic rather than resolving the way an ordinary sprain does is that the sacroiliac ligaments are large, thick, and constantly weightbearing, and in my experience they often do not fully settle down and heal the way a non-weightbearing ligament might. That reasoning is my own clinical judgment, built from years of hands-on practice, and it is offered here in that spirit: as an experienced clinician's working explanation for a pattern I see often, not as a conclusion drawn from a specific published study.

One tool I use clinically for suspected sacroiliac involvement is a sacroiliac belt, a simple strap worn snugly around the pelvis just below the hip bones, below the waistline rather than at it. The idea behind it is mechanical: by gently compressing the joint, the belt is thought to add a measure of external stability while the surrounding ligaments are irritated or lax. There is some direct supporting evidence for this mechanism. In a small pilot study, wearing a pelvic compression belt measurably changed how the gluteus maximus and hamstring muscles activated during single-leg standing in people with confirmed sacroiliac joint pain, compared with pain-free volunteers, consistent with the idea that external support changes how the surrounding muscles are recruited to protect the joint.⁴⁸ That is meaningful mechanistic evidence, but it comes from a small pilot study measuring muscle activity, not a large trial measuring pain relief or function over time, and the broader evidence on how much sacroiliac belts reduce pain in everyday use is thinner than their popularity in clinical practice would suggest. If your provider recommends one, think of it as a low-risk, mechanically sensible support to try alongside the rest of your care, not a stand-alone fix.

None of this is a self-diagnosis tool. Sacroiliac joint pain overlaps heavily with the other causes of low back pain covered elsewhere in this guide, and confirming it takes a hands-on exam by a provider trained to look for it, sometimes alongside a diagnostic injection. What this section is meant to do is put a name to a pattern you may already be living with: chronic low back pain paired with glutes that never seem to release. If that describes you, it is worth raising the sacroiliac joint specifically at your next visit, by name, rather than continuing to treat the pain as generic and untraceable.

The Hip Connection: Why the Joint Below Often Drives the Pain

Your lumbar spine is built for stability. Your hips are built for mobility. That division of labour matters more than almost anything else in this guide, because when a joint that is supposed to move stops moving, the joint next to it starts moving more than it was designed to in order to make up the difference. If your hips have lost rotation or extension, the low back is what bends, twists, and absorbs in their place, every time you stand up, reach, or take a step.

That is why hip range of motion belongs in a low back evaluation, and why the movement work later in this guide spends time on your hips when your complaint is squarely in your back.

What the evidence actually shows, stated carefully. A 2024 systematic review examined hip biomechanics in people with non-specific low back pain, looking at hip range of motion, strength, movement patterns, and muscle activity. The authors' honest summary is that the relationship is real enough to keep investigating and still contested in its details, and they say so plainly rather than overselling it.⁴⁹ So I am not going to tell you that stiff hips caused your back pain. What I will tell you is that hips are close enough, mechanically linked enough, and implicated often enough that leaving them unexamined is a mistake I see made constantly.

There is a second line of evidence here, and it is stronger than the biomechanics one. The Functional Movement Screen is a standardized set of seven movement patterns used to find where motion is missing rather than where pain is felt. A 2024 systematic review and meta-analysis, rated Level 1 evidence, compared screen scores between people with low back pain and people without it. The low back pain group scored significantly lower overall, by 1.81 points out of 21, and scored lower on six of the seven individual patterns.⁵⁰

Look at which patterns those were: deep squat, hurdle step, inline lunge, active straight leg raise, trunk stability push-up, and rotational stability. Four of those six are patterns that load the hip directly, and the active straight leg raise is essentially a hip mobility test. People with low back pain do not just move worse in general. They move worse specifically in the patterns that ask the hip to do its job.

And the honest limit, because it matters. This same screen has been studied as a tool for predicting who will get injured, and there it performs poorly. A meta-analysis of thirteen studies found only five of twelve reported an association between a low score and later injury, with substantial disagreement between studies, and the authors' conclusion was equivocal.⁵¹ So this is a screen that describes how you move right now. It is not a crystal ball, and anyone selling it to you as injury prediction is overstating it. I use it for what it does well: showing which motion is missing, so the treatment goes there.

What this means practically. If your low back pain has not responded to treatment aimed only at your back, get your hip motion assessed before concluding that nothing works. The hip is the single most common place where the actual restriction lives while the pain shows up somewhere else.

4. How Low Back Pain Is Diagnosed

For most cases of acute low back pain, extensive testing is unnecessary and can be counterproductive by increasing fear and anxiety. Routine imaging is specifically recommended against in national guidelines for recent-onset low back pain.⁶

Clinical Examination

Your healthcare provider will typically take a detailed history covering onset, aggravating and relieving factors, and red flags, and assess your range of motion in flexion, extension, side-bending, and rotation. If leg symptoms are present, they will perform neurological testing of strength, reflexes, and sensation, along with special tests such as the straight leg raise and femoral nerve stretch. They will also assess your gait and posture and palpate the spine, pelvis, and surrounding muscles.

Imaging: When Is It Needed?

Usually **not** needed in the first 4 to 6 weeks unless red flags are present.

Imaging is generally indicated when red flags are present (see Chapter 12), when there are progressive neurological deficits such as weakness or numbness, when symptoms persist beyond 6 weeks despite conservative care, when there is a history of significant trauma, or when there is suspicion of infection, cancer, or inflammatory disease.

Types of imaging: X-ray shows bones and alignment and is useful for fractures, spondylolisthesis, and severe arthritis. MRI shows soft tissues including discs, nerves, ligaments, and muscles, and is the best test for visualizing disc herniations and spinal stenosis. CT scan provides detailed bony images and is useful when MRI is contraindicated. Bone scan detects infections, tumors, or occult fractures.

5. The Healing Timeline: What to Expect

Acute Phase (0 to 6 Weeks)

Pain is typically most intense at the start and then improves markedly. In a meta-analysis of 33 inception cohorts covering 11,166 people, average pain in acute low back pain cohorts fell from 52 out of 100 at the start to 23 out of 100 at six weeks.⁷

Focus: Stay active, avoid bed rest, manage pain, reduce fear.

Subacute Phase (6 to 12 Weeks)

Improvement continues but slows. In the same meta-analysis, average pain reached 12 out of 100 at 26 weeks in the acute cohorts.⁷

Focus: Structured exercise, addressing biomechanics, returning to normal activities.

Chronic Phase (Beyond 12 Weeks)

Pain persists beyond expected healing time. Among people who already had persistent pain at the start, average pain was 51 out of 100 at baseline and still 23 out of 100 at one year, so a slower and more layered approach is appropriate here.⁷

Focus: Multidisciplinary, biopsychosocial approach. This is where integrative therapies matter most.

The Natural History

Most people improve markedly in the first six weeks, and then improvement slows. Low to moderate levels of pain and disability are still present at one year for many people, especially those whose pain was already persistent.⁷

Recurrence is common, though the research does not give a precise number. A systematic review of recurrence found the studies too varied and too low in quality to pool. The single study judged to be a true inception cohort reported a one-year recurrence of **33%**, and a history of previous episodes was the only factor that consistently predicted recurrence.⁴

The practical takeaway is unchanged: prevention and maintenance matter, not just treating the current episode.

6. Supplements for Spinal Health and Pain Relief: What the Evidence Actually Says

This is the chapter where honesty matters most, because supplements cost money and it is easy to overstate them.

Here is the headline: **most supplement research relevant to this chapter was done in knee osteoarthritis or in exercise-induced muscle soreness, not in low back pain.** That does not make the products useless, but it does mean the evidence is indirect. Where a supplement has been tested specifically in low back pain and did not work, this guide says so.

Always discuss supplements with your healthcare provider, especially if you take blood thinners or have kidney disease.

6.1 Omega-3 Fatty Acids (EPA and DHA)

What it does: Omega-3 fats are incorporated into cell membranes and influence the production of inflammatory signaling molecules.

Evidence: A 2006 report in *Surgical Neurology* asked 250 neurosurgical patients with nonsurgical neck or back pain to take 1,200 to 2,400 mg per day of omega-3 fatty acids. Of the 125 who returned the questionnaire after an average of 75 days, 59% had stopped their prescription NSAIDs and 60% said their overall pain had improved.⁸

Important caveat: That study was an uncontrolled questionnaire survey with a 50% response rate, not a randomized trial. It cannot establish that omega-3 works as well as ibuprofen for back pain, and it should be read as encouraging but preliminary.

Dosage commonly used: 1,200 to 2,400 mg combined EPA and DHA daily. That is the range used in the one study run in people with nonsurgical neck and back pain, where most participants took 1,200 mg daily and about a fifth took 2,400 mg. That study was an uncontrolled questionnaire follow-up rather than a placebo-controlled trial, so treat it as a starting point rather than a proven dose. Higher intakes appear in the joint-pain literature, but that work was done in rheumatoid arthritis and other inflammatory conditions rather than in the mechanical and degenerative problems this guide covers, so I have not carried its dose across.⁸

Best form: Triglyceride or re-esterified triglyceride form **Caution:** May increase bleeding risk at high doses. Discuss with your surgeon before any operation.

6.2 Turmeric and Curcumin

What it does: Curcumin acts on several inflammatory signaling pathways, including NF-kB and COX-2.

Evidence: A meta-analysis of randomized trials of turmeric extracts and curcumin in arthritis found reduced pain scores compared with placebo, and no significant difference in pain compared with pain medication. The authors noted that the number of trials, total sample size, and study quality were not sufficient for definitive conclusions.⁹ In a multicenter randomized trial of 367 people with knee osteoarthritis, *Curcuma domestica* extract was as effective as ibuprofen over four weeks, with fewer gastrointestinal side effects.¹⁰ A separate systematic review of 11 trials found curcuminoid formulations statistically better than placebo for knee osteoarthritis pain and function, while again cautioning that the evidence base is not adequate in size or quality for firm clinical recommendations.¹¹

Scope note: All of this evidence is in arthritis, primarily of the knee. Curcumin has not been established as a treatment for low back pain specifically.

Dosage commonly used: 500 to 1,000 mg curcuminoids daily, with piperine or in a liposomal or phytosome form **Caution:** May interact with blood thinners. Avoid in gallbladder disease.

6.3 Vitamin D

What it does: Vitamin D is essential for bone health, muscle function, and immune regulation.

Evidence, and an important correction: Vitamin D is frequently promoted for back pain, but the trial evidence does not support it. A systematic review and meta-analysis of eight clinical trials concluded that **vitamin D supplementation is not more effective than placebo, no intervention, or other conservative treatments for low back pain**, and that the prescription of vitamin D for low back pain cannot be recommended until better trials suggest otherwise. The result held regardless of the type of low back pain or the form of vitamin D used. The authors rated the overall quality of evidence as very low.¹²

What this means for you: Correcting a genuine vitamin D deficiency is worth doing for bone and general health, and it is reasonable to have your level checked. Do not expect it to function as a back pain treatment.

Dosage commonly used for deficiency: Guided by blood levels, under your provider's direction **Best form:** D3 (cholecalciferol) **Caution:** Have levels checked periodically rather than supplementing blindly.

6.4 Magnesium

What it does: Magnesium is involved in hundreds of enzymatic reactions, in energy production, and in normal muscle and nerve function.

Evidence: There is no reliable trial evidence that magnesium supplementation treats low back pain, and this guide no longer claims otherwise. The closest relevant finding is a systematic review of magnesium supplementation and **exercise-induced muscle soreness** in physically active people. It included only four eligible studies, which reported reduced muscle soreness and better recovery. That is a small and narrow evidence base, and it is about post-exercise soreness, not back pain.¹³

Dosage commonly used: 200 to 400 mg elemental magnesium daily This is a general supplemental intake range rather than a dose drawn from a trial in this condition, which is why no study is cited beside it. **Caution:** Can cause loose stools. Avoid in severe kidney disease. Discuss with your provider.

6.5 Collagen or Gelatin with Vitamin C

What it does: Supplies the amino acids (glycine, proline, hydroxyproline) used in collagen synthesis. Vitamin C is required for collagen cross-linking.

Evidence: In a randomized crossover study, participants who took 15 g of vitamin C-enriched gelatin one hour before a short bout of exercise showed **double** the blood marker of collagen synthesis compared with placebo. Serum from supplemented participants also increased collagen content and improved mechanics in engineered ligament tissue. The authors concluded that gelatin added to an intermittent exercise program improves collagen synthesis and could play a beneficial role in injury prevention and tissue repair.¹⁴

Scope note: This study measured collagen synthesis markers and engineered tissue, not back pain outcomes. It is a mechanism study, and a good one, but it does not demonstrate pain relief.

Dosage used in the study: 15 g gelatin with 50 mg vitamin C, about 60 minutes before exercise **Caution:** Generally very well tolerated.

6.6 Boswellia Serrata (Frankincense)

What it does: Inhibits 5-lipoxygenase, an enzyme in one branch of the inflammatory cascade.

Evidence: In a systematic review and meta-analysis of 11 randomized trials, boswellia formulations were statistically significantly better than placebo for pain relief and functional improvement in **knee osteoarthritis**, with no significant difference from placebo in safety outcomes. The authors were explicit that the current body of evidence is not adequate in size or quality to make meaningful clinical practice recommendations.¹¹

Scope note: Knee osteoarthritis, not low back pain.

Dosage commonly used: 300 to 500 mg standardized extract twice daily **Caution:** May interact with anti-inflammatory medications.

6.7 Ginger

What it does: Contains gingerols and shogaols, compounds with anti-inflammatory and analgesic activity in laboratory models.

Evidence: In two double-blind, placebo-controlled randomized experiments (34 and 40 volunteers), 11 days of 2 g daily raw or heat-treated ginger reduced muscle pain 24 hours after pain-inducing eccentric exercise by roughly **23% to 25%** compared with placebo.¹⁵

Scope note: Exercise-induced muscle pain in small samples, not low back pain.

Dosage commonly used: 500 to 1,000 mg standardized extract daily **Caution:** May interact with blood thinners.

6.8 Methylsulfonylmethane (MSM)

What it does: An organic sulfur compound studied for anti-inflammatory and antioxidant effects.

Evidence: In a pilot randomized trial, MSM at 3 g twice daily produced significant decreases in pain and physical function impairment on the WOMAC scale in **knee osteoarthritis**, without major adverse events. The authors stated that benefits and long-term safety could not be confirmed from a pilot trial.¹⁶ A later randomized, double-blind, placebo-controlled trial in 88 healthy Japanese adults with **mild knee pain** found that 2,000 mg per day of MSM for 12 weeks improved knee-related quality of life scores compared with placebo.¹⁷

Scope note: Both trials studied the knee. There is no low back pain trial evidence for MSM presented here.

Dosage used in trials: 2,000 to 6,000 mg daily **Caution:** Generally well tolerated.

6.9 Palmitoylethanolamide (PEA)

What it does: A naturally occurring fatty acid amide studied for its effects on mast cells and glial cells, the immune-like cells involved in neuroinflammation.

Evidence: A pooled data meta-analysis of 12 studies of micronized and ultra-micronized PEA in chronic and neuropathic pain found a progressive reduction in pain intensity significantly greater than in controls, roughly 1 point every 2 weeks on the pain scale versus 0.2 points in controls. A pain score of 3 or less was reached by 81% of PEA-treated patients compared with 40.9% of controls.¹⁸

Important caveat: Only 3 of the 12 pooled studies were double-blind. Pooling open-label studies with controlled trials inflates apparent benefit, so treat this as promising rather than settled, and note the review was not specific to low back pain.

Dosage commonly used: 300 to 600 mg twice daily, micronized form The upper end of that range is the dose used in a randomized trial of ultra-micronized PEA at 600 mg twice daily in older adults with chronic back, joint, or limb pain.⁵² **Caution:** Well tolerated in the studies reported.

6.10 Glucosamine and Chondroitin

What it does: Components of cartilage, widely marketed for joint and disc health.

Evidence, and a clear negative result: In a randomized, double-blind, placebo-controlled trial of people with **chronic low back pain and degenerative lumbar osteoarthritis**, six months of oral glucosamine did

not reduce pain-related disability compared with placebo, either at the end of treatment or at one-year follow-up.¹⁹

What this means for you: This is the single most directly relevant supplement trial in this chapter, and it is negative. Glucosamine is not recommended here for low back pain. If you take it for a knee or another joint, that is a separate conversation with your provider.

What Was Removed From This Chapter, and Why

An earlier draft of this guide recommended **proteolytic enzymes (bromelain and serrapeptase)** on the strength of a cited review. That citation could not be traced to a real paper supporting a clinical benefit for back pain or soft tissue swelling, so the section was removed rather than left standing on an unverifiable claim. If your provider recommends these products based on evidence they can show you, that is their call to make with you.

7. Movement and Exercise for Low Back Recovery

The most important message: bed rest is not the answer. Gentle, graded movement is the foundation of recovery. Guidelines consistently recommend advice to remain active, along with patient education and supervised exercise.⁶

Exercise therapy for chronic low back pain is supported by moderate-certainty evidence that it probably reduces pain compared with no treatment, usual care, or placebo. The effect on functional limitations is small, and the advantage over other conservative treatments is small and often below the threshold clinicians consider clinically important.²⁰ In plain terms: exercise helps, it is one of the better-supported options you have, and it is not magic. Consistency matters more than the specific method.

Before You Begin: The Pre-Treatment Warm-Up

Best for: Everyone, before any of the exercises below. This is not the treatment. It is what makes the treatment possible to do without guarding.

This is the same soft-tissue and mobility sequence used in office at Root Health before loading work, tools and doses unchanged from the sheet patients receive. Use a percussion massage device, a foam roller, or your own hands, whichever you have, and respect a 6 out of 10 pressure ceiling throughout.

Cat-Camel: 1. On hands and knees, arch your back up, then let it sag. 2. 1 set of 10 repetitions.

Hamstring Stretch with a Strap: 1. Lying on your back, loop a strap or towel around one foot. 2. Straighten the leg toward the ceiling with a soft knee. 3. 1 set of 10 reps each side.

Ball Bounce: 1. Seated on an exercise ball, palms up. 2. Bounce gently up and down, then side to side. 3. 1 set of 10 each direction.

Glute Release: 1. Lying on your back, place a massage ball or foam roller under one glute. 2. Move that leg up and down to work the tissue. 3. Work each side.

TFL and Hip Flexor Release: 1. With a percussion device, run it over the TFL and hip flexors. 2. 1 to 2 minutes each side.

Psoas Release: 1. With a massage ball or a lacrosse ball, place it over the front of the abdomen where the psoas runs. 2. Rock gently back and forth, then up and down. 3. 20 to 30 seconds each side.

This one is unusual and it is deliberate: the psoas connects the lumbar spine to the leg, and it rarely gets touched by anything else.

Lower Back Rest: 1. Lie on your back with a Posture Curve or a rolled towel under your lower back, knees bent. 2. Straighten the legs if comfortable. 3. Rest for 30 to 60 seconds.

Calf Release: 1. With a roller, find a sore spot on the calf. 2. Move the foot up, down, left, right, and in circles. 3. 2 to 3 repetitions per area.

Why it is here: the calves are part of the chain that decides how your gait loads your spine, which is also why this sequence ends there rather than stopping at the back. Do this before the exercises below, not instead of them. The evidence in the rest of this chapter is for the movement. This routine is what gets you loose enough to do it.

7.1 Walking

Best for: Everyone, especially during acute flare-ups.

Protocol: Start with 10 to 15 minutes on flat, even surfaces, and gradually increase toward 30 to 45 minutes daily. Maintain an upright, relaxed posture, swing your arms naturally, and increase your pace and distance as tolerated.

Evidence: A systematic review and meta-analysis of randomized trials found that pain, disability, quality of life, and fear-avoidance improved **similarly** with walking and with other physical exercise in chronic low back pain. Adding walking on top of exercise did not produce further short-term improvement. The authors concluded that walking may be a less expensive alternative to other physical activity.²¹

Why this is good news: You do not need a gym or equipment to get a large share of the available benefit.

7.2 McKenzie Extension Exercises

Best for: Disc-related pain that worsens with sitting and bending and eases with standing and walking.

Prone on Elbows: 1. Lie face down with elbows under your shoulders 2. Relax your lower back completely 3. Hold for 2 to 3 minutes, breathing deeply 4. Repeat 3 to 5 times daily

Press-Up: 1. Lie face down, hands under shoulders 2. Keep hips and pelvis on the floor 3. Press your upper body up, extending your arms 4. Hold for 2 seconds, lower slowly 5. Start with 5 reps, progress to 10 6. Perform every 2 hours during an acute phase

What to look for: Many people with disc-related pain notice that symptoms move out of the leg and toward the midline of the back with repeated extension. Clinicians call this centralization and generally treat it as a favorable sign. If a movement consistently increases your leg symptoms, stop and check with your provider.

7.3 Core Stabilization (The “Big Three”)

These three exercises, popularized by Dr. Stuart McGill, target the deep core stabilizers without heavy spinal loading.

Modified Curl-Up: 1. Lie on your back, one knee bent, one straight 2. Place hands under the small of your back to maintain your neutral lumbar curve 3. Lift your head and shoulders slightly off the floor, not a full sit-up

4. Hold for 10 seconds, breathing normally 5. Perform 5 reps, progressing hold time

Side Bridge (Side Plank): 1. Lie on your side, propped on your forearm 2. Lift your hips so your body forms a straight line 3. Hold for 10 seconds 4. Perform 3 reps per side

Bird Dog: 1. Start on hands and knees 2. Extend one arm forward and the opposite leg back 3. Keep your core engaged and your back flat 4. Hold for 10 seconds, return to start 5. Perform 5 reps per side

Evidence: A review of five studies including 414 patients found core stability exercise more effective than general exercise for reducing pain and improving back-specific function **at 3 months**, with no significant difference remaining at 6 months.²² Separately, a trial of specific stabilizing exercise after a first episode of low back pain reported markedly fewer recurrences in the exercise group: **30% versus 84%** at one year, and **35% versus 75%** at two to three years.²³

Reading that honestly: The short-term advantage over general exercise fades. The recurrence-prevention finding is striking but comes from a single small trial. Both point the same direction, which is that doing the work is what counts.

7.4 Hip Mobility and Glute Strengthening

Weak or inhibited glutes tend to leave the lower back doing more of the work.

Glute Bridge: 1. Lie on your back, knees bent, feet flat 2. Lift your hips until knees, hips, and shoulders form a straight line 3. Squeeze your glutes at the top 4. Hold for 3 seconds, lower slowly 5. Perform 2 to 3 sets of 10 to 15 reps

Clamshell: 1. Lie on your side, knees bent at 45 degrees 2. Keeping your feet together, lift the top knee like a clamshell opening 3. Hold for 2 seconds, lower slowly 4. Perform 2 to 3 sets of 15 reps per side

Why it makes sense: The gluteus maximus and medius are primary hip extensors and pelvic stabilizers. Strengthening them is a reasonable, low-risk part of a general exercise program.

7.5 Hamstring and Hip Flexor Stretching

Supine Hamstring Stretch: 1. Lie on your back, loop a towel or strap around one foot 2. Gently straighten the leg toward the ceiling 3. Hold for 30 seconds, repeat 3 times per leg

Half-Kneeling Hip Flexor Stretch: 1. Kneel on one knee, other foot forward in a lunge 2. Tuck your pelvis under (posterior tilt) 3. Gently shift weight forward 4. Hold for 30 seconds, repeat 3 times

7.6 Yoga and Pilates

Best for: Chronic low back pain, stress reduction, flexibility, and body awareness.

Evidence for yoga: A Cochrane review of 21 trials including 2,223 participants found low- to moderate-certainty evidence that yoga, compared with **no exercise**, produces small improvements in back-related function and pain that the reviewers described as clinically unimportant. Compared with **other back-focused exercise**, there is probably little or no difference in function at three months. Yoga was associated with more adverse events than no exercise, and a similar risk to other exercise.²⁴

Evidence for Pilates: A Cochrane review of 10 trials with 510 participants found low- to moderate-quality evidence that Pilates is more effective than **minimal intervention** for pain and disability, and only a small effect on function compared with **other exercises**. There was no high-quality evidence for any comparison, and no conclusive evidence that Pilates is superior to other forms of exercise.²⁵

The practical conclusion: Choose the movement practice you will actually keep doing. The research does not crown a winner, and both reviews point toward preference, cost, and availability as the deciding factors.

Gentle options to start with: Cat-cow, child's pose, sphinx pose, supine twist, happy baby. Focus on gentle, mindful movement rather than aggressive stretching.

8. Integrative and Physical Therapies

8.1 Physical Therapy

Physical therapy remains a cornerstone of low back pain management. A skilled physical therapist will assess your movement patterns, muscle imbalances, and joint mobility, prescribe individualized exercise progressions, and use manual therapy such as joint mobilization and soft tissue work. They will also educate you on body mechanics and posture, and help you address fear of movement while gradually reintroducing activity.

Evidence: See the exercise therapy evidence in Chapter 7, which found moderate-certainty evidence that exercise probably reduces pain in chronic low back pain compared with no treatment, usual care, or placebo.²⁰ Guidelines recommend supervised exercise and manual therapy when treatment is needed.⁶

8.2 Chiropractic Care and Spinal Manipulation

Spinal manipulation involves controlled forces applied to spinal joints to improve mobility and reduce pain.

Evidence: A systematic review and meta-analysis in *JAMA* found moderate-quality evidence that spinal manipulative therapy was associated with **modest** improvements in pain and function for **acute** low back pain at up to 6 weeks, alongside transient minor musculoskeletal side effects such as temporary soreness. Heterogeneity between studies was large.²⁶ The American College of Physicians guideline includes spinal manipulation among recommended nonpharmacologic options.²⁷ The 2018 *Lancet* low back pain series similarly emphasizes aligning care with the evidence, reducing the focus on spinal abnormalities, and promoting activity and function.²⁸

Realistic expectation: Modest short-term benefit for acute pain, as part of a broader active plan.

8.3 Acupuncture

Acupuncture involves inserting thin needles into specific points.

Evidence: An individual patient data meta-analysis concluded that acupuncture is effective for chronic musculoskeletal pain, headache, and osteoarthritis pain, that the effects persist over time, and that they cannot be explained solely by placebo effects. The authors concluded that referral for a course of acupuncture is a reasonable option for a patient with chronic pain.²⁹ The American College of Physicians guideline includes acupuncture among recommended nonpharmacologic treatments.²⁷

Where the guidelines disagree: The Danish national clinical guidelines for recent-onset low back pain and lumbar radiculopathy recommend **against** acupuncture.⁶ Reasonable expert bodies read this literature differently. It is worth knowing that before you spend money on a course of treatment.

8.4 Massage Therapy

Evidence: A randomized controlled trial in *Annals of Internal Medicine* comparing relaxation massage, structural massage, and usual care in chronic back pain found that massage may be effective, **with benefits lasting at least 6 months**, and no clinically meaningful difference between the two massage styles.³⁰ A Cochrane review of 25 trials with 3,096 participants was more cautious, stating the reviewers had very little confidence that massage is an effective treatment for low back pain, with pain improvements seen only at short-term follow-up. Adverse effects were minor.³¹

Realistic expectation: Pleasant, low-risk, and possibly helpful in the short term. Not a structural fix, and the highest-quality synthesis is lukewarm.

8.5 Heat and Cold Therapy

Cold (first 48 hours of an acute flare): Numbs pain and may reduce local swelling. Apply ice wrapped in cloth for 15 to 20 minutes.

Heat (after the acute phase, or for muscle spasm): Increases local blood flow, relaxes muscles, and reduces stiffness. Use a heating pad or warm bath for 15 to 20 minutes.

Evidence: A Cochrane review of nine trials with 1,117 participants found moderate evidence in a small number of trials that **heat wrap therapy** provides a small short-term reduction in pain and disability in acute and subacute low back pain, and that adding exercise reduces pain further and improves function. The evidence for **cold** was much more limited, based on only three poor-quality studies, and the reviewers concluded that no conclusions can be drawn about cold for low back pain.³²

Practical translation: Heat has modest support. Cold is not well studied, so use it if it feels good and skip it if it does not.

8.6 Dry Needling

Dry needling targets myofascial trigger points with thin needles.

Evidence: A systematic review and meta-analysis of 16 randomized trials found dry needling more effective than sham needling and acupuncture for reducing pain and disability immediately after the intervention, with effects at follow-up equal to acupuncture. The reviewers stressed that most included studies had high or unclear risk of bias, and that current evidence is not strong enough for a firm conclusion.³³

8.7 Low-Level Laser Therapy (LLLT)

LLLT uses specific wavelengths of light applied to tissue.

Evidence, with a correction: An earlier draft of this guide attributed a favorable LLLT finding to a 2015 systematic review in *The Lancet*. That was wrong on two counts. The *Lancet* review of LLLT was published in

2009 and studied **neck pain**, where it found pain reduction immediately after treatment in acute neck pain and up to 22 weeks after treatment in chronic neck pain.³⁴

For **low back pain** specifically, a separate systematic review and meta-analysis of seven randomized trials including 394 patients found that LLLT significantly reduced pain scores compared with placebo, with **no** significant effect on disability or spinal range of motion.³⁵ LLLT also appears among the nonpharmacologic options in the American College of Physicians guideline.²⁷

Realistic expectation: Possible short-term pain relief, no demonstrated effect on function.

8.8 Cupping Therapy

Evidence: A systematic review and meta-analysis of 11 trials with 921 participants found that cupping significantly improved pain at the 2 to 8 week end of intervention, but that the improvement was **not** maintained at one month or at 3 to 6 months. The reviewers noted notable heterogeneity between studies and called for standardized protocols.³⁶

Realistic expectation: Short-lived benefit, if any. Cupping commonly leaves circular bruise-like marks for several days.

8.9 TENS (Transcutaneous Electrical Nerve Stimulation)

TENS delivers low-voltage electrical currents through skin electrodes.

Evidence, and a correction: An earlier draft described TENS as moderately effective. A Cochrane review of four high-quality randomized trials with 585 patients found **conflicting** evidence on whether TENS reduces chronic back pain intensity and **consistent** evidence in two trials with 410 patients that it did **not** improve back-specific function. The reviewers concluded that the evidence does not support the use of TENS in the routine management of chronic low back pain.³⁷

What this means for you: TENS units are cheap and safe for home use, and some people like them. Do not expect them to change your function, and do not let one substitute for active treatment.

9. The Anti-Inflammatory Diet for Spinal Recovery

What you eat influences inflammation, tissue repair, and overall health. The pattern below is a well-established general healthy-eating approach. It is presented here as sensible nutrition that supports recovery, not as a proven treatment for low back pain.

9.1 Foods to Emphasize

Omega-3 rich foods: Wild-caught salmon, sardines, mackerel, anchovies, walnuts, flaxseeds, chia seeds

Colorful vegetables and fruits: Leafy greens such as spinach, kale, and Swiss chard, berries such as blueberries, strawberries, and blackberries, cruciferous vegetables such as broccoli, cauliflower, and Brussels sprouts, and orange and yellow vegetables such as carrots and sweet potatoes.

Anti-inflammatory spices: Turmeric (with black pepper), ginger, garlic, cinnamon, rosemary

Quality protein: Grass-fed meats, pasture-raised poultry, wild fish, eggs, legumes

Bone broth and collagen sources: Supply the amino acids used in connective tissue repair

Healthy fats: Extra virgin olive oil, avocados, nuts and seeds

9.2 Foods to Minimize

Pro-inflammatory fats: Trans fats, excessive omega-6 vegetable oils, deep-fried foods

Refined carbohydrates and sugars: White bread, pastries, sugary cereals, soda, candy

Processed foods: Deli meats, fast food, packaged snacks

Excessive alcohol: Interferes with sleep and adds empty calories

9.3 Hydration

Spinal discs are largely made of water, and staying well hydrated is part of general health. There is no need to chase a specific ounce target. Drink regularly through the day, keep water with you, and use thirst and pale urine as your guide. If you have heart or kidney conditions, follow your provider's fluid guidance instead.

Tips: Have water with each meal, keep a bottle where you work, and watch caffeine and alcohol late in the day, mostly for sleep quality.

9.4 Sample Anti-Inflammatory Day

Breakfast: Smoothie with spinach, blueberries, collagen powder, flaxseed, ginger, almond butter, and unsweetened almond milk. Or oatmeal with walnuts, chia seeds, cinnamon, and berries.

Lunch: Large salad with mixed greens, grilled salmon, avocado, olive oil and lemon dressing. Side of vegetable soup.

Afternoon snack: Apple with almond butter, or carrot sticks with hummus.

Dinner: Grilled chicken or tofu with turmeric and garlic, roasted broccoli and sweet potato, side salad.

Evening: Ginger or turmeric tea. Small square of dark chocolate (70% cacao or higher).

10. Lifestyle Modifications: Sleep, Stress, and Ergonomics

10.1 Sleep Optimization

Sleep is when your body repairs tissue and regulates inflammation, and poor sleep makes pain worse.

Evidence: A brain imaging study published in the *Journal of Neuroscience* showed that sleep deprivation enhances pain responsivity in the primary sensory regions of the cortex while blunting activity in regions that modulate pain, including the striatum and insula. In a separate general-population sample, even subtle night-to-night changes in sleep predicted day-to-day changes in pain, in both directions.³⁸

Strategies: Maintain a consistent sleep schedule, and sleep on your side with a pillow between your knees, or on your back with a pillow under your knees. Keep your room cool, around 65 to 68 degrees Fahrenheit, limit screens for an hour before bed, and avoid caffeine in the afternoon. If you use magnesium at night, discuss the dose with your provider.

10.2 Stress Management

Chronic stress amplifies pain perception and makes coping harder.

Evidence: A systematic review and meta-analysis of 47 trials with 3,515 participants found that mindfulness meditation programs had moderate evidence of improved anxiety and depression, and moderate evidence of improved **pain** (effect size 0.33). The authors concluded that meditation programs can produce small to moderate reductions in multiple dimensions of psychological stress.³⁹

Effective techniques: Diaphragmatic breathing for 5 minutes twice daily, mindfulness meditation, progressive muscle relaxation, and time outdoors and in nature.

10.3 Ergonomics

Sitting: Use a chair with lumbar support, keep your feet flat with hips slightly higher than knees, take standing or walking breaks every 30 to 45 minutes, and consider a sit-stand desk.

Standing: Shift your weight periodically and use an anti-fatigue mat.

Lifting: Bend at the knees, keep the load close to your body, and avoid twisting while lifting.

Driving: Adjust the seat so your knees are slightly bent, use lumbar support, and take breaks every 1 to 2 hours.

10.4 Weight Management

Evidence: A meta-analysis found that overweight and obesity increase the risk of low back pain, with the strongest associations for seeking care for low back pain (odds ratio 1.56) and for chronic low back pain (odds ratio 1.43).²

An honest caveat: That association does not automatically mean losing weight will treat existing back pain. A systematic review of 22 weight-loss trials with 3,602 participants found small to moderate improvements in pain and disability for **osteoarthritis**, but described the evidence for weight-loss interventions targeting **spinal pain** as limited and inconclusive.⁴⁰

What this means for you: Reaching a healthy weight is worth doing for many good reasons, including reduced risk of a future back pain episode. Treat improvement in current back pain as a possible bonus rather than a guaranteed outcome.

10.5 Smoking Cessation

Evidence: In a study of identical twin pairs highly discordant for cigarette smoking, which controls for genetics and much of shared upbringing, smokers had **18% greater** mean disc degeneration scores across the lumbar spine than their non-smoking twins. The effect appeared across the entire lumbar spine, implicating a systemic mechanism.³ Smokers are also among the groups at greatest risk of reporting low back pain.¹

Quitting is one of the highest-value things you can do for spinal health and for general health. Your provider can help you with a plan.

11. Mind-Body Approaches to Chronic Pain

For chronic low back pain, addressing the mind-body connection is not optional. It is essential. This is not a suggestion that your pain is imaginary. It is a recognition that the nervous system that produces pain is the same one shaped by stress, sleep, fear, and attention.

11.1 Cognitive Behavioral Therapy (CBT)

CBT helps you identify and change unhelpful thoughts and behaviors related to pain, including catastrophizing and fear avoidance.

Evidence: In a randomized trial of 342 adults with chronic low back pain, CBT produced greater improvement in back pain and functional limitations than usual care at 26 weeks. Outcomes were also assessed at 52 weeks, and 84.8% of participants completed the study at that point.⁴¹ CBT is among the nonpharmacologic treatments recommended in the American College of Physicians guideline.²⁷

11.2 Mindfulness-Based Stress Reduction (MBSR)

MBSR teaches present-moment awareness and acceptance of pain without judgment.

Evidence: In the same randomized trial, MBSR also produced greater improvement in back pain and functional limitations than usual care at 26 weeks, with **no significant difference in outcomes between MBSR and CBT**. The authors concluded that MBSR may be an effective treatment option for chronic low back pain.⁴¹

The useful takeaway: Two very different-looking approaches worked about equally well. Pick the one you will attend.

11.3 Biofeedback

Biofeedback uses sensors to help you learn to influence physiological responses such as muscle tension and heart rate.

Evidence: A meta-analysis found a small-to-medium effect of biofeedback on pain intensity reduction in chronic back pain (Hedges' $g = 0.60$), which remained stable over an average 8-month follow-up, along with improvements in depression and disability. The authors concluded biofeedback can improve pain-related outcomes in the short and long term, both alone and as an add-on.⁴² Electromyography biofeedback also appears among the recommended nonpharmacologic options in the American College of Physicians guideline.²⁷

11.4 Acceptance and Commitment Therapy (ACT)

ACT focuses on accepting pain while committing to valued activities, reducing the exhausting struggle against pain.

Evidence: A systematic review and meta-analysis of acceptance-based interventions in chronic pain found an effect size of 0.37 on pain and 0.32 on depression in controlled studies. The authors reported that MBSR and ACT were **not superior to CBT** but can be good alternatives, with small to medium effects on physical and mental health.⁴³

11.5 Graded Exposure and Pacing

Graded exposure involves gradually reintroducing feared or avoided movements. Pacing involves breaking activities into manageable chunks so you are not swinging between overdoing it and shutting down.

Evidence: In a randomized controlled trial in patients with chronic back pain, graded in vivo exposure produced significantly greater improvements than graded activity on measures of fear of pain and movement, fear-avoidance beliefs, pain-related anxiety, and pain self-efficacy, and greater improvements than a wait-list control across those measures plus pain catastrophizing and pain experience. Gains were maintained at one-month follow-up. Differences on pain-related disability itself were at the level of a trend.⁴⁴

Reading that honestly: Graded exposure reliably reduces fear. Its direct effect on disability in that trial was less clear-cut. Reducing fear is still worth a great deal, because fear is what keeps people from moving.

12. The In-Clinic Pre-Treatment Protocol

Understand what this is for, because the distinction matters.

None of this rebuilds tissue. The loading and strengthening work in the previous chapters does that. What this protocol does is reduce the muscle tone, stiffness and tissue sensitivity that make the strength work harder to tolerate and less pleasant to do, particularly in the early weeks when things are at their most irritable.

It is preparation, not treatment. I use it in the clinic before the working part of a session, and I give it to patients to use at home before theirs. Patients who do it tend to tolerate the loading better and stick with it longer, and sticking with it is the actual mechanism by which recovery happens.

If you have to choose between doing this and doing your strength work, do your strength work.

Heat and vibration back wrap, set to vibrate and heat, while performing:

- Cat-camel stretch, 1 set of 10.
- Hamstring stretch with a stretch strap, 1 set of 10 each side.
- Ball bounce on an exercise ball, up and down then side to side, 1 set of 10 each, palms up.

Vibrating massage ball or foam roller. Lie on your back with the sphere on the glute muscles, moving the leg up and down.

Percussion massager, round or flat head. Over the tensor fasciae latae and hip flexors, 1 to 2 minutes each side.

Massage ball over the abdomen, working the psoas, rocking back and forth and up and down, 20 to 30 seconds each side.

Posture Curve under the lower back, knees bent to start and straightening if comfortable, resting 30 to 60 seconds.

Vibrating sphere or foam roller on sore calf areas, moving the foot up, down, left, right and in circles, 2 to 3 reps per area.

Safety, and I mean these.

- Avoid all of this over an acutely injured area in the first 48 to 72 hours.
- Reduce pressure over anything actively inflamed.
- Skip percussion therapy entirely if you have a pacemaker or another implanted electronic device.
- Avoid aggressive pressure over varicose veins or compromised skin.

- Start light. Pain must not exceed 6 out of 10. Back off immediately for sharp or shooting pain.
- Stop and seek advice if you get numbness, tingling, or weakness.

One absolute rule before you roll or percuss a calf. Never work on a calf that is swollen, warm, or tight in one leg only, particularly after travel, surgery, or a period of immobilisation. That is a blood clot concern. It needs assessing the same day and it must never be massaged.

13. Red Flags: When to Seek Immediate Care

Seek immediate care if you experience:

- **Cauda equina syndrome:** Loss of bowel or bladder control, urinary retention, or numbness in the saddle region (inner thighs, groin, buttocks). **This is a surgical emergency.**
- Progressive weakness in the legs or feet
- Severe, unrelenting pain not improved by rest
- Fever with back pain
- A history of cancer with new onset back pain
- Unexplained weight loss with back pain
- Recent significant trauma (fall, car accident)
- Intravenous drug use with back pain and fever (possible infection)

Contact your provider promptly if:

- Symptoms worsen after 2 weeks of conservative care
- Pain prevents sleep despite medication
- New numbness, tingling, or weakness develops
- Symptoms persist beyond 6 to 8 weeks without improvement

Guidelines recommend that care include clear information about prognosis and warning signs, along with advice to remain active.⁶

14. Building Your Personal Recovery Plan

Use this as a template to discuss with your provider, not as a prescription. Adjust anything that does not fit your situation, and skip any supplement your provider advises against.

Week 1 to 2: Calm and Protect

- **Movement:** Short walks, gentle McKenzie extensions if they help your symptoms, breathing exercises
- **Therapies:** Heat for muscle spasm, initial physical therapy or chiropractic evaluation
- **Diet:** Begin anti-inflammatory eating, stay well hydrated
- **Lifestyle:** Prioritize sleep, reduce stress, adjust your workstation
- **Supplements (optional, with your provider):** Omega-3 is the most reasonable starting point given its safety profile

Week 3 to 6: Rebuild

- **Movement:** Progress to the core Big Three, glute strengthening, hip mobility
- **Therapies:** Regular physical therapy, and yoga or Pilates based on your preference
- **Diet:** Continue the anti-inflammatory pattern
- **Lifestyle:** Implement stress management, optimize sleep hygiene

Week 7 to 12: Strengthen and Restore

- **Movement:** Advanced core work, functional movement, return to normal activities
- **Therapies:** Maintenance care as needed, biofeedback if pain is becoming persistent
- **Diet:** Anti-inflammatory eating as a lifestyle, not a diet
- **Lifestyle:** Maintain a healthy weight, keep exercising, keep managing stress

Beyond 12 Weeks: Maintenance and Prevention

- Continue core and general exercise 3 to 4 times per week
- Maintain the anti-inflammatory eating pattern
- Regular walking or low-impact cardio
- Address flare-ups early rather than waiting
- For persistent pain, keep up a mind-body practice (CBT, MBSR, or ACT)

Final Thoughts

Low back pain is not a life sentence. Your spine is strong, adaptable, and designed to move. The evidence points clearly toward active, whole-person self-care rather than passive treatment alone.

That means moving your body regularly and mindfully, nourishing yourself with good food, sleeping well, managing stress, and addressing the psychological and social dimensions of pain. It means understanding that pain does not always equal damage, and that you can hurt without being harmed.

The biopsychosocial model is not just a theory. It is a practical roadmap. When you address your biology (exercise, nutrition, sleep), your psychology (thoughts, emotions, fear), and your social context (work, relationships, support), you create the conditions for genuine, lasting improvement.

There will be good days and difficult days. Healing is rarely linear. But with consistency, patience, and the right tools, you can recover from this episode and build a stronger, more resilient back along the way.

You are not broken. You are healing.

15. References

Every reference below was retrieved and checked against the published record. PMID numbers are included so you or your provider can look up any claim in this guide directly.

1. Hartvigsen J, Hancock MJ, Kongsted A, et al. What low back pain is and why we need to pay attention. *Lancet*. 2018;391(10137):2356-2367. PMID: 29573870
2. Shiri R, Karppinen J, Leino-Arjas P, Solovieva S, Viikari-Juntura E. The association between obesity and low back pain: a meta-analysis. *Am J Epidemiol*. 2010;171(2):135-154. PMID: 20007994
3. Battié MC, Videman T, Gill K, et al. 1991 Volvo Award in clinical sciences. Smoking and lumbar intervertebral disc degeneration: an MRI study of identical twins. *Spine (Phila Pa 1976)*. 1991;16(9):1015-1021. PMID: 1948392
4. da Silva T, Mills K, Brown BT, Herbert RD, Maher CG, Hancock MJ. Risk of recurrence of low back pain: a systematic review. *J Orthop Sports Phys Ther*. 2017;47(5):305-313. PMID: 28355981
5. Brinjikji W, Luetmer PH, Comstock B, et al. Systematic literature review of imaging features of spinal degeneration in asymptomatic populations. *AJNR Am J Neuroradiol*. 2015;36(4):811-816. PMID: 25430861
6. Stochkendahl MJ, Kjaer P, Hartvigsen J, et al. National clinical guidelines for non-surgical treatment of patients with recent onset low back pain or lumbar radiculopathy. *Eur Spine J*. 2018;27(1):60-75. PMID: 28429142
7. da C Menezes Costa L, Maher CG, Hancock MJ, McAuley JH, Herbert RD, Costa LO. The prognosis of acute and persistent low-back pain: a meta-analysis. *CMAJ*. 2012;184(11):E613-E624. PMID: 22586331
8. Maroon JC, Bost JW. Omega-3 fatty acids (fish oil) as an anti-inflammatory: an alternative to nonsteroidal anti-inflammatory drugs for discogenic pain. *Surg Neurol*. 2006;65(4):326-331. PMID: 16531187
9. Daily JW, Yang M, Park S. Efficacy of turmeric extracts and curcumin for alleviating the symptoms of joint arthritis: a systematic review and meta-analysis of randomized clinical trials. *J Med Food*. 2016;19(8):717-729. PMID: 27533649
10. Kuptniratsaikul V, Dajpratham P, Taechaarpornkul W, et al. Efficacy and safety of *Curcuma domestica* extracts compared with ibuprofen in patients with knee osteoarthritis: a multicenter study. *Clin Interv Aging*. 2014;9:451-458. PMID: 24672232
11. Bannuru RR, Osani MC, Al-Eid F, Wang C. Efficacy of curcumin and Boswellia for knee osteoarthritis: systematic review and meta-analysis. *Semin Arthritis Rheum*. 2018;48(3):416-429. PMID: 29622343
12. Zadro JR, Shirley D, Ferreira M, et al. Is vitamin D supplementation effective for low back pain? A systematic review and meta-analysis. *Pain Physician*. 2018;21(2):121-145. PMID: 29565945
13. Tarsitano MG, Quinzi F, Folino K, et al. Effects of magnesium supplementation on muscle soreness in different type of physical activities: a systematic review. *J Transl Med*. 2024;22(1):629. PMID: 38970118

14. Shaw G, Lee-Barthel A, Ross ML, Wang B, Baar K. Vitamin C-enriched gelatin supplementation before intermittent activity augments collagen synthesis. *Am J Clin Nutr.* 2017;105(1):136-143. PMID: 27852613
15. Black CD, Herring MP, Hurley DJ, O'Connor PJ. Ginger (*Zingiber officinale*) reduces muscle pain caused by eccentric exercise. *J Pain.* 2010;11(9):894-903. PMID: 20418184
16. Kim LS, Axelrod LJ, Howard P, Buratovich N, Waters RF. Efficacy of methylsulfonylmethane (MSM) in osteoarthritis pain of the knee: a pilot clinical trial. *Osteoarthritis Cartilage.* 2006;14(3):286-294. PMID: 16309928
17. Toguchi A, Noguchi N, Kanno T, Yamada A. Methylsulfonylmethane improves knee quality of life in participants with mild knee pain: a randomized, double-blind, placebo-controlled trial. *Nutrients.* 2023;15(13):2995. PMID: 37447322
18. Paladini A, Fusco M, Cenacchi T, Schievano C, Piroli A, Varrassi G. Palmitoylethanolamide, a special food for medical purposes, in the treatment of chronic pain: a pooled data meta-analysis. *Pain Physician.* 2016;19(2):11-24. PMID: 26815246
19. Wilkens P, Scheel IB, Grundnes O, Hellum C, Storheim K. Effect of glucosamine on pain-related disability in patients with chronic low back pain and degenerative lumbar osteoarthritis: a randomized controlled trial. *JAMA.* 2010;304(1):45-52. PMID: 20606148
20. Hayden JA, Ellis J, Ogilvie R, Malmivaara A, van Tulder MW. Exercise therapy for chronic low back pain. *Cochrane Database Syst Rev.* 2021;9(9):CD009790. PMID: 34580864
21. Vanti C, Andreatta S, Borghi S, Guccione AA, Pillastrini P, Bertozzi L. The effectiveness of walking versus exercise on pain and function in chronic low back pain: a systematic review and meta-analysis of randomized trials. *Disabil Rehabil.* 2019;41(6):622-632. PMID: 29207885
22. Coulombe BJ, Games KE, Neil ER, Eberman LE. Core stability exercise versus general exercise for chronic low back pain. *J Athl Train.* 2017;52(1):71-72. PMID: 27849389
23. Hides JA, Jull GA, Richardson CA. Long-term effects of specific stabilizing exercises for first-episode low back pain. *Spine (Phila Pa 1976).* 2001;26(11):E243-E248. PMID: 11389408
24. Wieland LS, Skoetz N, Pilkington K, et al. Yoga for chronic non-specific low back pain. *Cochrane Database Syst Rev.* 2022;11(11):CD010671. PMID: 36398843
25. Yamato TP, Maher CG, Saragiotto BT, et al. Pilates for low back pain. *Cochrane Database Syst Rev.* 2015;2015(7):CD010265. PMID: 26133923
26. Paige NM, Miake-Lye IM, Booth MS, et al. Association of spinal manipulative therapy with clinical benefit and harm for acute low back pain: systematic review and meta-analysis. *JAMA.* 2017;317(14):1451-1460. PMID: 28399251. (Errata: *JAMA.* 2017;317(21):2239, an error in 2 figures, PMID 28586870; and *JAMA.* 2017;318(20):2048, incorrect data in text, PMID 29183047. Screened 2026-09-05; corrected, not retracted. The correction text is subscription-gated and was not read. This guide reproduces no numeric value from this paper, so no figure here depends on it.)

27. Qaseem A, Wilt TJ, McLean RM, Forciea MA. Noninvasive treatments for acute, subacute, and chronic low back pain: a clinical practice guideline from the American College of Physicians. *Ann Intern Med.* 2017;166(7):514-530. PMID: 28192789
28. Foster NE, Anema JR, Cherkin D, et al. Prevention and treatment of low back pain: evidence, challenges, and promising directions. *Lancet.* 2018;391(10137):2368-2383. PMID: 29573872
29. Vickers AJ, Vertosick EA, Lewith G, et al. Acupuncture for chronic pain: update of an individual patient data meta-analysis. *J Pain.* 2018;19(5):455-474. PMID: 29198932
30. Cherkin DC, Sherman KJ, Kahn J, et al. A comparison of the effects of 2 types of massage and usual care on chronic low back pain: a randomized, controlled trial. *Ann Intern Med.* 2011;155(1):1-9. PMID: 21727288
31. Furlan AD, Giraldo M, Baskwill A, Irvin E, Imamura M. Massage for low-back pain. *Cochrane Database Syst Rev.* 2015;2015(9):CD001929. PMID: 26329399
32. French SD, Cameron M, Walker BF, Reggars JW, Esterman AJ. Superficial heat or cold for low back pain. *Cochrane Database Syst Rev.* 2006;2006(1):CD004750. PMID: 16437495
33. Hu HT, Gao H, Ma RJ, Zhao XF, Tian HF, Li L. Is dry needling effective for low back pain? A systematic review and PRISMA-compliant meta-analysis. *Medicine (Baltimore).* 2018;97(26):e11225. PMID: 29952980
34. Chow RT, Johnson MI, Lopes-Martins RA, Bjordal JM. Efficacy of low-level laser therapy in the management of neck pain: a systematic review and meta-analysis of randomised placebo or active-treatment controlled trials. *Lancet.* 2009;374(9705):1897-1908. PMID: 19913903. (Erratum: *Lancet.* 2010;375(9718):894. Screened 2026-09-05; corrected, not retracted.)
35. Huang Z, Ma J, Chen J, Shen B, Pei F, Kraus VB. The effectiveness of low-level laser therapy for nonspecific chronic low back pain: a systematic review and meta-analysis. *Arthritis Res Ther.* 2015;17:360. PMID: 26667480
36. Zhang Z, Pasapula M, Wang Z, Edwards K, Norrish A. The effectiveness of cupping therapy on low back pain: a systematic review and meta-analysis of randomized control trials. *Complement Ther Med.* 2024;80:103013. PMID: 38184285
37. Khadilkar A, Odebiyi DO, Brosseau L, Wells GA. Transcutaneous electrical nerve stimulation (TENS) versus placebo for chronic low-back pain. *Cochrane Database Syst Rev.* 2008;2008(4):CD003008. PMID: 18843638
38. Krause AJ, Prather AA, Wager TD, Lindquist MA, Walker MP. The pain of sleep loss: a brain characterization in humans. *J Neurosci.* 2019;39(12):2291-2300. PMID: 30692228
39. Goyal M, Singh S, Sibinga EM, et al. Meditation programs for psychological stress and well-being: a systematic review and meta-analysis. *JAMA Intern Med.* 2014;174(3):357-368. PMID: 24395196
40. Robson EK, Hodder RK, Kamper SJ, et al. Effectiveness of weight-loss interventions for reducing pain and disability in people with common musculoskeletal disorders: a systematic review with meta-analysis. *J Orthop Sports Phys Ther.* 2020;50(6):319-333. PMID: 32272032

41. Cherkin DC, Sherman KJ, Balderson BH, et al. Effect of mindfulness-based stress reduction vs cognitive behavioral therapy or usual care on back pain and functional limitations in adults with chronic low back pain: a randomized clinical trial. *JAMA*. 2016;315(12):1240-1249. PMID: 27002445
42. Sielski R, Rief W, Glombiewski JA. Efficacy of biofeedback in chronic back pain: a meta-analysis. *Int J Behav Med*. 2017;24(1):25-41. PMID: 27307013
43. Veehof MM, Oskam MJ, Schreurs KMG, Bohlmeijer ET. Acceptance-based interventions for the treatment of chronic pain: a systematic review and meta-analysis. *Pain*. 2011;152(3):533-542. PMID: 21251756
44. Woods MP, Asmundson GJG. Evaluating the efficacy of graded in vivo exposure for the treatment of fear in patients with chronic back pain: a randomized controlled clinical trial. *Pain*. 2008;136(3):271-280. PMID: 17716819
45. DePalma MJ, Ketchum JM, Saullo T. What is the source of chronic low back pain and does age play a role? *Pain Med*. 2011;12(2):224-233. PMID: 21266006
46. Laslett M. Evidence-based diagnosis and treatment of the painful sacroiliac joint. *J Man Manip Ther*. 2008;16(3):142-152. PMID: 19119403
47. Fortin JD, Dwyer AP, West S, Pier J. Sacroiliac joint: pain referral maps upon applying a new injection/arthrography technique. Part I: Asymptomatic volunteers. *Spine (Phila Pa 1976)*. 1994;19(13):1475-1482. PMID: 7939978
48. Jung HS, Jeon HS, Oh DW, Kwon OY. Effect of the pelvic compression belt on the hip extensor activation patterns of sacroiliac joint pain patients during one-leg standing: a pilot study. *Man Ther*. 2013;18(2):143-148. PMID: 23111368
49. Pizol GZ, Miyamoto GC, Cabral CMN. Hip biomechanics in patients with low back pain, what do we know? A systematic review. *BMC Musculoskelet Disord*. 2024;25(1):415. PMID: 38807086
50. Alkhathami KM, Alqahtani B. Comparing the scores of the Functional Movement Screen in individuals with low back pain versus healthy individuals: a systematic review and meta-analysis. *Int J Sports Phys Ther*. 2024;19(7):834-848. PMID: 38966824
51. Trinidad-Fernandez M, Gonzalez-Sanchez M, Cuesta-Vargas AI. Is a low Functional Movement Screen score (14/21 or below) associated with injuries in sport? A systematic review and meta-analysis. *BMJ Open Sport Exerc Med*. 2019;5(1):e000501. PMID: 31673399
52. Germini F, Coerezza A, Andreinetti L, et al. N-of-1 Randomized Trials of Ultra-Micronized Palmitoylethanolamide in Older Patients with Chronic Pain. *Drugs Aging*. 2017;34(12):941-952. PMID: 29210011

Working with Root Health

If your back pain has not turned the corner, and you would like an evaluation that looks at your hips and your loading tolerance rather than stopping at the spot that hurts, I offer a free Root Discovery Session at Root Health to talk through what may be contributing and what a reasonable next step looks like. You can schedule one at rootcauseunlocked.com/discovery, or reach the office at (919) 662-0520.

Consultations and ongoing care are both available wherever you are located. Being outside North Carolina is not a barrier to working with me. Tell me where you are when you reach out and I will lay out exactly how we would work together from there.

If you would rather start on your own, the free Root Cause Unlocked tracker has a mode built for musculoskeletal recovery, at rootcauseunlocked.com/tracker. Eight weeks of it turns “I think it might be helping” into something you and I can both read.

This guide was compiled from peer-reviewed medical literature and clinical best practices. Every citation was verified against the published record. It is intended to give you knowledge and options to discuss with your healthcare provider. It is not a substitute for individual medical care, and it does not promise a cure.

© 2026 · For personal educational use only

Disclaimer: This ebook is for educational purposes only and does not constitute medical advice. Always consult with a qualified healthcare provider before starting any new treatment, supplement, exercise program, or dietary change. Do not start, stop, or change any prescribed medication without talking to the provider who prescribed it. If you experience progressive weakness, loss of bowel or bladder control, or severe worsening pain, seek emergency medical attention immediately.

A note on the evidence in this guide. Every claim below that carries a numbered marker has been checked against the actual published study. Where a statement is general clinical background rather than a specific research finding, it carries no marker. Claims that could not be traced to a real, on-point study were removed rather than softened.

1. What Is Sciatica?

Sciatica refers to pain that radiates along the path of the sciatic nerve, the largest and longest nerve in your body. This nerve branches from your lower back through your hips and buttocks and down each leg. Typically, sciatica affects only one side of the body.^{1,2}

Contrary to popular belief, sciatica is not a condition itself but rather a symptom of an underlying problem affecting the sciatic nerve or its nerve roots. Understanding this distinction matters, because effective treatment means addressing the source of the irritation, not just masking the pain.²

Common Symptoms

Common symptoms include sharp, burning, or shooting pain that travels from the lower back down the leg, and numbness, tingling, or “pins and needles” in the leg or foot. Muscle weakness in the affected leg can occur, and pain often worsens with sitting, coughing, sneezing, or straining. Some people also notice difficulty moving the leg or foot.

2. Understanding the Anatomy

The sciatic nerve is formed by the convergence of nerve roots from the L4, L5, S1, S2, and S3 spinal segments. It exits the pelvis through the greater sciatic foramen and travels down the posterior thigh, branching into smaller nerves that serve the lower leg and foot.

Because the nerve is so long, irritation can occur at multiple points along its path. At the spinal level, herniated discs, spinal stenosis, or spondylolisthesis can compress a nerve root. At the pelvic level, piriformis syndrome occurs when the piriformis muscle compresses the nerve. And at the hamstring level, entrapment can occur from fibrous bands or hamstring tendons.

Understanding where your symptoms originate helps guide the most effective treatment approach.

3. Common Causes and Risk Factors

Primary Causes

Lumbar Herniated Disc (the most common cause): the gel-like center of an intervertebral disc pushes through the tough outer layer, pressing on a nerve root. Lumbar disc herniation is the most frequently identified structural cause of sciatica, although published estimates of how often it is responsible vary widely between studies and definitions.^{1,2}

Degenerative Disc Disease: as discs age, they lose height and water content, reducing the space where nerves exit the spine.

Lumbar Spinal Stenosis: narrowing of the spinal canal, often age-related, compresses the nerve roots.

Spondylolisthesis: one vertebra slips forward over another, narrowing the nerve exit space.

Piriformis Syndrome: the piriformis muscle, deep in the buttock, spasms or tightens and compresses the sciatic nerve. This is a recognized but less common source of sciatic-type pain.

Risk Factors

Several factors increase the risk of developing sciatica. Age is one, since it is most common between 30 and 50 years.¹ Prolonged sitting or a sedentary lifestyle, and occupations requiring heavy lifting or twisting, both add mechanical stress. Obesity increases spinal load and is an established risk factor for sciatica.^{46,47} Diabetes increases nerve damage risk, smoking is associated with reduced blood flow to spinal discs, and pregnancy carries its own risk through hormonal changes and shifts in weight distribution.

4. How Sciatica Is Diagnosed

A proper diagnosis typically involves:

Clinical Examination

A clinical examination typically includes a straight leg raise test (Lasègue's sign), a neurological examination of strength, reflexes, and sensation, gait analysis, and palpation of the spine and pelvis.

Imaging (when needed)

When imaging is needed, MRI is the preferred study for visualizing soft tissues, discs, and nerve compression. CT scanning is useful for bony detail, while X-rays have limited utility for soft tissue but are helpful for ruling out fractures or alignment issues. EMG and nerve conduction studies assess nerve function when the diagnosis is unclear.

Important: many people have MRI findings such as disc bulges without any symptoms, and clinical guidelines advise against routine imaging for back pain in the absence of red flags or a plan for surgery. Imaging should correlate with your clinical presentation, not drive treatment decisions on its own.³

5. The Healing Timeline: What to Expect

Understanding the natural history of sciatica helps set realistic expectations.

Acute phase (0 to 6 weeks): pain is usually most intense. A large share of cases improve substantially during this window with conservative care alone. Prospective follow-up of sciatica associated with disc pathology has shown that symptoms and the disc herniation itself often resolve over time without surgery.⁴

Subacute phase (6 to 12 weeks): gradual improvement continues. This is when targeted rehabilitation becomes most important.

Chronic phase (beyond 12 weeks): if pain persists, a more comprehensive, multidisciplinary approach is warranted.

Key insight. Randomized evidence comparing early surgery with prolonged conservative care for sciatica from a lumbar disc herniation found that early surgery gave faster relief of leg pain, but that outcomes at one year were similar between the two groups.⁶ The SPORT randomized trial and a systematic review of surgery versus conservative management reached a compatible conclusion: surgery tends to speed up recovery, while long-term outcomes converge.^{5,7} Ten-year follow-up from the Maine Lumbar Spine Study likewise found that patients treated surgically and nonsurgically reported broadly similar long-term outcomes on several measures.⁸

This is why an unhurried, well-structured conservative plan is a reasonable first step for most people who do not have red-flag findings.

6. Evidence-Based Supplements for Nerve Health and Inflammation

The supplements below have research support of varying strength. Read the scope of each finding carefully. Much of the strongest supplement research in this area comes from joint pain or diabetic nerve pain, not from sciatica specifically, and that distinction is stated wherever it applies.

Always discuss supplements with your healthcare provider, especially if you take medications, since nothing here is a cure or a substitute for prescribed treatment.

6.1 Omega-3 Fatty Acids (EPA and DHA)

What it does: EPA and DHA are incorporated into cell membranes and serve as precursors to resolvins and protectins, a family of mediators involved in actively resolving inflammation rather than only suppressing it. Marine omega-3 fatty acids influence multiple steps in the inflammatory process.⁹

What the evidence shows: the mechanistic and biochemical case for omega-3 fatty acids in inflammation is well described.⁹ There is no high-quality trial evidence showing that omega-3 supplementation relieves sciatica, so treat this as general anti-inflammatory support rather than a sciatica treatment.

Typical intake: 1,200 to 2,400 mg combined EPA and DHA daily. That is the range used in the one study run in people with nonsurgical neck and back pain, where most participants took 1,200 mg daily and about a fifth took 2,400 mg. That study was an uncontrolled questionnaire follow-up rather than a placebo-controlled trial, so treat it as a starting point rather than a proven dose. Higher intakes appear in the joint-pain literature, but that work was done in rheumatoid arthritis and other inflammatory conditions rather than in the mechanical and degenerative problems this guide covers, so I have not carried its dose across.⁵³ **Form:** triglyceride or re-esterified triglyceride forms are commonly chosen for absorption. **Caution:** may increase bleeding risk at higher doses. Tell your surgeon if a procedure is planned.

6.2 Turmeric and Curcumin

What it does: curcumin, the active compound in turmeric, acts on NF-κB and on inflammatory enzymes including COX-2.

What the evidence shows: a systematic review and meta-analysis of randomized clinical trials found that turmeric extracts and curcumin reduced arthritis symptoms, while noting that the number and size of the trials were limited.¹⁰ A multicenter randomized trial in knee osteoarthritis found a *Curcuma domestica* extract comparable to ibuprofen for pain and function over four weeks.¹¹ Both trials were conducted in joint arthritis populations; neither one addresses sciatica or nerve pain.

Typical intake: 500 to 1,000 mg curcuminoids daily, often taken with piperine (black pepper extract) or in a liposomal or phytosome form for absorption. **Caution:** may interact with blood thinners. Discuss before use if you have gallbladder disease.

6.3 Vitamin D

What it does: supports bone health, muscle function, and immune regulation.

What the evidence shows: a study of patients with persistent, nonspecific musculoskeletal pain found a high prevalence of severe vitamin D deficiency in that group.¹² This shows how common deficiency was in that group; it does not demonstrate that correcting it relieves pain. A systematic review of vitamin D supplementation in diabetic peripheral neuropathy reported improvement in neuropathic pain scores, while flagging the low quality of the available studies.¹³ Checking your level is reasonable. Expecting vitamin D alone to resolve sciatica is not.

Typical intake: dosing should be guided by a blood level. Many clinicians target a level in the sufficient range and dose accordingly. **Form:** D3 (cholecalciferol). **Caution:** have levels checked periodically. Toxicity is uncommon but possible at very high intakes.

6.4 Magnesium

What it does: magnesium acts as an NMDA receptor antagonist, a mechanism relevant to central sensitization and pain amplification, and it participates in normal muscle and nerve function.

What the evidence shows: a review of magnesium and pain summarizes the NMDA-related mechanism and the mixed clinical trial results across pain conditions.¹⁴ No trial evidence addresses sciatica specifically.

Typical intake: 200 to 400 mg elemental magnesium daily. Glycinate and threonate are commonly chosen for tolerability; citrate is often used when constipation is also an issue. This is a general supplemental intake range rather than a dose drawn from a trial in this condition, which is why no study is cited beside it. **Caution:** can cause loose stools. Avoid in severe kidney disease unless your physician directs otherwise.

6.5 Methylcobalamin (Active Vitamin B12)

What it does: vitamin B12 is required for myelin maintenance, the protective sheath around nerves.

What the evidence shows: a systematic review of clinical controlled trials of vitamin B12 in diabetic neuropathy found that supplementation improved some symptom measures, but concluded the evidence base was limited and of uneven quality.¹⁵ The trials reviewed were in diabetic neuropathy; none examined sciatica.

Typical intake: 1,000 to 2,000 mcg daily, sublingual or oral. **Caution:** generally very well tolerated.

6.6 Alpha-Lipoic Acid (ALA)

What it does: an antioxidant studied extensively in diabetic nerve damage, where oxidative stress is part of the mechanism.

What the evidence shows: the SYDNEY trial found that intravenous alpha-lipoic acid improved sensory symptoms in diabetic polyneuropathy.¹⁶ The follow-up SYDNEY 2 trial found that oral alpha-lipoic acid at 600 mg daily improved symptomatic diabetic polyneuropathy, with higher doses adding side effects rather than benefit.¹⁷ A meta-analysis of alpha-lipoic acid in symptomatic diabetic polyneuropathy supported short-term symptom improvement.¹⁸ Every one of these trials was conducted in diabetic polyneuropathy; sciatica was not studied.

Typical intake: 600 mg daily is the dose used in the oral trial. **Caution:** may lower blood sugar. If you take diabetes medication, coordinate with your prescriber before adding it.

6.7 Palmitoylethanolamide (PEA)

What it does: a naturally occurring fatty acid amide that acts on mast cells and glial cells, pathways involved in neuroinflammation.

What the evidence shows: a pooled data meta-analysis reported that PEA reduced pain intensity across chronic pain conditions with a favorable tolerability profile.¹⁹ A later systematic review and meta-analysis of double-blind randomized controlled trials also found a reduction in pain, while noting heterogeneity across studies.²⁰ Little of this trial evidence was gathered in sciatica specifically.

Typical intake: 300 to 600 mg twice daily, micronized form commonly used. The upper end of that range is the dose used in a randomized trial of ultra-micronized PEA at 600 mg twice daily in older adults with chronic pain.⁵⁰ **Caution:** generally well tolerated with few reported side effects.

6.8 Boswellia Serrata (Frankincense)

What it does: inhibits 5-lipoxygenase, an enzyme in the inflammatory cascade, reducing leukotriene production.

What the evidence shows: a systematic review and meta-analysis found Boswellia and Boswellia extract effective for pain, stiffness, and function in osteoarthritis patients.²¹ The population studied was osteoarthritis, so this does not establish a benefit for nerve pain.

Typical intake: 300 to 500 mg standardized extract twice daily. **Caution:** discuss with your provider if you take anti-inflammatory medications.

6.9 Ginger

What it does: contains gingerols and shogaols, compounds with anti-inflammatory activity.

What the evidence shows: a randomized study found that daily ginger supplementation reduced muscle pain caused by eccentric exercise.²² A systematic review of ginger as an analgesic and ergogenic aid in sport found modest analgesic effects and limited, inconsistent performance effects.²³ None of this is nerve pain research, and no ginger trial has tested a sciatica claim.

Typical intake: 500 to 1,000 mg standardized extract daily, or fresh ginger in cooking. The randomized trial of a standardized ginger extract in knee osteoarthritis dosed it twice daily; the milligram amount is not stated in its abstract.⁵² **Caution:** may interact with blood thinners and can cause stomach upset at higher doses.

6.10 Collagen (Type II)

What it does: provides amino acid building blocks associated with cartilage and connective tissue.

What the evidence shows: a clinical trial of undenatured type II collagen in knee osteoarthritis reported reduced joint pain compared with a glucosamine and chondroitin combination.²⁴ The finding is specific to the knee joint; no evidence shows collagen supplementation repairs intervertebral discs or relieves sciatica.

Typical intake: 40 mg undenatured type II collagen daily, or 10 g hydrolyzed collagen as a general connective tissue support. The 40 mg figure is the exact daily dose used in the multicenter randomized trial of undenatured type II collagen in knee osteoarthritis.⁵¹ **Caution:** generally very safe. Source quality varies.

7. Targeted Exercises for Sciatica Relief

Critical rule: stop any exercise that increases radiating leg pain, numbness, or weakness. Mild discomfort in the back or buttock is usually acceptable, but pain traveling farther down the leg indicates nerve irritation and is a signal to back off.

Exercise as a category has broad support in chronic low back pain and chronic pain generally, with modest effect sizes and a good safety profile.^{28,30}

Before You Begin: The Pre-Treatment Warm-Up

Best for: Everyone, before any of the exercises below. This is soft-tissue release, not a substitute for the nerve-specific work that follows, and it is the same sequence used in office at Root Health, tools and doses unchanged. Use a percussion massage device, a foam roller, or your own hands, whichever you have, and respect a 6 out of 10 pressure ceiling throughout. The same critical rule from above applies here too: stop if anything increases radiating leg pain, numbness, or weakness.

Cat-Camel: 1. On hands and knees, arch your back up, then let it sag. 2. 1 set of 10 repetitions.

Glute Release, All Three Muscles: 1. With a percussion device, cover the glute maximus, medius, and minimus. 2. 1 to 2 minutes per side.

TFL and Hip Flexor Release: 1. With the same device, run it over the TFL and hip flexors. 2. 1 to 2 minutes each side.

Deep Hip Rotator and IT Band Release: 1. Lying on your back with a massage ball or foam roller under the glutes, move the leg up and down, or sit on the roller and search for tender spots. 2 to 3 minutes per side. 2. Side-lying, roll the outer thigh from just below the hip to above the knee for the IT band. 1 to 2 minutes per side.

This is the region worth lingering on if your pain sits mostly in the buttock: the piriformis and the other deep rotators run directly alongside the sciatic nerve, and section 7.3 below gives you the stretch that follows naturally from this release work.

Psoas Release: 1. With a massage ball or a lacrosse ball, place it over the front of the abdomen where the psoas runs. 2. Rock gently back and forth, then up and down. 3. 20 to 30 seconds each side.

This one is unusual and it is deliberate: the psoas connects the lumbar spine to the leg, and it rarely gets touched by anything else.

Why it is here: this is preparation, not treatment. It is what gets the tissue loose enough that the exercises below, which are the actual evidence-backed part of this chapter, are tolerable to do. If a step here increases leg symptoms rather than easing them, skip it and move on to the exercises.

7.1 McKenzie Extension Exercises

Best for: disc-related sciatica, especially when pain worsens with sitting and bending and improves with standing and walking.

Prone on Elbows 1. Lie face down with elbows under your shoulders. 2. Relax your lower back and hips completely. 3. Hold for 2 to 3 minutes, breathing deeply. 4. Repeat 3 to 5 times daily.

Press-Up (Prone Press-Up) 1. Lie face down, hands under shoulders as if doing a push-up. 2. Keep your hips and pelvis in contact with the floor. 3. Slowly press your upper body up, extending your arms. 4. Hold at the top for 2 seconds, then lower. 5. Start with 5 repetitions and progress to 10. 6. During an acute flare, many clinicians have patients repeat this at intervals through the day.

Why it works: the McKenzie approach, formally called Mechanical Diagnosis and Therapy, classifies patients by their response to repeated end-range movements and prescribes the direction that reduces symptoms. A literature review with meta-analysis found the McKenzie method more effective than comparison interventions for pain and disability in low back pain, with the authors noting that the effect sizes were small and the evidence quality limited.²⁵

7.2 Nerve Glides (Neural Flossing)

Best for: restricted nerve mobility anywhere along the pathway.

Sciatic Nerve Glide (Supine) 1. Lie on your back, legs straight. 2. Bring the affected leg up toward the ceiling, holding behind the thigh. 3. Slowly flex your ankle (pull toes toward you) while extending the knee. 4. Then point the toes while slightly bending the knee. 5. This creates a gliding motion along the nerve. 6. Perform 10 slow, controlled repetitions. 7. Never push into pain. A mild pulling or tingling is acceptable, sharp pain is not.

Why it works: neural mobilization aims to restore normal gliding of the nerve relative to surrounding tissue. A systematic review of neural mobilization in low back and radicular pain found evidence of benefit for pain and disability, with the authors cautioning that study quality varied.²⁶

7.3 Piriformis Stretching

Best for: piriformis-related symptoms, or when pain is primarily in the buttock.

Supine Piriformis Stretch (Figure 4) 1. Lie on your back, knees bent, feet flat. 2. Cross the affected leg over the opposite knee, creating a figure 4. 3. Gently pull the uncrossed thigh toward your chest. 4. You should feel a stretch in the buttock of the crossed leg. 5. Hold for 30 seconds, repeat 3 times.

Seated Piriformis Stretch 1. Sit in a chair and cross the affected leg over the opposite knee. 2. Keep your back straight and gently lean forward. 3. Hold for 30 seconds, repeat 3 times.

Why it works: the piriformis lies close to the sciatic nerve. When it is tight or in spasm it can contribute to buttock and posterior leg symptoms. Stretching aims to reduce that tension.

7.4 Core Stabilization

Best for: long-term support and prevention.

Dead Bug 1. Lie on your back, arms extended toward the ceiling, knees bent at 90 degrees. 2. Slowly lower one arm behind you while extending the opposite leg. 3. Keep your lower back pressed firmly into the floor. 4. Return to start and alternate sides. 5. Perform 8 to 10 reps per side, 2 to 3 sets.

Bird Dog 1. Start on hands and knees in a tabletop position. 2. Extend one arm forward and the opposite leg back. 3. Keep your core engaged and your back flat. 4. Hold for 5 seconds, then return to start. 5. Perform 8 to 10 reps per side, 2 to 3 sets.

Why it works: spinal stability comes from coordinated activity of many trunk muscles rather than from any single muscle, and stability demands change with the task.²⁷ In practice, exercise programs for chronic low back pain produce modest improvements in pain and function, and no single exercise type has been shown to be clearly superior to the others.²⁸

7.5 Hamstring and Hip Flexor Stretching

Best for: reducing tension across the lumbar spine and pelvis.

Standing Hamstring Stretch 1. Place the affected leg on a low step or chair. 2. Keep the knee slightly bent, not locked. 3. Hinge forward at the hips, keeping your back straight. 4. Hold for 30 seconds, repeat 3 times.

Half-Kneeling Hip Flexor Stretch 1. Kneel on one knee, other foot forward in a lunge position. 2. Tuck your pelvis under (posterior tilt). 3. Gently shift your weight forward. 4. Hold for 30 seconds, repeat 3 times.

Why it works: tight hamstrings and hip flexors change pelvic mechanics and can add stress to the lumbar spine. Note that in sciatica, a hamstring stretch can also tension the nerve itself, so ease into it and stop if leg symptoms increase.

7.6 Walking

Best for: almost everyone. It is foundational.

Protocol

Start with 10 to 15 minutes at a comfortable pace, then gradually build toward 30 to 45 minutes daily as tolerated. Maintain an upright posture, avoid hills initially if they aggravate symptoms, and stick to flat, even surfaces during acute phases.

Why it works: walking maintains mobility and load tolerance without high spinal stress. A systematic review and meta-analysis comparing walking with other exercise in chronic low back pain found no significant difference between them for pain and disability, which means walking is a reasonable, low-barrier option.²⁹

8. Physical Therapies That Work

8.1 Physical Therapy

Physical therapy is a cornerstone of conservative sciatica management. A skilled physical therapist will assess your specific movement patterns and identify aggravating factors, prescribe individualized exercises based on your presentation, use manual therapy techniques to improve joint mobility, educate you on body mechanics and posture, and progress you through a structured rehabilitation program.

Evidence: a systematic review of conservative management for lumbar disc herniation with associated radiculopathy found support for advice to stay active and for structured exercise programs, while noting that no single conservative protocol has been shown to be clearly superior.³¹ The American College of Physicians guideline recommends nonpharmacologic treatment, including exercise, as first-line care for low back pain.³²

8.2 Chiropractic Care and Spinal Manipulation

Spinal manipulation involves controlled, specific forces applied to spinal joints with the aim of improving mobility and reducing symptoms.

Evidence: a systematic review and meta-analysis found that spinal manipulative therapy was associated with modest improvements in pain and function in acute low back pain, with transient minor musculoskeletal soreness the most common adverse effect.³³ The American College of Physicians guideline includes spinal manipulation among the nonpharmacologic options for acute and subacute low back pain.³² A network meta-analysis of management strategies specifically for sciatica compared a wide range of options and found the evidence base uneven, with no clear ranking of conservative treatments against one another.³⁴

Be aware that most of the manipulation evidence above is for low back pain, and that sciatica with true nerve root involvement is a distinct and less-studied population.

What to expect: care typically combines manipulation or mobilization with exercise prescription and activity advice. Ask your provider for a defined plan of care with a scheduled re-evaluation point, so that progress is measured rather than assumed.

8.3 Acupuncture

Acupuncture involves inserting thin needles at specific points with the aim of modulating pain processing.

Evidence: an updated individual patient data meta-analysis found acupuncture superior to both sham acupuncture and no acupuncture for chronic musculoskeletal pain, with effects that persisted over time, although the difference from sham was modest.³⁵ A systematic review and meta-analysis focused on sciatica found that acupuncture improved pain outcomes compared with control interventions, with the authors noting that the included trials were of limited methodological quality.³⁶

What to expect: most protocols involve one to two sessions per week over several weeks. Benefits tend to accumulate across sessions.

8.4 Massage Therapy and Myofascial Release

Therapeutic massage addresses muscle tension and trigger points that may contribute to symptoms.

Evidence: a randomized controlled trial comparing structural massage, relaxation massage, and usual care for chronic low back pain found that both massage types produced better function and reduced symptoms than usual care, with benefits largely persisting at six months.³⁷ That trial studied chronic low back pain, not sciatica specifically.

Focus areas

Focus areas typically include the gluteal muscles, including the piriformis and gluteus medius and minimus, the lumbar paraspinals, the hamstrings, and the hip flexors.

8.5 Heat and Cold Therapy

Cold therapy (often used in the first 48 to 72 hours of an acute flare)

Cold therapy may help numb pain in the acute period. Apply an ice pack wrapped in cloth for 15 to 20 minutes, and allow the skin to return to normal temperature between applications.

Heat therapy (after the acute phase, or for muscle spasm)

Heat increases local blood flow and can reduce the sensation of stiffness. Use a heating pad or warm bath for 15 to 20 minutes; many people find it useful before stretching or exercise.

Evidence: a Cochrane review of superficial heat or cold for low back pain found moderate evidence that heat wrap therapy reduces pain and disability in acute and subacute low back pain, and found insufficient evidence to evaluate cold therapy.³⁸ In other words, heat has better support than ice, and neither has been shown to change the course of sciatica.

8.6 TENS (Transcutaneous Electrical Nerve Stimulation)

TENS units deliver low-voltage electrical currents through the skin with the aim of modulating pain signals.

Evidence: a Cochrane review of TENS for acute pain concluded that the available evidence was inconclusive because of the small number and low quality of eligible trials.³⁹ TENS is generally safe and can be tried at home, but you should treat it as a comfort measure with uncertain evidence rather than an established treatment.

Usage: place electrodes along the pain pathway. Sessions of 20 to 30 minutes are typical. Follow the device manufacturer's instructions and your provider's guidance.

8.7 Epidural Steroid Injections

For severe, persistent sciatica, an epidural injection delivers medication near the inflamed nerve root.

Evidence: a systematic review of randomized controlled trials comparing saline, local anesthetics, and steroids in epidural and facet joint injections found that epidural injections can provide meaningful relief in spinal pain, and notably that the injectate itself, including local anesthetic alone, often accounted for a substantial part of the effect.⁴⁰ A separate randomized trial in lumbar spinal stenosis, a different condition from disc-related sciatica, found that adding a glucocorticoid to lidocaine offered minimal or no additional benefit at six weeks.⁴¹

Considerations: injections are usually limited in number per year because of potential side effects. Discuss the risks, including effects on blood sugar and bone density, with your physician. Injections are best understood as a possible bridge that makes rehabilitation feasible, not as a stand-alone solution.

9. The Anti-Inflammatory Diet for Nerve Recovery

What you eat influences overall health and inflammatory load. The dietary pattern below is a sensible, well-tolerated way of eating. Be clear-eyed about what it is: general health support, not a proven sciatica treatment.

9.1 Foods to Emphasize

Emphasize omega-3 rich foods such as wild-caught salmon, sardines, mackerel, and anchovies, along with walnuts, flaxseeds, and chia seeds, aiming for 2 to 3 servings of fatty fish per week. Fill out your plate with colorful vegetables and fruits: leafy greens such as spinach, kale, and Swiss chard, berries such as blueberries, strawberries, and blackberries, cruciferous vegetables such as broccoli, cauliflower, and Brussels sprouts, and orange and yellow vegetables such as carrots, sweet potatoes, and peppers, with the goal of a large and varied intake spread across the day. Anti-inflammatory spices such as turmeric (paired with black pepper), ginger, garlic, cinnamon, and rosemary are worth incorporating regularly, along with healthy fats such as extra virgin olive oil, avocados, and nuts and seeds. Quality protein sources include fish, poultry, meat, eggs, and legumes as tolerated, and bone broth and other collagen sources provide a traditional source of glycine and proline.

9.2 Foods to Minimize

Minimize pro-inflammatory fats such as trans fats (partially hydrogenated oils), deep-fried foods, and very high intakes of refined seed oils relative to omega-3 sources. Cut back on refined carbohydrates and added sugars, including white bread, pastries, and sugary cereals, soda and sweetened beverages, and candy and desserts. Highly processed foods such as deli meats, hot dogs, bacon, fast food, and packaged snacks are also worth minimizing. Excessive alcohol interferes with sleep, which matters for recovery, and heavy intake is associated with poorer nutritional status. Some individuals notice a pattern with nightshades such as tomatoes, potatoes, eggplant, and peppers, though most people are not sensitive to them; this is worth testing individually rather than assuming.

9.3 Hydration

Intervertebral discs are highly hydrated tissues that rely on diffusion for nutrition, so staying well hydrated is sensible.

A practical rule of thumb: drink enough that you are rarely thirsty and your urine stays pale through the day. Individual needs vary with body size, climate, and activity.

Tips

Start your day with a glass or two of water, and drink before meals. Be aware that caffeine and alcohol increase fluid losses, and add electrolytes if you sweat heavily.

9.4 Sample Anti-Inflammatory Day

A sample anti-inflammatory day might begin with a smoothie of spinach, blueberries, flaxseed, ginger, almond butter, and unsweetened almond milk, or oatmeal with walnuts, chia seeds, cinnamon, and berries. Lunch could be a large salad with mixed greens, grilled salmon, avocado, and an olive oil and lemon dressing, with a side of vegetable soup. An afternoon snack of a handful of walnuts and an apple, or carrot and celery sticks with hummus, helps keep energy steady. Dinner might feature grilled chicken or tofu with turmeric and garlic, roasted broccoli and sweet potato, and a small side salad with olive oil. In the evening, if you like, a cup of ginger or turmeric tea and a small square of dark chocolate round out the day.

10. Lifestyle Modifications and Pain Management

10.1 Sleep Optimization

Sleep and pain influence each other in both directions. Poor sleep tends to worsen pain, and pain disrupts sleep.⁴³

Strategies

Maintain a consistent sleep schedule, with the same bedtime and wake time, and sleep on your side with a pillow between your knees, or on your back with a pillow under your knees. Many people find stomach sleeping aggravating for the low back. Keep your room cool, limit screens in the hour before bed, and consider that some people find magnesium glycinate in the evening helpful (see section 6.4).

Evidence: an experimental neuroimaging study found that a night of sleep deprivation lowered pain thresholds and altered pain-related activity in the brain, including heightened somatosensory cortex reactivity and blunted activity in regions involved in pain modulation.⁴² A review of the sleep and pain relationship concluded that sleep disturbance is a strong predictor of pain, arguably more so than the reverse.⁴³

10.2 Stress Management

Chronic stress affects pain perception and coping capacity.

Effective techniques

Effective techniques include diaphragmatic breathing for 5 minutes, twice daily, and mindfulness meditation: a randomized clinical trial found that mindfulness-based stress reduction improved back pain and functional limitations compared with usual care, with results similar to cognitive behavioral therapy.⁴⁴ Progressive muscle relaxation, systematically tensing and releasing muscle groups, is another option, as is gentle yoga; a randomized noninferiority trial found yoga noninferior to physical therapy for pain and function in chronic low back pain.⁴⁵

Note that both of those trials studied chronic low back pain rather than sciatica specifically.

10.3 Ergonomics and Body Mechanics

Sitting

Use a chair with lumbar support, or add a lumbar roll, keep your feet flat on the floor or on a footrest, and position your hips slightly higher than your knees. Take standing or walking breaks every 30 to 45 minutes, and consider a sit-stand desk.

Standing

Shift your weight periodically, use an anti-fatigue mat, and rest one foot on a small step to reduce lumbar extension.

Lifting

Bend at the knees and hips, not the waist, and keep the load close to your body. Avoid twisting while lifting, and ask for help with heavy items.

Driving

Adjust the seat so your knees are slightly bent, use a lumbar support, and take breaks every 1 to 2 hours on long trips.

10.4 Weight Management

Evidence: a meta-analysis found that overweight and obesity are associated with an increased prevalence and incidence of low back pain.⁴⁶ A separate meta-analysis found that obesity is a risk factor for sciatica specifically.⁴⁷ These are association findings. A systematic review of weight loss programs for low back pain found the evidence that intentional weight loss reduces back pain to be limited and inconsistent.⁴⁸

The honest summary: carrying less weight reduces spinal load and is good for your general health, but do not expect weight loss on its own to resolve sciatica.

Approach

Focus on the dietary pattern outlined above, aim for gradual and sustainable change, and combine it with walking and gentle exercise as tolerated.

10.5 Smoking Cessation

Smoking is associated with poorer spinal health and reduced blood supply to spinal tissues, and it is associated with worse outcomes across many musculoskeletal conditions. If you smoke, stopping is one of the highest-value changes you can make for your overall health. Ask your provider about cessation support.

11. The In-Clinic Pre-Treatment Protocol

Understand what this is for, because the distinction matters.

None of this rebuilds tissue. The loading and strengthening work in the previous chapters does that. What this protocol does is reduce the muscle tone, stiffness and tissue sensitivity that make the strength work harder to tolerate and less pleasant to do, particularly in the early weeks when things are at their most irritable.

It is preparation, not treatment. I use it in the clinic before the working part of a session, and I give it to patients to use at home before theirs. Patients who do it tend to tolerate the loading better and stick with it longer, and sticking with it is the actual mechanism by which recovery happens.

If you have to choose between doing this and doing your strength work, do your strength work.

Heat and vibration back wrap, set to vibrate and heat, while performing:

- Cat-camel stretch, 1 set of 10.
- Hamstring stretch with a stretch strap, 1 set of 10 each side.
- Ball bounce on an exercise ball, up and down then side to side, 1 set of 10 each, palms up.

Vibrating massage ball or foam roller. Lie on your back with the sphere on the glute muscles, moving the leg up and down.

Percussion massager, round or flat head. Over the tensor fasciae latae and hip flexors, 1 to 2 minutes each side.

Massage ball over the abdomen, working the psoas, rocking back and forth and up and down, 20 to 30 seconds each side.

Posture Curve under the lower back, knees bent to start and straightening if comfortable, resting 30 to 60 seconds.

Vibrating sphere or foam roller on sore calf areas, moving the foot up, down, left, right and in circles, 2 to 3 reps per area.

Safety, and I mean these.

- Avoid all of this over an acutely injured area in the first 48 to 72 hours.
- Reduce pressure over anything actively inflamed.
- Skip percussion therapy entirely if you have a pacemaker or another implanted electronic device.
- Avoid aggressive pressure over varicose veins or compromised skin.

- Start light. Pain must not exceed 6 out of 10. Back off immediately for sharp or shooting pain.
- Stop and seek advice if you get numbness, tingling, or weakness.

One absolute rule before you roll or percuss a calf. Never work on a calf that is swollen, warm, or tight in one leg only, particularly after travel, surgery, or a period of immobilisation. That is a blood clot concern. It needs assessing the same day and it must never be massaged.

12. Red Flags: When to Seek Immediate Care

While most sciatica improves with conservative care, certain symptoms require urgent medical evaluation.^{2,3}

Seek immediate care if you experience:

- **Cauda equina syndrome signs:** loss of bowel or bladder control, urinary retention or new difficulty urinating, or numbness in the saddle area (inner thighs, genitals, buttocks). **This is a surgical emergency. Go to an emergency department immediately, do not wait for an office appointment.**
- **Progressive weakness:** foot drop, meaning an inability to lift the front of your foot, or worsening leg weakness
- **Severe, unrelenting pain:** pain that does not improve with rest or with your usual measures
- **Fever with back pain:** may indicate infection
- **History of cancer:** new onset back pain, particularly with unexplained weight loss
- **Recent significant trauma:** a fall or car accident, especially if you are over 50 or have osteoporosis
- **Unexplained weight loss** accompanying back pain

Contact your healthcare provider promptly if:

- Symptoms worsen after 2 weeks of conservative care
- Pain is severe enough to prevent sleep
- You develop new areas of numbness or tingling
- Symptoms persist beyond 6 to 8 weeks without improvement

13. Building Your Personal Recovery Plan

Use this framework as a starting point for a conversation with your healthcare team, not as a substitute for one. Timelines are typical, not guaranteed, and your plan should be adjusted based on how you actually respond.

Weeks 1 and 2: Calm Things Down (Acute Phase)

The goal in weeks 1 and 2 is to calm things down. Discuss any supplement additions with your provider first. For movement, stick to short walks, prone on elbows, and gentle nerve glides if tolerated. Therapies at this stage include heat or ice for comfort, temporary modification of aggravating activities, and an initial evaluation by a physical therapist, chiropractor, or physician. On the diet side, begin the anti-inflammatory pattern and keep hydration steady, and for lifestyle, prioritize sleep, reduce stress load, and adjust your workstation.

Weeks 3 to 6: Rebuild (Subacute Phase)

Weeks 3 to 6 are about rebuilding. Review with your provider what supplements are helping and what are not. For movement, progress your directional exercises, add core stabilization, and increase walking duration. Therapies at this stage typically mean regular physical therapy or chiropractic care, with consideration of a trial of acupuncture or massage. Diet-wise, aim for the full anti-inflammatory pattern, and for lifestyle, work on building a repeatable stress management routine while keeping sleep hygiene consistent.

Weeks 7 to 12: Strengthen and Prevent (Rehabilitation Phase)

Weeks 7 to 12 focus on strengthening and prevention. Reassess the need for ongoing supplement use with your provider. Movement progresses to advanced core work, functional movement training, and a gradual return to normal activities. Therapies taper to as-needed care as you transition to a home exercise program. Diet becomes anti-inflammatory eating as a lifestyle, and for lifestyle, focus on maintaining your weight, keeping the exercise routine, and managing stress.

Beyond 12 Weeks: Maintenance

Beyond 12 weeks, the focus shifts to maintenance: continue core exercises several times per week, maintain the anti-inflammatory dietary pattern, and keep walking or another low-impact activity in the routine. Schedule periodic check-ins with your healthcare team, and address flare-ups early with your established toolkit.

If you are not improving on this timeline, that is information, not failure. Go back to your provider for re-evaluation.⁴⁹

Final Thoughts

Sciatica can be frightening and debilitating, but most cases improve significantly with conservative care. Your body has a real capacity to heal when given the right support: sensible nutrition, targeted movement, stress management, sleep, and patience.

Healing is rarely linear. You may have good days and bad days. The key is consistency with your plan and open communication with your healthcare team. If progress stalls, seek re-evaluation. Every person's sciatica is different, and your treatment should be tailored to your specific situation.

You are not alone in this, and with the right tools and support, meaningful improvement is a realistic goal.

References

Every reference below was retrieved and verified against the PubMed record. The PMID is provided so you can look up any of them yourself.

1. Konstantinou K, Dunn KM. Sciatica: review of epidemiological studies and prevalence estimates. *Spine (Phila Pa 1976)*. 2008;33(22):2464-2472. PMID 18923325.
2. Koes BW, van Tulder MW, Peul WC. Diagnosis and treatment of sciatica. *BMJ*. 2007;334(7607):1313-1317. PMID 17585160.
3. Chou R, Qaseem A, Snow V, et al. Diagnosis and treatment of low back pain: a joint clinical practice guideline from the American College of Physicians and the American Pain Society. *Ann Intern Med*. 2007;147(7):478-491. PMID 17909209. (Erratum: *Ann Intern Med*. 2008;148(3):247-248. PMID 18257154. Screened 2026-09-05; corrected, not retracted.)
4. Bush K, Cowan N, Katz DE, Gishen P. The natural history of sciatica associated with disc pathology. A prospective study with clinical and independent radiologic follow-up. *Spine (Phila Pa 1976)*. 1992;17(10):1205-1212. PMID 1440010.
5. Jacobs WC, van Tulder M, Arts M, et al. Surgery versus conservative management of sciatica due to a lumbar herniated disc: a systematic review. *Eur Spine J*. 2011;20(4):513-522. PMID 20949289.
6. Peul WC, van Houwelingen HC, van den Hout WB, et al. Surgery versus prolonged conservative treatment for sciatica. *N Engl J Med*. 2007;356(22):2245-2256. PMID 17538084.
7. Weinstein JN, Tosteson TD, Lurie JD, et al. Surgical vs nonoperative treatment for lumbar disk herniation: the Spine Patient Outcomes Research Trial (SPORT): a randomized trial. *JAMA*. 2006;296(20):2441-2450. PMID 17119140.
8. Atlas SJ, Keller RB, Wu YA, Deyo RA, Singer DE. Long-term outcomes of surgical and nonsurgical management of sciatica secondary to a lumbar disc herniation: 10 year results from the Maine Lumbar Spine Study. *Spine (Phila Pa 1976)*. 2005;30(8):927-935. PMID 15834338.
9. Calder PC. Omega-3 fatty acids and inflammatory processes: from molecules to man. *Biochem Soc Trans*. 2017;45(5):1105-1115. PMID 28900017.
10. Daily JW, Yang M, Park S. Efficacy of turmeric extracts and curcumin for alleviating the symptoms of joint arthritis: a systematic review and meta-analysis of randomized clinical trials. *J Med Food*. 2016;19(8):717-729. PMID 27533649.
11. Kuptniratsaikul V, Dajpratham P, Taechaarpornkul W, et al. Efficacy and safety of *Curcuma domestica* extracts compared with ibuprofen in patients with knee osteoarthritis: a multicenter study. *Clin Interv Aging*. 2014;9:451-458. PMID 24672232.
12. Plotnikoff GA, Quigley JM. Prevalence of severe hypovitaminosis D in patients with persistent, nonspecific musculoskeletal pain. *Mayo Clin Proc*. 2003;78(12):1463-1470. PMID 14661675.
13. Yamine K, Wehbe R, Assi C. A systematic review on the efficacy of vitamin D supplementation on diabetic peripheral neuropathy. *Clin Nutr*. 2020;39(10):2970-2974. PMID 32089370.
14. Shin HJ, Na HS, Do SH. Magnesium and pain. *Nutrients*. 2020;12(8). PMID 32718032.
15. Sun Y, Lai MS, Lu CJ. Effectiveness of vitamin B12 on diabetic neuropathy: systematic review of clinical controlled trials. *Acta Neurol Taiwan*. 2005;14(2):48-54. PMID 16008162.
16. Ametov AS, Barinov A, Dyck PJ, et al. The sensory symptoms of diabetic polyneuropathy are improved with alpha-lipoic acid: the SYDNEY trial. *Diabetes Care*. 2003;26(3):770-776. PMID 12610036. (Erratum: *Diabetes Care*. 2003;26(7):2227. Screened 2026-09-05; corrected, not retracted.)
17. Ziegler D, Ametov A, Barinov A, et al. Oral treatment with alpha-lipoic acid improves symptomatic diabetic polyneuropathy: the SYDNEY 2 trial. *Diabetes Care*. 2006;29(11):2365-2370. PMID 17065669.

18. Ziegler D, Nowak H, Kempler P, Vargha P, Low PA. Treatment of symptomatic diabetic polyneuropathy with the antioxidant alpha-lipoic acid: a meta-analysis. *Diabet Med*. 2004;21(2):114-121. PMID 14984445.
19. Paladini A, Fusco M, Cenacchi T, et al. Palmitoylethanolamide, a special food for medical purposes, in the treatment of chronic pain: a pooled data meta-analysis. *Pain Physician*. 2016;19(2):11-24. PMID 26815246.
20. Lang-Illievich K, Klivinyi C, Lasser C, et al. Palmitoylethanolamide in the treatment of chronic pain: a systematic review and meta-analysis of double-blind randomized controlled trials. *Nutrients*. 2023;15(6). PMID 36986081.
21. Yu G, Xiang W, Zhang T, et al. Effectiveness of Boswellia and Boswellia extract for osteoarthritis patients: a systematic review and meta-analysis. *BMC Complement Med Ther*. 2020;20(1):225. PMID 32680575.
22. Black CD, Herring MP, Hurley DJ, O'Connor PJ. Ginger (*Zingiber officinale*) reduces muscle pain caused by eccentric exercise. *J Pain*. 2010;11(9):894-903. PMID 20418184.
23. Wilson PB. Ginger (*Zingiber officinale*) as an analgesic and ergogenic aid in sport: a systemic review. *J Strength Cond Res*. 2015;29(10):2980-2995. PMID 26200194.
24. Crowley DC, Lau FC, Sharma P, et al. Safety and efficacy of undenatured type II collagen in the treatment of osteoarthritis of the knee: a clinical trial. *Int J Med Sci*. 2009;6(6):312-321. PMID 19847319.
25. Lam OT, Strenger DM, Chan-Fee M, Pham PT, Preuss RA, Robbins SM. Effectiveness of the McKenzie method of mechanical diagnosis and therapy for treating low back pain: literature review with meta-analysis. *J Orthop Sports Phys Ther*. 2018;48(6):476-490. PMID 29602304.
26. Peacock M, Douglas S, Nair P. Neural mobilization in low back and radicular pain: a systematic review. *J Man Manip Ther*. 2023;31(1):4-12. PMID 35583521.
27. McGill SM. Low back stability: from formal description to issues for performance and rehabilitation. *Exerc Sport Sci Rev*. 2001;29(1):26-31. PMID 11210443.
28. Searle A, Spink M, Ho A, Chuter V. Exercise interventions for the treatment of chronic low back pain: a systematic review and meta-analysis of randomised controlled trials. *Clin Rehabil*. 2015;29(12):1155-1167. PMID 25681408.
29. Vanti C, Andreatta S, Borghi S, Guccione AA, Pillastrini P, Bertozzi L. The effectiveness of walking versus exercise on pain and function in chronic low back pain: a systematic review and meta-analysis of randomized trials. *Disabil Rehabil*. 2019;41(6):622-632. PMID 29207885.
30. Geneen LJ, Moore RA, Clarke C, Martin D, Colvin LA, Smith BH. Physical activity and exercise for chronic pain in adults: an overview of Cochrane Reviews. *Cochrane Database Syst Rev*. 2017;4(4):CD011279. PMID 28436583.
31. Hahne AJ, Ford JJ, McMeeken JM. Conservative management of lumbar disc herniation with associated radiculopathy: a systematic review. *Spine (Phila Pa 1976)*. 2010;35(11):E488-E504. PMID 20421859.
32. Qaseem A, Wilt TJ, McLean RM, Forciea MA. Noninvasive treatments for acute, subacute, and chronic low back pain: a clinical practice guideline from the American College of Physicians. *Ann Intern Med*. 2017;166(7):514-530. PMID 28192789.
33. Paige NM, Miake-Lye IM, Booth MS, et al. Association of spinal manipulative therapy with clinical benefit and harm for acute low back pain: systematic review and meta-analysis. *JAMA*. 2017;317(14):1451-1460. PMID 28399251. (Errata: *JAMA*. 2017;317(21):2239, an error in 2 figures, PMID 28586870; and *JAMA*. 2017;318(20):2048, incorrect data in text, PMID 29183047. Screened 2026-09-05; corrected, not retracted. The correction text is subscription-gated and was not read. This guide reproduces no numeric value from this paper, so no figure here depends on it.)
34. Lewis RA, Williams NH, Sutton AJ, et al. Comparative clinical effectiveness of management strategies for sciatica: systematic review and network meta-analyses. *Spine J*. 2015;15(6):1461-1477. PMID 24412033.
35. Vickers AJ, Vertosick EA, Lewith G, et al. Acupuncture for chronic pain: update of an individual patient data meta-analysis. *J Pain*. 2018;19(5):455-474. PMID 29198932.

36. Han KH, Cho KH, Han C, et al. The effectiveness and safety of acupuncture treatment on sciatica: a systematic review and meta-analysis. *Complement Ther Med*. 2022;71:102872. PMID 35985442.
37. Cherkin DC, Sherman KJ, Kahn J, et al. A comparison of the effects of 2 types of massage and usual care on chronic low back pain: a randomized, controlled trial. *Ann Intern Med*. 2011;155(1):1-9. PMID 21727288.
38. French SD, Cameron M, Walker BF, Reggars JW, Esterman AJ. Superficial heat or cold for low back pain. *Cochrane Database Syst Rev*. 2006;(1):CD004750. PMID 16437495.
39. Johnson MI, Paley CA, Howe TE, Sluka KA. Transcutaneous electrical nerve stimulation for acute pain. *Cochrane Database Syst Rev*. 2015;(6):CD006142. PMID 26075732.
40. Manchikanti L, Nampiaparampil DE, Manchikanti KN, et al. Comparison of the efficacy of saline, local anesthetics, and steroids in epidural and facet joint injections for the management of spinal pain: a systematic review of randomized controlled trials. *Surg Neurol Int*. 2015;6(Suppl 4):S194-S235. PMID 26005584.
41. Friedly JL, Comstock BA, Turner JA, et al. A randomized trial of epidural glucocorticoid injections for spinal stenosis. *N Engl J Med*. 2014;371(1):11-21. PMID 24988555. (Erratum: *N Engl J Med*. 2014;371(4):390. Screened 2026-09-05; corrected, not retracted.)
42. Krause AJ, Prather AA, Wager TD, Lindquist MA, Walker MP. The pain of sleep loss: a brain characterization in humans. *J Neurosci*. 2019;39(12):2291-2300. PMID 30692228.
43. Finan PH, Goodin BR, Smith MT. The association of sleep and pain: an update and a path forward. *J Pain*. 2013;14(12):1539-1552. PMID 24290442.
44. Cherkin DC, Sherman KJ, Balderson BH, et al. Effect of mindfulness-based stress reduction vs cognitive behavioral therapy or usual care on back pain and functional limitations in adults with chronic low back pain: a randomized clinical trial. *JAMA*. 2016;315(12):1240-1249. PMID 27002445.
45. Saper RB, Lemaster C, Delitto A, et al. Yoga, physical therapy, or education for chronic low back pain: a randomized noninferiority trial. *Ann Intern Med*. 2017;167(2):85-94. PMID 28631003.
46. Shiri R, Karppinen J, Leino-Arjas P, Solovieva S, Viikari-Juntura E. The association between obesity and low back pain: a meta-analysis. *Am J Epidemiol*. 2010;171(2):135-154. PMID 20007994.
47. Shiri R, Lallukka T, Karppinen J, Viikari-Juntura E. Obesity as a risk factor for sciatica: a meta-analysis. *Am J Epidemiol*. 2014;179(8):929-937. PMID 24569641.
48. Chen LH, Weber K, Mehrabkhani S, Baskaran S, Abbass T, Macedo LG. The effectiveness of weight loss programs for low back pain: a systematic review. *BMC Musculoskelet Disord*. 2022;23(1):488. PMID 35606809.
49. Luijsterburg PA, Verhagen AP, Ostelo RW, et al. Effectiveness of conservative treatments for the lumbosacral radicular syndrome: a systematic review. *Eur Spine J*. 2007;16(7):881-899. PMID 17415595.
50. Germini F, Coerezza A, Andreinetti L, et al. N-of-1 Randomized Trials of Ultra-Micronized Palmitoylethanolamide in Older Patients with Chronic Pain. *Drugs Aging*. 2017;34(12):941-952. PMID: 29210011
51. Lugo JP, Saiyed ZM, Lane NE. Efficacy and tolerability of an undenatured type II collagen supplement in modulating knee osteoarthritis symptoms: a multicenter randomized, double-blind, placebo-controlled study. *Nutr J*. 2016;15:14. PMID: 26822714
52. Altman RD, Marcussen KC. Effects of a ginger extract on knee pain in patients with osteoarthritis. *Arthritis Rheum*. 2001;44(11):2531-2538. PMID: 11710709
53. Maroon JC, Bost JW. Omega-3 fatty acids (fish oil) as an anti-inflammatory: an alternative to nonsteroidal anti-inflammatory drugs for discogenic pain. *Surg Neurol*. 2006;65(4):326-331. PMID: 16531187

This guide was compiled from peer-reviewed medical literature and clinical best practices. It is intended to give you knowledge and options to discuss with your healthcare provider. It does not promise a cure and it is not a substitute for individualized medical care.

Working with Root Health

If your leg symptoms are not improving, or you are being asked to decide about surgery and want the conservative case laid out honestly first, I offer a free Root Discovery Session at Root Health to talk through where you are and what a reasonable next step looks like. You can schedule one at rootcauseunlocked.com/discovery, or reach the office at (919) 662-0520.

Consultations and ongoing care are both available wherever you are located. Being outside North Carolina is not a barrier to working with me. Tell me where you are when you reach out and I will lay out exactly how we would work together from there.

If you would rather start on your own, the free Root Cause Unlocked tracker has a mode built for musculoskeletal recovery, at rootcauseunlocked.com/tracker. Eight weeks of it turns “I think it might be helping” into something you and I can both read.

© 2026 · For personal educational use only

Disclaimer: *This ebook is for educational purposes only and does not constitute medical advice. Always consult with a qualified healthcare provider before starting any new treatment, supplement, exercise program, or dietary change. Do not start, stop, or change any prescribed medication without speaking to the prescriber. This guide does not diagnose your condition. If you experience the warning signs described in Chapter 11, seek medical evaluation promptly.*

A note on the evidence in this guide. *Every numbered citation points to a specific published study that was checked against the source paper before it was used here. Where the research on a topic is strong, this guide says so. Where the research is thinner than clinical experience suggests, or where a claim rests mainly on years of hands on practice rather than a body of trials, this guide says that too, plainly, rather than dressing clinical judgment up as settled science.*

1. Understanding Hip Pain

If your hip has been bothering you, you have probably already noticed that it does not behave the way you expected. Maybe it aches deep in the groin when you stand up from a chair. Maybe it is a sharp, nagging pain on the outside of your hip that flares up the moment you lie on that side at night. Maybe it started as low back pain and slowly migrated into your buttock and hip. Hip pain is not one condition. It is a region, and a great many different structures in that region can become the source of your discomfort.

This guide takes an integrative, root cause approach to hip pain, the same approach I use in my practice at Root Health in Garner, North Carolina. That approach starts from a simple premise: the place where you feel pain is not always the place where the problem began. A tight, weak, or poorly coordinated muscle group around the hip and pelvis can quietly overload a tendon, a joint, or a nerve until it finally speaks up as pain. Understanding those relationships, rather than chasing the painful spot alone, is what this guide is built around.

You will also notice this guide spends real time on gluteus medius and gluteus minimus tendinopathy, often still called greater trochanteric pain syndrome or, incorrectly, hip bursitis. In my clinical experience, these two muscles are involved in the majority of hip presentations I see, whether the pain is on the outside of the hip itself or in a related pattern elsewhere in the leg or low back. That observation lines up closely with what the research literature now says about lateral hip pain, which is covered in detail in Chapter 4.

Hip pain is also, encouragingly, very often manageable with active, structured care rather than a quick procedure. Most of the conditions in this guide respond to a combination of load management, targeted strengthening, and attention to the muscles and movement patterns around the hip and pelvis, not to rest and waiting. That is the direction this guide points you in throughout.

One more thing before you continue. This guide organizes your thinking about hip pain. It does not diagnose you. A qualified clinician, examining your specific hip, history, and movement pattern, is the only way to know for certain what is driving your pain. Use this guide to understand your options and to ask better questions at your next appointment, not as a substitute for that appointment.

2. Anatomy of the Hip: The Structures at Play

The hip joint itself is a deep, stable ball and socket joint, the femoral head seated inside the acetabulum of the pelvis. That depth and stability is part of why the hip rarely dislocates the way a shoulder does. It is also part of why hip pain is so often generated not by the joint surfaces themselves but by the dense network of muscles, tendons, and ligaments that surround and move that joint.

The Bony and Joint Structures

The femur, the pelvis, and the sacrum form the bony architecture of the hip and pelvic girdle. Where the sacrum meets the pelvis on either side, you have the sacroiliac, or SI, joints, small joints that transmit load between your spine and your legs. A cartilage rim called the labrum lines the acetabulum and deepens the socket, and a layer of articular cartilage covers the joint surfaces themselves.

The Muscle Groups I Focus On

Rather than list every muscle that crosses the hip, it is more useful to understand the functional groups I evaluate in nearly every hip patient, because these are the groups most often out of balance.

The hip flexors: psoas major and iliacus. Together often called the iliopsoas, these muscles run from the lumbar spine and the inside of the pelvis down to the top of the femur, and their job is to flex the hip and help stabilize the lumbar spine and pelvis. In my clinical experience, a tight or poorly functioning psoas and iliacus is a frequent contributor to hip dysfunction, particularly in people who sit for long stretches of the day. It is worth being direct about the evidence here: the idea of a “tight psoas” is one of the most widely repeated concepts in manual and movement based practice, and it has real clinical currency, but it is not backed by a deep body of high quality trial evidence specifically linking psoas tightness to hip pain. This guide treats it as clinical reasoning, not an established fact, and says so again in Chapter 6.

The external rotators of the hip. A small group of deep muscles, including the piriformis, that rotate the thigh outward and help stabilize the femoral head in the socket during walking and standing. When these muscles are chronically short or overactive, my clinical experience is that they can alter hip mechanics and contribute to lateral and posterior hip discomfort. As with the psoas, this is a clinical pattern more than a heavily researched one, and it is presented that way in Chapter 6.

The tensor fasciae latae, or TFL. A muscle at the front of the outer hip that blends into the iliotibial band running down the outside of the thigh. It is frequently implicated in lateral hip and knee complaints in clinical practice. A 2020 systematic review and meta-analysis in *Sports Medicine* examined whether the TFL actually differs in structure or activation between people with lower limb musculoskeletal conditions, including lateral hip pain, and people without. The reviewers found only weak, inconsistent evidence for the common clinical assumptions about the TFL's role, and concluded that those assumptions are not well investigated and are poorly supported by current research.¹ That does not mean the TFL is irrelevant. It means the confident claims sometimes made about it outrun what has actually been studied, which is worth knowing before anyone recommends an aggressive TFL-focused protocol to you.

The gluteus medius and gluteus minimus. These two muscles sit on the outer surface of the pelvis, beneath the larger gluteus maximus, and their primary job is to stabilize your pelvis every time you take a step on one leg. My clinical experience is that these two muscles are involved in almost every hip case I evaluate, and as you will see in Chapter 4, that observation is strongly supported by the research literature on lateral hip pain specifically.

How These Groups Work Together

None of these muscle groups act alone. The hip flexors, external rotators, TFL, and glute medius and minimus all cross the same joint and share the job of controlling your pelvis during standing, walking, and single leg tasks like climbing stairs. When one group is tight, weak, or slow to activate, the others are forced to compensate, and that compensation is frequently where pain begins. This interconnected view of the hip carries through the rest of this guide, and it is the foundation of the regional interdependence model discussed fully in Chapter 6.

3. Common Causes of Hip Pain

Hip pain is generally grouped by where you feel it: the front of the hip and groin, the outside of the hip, or the back of the hip and buttock. This is a useful starting framework, because different structures tend to produce pain in different locations, although there is real overlap.

Anterior, or front of hip and groin, pain is often related to the hip joint itself. In younger, more active adults this can include labral tears or femoroacetabular impingement, a mechanical conflict between the femur and the socket. In older adults, hip osteoarthritis is a common driver of anterior groin pain. Front of hip pain can also be referred from structures inside the abdomen or pelvis, which is one more reason self-diagnosis is not a good idea.

Lateral, or outside of hip, pain is what most people mean when they describe a nagging ache over the bony point of the hip that worsens with lying on that side, walking, or climbing stairs. This is the territory of greater trochanteric pain syndrome, covered in full in Chapter 4, and it is by far the most common presentation my approach is built around.

Posterior, or back of hip and buttock, pain can come from the hip joint itself, from deep gluteal structures including the piriformis and the sciatic nerve as it passes beneath it, or it can be referred pain traveling down from the lumbar spine. A 2021 review in *American Family Physician* laid out this same anterior, lateral, and posterior framework for evaluating adult hip pain, and specifically noted that posterior hip pain includes referred pain from lumbar spinal pathology as well as deep gluteal syndrome with sciatic nerve entrapment.² That overlap between the low back and the hip is a recurring theme in this guide and is central to Chapter 6.

A Common Misdiagnosis Worth Naming Early

For decades, persistent pain over the outer hip was routinely labeled “trochanteric bursitis,” implying inflammation of the small fluid filled sac, the bursa, that cushions the tendons near the greater trochanter. Current evidence tells a different story, and it is important enough that the next chapter is devoted to it entirely.

A Brief Note on the Sacroiliac Joint

Some readers of this guide will have pain that is genuinely coming from the sacroiliac joint rather than the hip itself. SI joint pain typically sits low and to the side of the spine, over the dimples at the top of the buttock, and it is a real and commonly under-recognized contributor to what people describe as low back and hip pain, affecting an estimated 15 to 25 percent of people with axial low back pain according to a widely cited review in *Anesthesia & Analgesia*.³ Because SI joint dysfunction is experienced by most people as chronic low back and buttock pain rather than as hip pain specifically, the full discussion of how it develops and how it is managed, including the role of sacroiliac support belts, lives in my companion guide, *The Complete Patient's Guide to Low Back Pain*. If your pain sits more over your low back and buttock than over the side of your hip, or if it has never resolved despite treatment aimed at your back, that guide and a proper evaluation for SI joint involvement are worth pursuing alongside this one.

4. The Real Story Behind Lateral Hip Pain: Gluteal Tendinopathy

If you have lateral hip pain, the single most important thing this guide can tell you is this: it is very likely not bursitis.

For years, the working assumption in medicine was that pain over the outer hip, especially pain that worsened with lying on that side, was caused by inflammation of the trochanteric bursa, a small cushioning sac near the top of the femur. Research over the past two decades has substantially overturned that assumption. A landmark study published in *Arthritis and Rheumatism* examined 24 patients with the classic clinical picture of greater trochanteric pain syndrome using MRI. Gluteus medius tendon pathology, either tearing or tendinitis, was found in the large majority of patients, while true bursal distension was uncommon and, notably, did not occur in the absence of gluteus medius pathology.⁴ In other words, when the bursa was inflamed at all, the tendon was already involved too. The bursa was a bystander, not the primary problem.

A comprehensive 2015 review in *Sports Medicine* confirmed and extended this picture, stating plainly that tendinopathy of the gluteus medius and gluteus minimus tendons is now recognized as the primary local source of lateral hip pain.⁵ This condition is most common in midlife, occurs in both athletes and people who do not regularly exercise, and affects women more than men. It interferes with sleep, particularly side lying, and with common weight bearing tasks, which is exactly the pattern many of my patients describe: pain that wakes them at night and pain that builds with walking or stairs.

This is precisely the pattern my clinical model points to directly. I have long observed that the gluteus medius and gluteus minimus are almost always involved in hip presentations, and the research on greater trochanteric pain syndrome now bears that observation out with real specificity. What was once dismissed as a vague, catch all “bursitis” diagnosis has a much more precise and much more treatable explanation: overloaded, irritated hip abductor tendons.

Why This Distinction Actually Matters to You

Calling this condition bursitis matters because it changes how people try to treat it. Bursitis suggests inflammation that might respond to rest and anti-inflammatory measures alone. Tendinopathy is a different problem, one driven by how the tendon is being loaded over time, and it generally responds best to a structured, progressive loading program rather than rest. Chapter 7 walks through what that looks like in practice.

The Role of Compression

One of the more clinically useful findings from the tendinopathy research is the role of compression. The 2015 *Sports Medicine* review found that the combination of excessive compressive load and high tensile load on the gluteal tendons is thought to be the most damaging combination, and that joint position, particularly excessive hip adduction, where the thigh crosses toward or past the midline of the body, is a key factor in this condition.⁵ Every time your femur moves into an adducted position, whether from standing with your weight shifted onto one hip, sitting with your legs crossed, or sleeping curled on your side without support between your knees, the iliotibial band and the gluteal tendons are compressed against the bony point of the hip.

This has a direct and somewhat counterintuitive practical implication, addressed fully in Chapter 7: some of the stretches people instinctively reach for to relieve lateral hip pain, particularly stretches that pull the leg across the body into adduction, can actually provoke the very compression that is aggravating the tendon in the first place. Stretching alone, and the wrong kind of stretching in particular, is not a reliable fix for this condition.

What About Injections?

Corticosteroid injections are commonly offered for greater trochanteric pain syndrome, and they can and do provide real short term relief for many people. The strongest evidence available on how injections compare with a structured exercise program comes from a large, well conducted trial known as the LEAP trial, published in *BMJ* in 2018. Researchers randomly assigned 204 adults with confirmed gluteal tendinopathy to one of three groups: a physiotherapy led program of education plus exercise over eight weeks, a single corticosteroid injection, or a wait and see approach with no active treatment.⁶

The results were notable. At eight weeks, both the education plus exercise group and the corticosteroid injection group reported significantly better global improvement and less pain than the wait and see group, which is exactly what you would expect: doing something beats doing nothing. But the education plus exercise group did not just match the injection group, it outperformed it, both on the global rating of improvement and on reported pain scores at eight weeks. At the one year follow up, the education plus exercise group continued to show better global outcomes than the corticosteroid injection group, while the two groups' pain scores had converged somewhat by that point, with both still ahead of the wait and see group.⁶

The honest, plainly stated takeaway from this trial is not that injections do nothing. They clearly helped compared with doing nothing at all. The takeaway is that a structured program of education and exercise did at least as well as an injection in the short term and did better on the measure of overall improvement a year later. An injection is not a solution on its own, and it should not be presented to you as one. If you are offered a corticosteroid injection for lateral hip pain, it is entirely reasonable to ask your provider whether a loading program is being paired with it, because the best evidence available says that combination, or exercise alone, is where the durable improvement comes from.

5. How Hip Pain Is Diagnosed

A thorough evaluation for hip pain typically combines your history, a physical examination, and, when needed, imaging.

History and Physical Examination

Your provider will want to know where exactly your pain is located, what makes it better or worse, whether it changes with specific positions like lying on your side or crossing your legs, and whether you have any of the warning signs covered in Chapter 11. Physical examination typically includes assessing your hip's range of motion, your strength in hip abduction and other directions, your gait, and specific provocative tests. One classic finding worth knowing about is the Trendelenburg sign, in which the pelvis drops on the side opposite the affected hip when you stand on the affected leg alone, a sign of gluteus medius weakness or dysfunction. In the original MRI study of greater trochanteric pain syndrome discussed in Chapter 4, this sign was the most accurate of the physical tests examined for predicting an actual gluteus medius tendon tear, correctly identifying about 73 percent of tears and correctly ruling out the condition in about 77 percent of people who did not have a tear.⁴

Imaging: When It Helps

Not everyone with hip pain needs imaging right away. A 2021 review in *American Family Physician* on evaluating hip pain in adults notes that plain radiography of the hip and pelvis is usually the appropriate initial imaging test, with ultrasound or MRI reserved for cases where the diagnosis remains unclear or where findings would change management, such as suspected labral tears, femoroacetabular impingement, or gluteus medius tendon tears, which often have good outcomes with surgery when appropriate and where earlier imaging and referral can matter.² For most cases of lateral hip pain consistent with gluteal tendinopathy, a clinical diagnosis based on history and examination is often sufficient to begin a structured treatment program, without needing advanced imaging first.

Differential Diagnosis: Why Location Alone Is Not Enough

As Chapter 3 introduced, hip pain located at the front, side, or back of the hip tends to point toward different structures, but there is meaningful overlap, and referred pain from the lumbar spine can mimic pain originating in the hip itself.² This is exactly why this guide does not attempt to diagnose you, and why a proper clinical evaluation, one that also assesses your lumbar spine and pelvis rather than only the hip in isolation, is genuinely necessary before starting a treatment plan.

6. Regional Interdependence: Why the Hip Rarely Works Alone

This chapter is the connective tissue of my clinical model, and it deserves to be read as its own idea rather than a footnote to the others.

The formal term for this way of thinking is regional interdependence, and it has a real foothold in the physical therapy and rehabilitation literature. A widely cited paper on the model, published in the *Journal of Manual and Manipulative Therapy*, describes the underlying premise directly: seemingly unrelated impairments in remote anatomical regions of the body may contribute to and be associated with a patient's primary report of symptoms, and interventions directed at one region of the body will often have effects at remote and seemingly unrelated areas.⁷ In plainer language, the place that hurts and the place that is driving the problem are frequently not the same place.

This is the lens I apply to hip dysfunction. A tight or weak psoas and iliacus, tight external rotators, and an overactive TFL are not simply local nuisances. They change how your femur sits in the socket and how much work your gluteus medius and minimus have to do to stabilize your pelvis with every step. When those upstream muscles are not doing their share, the tendons at the outer hip end up absorbing load they were not built to handle alone, which loops directly back to the gluteal tendinopathy story in Chapter 4.

Some supporting research for this interconnected view comes from outside the hip specifically. A 2025 study in *Scientific Reports* examined hamstring strength and hip mobility, including hip extension and hip internal rotation range of motion, in young adults with and without clinically defined lumbar instability, and found that both hamstring strength and hip mobility were strongly associated with pain and disability, independent of each other.⁸ This is not a study of hip pain patients specifically, and its authors were careful to describe their findings as hypothesis generating rather than diagnostic, but it is a genuine, measured example of hip function and low back symptoms moving together, which is exactly the kind of relationship the regional interdependence model predicts.

Functional Leg Length Inequality

The fifth pillar of my clinical model is functional leg length inequality, a difference in effective leg length that arises from muscular, postural, or pelvic factors rather than an actual difference in bone length. In my clinical experience, this kind of asymmetry can contribute not only to hip pain but to low back, knee, ankle, and foot pain as well, because every step is being taken on an uneven foundation.

This is genuinely an area where, as with the psoas, the trial evidence is thinner than the clinical pattern would suggest, and it is worth saying so directly rather than overselling it. A systematic review published in *Archives of Physical Medicine and Rehabilitation* examined shoe lift interventions for leg length discrepancy in adults with painful musculoskeletal conditions and found that leg length discrepancy was indeed associated with low back pain, scoliosis, and osteoarthritis of the hip and knee in the literature, but that the overall quality of the included studies was very low to poor, and that only one included study was a randomized controlled trial.⁹ Within that limited evidence, the results were still worth noting: in the studies that reported it, the large majority of people treated with a shoe lift experienced partial or complete pain relief, and the one randomized trial found meaningful pain reduction in the treated group.⁹ A more recent, narrower systematic review in the *Journal of the American Podiatric Medical Association* looked specifically at foot orthoses and low back pain in people with leg length inequality and found that every included study reported a significant reduction in low back pain with orthosis use, although the evidence on whether orthoses actually changed walking mechanics was inconsistent.¹⁰ A separate review connecting

foot and ankle mechanics to chronic low back pain likewise noted that excessive pronation has been linked to leg length discrepancies that contribute to pelvic tilt and low back pain, describing the connection as part of a broader kinetic chain running from the foot to the low back.¹¹

Taken together, this is a real, plausible, and clinically observed pattern that does not yet have the kind of large, rigorous trial evidence behind it that the gluteal tendinopathy research does. My use of functional leg length assessment and correction, where appropriate, reflects clinical judgment built on this pattern and on years of hands on evaluation, not on a settled body of high quality research. That distinction matters, and this guide is not going to claim more certainty than the evidence supports.

The Practical Upshot

Regional interdependence is the reason a thorough hip evaluation looks beyond the hip itself, checking your lumbar spine, your pelvis, your leg length, and the muscle groups covered in Chapter 2. It is also the reason that a treatment plan built only around the painful spot, without addressing what is feeding into it, so often stalls out. The exercise and movement approach in the next chapter is built with this whole picture in mind.

7. Movement and Exercise for Hip Recovery

The most important message in this chapter: load management and progressive strengthening have the strongest evidence behind them of anything discussed in this guide, and they should be the center of your recovery plan, not an afterthought.

Understanding Load, Not Just Pain

A useful way to think about tendinopathy, including gluteal tendinopathy, comes from a foundational 2009 paper in the *British Journal of Sports Medicine*, which proposed that tendon pathology exists on a continuum rather than as a single fixed injury.¹² Depending on how a tendon has been loaded over time, it can move along that continuum toward reactive irritation or toward more degenerative change, and different points on that continuum call for different treatment intensity. The practical takeaway for you is straightforward: your tendon's pain is telling you something about how it has been loaded, not simply that it is damaged, and the right amount and right kind of loading, not none at all, is usually the way forward.

The Evidence Centerpiece: Education Plus Exercise

As discussed fully in Chapter 4, the LEAP trial found that an eight week, physiotherapy led program combining education about load management with a progressive exercise program outperformed a single corticosteroid injection, both in the short term and at one year follow up, for people with confirmed gluteal tendinopathy.⁶ That program is the single best supported intervention for this condition among everything covered in this guide, and it should be the anchor of any recovery plan for lateral hip pain.

What to Avoid: Compressive Stretches

Because excessive hip adduction compresses the gluteal tendons against the greater trochanter, and because that compression is implicated as a driver of gluteal tendinopathy, some commonly recommended hip stretches can do more harm than good if you have this condition.⁵ Be cautious with any stretch that pulls your leg across the midline of your body, such as pulling your knee toward your opposite shoulder while lying on your back, or a standing IT band stretch where you cross one leg behind the other and lean away. Sitting with your legs crossed for long periods and standing with your weight sunk into one hip, a posture sometimes called hip hitching, create the same compressive load and are worth minimizing during a flare. This does not mean all stretching is off limits, but it does mean stretching should be selected thoughtfully rather than reached for automatically, ideally with guidance from a clinician familiar with this condition.

Gentle Isometrics to Begin

For an irritable tendon, isometric holds, contracting the muscle without moving the joint, are often a reasonable and well tolerated starting point before progressing to more dynamic strengthening.

Isometric Hip Abduction Against a Wall: 1. Stand with the outside of your affected hip a few inches from a wall 2. Press your leg outward into the wall without letting your body move 3. Hold for 20 to 30 seconds, breathing normally 4. Repeat 3 to 5 times

Progressive Strengthening

Once the tendon tolerates isometric loading comfortably, progressive strengthening of the hip abductors, particularly the gluteus medius and gluteus minimus, becomes the priority.

Sidelying Hip Abduction: 1. Lie on your unaffected side with both knees slightly bent 2. Keeping your hips stacked and your top hip from rolling backward, lift your top leg toward the ceiling 3. Avoid lifting the leg too high, which shifts the work away from the target muscles 4. Perform 2 to 3 sets of 10 to 15 reps, progressing to a resistance band around the thighs as tolerated

Standing Hip Hike: 1. Stand on your affected leg on a small step, letting the other foot hang off the edge 2. Slowly lower the hanging hip toward the floor, then use your standing hip's outer muscles to lift it back level 3. Perform 2 to 3 sets of 10 reps

Bridge with Band: 1. Lie on your back, knees bent, a resistance band around your thighs just above the knees 2. Press your knees gently outward against the band while lifting your hips into a bridge 3. Hold briefly at the top, lower slowly 4. Perform 2 to 3 sets of 10 to 15 reps

Addressing the Whole Chain

Consistent with the regional interdependence model in Chapter 6, a complete program also addresses hip flexor mobility and general hip and pelvic control, not the outer hip alone.

Half Kneeling Hip Flexor Stretch, Modified: 1. Kneel on one knee with the other foot forward in a lunge position 2. Tuck your pelvis under and gently shift your weight forward, feeling a stretch through the front of the kneeling hip 3. Keep the front knee tracking over the foot, and avoid letting the kneeling side hip drift outward into adduction 4. Hold for 20 to 30 seconds, repeat 2 to 3 times per side

Bird Dog: 1. Start on hands and knees 2. Extend one arm forward and the opposite leg straight back, keeping your pelvis level 3. Hold for 5 to 10 seconds, return to start 4. Perform 2 to 3 sets of 8 reps per side

Walking

Regular walking remains one of the simplest and most useful tools for most people with hip pain, as it loads the hip abductors rhythmically and predictably. If walking aggravates your specific pain pattern, shortening your stride slightly and avoiding uneven or heavily cambered surfaces can help, and this is a good topic to bring to your provider if walking consistently flares your symptoms.

A Word on Pacing

Progress in tendon rehabilitation is rarely linear. A tendon that has been irritated for months did not become irritated overnight, and it will not resolve overnight either. Expect some day to day variation, and treat a single bad day as information rather than a sign that the whole plan has failed.

8. Integrative and Physical Therapies

8.1 Physical Therapy

A skilled physical therapist can build and progress the kind of graded loading program described in Chapter 7, assess the movement patterns and muscle imbalances discussed in Chapter 6, and adjust your program as your tendon's tolerance changes. Given the strength of the evidence behind structured education and exercise for gluteal tendinopathy specifically, physical therapy is a reasonable first line recommendation for most people with lateral hip pain.⁶

8.2 Chiropractic Care

Chiropractic evaluation and care can address joint mobility in the hip, pelvis, and lumbar spine, and can incorporate the regional interdependence assessment described in Chapter 6, including screening for functional leg length asymmetry and pelvic mechanics that may be contributing to your hip pain. This guide does not have a hip specific randomized trial of chiropractic care to cite here, and it is more honest to say so than to borrow evidence from a different condition. What chiropractic care can reasonably offer is a whole person evaluation of the kinetic chain feeding into your hip, paired with hands on treatment and a home exercise program.

8.3 Corticosteroid Injections

As detailed in Chapter 4, corticosteroid injections can provide genuine short term relief for gluteal tendinopathy, but the best available randomized evidence found that an eight week education and exercise program matched or outperformed a single injection at every time point measured, including one year later.⁶ An injection may be a reasonable part of a plan, particularly to reduce pain enough to begin exercise more comfortably, but it functions best as a bridge to active rehabilitation rather than as a standalone treatment.

8.4 Shockwave Therapy

Extracorporeal shockwave therapy, which delivers acoustic pulses to irritated tendon tissue, is used clinically for gluteal tendinopathy that has not responded to a structured exercise program. It is a reasonable option to discuss with your provider if first line care has stalled, though it is generally considered a second line option after a genuine trial of load management and exercise rather than a replacement for it.

8.5 Heat and Cold

Heat and cold are simple, low risk, and can offer symptomatic comfort. Cold applied for 15 to 20 minutes may help calm an acutely irritable tendon after a flare, while heat can ease general stiffness and muscle tension in the surrounding hip flexors and external rotators. Neither replaces a loading program, but either can make daily life more comfortable while you work through one.

8.6 Massage and Soft Tissue Work

Massage and soft tissue work targeting the hip flexors, external rotators, and TFL can provide short term relief of tension and may support your ability to tolerate the exercise program in Chapter 7. As with chiropractic care, this guide does not have hip

specific trial evidence to cite for massage, and presents it here as a reasonable, low risk adjunct rather than a primary treatment.

9. Nutrition for Joint and Tendon Health

What you eat will not repair a tendon on its own, and this guide is not going to claim otherwise. A general pattern that supports connective tissue repair and manages systemic inflammation is nonetheless a sensible part of a whole person recovery plan.

Foods to Emphasize

Favor foods that supply the building blocks of tendon and cartilage repair alongside a broadly anti-inflammatory pattern of eating: quality protein sources such as wild caught fish, pasture raised poultry, eggs, and legumes; colorful vegetables and fruits, particularly leafy greens and berries; omega-3 rich foods like salmon, sardines, walnuts, and flaxseed; and healthy fats such as olive oil and avocado. Bone broth and collagen containing foods supply amino acids used in connective tissue repair, though as with any supplement or food based approach, expect this to support your overall recovery environment rather than to resolve a specific tendon problem on its own.

Foods to Minimize

Refined carbohydrates and added sugars, deep fried foods, heavily processed meats, and excessive alcohol are worth minimizing, both for general inflammatory tone and because they tend to displace more nutrient dense choices from your plate.

Hydration

Adequate hydration supports tissue health broadly. There is no need to chase a specific number. Drink regularly through the day and let thirst and pale colored urine guide you, unless your provider has given you specific fluid guidance for a heart or kidney condition.

Weight and Joint Load

Carrying additional body weight increases the mechanical load passing through your hips with every step, which matters for both hip osteoarthritis and tendon based conditions like gluteal tendinopathy. Reaching and maintaining a healthy weight is worth pursuing for hip health among many other reasons, though it should be approached as one part of a broader plan rather than a precondition for starting the exercise program in Chapter 7.

10. Lifestyle Modifications: Sleep, Weight, and Footwear

10.1 Sleep Positioning

Because gluteal tendinopathy is aggravated by compression from hip adduction, how you sleep matters directly. If you sleep on your side, place a pillow between your knees to keep your top hip from adducting and compressing the irritated tendon against the mattress, and consider sleeping with a pillow supporting the top leg more broadly if lying on the affected side is particularly painful. Some people find temporary relief sleeping on their back with a pillow under their knees, or on the unaffected side.

10.2 Standing and Sitting Posture

Avoid standing for long periods with your weight shifted into one hip, the hip hitching posture mentioned in Chapter 7, which compresses the outer hip tendons on the weight bearing side. When sitting, avoid prolonged leg crossing for the same reason, and take regular breaks to change position rather than holding any one posture for hours.

10.3 Footwear and Leg Length

If a functional leg length asymmetry has been identified as part of your evaluation, as discussed in Chapter 6, your provider may recommend a heel lift or orthotic on one side. The evidence here, again, is real but modest: a systematic review of shoe lifts for leg length discrepancy found low quality but broadly positive evidence for pain relief, alongside a clear call for higher quality research to determine which patients benefit most and by how much correction.⁹ This is a reasonable intervention to try under professional guidance, with realistic expectations set from the outset. Supportive, well fitted footwear generally, with adequate cushioning for daily walking, is a reasonable and low risk addition regardless of leg length findings.

10.4 Activity Modification During a Flare

During a significant flare, temporarily reducing high compressive or high impact activities, such as running on cambered roads, deep squatting past 90 degrees, or prolonged standing on one leg, can calm symptoms while you continue the lower irritation isometric and strengthening work from Chapter 7. This is a temporary adjustment, not a long term restriction, and the goal is always to return to full activity through graded loading rather than through indefinite avoidance.

11. The In-Clinic Pre-Treatment Protocol

Understand what this is for, because the distinction matters.

None of this rebuilds tissue. The loading and strengthening work in the previous chapters does that. What this protocol does is reduce the muscle tone, stiffness and tissue sensitivity that make the strength work harder to tolerate and less pleasant to do, particularly in the early weeks when things are at their most irritable.

It is preparation, not treatment. I use it in the clinic before the working part of a session, and I give it to patients to use at home before theirs. Patients who do it tend to tolerate the loading better and stick with it longer, and sticking with it is the actual mechanism by which recovery happens.

If you have to choose between doing this and doing your strength work, do your strength work.

Percussion device (Hypervolt), round or flat tip.

- Tensor fasciae latae and hip flexors, 1 to 2 minutes.
- All three glutes, 1 to 2 minutes per side.
- Adductors, gently, 1 minute per side.

Vibrating sphere (Hypersphere Mini) or roller (Vyper or foam roller).

- Glutes, 2 to 3 minutes per side. Lie on your back and move the leg up and down, or sit and roll.
- Hip flexor, psoas and iliacus, 1 to 2 minutes per side.
- IT band, side lying, from hip to just above the knee, 1 to 2 minutes per side.

Safety, and I mean these.

- Avoid all of this over an acutely injured area in the first 48 to 72 hours.
- Reduce pressure over anything actively inflamed.
- Skip percussion therapy entirely if you have a pacemaker or another implanted electronic device.
- Avoid aggressive pressure over varicose veins or compromised skin.
- Start light. Pain must not exceed 6 out of 10. Back off immediately for sharp or shooting pain.
- Stop and seek advice if you get numbness, tingling, or weakness.

12. Red Flags: When to Seek Immediate Care

Most hip pain is not an emergency, and the great majority of what is covered in this guide responds to the kind of structured, active care described in earlier chapters. A small number of presentations, however, warrant prompt medical evaluation, and it is worth knowing them.

Seek prompt medical evaluation if you experience:

- Hip pain following a fall or direct trauma, particularly if you are older or have known osteoporosis risk, because this raises concern for a hip fracture
- Fever accompanying joint pain, which can indicate infection
- Unexplained weight loss along with hip or pelvic pain
- Night pain that is not relieved by any change in position, which is different from pain that simply worsens with lying on the affected side
- Progressive weakness, numbness, or other neurological symptoms in the leg
- Groin pain severe enough that you are unable to bear weight on the leg at all

These situations call for prompt evaluation, not panic, and most people who experience one of them do not turn out to have a serious underlying condition. But they are exactly the situations where this guide, or any self-directed exercise program, should be set aside in favor of a timely appointment or, for a suspected fracture or infection, urgent care.

13. Building Your Personal Recovery Plan

Use this as a starting framework to discuss with your provider, not as a fixed prescription. Your specific findings, particularly which structures are involved and how irritable your symptoms currently are, should shape how quickly you move through these stages.

Weeks 1 to 2: Calm and Assess

Begin with a proper evaluation that looks at your hip, your lumbar spine, your pelvis, and your movement pattern as a whole, consistent with the regional interdependence approach in Chapter 6. Start gentle isometric hip abduction work if tolerated, adjust your sleep position to avoid compressive adduction, and minimize hip hitching and prolonged leg crossing. Avoid aggressive stretching into hip adduction during this period.

Weeks 3 to 6: Build Tolerance

Progress from isometrics into the sidelying abduction, standing hip hike, and banded bridge work described in Chapter 7, adding hip flexor mobility and general core and pelvic control work like bird dog. Continue addressing sleep positioning and standing posture. If pain is significant, this is a reasonable point to discuss a corticosteroid injection with your provider as a bridge to more comfortable exercise participation, understanding from the LEAP trial evidence that the injection works best paired with, not instead of, the exercise program.⁶

Weeks 7 to 12: Strengthen and Restore

Progress resistance in your hip abductor strengthening, reintroduce walking distance and pace gradually, and begin addressing any identified functional leg length asymmetry with your provider's guidance. This is also a reasonable point to consider shockwave therapy if progress has plateaued despite consistent loading work.

Beyond 12 Weeks: Maintenance

Continue hip abductor and general lower body strength work two to three times per week indefinitely, since deconditioning is a common precursor to recurrence. Maintain attention to sleep positioning and posture as ongoing habits rather than temporary fixes, and revisit your provider if symptoms plateau or recur despite consistent adherence to your program.

Final Thoughts

Hip pain, and lateral hip pain in particular, has been misunderstood and mismanaged for a long time under the label of bursitis. The research is now clear that for most people with pain over the outer hip, the real story is a tendon that has been asked to do more than it was prepared for, usually because the muscles and movement patterns feeding into that tendon, the psoas and iliacus, the external rotators, the TFL, and the pelvis and low back beyond, were not sharing the load the way they should have been.

That reframing matters because it points toward a genuinely different and genuinely more hopeful path forward: not rest and injections alone, but understanding your load, retraining your gluteus medius and minimus, and addressing the whole kinetic chain around your hip rather than the painful spot in isolation. That is the root cause approach this guide has tried to walk you through, and it is the approach I use with my patients at Root Health every day.

You do not need to have all the answers before you start. You need an accurate evaluation, a plan that respects both the strong evidence behind load management and the honest limits of what is known about some of the other contributing factors, and the patience to let a tendon or a movement pattern change over weeks rather than days.

If you would like a thorough, root cause evaluation of your hip pain and the patterns that may be feeding into it, I offer a free Root Discovery Session at Root Health. You can schedule yours at rootcauseunlocked.com/discovery. You can also reach the office at (919) 662-0520.

Consultations and ongoing care are both available wherever you are located. Being outside North Carolina is not a barrier to working with me. Tell me where you are when you reach out and I will lay out exactly how we would work together from there.

14. References

Every reference below was retrieved and checked against the published record on PubMed. PMID numbers are included so you or your provider can look up any claim in this guide directly.

1. Besomi M, Maclachlan L, Mellor R, Vicenzino B, Hodges PW. Tensor fascia latae muscle structure and activation in individuals with lower limb musculoskeletal conditions: a systematic review and meta-analysis. *Sports Med*. 2020;50(5):965-985. PMID: 31898217
2. Chamberlain R. Hip pain in adults: evaluation and differential diagnosis. *Am Fam Physician*. 2021;103(2):81-89. PMID: 33448767. (Erratum: *Am Fam Physician*. 2021;103(5):263. PMID 33630548. Screened 2026-09-05; corrected, not retracted.)
3. Cohen SP. Sacroiliac joint pain: a comprehensive review of anatomy, diagnosis, and treatment. *Anesth Analg*. 2005;101(5):1440-1453. PMID: 16244008
4. Bird PA, Oakley SP, Shnier R, Kirkham BW. Prospective evaluation of magnetic resonance imaging and physical examination findings in patients with greater trochanteric pain syndrome. *Arthritis Rheum*. 2001;44(9):2138-2145. PMID: 11592379
5. Grimaldi A, Mellor R, Hodges P, Bennell K, Wajswelner H, Vicenzino B. Gluteal tendinopathy: a review of mechanisms, assessment and management. *Sports Med*. 2015;45(8):1107-1119. PMID: 25969366
6. Mellor R, Bennell K, Grimaldi A, Nicolson P, Kasza J, Hodges P, Wajswelner H, Vicenzino B. Education plus exercise versus corticosteroid injection use versus a wait and see approach on global outcome and pain from gluteal tendinopathy: prospective, single blinded, randomised clinical trial. *BMJ*. 2018;361:k1662. PMID: 29720374
7. Sueki DG, Cleland JA, Wainner RS. A regional interdependence model of musculoskeletal dysfunction: research, mechanisms, and clinical implications. *J Man Manip Ther*. 2013;21(2):90-102. PMID: 24421619
8. Kuligowski T, Dębiec-Bąk A, Cieślík B. Hamstring strength and hip mobility associated with pain and disability in lumbar instability. *Sci Rep*. 2025;16(1):2332. PMID: 41366102
9. Campbell TM, Ghaedi BB, Tanjong Ghogomu E, Welch V. Shoe lifts for leg length discrepancy in adults with common painful musculoskeletal conditions: a systematic review of the literature. *Arch Phys Med Rehabil*. 2018;99(5):981-993.e2. PMID: 29229292
10. Menez C, L'Hermette M, Lerebourg L, Coquart J. Effects of insoles on gait kinematics and low back pain in patients with leg length inequality: a systematic review. *J Am Podiatr Med Assoc*. 2023;113(2). PMID: 37134059
11. O'Leary CB, Cahill CR, Robinson AW, Barnes MJ, Hong J. A systematic review: the effects of podiatric deviations on nonspecific chronic low back pain. *J Back Musculoskelet Rehabil*. 2013;26(2):117-123. PMID: 23640312
12. Cook JL, Purdam CR. Is tendon pathology a continuum? A pathology model to explain the clinical presentation of load-induced tendinopathy. *Br J Sports Med*. 2009;43(6):409-416. PMID: 18812414

This guide was compiled from peer-reviewed medical literature and clinical experience. Every citation was verified against the published record. Where a claim reflects clinical judgment rather than trial evidence, this guide has said so directly. It is intended to give you knowledge and options to discuss with your healthcare provider. It is not a substitute for individual medical care, and it does not diagnose your condition or promise a cure.

This document is a DRAFT pending Dr. Seay's clinical review.

(c) 2026 Root Health · For personal educational use only

Read this part first

This guide is educational. It is not medical advice, it does not diagnose anything, and reading it does not make you my patient.

Some back pain is an emergency. Know these before anything else in this guide:

- New numbness in the saddle area (groin, inner thighs, the parts that would touch a bicycle seat), new trouble controlling bladder or bowels, or progressive weakness in both legs. This pattern can mean cauda equina compression. **Emergency room, now, not tomorrow.**
- Back pain with fever, back pain after significant trauma, or new back pain if you have a history of cancer, IV drug use, or a suppressed immune system. Same-day medical evaluation.
- A fall followed by groin or hip pain and inability to bear weight. That is fracture-until-proven-otherwise territory.
- Night pain that is unrelenting regardless of position, week after week, deserves a physician visit rather than a foam roller.

The tool safety rules, from the same sheets we use in clinic: do not work over an acute injury in its first 48 to 72 hours; reduce pressure over inflamed areas; skip percussion devices entirely if you have a pacemaker or other implanted electronic device; avoid aggressive pressure over varicose veins or compromised skin; be cautious if you take blood thinners. Start light. On the manual tools, the foam roller, roller stick, ball and trigger point tool, where the pressure is your own body weight, pain must not exceed 6 out of 10: uncomfortable but tolerable. **On a powered percussion device the rule is different and stricter.** Follow the manufacturer's instructions, which say to use it without producing pain or discomfort and to stop immediately if you feel any, apart from light muscle soreness. I am not asking you to push a powered tool to a number. Sharp or shooting pain means back off, whatever the tool.

Nothing here is a reason to stop a prescription or skip a recommended evaluation.

Why one guide covers the low back and the hip

The tools sort by region, and so does the biology. The low back and hip share their major movers (glutes, hip flexors, hamstrings, adductors), share load between them with every step, and irritate each other when either stops doing its share. Half of what presents as “low back pain” carries a hip contribution, and lateral hip pain is routinely mislabeled, including by the people who have it. This guide covers the region: low back pain, sciatica, and hip pain, including the gluteal tendinopathy that most “hip bursitis” actually is.

If you were told you have... (your diagnosis, translated)

Diagnostic labels for this region have changed, and most readers arrive carrying an older one. Here is where yours lives in this guide:

- **“Trochanteric bursitis,” “hip bursitis,” or “greater trochanteric pain syndrome”**: the lateral hip pain territory. The research now treats most of it as gluteal tendinopathy rather than an inflamed bursa, and the LEAP trial in part two is your evidence.
- **“Sciatica,” a “pinched nerve,” or a “bulging disc”**: the sciatica material in part two, plus the companion Sciatica ebook. A disc bulge on imaging is common in pain-free backs; the finding and the pain are not the same thing.
- **“Degenerative disc disease” or “arthritis of the spine”**: labels for common age-related change that sound far scarier than their prognosis. The low back program applies.
- **“Piriformis syndrome”**: a community label for deep buttock pain; the glute and hip work in this guide covers that territory.
- **“Spinal stenosis”** (leg symptoms when walking that ease when you sit or lean forward): this deserves its own clinical evaluation before assuming this guide’s program fits, especially past sixty.

The one idea that makes this kit honest

Here is the sentence that separates this guide from most of what sells in this category: **the tools buy you a window, and the movement spends it.**

The evidence for massage tools is real but modest, and it is short-term: foam rolling used as a warm-up produces small improvements in flexibility and sprint performance (PMID 31024339), and massage guns improve short-term range of motion, flexibility, and recovery measures, while doing nothing for strength or performance, and are actually not recommended immediately before demanding strength or balance work (PMID 37754971). That last point changes the order of your session rather than the contents of it, so let me be specific instead of leaving you to reconcile it: the rolling, the stretches, the heat and the ball work can all run straight into your movement work, and the percussion massager should not. Use it after a session, on a rest day, or leave roughly half an hour before you load. Heat has the same honest profile, and one finding worth reading twice: moderate evidence shows heat wrap therapy gives a small short-term reduction in low back pain and disability, **and adding exercise to the heat further reduces pain and improves function** (PMID 16437495). The tools prepare tissue and calm symptoms so that moving feels possible. The movement is the treatment. Everything in this guide is sequenced around that fact, which is exactly how these same protocols are used before treatment in our clinic.

Part one: the tissue prep protocols, exactly as used in clinic

These are the same protocols patients receive at Root Health, tools, targets, and doses unchanged. Do them before your movement work, not instead of it. Respect the pressure ceilings above, manual versus powered, throughout.

One timing rule carries over from the section above, and it is the only place the order matters: the percussion massager is the one tool not to use immediately before strength work. Everything else here can run straight into your session.

The low back sequence

Heat and vibration back wrap, set to vibrate and heat, worn while performing the following:

- **Cat-camel stretch:** on hands and knees, arch the back up, then let it sag. 1 set of 10 repetitions.
- **Hamstring stretch with stretch strap:** on your back, strap around the foot, leg straight up with a soft knee. 1 set of 10 reps each side.
- **Ball bounce:** seated on an exercise ball, bounce up and down, then side to side, palms up throughout. 1 set of 10 each.

Massage ball or foam roller: lie on your back, place it on the glute muscles, and move the leg up and down. Work each side.

Massage gun, round or flat head: run the device over the TFL and hip flexors for 1 to 2 minutes each side.

Massage ball: place over the abdomen (psoas territory) and rock back and forth, then up and down. 20 to 30 seconds each side. This one is unusual and it is deliberate: the psoas connects the lumbar spine to the leg, and it rarely gets touched by anything.

Lumbar support device or rolled towel: lie on your back with the device under the lower back, knees bent to start (straighten if comfortable), and rest 30 to 60 seconds.

Massage ball or foam roller on the calf: find a sore spot, then move the foot up, down, left, right, and in circles. 2 to 3 repetitions per area. The calves are part of the chain that decides how your gait loads your spine.

The hip sequence

Massage gun, round or flat head:

- TFL and hip flexors, 1 to 2 minutes.
- All three glute muscles (maximus, medius, minimus), 1 to 2 minutes per side.
- Adductors (inner thigh), gently, 1 minute per side.

Massage ball or foam roller:

- Glutes: lie on your back with the sphere on the glutes and move the leg up and down, or sit on the roller and search for tender spots. 2 to 3 minutes per side.
- Hip flexor (psoas, iliacus): roll over the front of the hip, 1 to 2 minutes per side.
- IT band: side-lying, roll the outer thigh from below the hip to above the knee, 1 to 2 minutes per side.

Stretches: cat-camel, 1 set of 10. Hamstring stretch with the strap, 1 set of 10 each side. Modified pigeon pose for the deep hip rotators, 20 to 30 seconds per side, only if comfortable.

One caution specific to lateral hip pain: if your pain lives on the outside of the hip and hurts to lie on, go easy on direct sustained pressure there (including sitting on the sphere and aggressive IT band rolling). Compressive load over an irritated gluteal tendon can flare it. Treat above and around it, and let the loading program in part two do the heavy lifting.

Part two: the movement evidence, which is the actual treatment

This is the least glamorous and best-supported section of the guide. Nobody profits from telling you to walk, which, given how this guide reads funding everywhere else, is worth something. The 2026 hip review below lists its funding as none at all.

For chronic low back pain, exercise works, and resting does not. The Cochrane review of exercise therapy for chronic low back pain found moderate-certainty evidence that exercise is effective for pain compared with no treatment, usual care, or placebo, and that exercise probably beats advice or education alone (PMID 34580864). Which style of exercise matters far less than doing one: the review found no clear winner between approaches, so the best program is the one you will actually repeat. For a new episode of acute back pain, the advice with moderate-quality evidence behind it is to stay active rather than rest in bed (PMID 20556780). Bed rest is not treatment. It is deconditioning with a mattress.

For sciatica, time is on your side more than most people are told. In the randomized trial comparing early surgery against prolonged conservative care for severe sciatica, early surgery relieved leg pain faster, but disability outcomes were no different across the first year, and only 39 percent of the conservative group ended up needing surgery at all (PMID 17538084). At five years, outcomes between the groups were still equivalent (PMID 23793663). Severe or progressive

weakness, or any of the red flags on page one, changes that calculus immediately. Short of those, a patient and progressive rehab approach is not the timid option; it is the evidence-based one, with surgery remaining available if genuinely needed.

For hip osteoarthritis, the honest answer got harder in 2026, and you should have it anyway. The Cochrane review on this question was rebuilt in July 2026, and it is the most sobering piece of evidence in this guide. Across 18 studies and 1,368 participants, exercise compared against a placebo or attention control may have little to no effect on pain, and may improve physical function only slightly, both on low-certainty evidence. Compared against no treatment or usual care, exercise probably does reduce pain and improve function, but the reviewers say plainly that the size of those improvements is unlikely to be clinically meaningful. And where exercise was added on top of another treatment, it probably made little to no difference to pain or function at all (PMID 42486497).

That is a smaller and less certain benefit than the previous edition of this review reported. The hip is not the knee, the evidence base here is much thinner, and I would rather hand you the current number than a more flattering one that has been superseded.

So why is exercise still the first thing I do for an arthritic hip, and the first thing every guideline recommends? Three reasons, and none of them require overselling. The alternatives for hip osteoarthritis are worse: injections, medications with real organ costs, and eventually a replacement. Exercise in that same review probably reduced the risk of adverse events slightly rather than raising it, which almost nothing else on the list can claim. And the trial averages you are reading are built from people who did the program and people who did not, which is precisely why the rest of this part is written to get you to actually do it. A small average benefit with no downside, in a condition where the other options carry real ones, is still the right first move. It is just not the miracle the internet sells, and you deserve the real number.

For lateral hip pain (gluteal tendinopathy), the trial worth knowing beat the injection. In a three-arm randomized trial published in the BMJ, fourteen sessions of load-management education plus exercise produced higher rates of global improvement and lower pain at eight weeks than either a corticosteroid injection or a wait-and-see approach, and at 52 weeks the education-plus-exercise group still reported better global improvement than the injection group (PMID 29720374). If you have been told your outer-hip pain is “bursitis” and offered a needle first, this trial is the conversation-changer: the boring program of progressive loading and not compressing the tendon (less leg-crossing, less hanging on one hip, side-sleeping with a pillow between the knees) outperformed it.

The program

Start with the finding that takes the pressure off. The Cochrane review of exercise for chronic low back pain looked for a winning style and did not find one (PMID 34580864). No approach beat the others. Read that as permission rather than as a disappointment: you are not one correct exercise away from your result, and a boring program you repeat for three months outperforms a perfect one you abandon in week two. If you can only do part of what follows, do that part.

The core, for everybody in this guide

1. **Walk most days.** Start at a distance that is comfortable today, and add roughly ten percent a week. Hold at the same distance for a week whenever a week feels hard. This is the single item with the broadest support in the whole guide and it costs nothing.
2. **Strength, two or three times a week, on non-consecutive days.** Two to three sets of 8 to 12 repetitions, slow enough that you could stop at any point in the range.
3. **Hinge well.** Learn to bend from the hips with a stiff-but-not-rigid spine, and use it for everything you pick up. This is a skill, not an exercise, and it pays out every day.
4. **Do not rest it.** For a new episode, staying active beats bed rest on moderate-quality evidence (PMID 20556780). Move at whatever size you can.

Start here: pick the one that sounds like you

If your picture is chronic nonspecific low back pain:

- Glute work is your priority: bridges, then single-leg bridges, plus standing or side-lying hip abduction.
- Add a hinge pattern you can load, and the cat-camel from part one as daily movement rather than as treatment.
- Add walking, which for this picture is not the warm-up. It is the treatment.
- Expect a sawtooth, not a straight line. Good weeks and bad weeks inside an overall trend upward is what recovery actually looks like here.

If your picture is sciatica (leg pain, sometimes with numbness or tingling, often worse than the back pain itself):

- Time and load are on your side. The trials say most people get there without surgery, and that waiting does not cost you the option (PMID 17538084, PMID 23793663).
- Find the directions that centralise your symptoms, meaning the ones that pull the pain out of the leg and back toward the spine, and favour those. Moving pain toward the midline is progress even when the intensity has not changed yet.
- Walk in short, frequent bouts rather than one long one.
- **Progressive weakness, a foot that slaps or catches, or any saddle-area or bladder change overrides all of this and needs assessment now, not at six weeks.**

If your picture is an arthritic hip:

- Sit-to-stand from a chair, chair height as your difficulty dial, plus hip abduction and bridges.
- Stationary cycling or pool work if weight-bearing is the limit. The 2026 review found no evidence that one exercise type beats another, so pick what you will keep doing.
- Read the honest section above before you set expectations. The average benefit here is small. Consistency and stacking it with weight management and walking is how you get above that average.

If your picture is lateral hip pain (outside of the hip, hurts to lie on that side, often called bursitis):

- **This is the pathway where the evidence is strongest and most specific in this entire guide.**
Education plus exercise beat a corticosteroid injection and beat waiting, at eight weeks and still at 52 weeks (PMID 29720374). Fourteen supervised sessions was the trial's dose, which tells you this is a months-long project rather than a fortnight's.
- The load-management half matters as much as the exercise half, and it is mostly about not compressing the tendon: less leg-crossing, less standing hanging on one hip, and a pillow between the knees when you sleep on your side.
- Progressive hip abduction loading is the exercise half. Start isometric, holding rather than moving, if the tendon is angry.
- Go easy with the tools directly over the painful spot, per the caution in part one.

How to make it harder

Progress one thing at a time, roughly every one to two weeks, and only when the current version is comfortable: more repetitions first, up to about 12; then a harder version of the movement; then added load; then more sets. If you add difficulty and the next two sessions feel worse, go back one step for a week. That is the method working, not a setback.

How to know it is going right, and when to back off

The soreness rule. Discomfort during and after this work is expected and is not damage. The test is the next morning. Soreness that has settled by then is a normal training response and you carry on. Soreness that is still there, or worse, means the last session was too much, so drop back to the version you handled comfortably.

Reduce the load if a flare lasts more than a couple of days, or if pain starts waking you at night.

Stop and get examined for anything in the warning list at the top of this guide, for leg symptoms that are spreading or intensifying, for new weakness, or for pain trending worse across weeks rather than better.

Give it twelve weeks before you judge it, and check in at six. Six weeks of honest effort with nothing to show is a reason to be examined rather than a reason to push harder. Track it, and let the trend rather than any single day judge the program.

When to get evaluated instead of self-treating

Beyond the emergencies on page one: pain that is not trending better by six weeks of honest effort, leg symptoms that are spreading or intensifying, new weakness (a foot that slaps or catches, a knee that buckles), or a flare pattern you cannot connect to load. Chiropractic, physical therapy, and medical evaluation are complements here, not rivals, and a clinician who examines you beats any guide that cannot.

Part three: the supplement half, honestly graded

Why nutrient evidence looks thinner than it is

A large outcome trial costs tens to hundreds of millions of dollars, and that money only exists when a patent can pay it back. You cannot patent magnesium, an herb, or a food. So the evidence hierarchy this guide grades against is not neutral: it systematically over-documents patentable drugs and under-documents nearly everything else. That is measurable, not rhetorical. A Cochrane methodology review found that manufacturer sponsorship of drug and device studies produces more favourable results and conclusions than independent funding, through a bias that standard quality checks do not catch (PMID 28207928).

So when this guide says “no evidence,” it will tell you which of three things it means: *tested and failed* (a real trial ran and the answer was no), *never properly tested* (nobody could afford the trial, so absence of evidence is not evidence of absence), or *tested small and won* (real but modest trials, usually measuring markers or short-term pain, because that is what a small budget can afford). The asymmetry earns nutrients humility, not automatic credit: small nutrient trials carry their own publication bias toward positive results, just as drug trials tilt toward their sponsor. One rule, both directions, and I apply it to the supplement industry below exactly as hard as I apply it to drug companies.

Sort the tissue before you buy anything

This is the step that decides whether a supplement has any chance of helping you, and almost nobody takes it. “Low back and hip pain” is not one problem. It is at least three different tissues failing in different ways, and they do not want the same things. The aisle sells one joint capsule to all three of you.

The joint is the arthritic hip: a worn, irritated bearing surface in a joint that has turned inflammatory. That is the tissue nearly all joint-supplement research was actually done in, so for that group the evidence is borrowed only slightly.

The tendon is lateral hip pain, the ache on the outside of the hip that hurts to lie on, which you were probably told was bursitis. That is gluteal tendinopathy, and a tendon is a rope of collagen, not a bearing surface. It follows completely different rules, and it is the group most easily failed, because it gets lumped in with nonspecific back pain and told that nothing works.

The spine and its muscles is nonspecific low back pain, where the honest answer stays thin and I am not going to dress it up.

One rule holds across all three, and it is why this section sits behind the movement section rather than in front of it. Every trial worth citing below layered its supplement on top of a loading program. Nutrition supplies raw material. Load is what tells the body where to put it. A supplement taken instead of the program is money spent on a signal your body has no reason to act on.

The spine: tested and failed, said plainly

Glucosamine for low back pain. A double-blind randomized trial of 250 patients with chronic low back pain and degenerative lumbar changes found six months of glucosamine produced no reduction in pain-related disability versus placebo, at six months or at one year (PMID 20606148). If you take glucosamine for your back, that is the trial to know.

Vitamin D for low back pain. The systematic review and meta-analysis conclusion is direct: not more effective than placebo or other interventions, and “the prescription of vitamin D for LBP cannot be recommended” on current evidence (PMID 29565945). Correcting a genuinely deficient vitamin D level still matters for bone and general health, and if you have a measured deficiency you should fix it on those terms. Expecting it to treat back pain is the part the evidence does not support. Test, then treat what the test shows, rather than treating a symptom with a nutrient that has been tried for exactly that and did not deliver.

For nonspecific low back pain, there is still no supplement with trial support worth your money, and I will not invent one. What I will do, instead of ending the paragraph there, is point you at the two things in this section that do apply to you regardless of which tissue you have: the protein floor and the load. Both are further down, both are free, and both outperform anything you would have bought here.

The arthritic hip: borrowed evidence, labeled as borrowed

The supplements with real positive signals in this territory were tested in **osteoarthritis**, overwhelmingly knee osteoarthritis. That transfers reasonably to an arthritic hip, which is the same disease in a different joint, and poorly to nonspecific low back pain, and this guide refuses to blur that line.

The systematic review and meta-analysis of dietary supplements for osteoarthritis, covering 20 supplements across 69 trials, found *Boswellia serrata* extract and curcumin among seven showing large and clinically important short-term pain reductions, while rating the overall quality of that evidence very low. At medium term only two supplements held a clinically important effect on pain, green-lipped mussel extract and undenatured type II collagen, and at long term none did (PMID 29018060). The *Boswellia*-specific meta-analysis of seven trials and 545 patients found meaningful short-term improvement in pain, stiffness and function, and concluded a fair trial runs at least four weeks (PMID 32680575). The same review found glucosamine and chondroitin either ineffective or showing small and arguably clinically unimportant effects even for osteoarthritis, which pairs with the back pain null above.

Curcumin has a delivery problem the label will not tell you about, and it decides whether you wasted your money.

Plain curcumin is absorbed poorly. The osteoarthritis trial that showed a real effect used a specifically engineered high-absorption preparation delivering 180 mg of curcumin daily for eight weeks, measured at roughly 27 times the absorption of ordinary curcumin powder, and the patients on it had less pain and needed significantly less anti-inflammatory medication than placebo (PMID 25308211). The number that matters is not the milligrams of turmeric on the front of the bottle, it is whether the product uses a documented absorption-enhanced form. A large dose of a poorly absorbed powder is not a bigger dose, it is a more expensive one.

Oral hyaluronic acid is genuinely promising rather than proven. In a twelve-month double-blind placebo-controlled trial, 200 mg daily alongside daily quadriceps strengthening in both groups improved osteoarthritis symptom scores more than placebo, but reached significance only in subjects aged 70 or younger and only at the second and fourth months (PMID 23226979). That is sixty people, age-conditional, and industry-funded. If you are under 70 with an arthritic hip and already doing the strengthening, 200 mg daily for three months is a defensible experiment, and it is not a reason to skip the strengthening that both groups in that trial were doing.

Lateral hip pain: you have a tendon, and that changes everything

If your pain lives on the outside of the hip and hurts to lie on, stop reading the arthritis section. You have a collagen problem, and collagen has its own biology, its own timing, and its own evidence.

Start with what a tendon actually is. Roughly three quarters of the dry weight of tendon is type I collagen, a rope of protein wound from repeating glycine, proline and lysine. To build it, your body chemically modifies proline and lysine after the chain is assembled, and the enzymes that do that, prolyl hydroxylase and lysyl hydroxylase, cannot function without vitamin C and iron. The enzyme that then cross-links those ropes into something that carries load, lysyl oxidase, requires copper. Manganese and zinc are required elsewhere on the same assembly line. This is textbook connective tissue biochemistry rather than a supplement claim, and it is why a collagen program that supplies protein without supplying cofactors is doing half the job.

Now the timing, which is the part almost nobody gets right. Collagen synthesis in tendon is driven by mechanical load and rises in the hours after you load the tissue, so the useful question is not “should I take collagen” but “is the raw material in my bloodstream when my tendon is being told to rebuild.” In the study that established this, eight healthy men took vitamin C-enriched gelatin an hour before six minutes of rope skipping, and the 15 gram dose doubled a blood marker of new collagen formation (PMID 27852613). A later dose-response trial found 30 grams of hydrolyzed collagen with 50 mg of vitamin C an hour before heavy resistance exercise produced more collagen synthesis than 15 grams, while **15 grams was not statistically different from nothing at all** (PMID 38007183). Most products on the shelf, and most people taking them, are dosed at or below the amount that failed to beat placebo. Run it at a dose the research supports, about an hour before the session, not at bedtime.

Here is the ceiling on that claim, and I would rather you hear it from me. A blood marker is not a healed tendon. When an independent group tested whether collagen peptides change tendon structure, giving 15 grams daily through fifteen weeks of resistance training with MRI endpoints, they found no advantage over placebo on tendon size, stiffness or mechanical properties. Both groups improved, and the training did the work (PMID 37436929). The most recent independent systematic review, nineteen studies and 768 participants, found small statistically significant effects and graded the tendon findings specifically as very low certainty (PMID 39060741). A positive trial does exist in a population close to many of you: 139 athletic adults with activity-related knee pain took 5 grams of bioactive collagen peptides daily for twelve weeks and reported significantly less activity-related pain than placebo (PMID 28177710). That trial carries a correction notice and one of its authors works for a collagen research institute, which is exactly the sponsorship pattern the Cochrane review at the top of this section warns about. I cite it anyway, labeled, because pretending it does not exist would be its own dishonesty.

So the fair summary is this. The mechanism is real and well described. The acute response is real and dose-dependent. Symptom trials are positive but small and mostly industry-adjacent. The one hard structural endpoint tested independently came back null. That is a cheap, low-risk experiment with honest odds, not a treatment with proven structural benefit, and anyone selling it as the latter is ahead of the evidence.

The cofactor half is the smarter half. If substrate is only useful when the assembly line is staffed, the cofactors matter as much as the grams, and they are what consumer products ignore. A practitioner formula supplying vitamin C, copper, manganese, zinc and the amino acids proline and lysine together is built on real biochemistry rather than a marketing story. Two honest notes. Its amino acid content is modest, typically around 150 mg each of proline and lysine, a rounding error next to the 15 to 30 grams used in the collagen trials, so it complements a collagen powder rather than replacing it. And these

formulas often contain iron, which is a genuine cofactor here, but most men and most postmenopausal women should not take supplemental iron without a documented reason. Iron is one of the few nutrients where routinely taking more than you need is actively harmful.

Stage matters more than product, and this is the most useful paragraph in the section. In a three-arm randomized trial of 59 patients with Achilles tendinopathy, a supplement of mucopolysaccharides with type I collagen and vitamin C added benefit specifically in patients with early, reactive tendinopathy, and added nothing once the tendon had reached the degenerative stage with established matrix and vascular change (PMID 28053674). That trial was run in an Achilles, not a gluteal tendon, so treat the specific result as borrowed. What transfers is the principle: an early, angry, reactive tendon is still responsive tissue, and that is the window where nutritional support appears to add something, while a long-standing degenerative tendon has structural change that a capsule did not reverse. This is why “how long has this been going on” is one of the first questions I ask, and it should change what you spend money on. Weeks old, the supplement side is worth running. Two years old, put the money and the attention into the progressive loading program in part two, because that is what was actually shown to work at that stage, and in lateral hip pain specifically it outperformed a corticosteroid injection at 52 weeks (PMID 29720374).

The floor under all three tissues

The interventions with the best evidence in this entire section are not sold in the supplement aisle.

Protein is the one number worth knowing. The meta-analysis and meta-regression across 49 studies and 1,863 participants found protein supplementation meaningfully improves gains from resistance training, with benefits plateauing around 1.6 grams per kilogram of body weight per day (PMID 28698222). For a 180 pound person that is roughly 130 grams daily, and most people rehabbing an injury eat well under it. Collagen is a poor quality protein in isolation, low in the essential amino acids muscle needs, so treat a collagen dose as a targeted addition to your protein intake and never as your protein intake.

One disclosure on that paper, because I said I would run this rule in both directions. It carries a later correction noting that one of its authors sat on a sports supplement manufacturer's advisory board while it was written. The correction changed none of the data and the 1.6 figure stands, but holding the supplement industry to the standard I hold drug companies to means saying so when the study behind a number I like has a commercial tie on it, and not only when the number is one I dislike.

Sleep is when repair actually happens, energy sufficiency keeps repair funded, and at a weight-bearing joint any sustainable progress on excess load is mechanically meaningful with every step. Those outrank everything in a capsule and they earn me nothing.

What I would actually run, and how to judge it

Pick the tissue, not the product. Arthritic hip: an absorption-enhanced curcumin preparation or a Boswellia extract for eight to twelve weeks, optionally with 200 mg of oral hyaluronic acid daily if you are under 70, on top of the strengthening program. Lateral hip tendon pain: 15 to 30 grams of gelatin or hydrolyzed collagen with vitamin C about an hour before your loading sessions, cofactors covered, for twelve weeks, and weight it toward the early-and-reactive case rather than the long-standing one. Nonspecific low back pain: skip the aisle, hit the protein floor, and spend the money on the loading program instead. Underneath all three, 1.6 grams of protein per kilogram, sleep, and enough food.

Then judge it. Set a start date and an end date before you buy anything, change one thing at a time, and write down what you are measuring. Everything here is a defined trial with a stop date, not a subscription. If twelve weeks of honest effort produced nothing, that is a real answer worth having, and it sends you back to the loading program and to an examination instead of to the next bottle.

Interactions, and the ones people miss

Curcumin and Boswellia can interact with blood thinners, and curcumin at supplement doses can bother reflux-prone stomachs. Clear either with your prescriber if you take anticoagulants. Do not add supplemental iron, including inside a combination formula, without a reason to. Proprietary blends hide the dose of each ingredient behind a single total, which makes it impossible to know whether you are getting a researched amount of anything, so prefer products that disclose amounts. And whichever source you use, look for independent third-party testing, because the manufacturing side of this industry is even less regulated than the marketing side.

Curcumin and your liver. The National Center for Complementary and Integrative Health warns that the high-absorption curcumin formulations, the ones engineered so you absorb more of it, may harm the liver, and that liver damage has been reported in some people who took them. That warning applies to the product I name below, because high absorption is the entire reason I name it. The problem appears to be uncommon, and I would still rather you heard it from me than not at all. Stop the curcumin and call me if you notice unusual fatigue, nausea, loss of appetite, dark urine, or any yellowing of the skin or eyes. Do not take turmeric or curcumin supplements if you are pregnant, which the same source considers possibly unsafe. And treat eight weeks as a checkpoint rather than an open-ended course: we look at whether it actually helped and decide together whether to keep going.

Nothing in this section substitutes for being examined, and none of it applies to pain that is getting worse, waking you at night, or accompanied by any of the warning signs at the top of this guide.

Track it, so the program can be judged by data

The free Root Cause Unlocked tracker has an MSK mode built for exactly this guide: daily function, range of motion, load tolerated, and flare days, alongside the basics. Eight weeks of that turns “I think it might be helping” into an answer, and it makes any clinician visit dramatically more useful.

Free tracker: rootcauseunlocked.com/tracker?utm_source=guide&utm_medium=ebook&utm_campaign=msk-lbhip-recovery

Dr. Seay recommends: the exact products

The protocols in this guide are written generically on purpose, because the technique matters and the brand mostly does not. A rolled towel works. This section is the other half of that promise: if you would rather not stand in an aisle guessing, here is exactly what I would hand you, why that specific item and not a cheaper-looking equivalent, and where to get it. Read the disclosure at the end before you buy anything, because I want you deciding with your eyes open.

On the supplement side, through my dispensary

Everything below is available through the Root Health Fullscript dispensary. Search the product name exactly as written.

us.fullscript.com/welcome/drseay

For the arthritic hip: Theracurmin HP, by Integrative Therapeutics. Two capsules daily for eight weeks. I name this one specifically because it is the actual preparation used in the osteoarthritis trial discussed above, at the actual dose that trial used: two capsules deliver 180 mg of curcumin in the water-dispersible form measured at roughly 27 times the absorption of standard curcumin extract (PMID 25308211). This is the clearest case in this guide where the brand genuinely matters, because ordinary curcumin powder at the same milligram count is not the same product and did not produce those results.

Before you start this one, read the liver and pregnancy caution in the interactions section above. It applies specifically to high-absorption preparations, which is what this is.

For collagen cofactors, if lateral hip tendon pain is your picture: Collagenics, by Metagenics. Three tablets daily. This is the assembly line rather than the raw material: vitamin C, copper, manganese, zinc and the amino acids proline and lysine in one formula, which is the cofactor set the biochemistry section above describes. Read that section's iron caution first. Collagenics contains 6 mg of iron, and if you are a man or a postmenopausal woman without documented low iron, ask me or your prescriber before starting it rather than after.

For an early, reactive tendon: SynovX Tendon and Ligament, by Xymogen. Two capsules daily. This supplies the mucopolysaccharide and type I collagen combination with vitamin C tested in the tendinopathy trial above, where it added benefit in early-stage reactive tendinopathy and not in long-standing degenerative tendons (PMID 28053674). That trial was run in an Achilles, so match it to your stage rather than assuming it was proven in a hip. If your lateral hip pain is years old, spend the money on the loading program instead, and I would rather tell you that than sell you a bottle.

For collagen substrate: a plain hydrolyzed collagen or gelatin powder, 15 to 30 grams, with vitamin C. Here the brand genuinely does not matter and I will not pretend otherwise. What matters is the dose and the timing: aim toward the upper end of that range, take it about an hour before your loading session, and make sure vitamin C is on board. An unflavored bulk powder is the honest buy. Do not pay a premium for a proprietary blend delivering two grams.

If nonspecific low back pain is your picture, buy none of the above. The evidence does not support it and I am not going to sell you something to feel thorough. Put the money toward protein and toward the loading program.

Slots I have deliberately left open rather than filled: a Boswellia extract, an oral hyaluronic acid at 200 mg, and vitamin D. The Boswellia trials pooled several different extracts, so there is no single trial-matched product to point at the way there is with Theracurmin, and vitamin D should follow a test result rather than a guide. Ask me at your visit and I will pick these to fit you rather than guessing in print.

On the tool side

The protocols name categories, not brands, and every one can be done with inexpensive equipment. Any dense foam roller does the rolling work, a vibrating roller just adds vibration. A massage ball or lacrosse ball substitutes for a vibrating sphere. A rolled towel approximates the lumbar support device closely enough to try before you buy one. The percussion massager and the heat-plus-vibration wraps are the two items without a good household substitute, and they are also the most expensive, so they are the ones worth thinking hardest about.

If you want the specific equipment I use in the clinic, these are the links:

Percussion massager, vibrating roller, vibrating sphere and heat-plus-vibration wraps:

*[[HYPERICE_AFFILIATE_LINK]] **Foam roller, massage ball, stretch strap, lumbar support and exercise ball:***

[[GENERAL_TOOLS_AFFILIATE_LINK]]

Disclosure: Root Health earns a portion of dispensary purchases, and the tool links above are affiliate links, which means we may earn a commission if you buy through them at no additional cost to you. Nothing in this guide was included, excluded, ranked, or dosed based on what it earns. The generic substitutions listed above are real substitutions and I would rather you use a rolled towel than skip the work, and the highest-value items in this guide, walking, glute strengthening, protein, and sleep, earn us nothing at all.

Working with Root Health

Everything here is general education, written to be safe for a reader I have never examined. A personalized plan starts from your history, your exam, and your actual loading tolerance.

***Book a free Root Discovery Session: [rootcauseunlocked.com/discovery?](https://rootcauseunlocked.com/discovery?utm_source=guide&utm_medium=ebook&utm_campaign=msk-lbhip-recovery)
[utm_source=guide&utm_medium=ebook&utm_campaign=msk-lbhip-recovery](https://rootcauseunlocked.com/discovery?utm_source=guide&utm_medium=ebook&utm_campaign=msk-lbhip-recovery)***

Consultations and ongoing care are both available wherever you are located. Being outside North Carolina is not a barrier to working with me. Tell me where you are when you reach out and I will lay out exactly how we would work together from there.

***Get in touch: [rootcauseunlocked.com/contact?](https://rootcauseunlocked.com/contact?utm_source=guide&utm_medium=ebook&utm_campaign=msk-lbhip-recovery)
[utm_source=guide&utm_medium=ebook&utm_campaign=msk-lbhip-recovery](https://rootcauseunlocked.com/contact?utm_source=guide&utm_medium=ebook&utm_campaign=msk-lbhip-recovery)***

The companion ebooks go deeper on each condition in this region: *The Complete Patient's Guide to Low Back Pain*, *The Complete Patient's Guide to Sciatica*, and *The Complete Patient's Guide to Hip Pain*.

References

References 1 to 14 were verified against live PubMed records on 2026-08-09. References 15 to 23, added with the rebuilt supplement section, were verified against live PubMed records on 2026-09-04. Both passes included a publication type and corrections check for retractions and errata. On 2026-09-05 every reference was additionally screened for superseded editions, which is how reference 4 was moved to the 2026 Cochrane update; no other reference in this guide is superseded. That same screen surfaced a previously undisclosed correction on reference 23, noted there. Verification detail is documented in `Recovery-Guide-Citation-Ledger.md`.

1. Hayden JA, Ellis J, Ogilvie R, Malmivaara A, van Tulder MW. Exercise therapy for chronic low back pain. *Cochrane Database Syst Rev*. 2021;9(9):CD009790. PMID 34580864
2. Dahm KT, Brurberg KG, Jamtvedt G, Hagen KB. Advice to rest in bed versus advice to stay active for acute low-back pain and sciatica. *Cochrane Database Syst Rev*. 2010;2010(6):CD007612. PMID 20556780
3. French SD, Cameron M, Walker BF, Reggars JW, Esterman AJ. Superficial heat or cold for low back pain. *Cochrane Database Syst Rev*. 2006;2006(1):CD004750. PMID 16437495
4. Hall M, Lawford BJ, Hinman RS, Dobson F, Spiers L, Kimp A, French HP, Reichenbach S, Hernandez-Molina G, Bennell KL. Exercise for osteoarthritis of the hip. *Cochrane Database Syst Rev*. 2026;7:CD007912. PMID 42486497 (Current edition, 23 July 2026. It supersedes Fransen et al. 2014, PMID 24756895, which earlier drafts of this guide cited. Funding: none.)
5. Mellor R, Bennell K, Grimaldi A, et al. Education plus exercise versus corticosteroid injection use versus a wait and see approach on global outcome and pain from gluteal tendinopathy: prospective, single blinded, randomised clinical trial. *BMJ*. 2018;361:k1662. PMID 29720374
6. Peul WC, van Houwelingen HC, van den Hout WB, et al. Surgery versus prolonged conservative treatment for sciatica. *N Engl J Med*. 2007;356(22):2245-2256. PMID 17538084
7. Lequin MB, Verbaan D, Jacobs WC, et al. Surgery versus prolonged conservative treatment for sciatica: 5-year results of a randomised controlled trial. *BMJ Open*. 2013;3(5). PMID 23793663
8. Wiewelhove T, Döweling A, Schneider C, et al. A Meta-Analysis of the Effects of Foam Rolling on Performance and Recovery. *Front Physiol*. 2019;10:376. PMID 31024339
9. Ferreira RM, Silva R, Vigário P, et al. The Effects of Massage Guns on Performance and Recovery: A Systematic Review. *J Funct Morphol Kinesiol*. 2023;8(3). PMID 37754971
10. Wilkens P, Scheel IB, Grundnes O, Hellum C, Storheim K. Effect of glucosamine on pain-related disability in patients with chronic low back pain and degenerative lumbar osteoarthritis: a randomized controlled trial. *JAMA*. 2010;304(1):45-52. PMID 20606148
11. Zadro JR, Shirley D, Ferreira M, et al. Is Vitamin D Supplementation Effective for Low Back Pain? A Systematic Review and Meta-Analysis. *Pain Physician*. 2018;21(2):121-145. PMID 29565945
12. Liu X, Machado GC, Eyles JP, Ravi V, Hunter DJ. Dietary supplements for treating osteoarthritis: a systematic review and meta-analysis. *Br J Sports Med*. 2018;52(3):167-175. PMID 29018060
13. Yu G, Xiang W, Zhang T, Zeng L, Yang K, Li J. Effectiveness of Boswellia and Boswellia extract for osteoarthritis patients: a systematic review and meta-analysis. *BMC Complement Med Ther*. 2020;20(1):225. PMID 32680575
14. Lundh A, Lexchin J, Mintzes B, et al. Industry sponsorship and research outcome. *Cochrane Database Syst Rev*. 2017;2(2):MR000033. PMID 28207928

15. Shaw G, Lee-Barthel A, Ross ML, Wang B, Baar K. Vitamin C-enriched gelatin supplementation before intermittent activity augments collagen synthesis. *Am J Clin Nutr.* 2017;105(1):136-143. PMID 27852613
16. Lee J, Tang JCY, Dutton J, et al. The Collagen Synthesis Response to an Acute Bout of Resistance Exercise Is Greater when Ingesting 30 g Hydrolyzed Collagen Compared with 15 g and 0 g in Resistance-Trained Young Men. *J Nutr.* 2024;154(7):2076-2086. PMID 38007183
17. Balshaw TG, Funnell MP, McDermott EJ, et al. The Effect of Specific Bioactive Collagen Peptides on Tendon Remodeling during 15 wk of Lower Body Resistance Training. *Med Sci Sports Exerc.* 2023;55(11):2083-2095. PMID 37436929
18. Bischof K, Moitzi AM, Stafilidis S, Konig D. Impact of Collagen Peptide Supplementation in Combination with Long-Term Physical Training on Strength, Musculotendinous Remodeling, Functional Recovery, and Body Composition in Healthy Adults: A Systematic Review with Meta-analysis. *Sports Med.* 2024;54(11):2865-2888. PMID 39060741
19. Zdzieblik D, Oesser S, Gollhofer A, Konig D. Improvement of activity-related knee joint discomfort following supplementation of specific collagen peptides. *Appl Physiol Nutr Metab.* 2017;42(6):588-595. PMID 28177710 (Erratum: *Appl Physiol Nutr Metab.* 2017;42(11):1237. Corrected, not retracted. One author is affiliated with a collagen research institute, disclosed in the text.)
20. Nakagawa Y, Mukai S, Yamada S, et al. Short-term effects of highly-bioavailable curcumin for treating knee osteoarthritis: a randomized, double-blind, placebo-controlled prospective study. *J Orthop Sci.* 2014;19(6):933-939. PMID 25308211
21. Balius R, Alvarez G, Baro F, et al. A 3-Arm Randomized Trial for Achilles Tendinopathy: Eccentric Training, Eccentric Training Plus a Dietary Supplement Containing Mucopolysaccharides, or Passive Stretching Plus a Dietary Supplement Containing Mucopolysaccharides. *Curr Ther Res Clin Exp.* 2016;78:1-7. PMID 28053674 (Two authors are employed by the ingredient manufacturer, disclosed in the text.)
22. Tashiro T, Seino S, Sato T, Matsuoka R, Masuda Y, Fukui N. Oral administration of polymer hyaluronic acid alleviates symptoms of knee osteoarthritis: a double-blind, placebo-controlled study over a 12-month period. *ScientificWorldJournal.* 2012;2012:167928. PMID 23226979
23. Morton RW, Murphy KT, McKellar SR, et al. A systematic review, meta-analysis and meta-regression of the effect of protein supplementation on resistance training-induced gains in muscle mass and strength in healthy adults. *Br J Sports Med.* 2018;52(6):376-384. PMID 28698222 (Correction: *Br J Sports Med.* 2020;54(19):e7. PMID 32943392. The correction adds a competing interest, not a change to the data: one author served on the advisory board of a sports supplement manufacturer while the paper was written. Corrected, not retracted; the breakpoint used here stands. Disclosed in the text.)

Dr. Adam Seay, DC · Seay Wellness / Root Health, Garner, NC
rootcauseunlocked.com · (919) 662-0520

Educational content only. Not medical advice. Not a diagnosis. Not a substitute for examination. The emergency signs at the top of this guide outrank everything else in it. Talk with your clinician before starting new exercise or supplements, particularly if you take blood thinners or have osteoporosis, and respect the pressure ceilings above, which differ for manual tools and powered devices.