



ROOT CAUSE
UNLOCKED

DRAFT · CLINICAL REVIEW COPY

ROOT CAUSE UNLOCKED · ULTIMATE GUIDE

Unlock Your *Reflux*, Ultimate Guide

A Comprehensive Guide to Healing

Dr. Adam Seay, DC

Root Cause Unlocked, Garner, NC · rootcauseunlocked.com · (919) 662-0520

Unlock Your Reflux, Ultimate Guide

A Comprehensive Guide to Healing

DRAFT · CLINICAL REVIEW COPY

This edition is pending Dr. Seay's clinical sign-off. Every citation has been verified against its live PubMed record, including a check for retractions. The companion citation ledger lists each source and the claims that were investigated but left uncited.

BEFORE YOU START

Contents

Introduction

Chapter 1: The Misunderstood Stomach: Unraveling the Role of Stomach Acid

Chapter 2: The Hidden Invader: H. pylori and Its Role in Reflux

Chapter 3: When Food Fights Back: Sensitivities & Allergies

Chapter 4: The Mind-Gut Connection: Vagus Nerve Dysfunction & Poor Parasympathetic Tone

Chapter 5: The Overgrown Garden: SIBO (Small Intestinal Bacterial Overgrowth) and Histamine Intolerance

Chapter 6: The Foundation: The Impact of Poor Posture

Chapter 7: Your Personalized Action Plan: Bringing It All Together

Conclusion

Working with Root Health

Appendix and Resources

References

From Understanding to Doing

Read this part first

The honest starting point, and the five drivers

The complication you should understand before anything else

Driver 1: the barrier, which is mechanical and has no blood test

Driver 2: the signal, where the nervous system sets the volume

Helicobacter pylori

Iron studies and a complete blood count

Vitamin B12 and magnesium, the two the medication takes

Eosinophilic esophagitis

Celiac disease

Cardiac disease

Stomach acid, and the honest state of that question

The esophageal and gastric microbiome

Small intestinal bacterial overgrowth

The rest of the terrain

Functional ranges for the markers in this guide

Reading your results as a pattern

The conversation with your doctor

What these labs cannot tell you

When to seek clinical review

Your next step

References

Read this part first

Where food actually sits in the reflux picture

Weight

Timing and position

Smoking, with the honest nuance

The two-week test, run properly

What is worth knowing about the usual suspects

Part two and a half: how you eat, which is the vagus nerve at the table

A plant-forward Mediterranean pattern

Fiber, and the reason it belongs here

The histamine question, briefly

Bacterial overgrowth, if bloating dominates

A week that supports the plan

What I deliberately left out, and why

When to seek professional care

Working with Root Health

References

Read this part first

Why nutrient evidence looks thinner than it is

Where supplements actually fit in reflux

How to read this guide

Foundation 1: retrain the diaphragm, because it is half of your sphincter

Foundation 2: work the nervous system, which sets the volume

Alginate

Melatonin

Iberogast and similar multi-herb formulas

Curcumin

Zinc L-carnosine

Ginger

D-limonene

Probiotics, mainly as an adjunct to H. pylori treatment

Betaine HCl and the low stomach acid question

Deglycyrrhizinated licorice

Coming off a PPI, which is what most readers actually came here for

What I deliberately left out, and why

Realistic expectations

When to seek professional care

Where to find quality versions of these

Working with Root Health

References

Introduction

Maybe you dread mealtime, fearing the inevitable burn and discomfort that follows. Maybe you're constantly reaching for antacids, only to find temporary relief that never truly solves the problem. You're not alone. Millions of people worldwide suffer from reflux, a condition often mistakenly labeled as simply "heartburn" or "acid indigestion." And the conventional wisdom about reflux is often incomplete, or even outright wrong.

For too long, the standard approach to reflux has been to suppress stomach acid and stop there. The reasoning is that too much acid is the culprit, so medications like proton pump inhibitors (PPIs) shut acid production down. Those drugs work, and for people with erosive disease they are genuinely important. What they do not do is address the reasons you started refluxing. And here is the number that should reframe the whole conversation: up to half of people remain symptomatic despite adequate acid suppression. That is the argument, in one statistic, that reflux is multifactorial rather than purely an acid problem.

This ebook is a guide to the rest of that picture. I dig into the drivers that go unexamined when the answer stops at a prescription. Think of it as a detective mission, uncovering the keys that explain why your particular valve is failing.

What you are buying here, stated plainly. The two interventions with the strongest evidence in this entire book are losing weight if you carry extra, and raising the head of your bed. Neither costs anything and neither is exciting. Some of what follows sits on good trial evidence, particularly alginate, the deprescribing sequence, and the H. pylori work. Some of it sits on mechanism and clinical use rather than on outcome trials, and I have marked which is which as I go, chapter by chapter, rather than letting you assume it is all the same grade. Where a claim is contested, I say so, including the one this book is best known for.

I explore factors like:

- Stomach acid, and the low-acid claim, which needs handling more carefully than most reflux content gives it.
- The role of H. pylori infection.
- Food sensitivities and inflammation, and which famous trigger foods do not hold up.
- The impact of your vagus nerve and nervous system.
- SIBO (Small Intestinal Bacterial Overgrowth) and histamine intolerance.
- Something as fundamental as your posture and your diaphragm.

By the end you will have an understanding of what may be going on and a roadmap for investigating it in a sensible order. This is not about masking symptoms. It is about finding which of the five drivers is yours.

Chapter 1: The Misunderstood Stomach: Unraveling the Role of Stomach Acid

When you experience that burning sensation in your chest, the first explanation that comes to mind is usually “too much stomach acid.” That belief has led to a widespread approach of suppressing acid production, and for a great many people that strategy still leaves them symptomatic. So a second explanation circulates widely in functional medicine: that the real problem is the opposite, low stomach acid, a condition known as hypochlorhydria.

I am going to handle that claim carefully, because it is the single most repeated idea in reflux content and it is not as settled as it is usually presented.

The low stomach acid claim, and what actually stands behind it

Here is the honest position, stated before the mechanism, so nothing that follows misleads you.

Low stomach acid is a real, defined, testable condition. Where it has been demonstrated, replacing acid has a coherent rationale, and the strongest published case is made for **autoimmune gastritis**, in which the parietal cells that make acid are destroyed. In that population the argument is that B12 and iron get replaced while the missing acid does not, and that these patients are sometimes prescribed acid suppression for a stomach that is not producing acid.

What that supports and what it does not. It supports acid replacement in someone with demonstrated low acid production. **It is not evidence that most people with reflux have low stomach acid, and that broader claim does not have trial support behind it.** The lab guide in this kit lays out the workup, and the supplement guide lays out where betaine HCl sits and why it is not a first move.

So read this chapter as what it is: a mechanistically coherent account of why acid matters, offered as a hypothesis worth testing in your case rather than as the settled explanation for your symptoms. If you take one instruction from it, take this one. Demonstrate low acid before you treat it. Adding acid to an ulcer, to erosive esophagitis, or to active gastritis can hurt you, and those conditions produce exactly the symptoms that send people looking for betaine HCl in the first place.

Why stomach acid matters regardless of who has too little

Our stomach is an incredible organ, and stomach acid, or hydrochloric acid (HCl), is its mighty powerhouse. When you eat, your stomach should become highly acidic, with a pH between 1.5 and 3.5. This extremely acidic environment isn't just for show; it's absolutely essential for several key digestive and protective functions:

- **Breaking Down Food:** HCl helps to break down proteins into smaller, more manageable peptides, preparing them for further digestion in the small intestine. Without sufficient acid, food sits in your stomach longer, fermenting and creating gas.
- **Nutrient Absorption:** Stomach acid is critical for absorbing essential nutrients like Vitamin B12, iron, calcium, magnesium, and zinc.¹ If your acid is low, you might be eating a nutrient-rich diet but still be deficient in these nutrients.
- **Killing Pathogens:** Your stomach acid acts as your body's first line of defense against harmful bacteria, viruses, parasites, and fungi that hitch a ride on your food. It sterilizes your meals before they enter your delicate small intestine.
- **Triggering Digestive Cascade:** Adequate stomach acid signals to your pancreas and gallbladder to release digestive enzymes and bile. If this initial signal is weak, the entire digestive process downstream suffers.
- **Maintaining the Lower Esophageal Sphincter (LES):** the LES is a muscular valve at the bottom of your esophagus that opens to let food into your stomach and then closes to prevent stomach contents from flowing back up. **A proposed mechanism**, and this is the part of the low-acid model that carries the most weight in functional medicine, is that the LES tightens in response to a properly acidified stomach, so that when acid is low the valve does not close as it should. Note that this is a proposed mechanism rather than a demonstrated cause of reflux in the general population, and it is still argued in the literature. What is not argued is that the valve is only half the barrier. The diaphragm is the other half, which is why the posture and breathing work in Chapter 6 is not a wellness postscript.

So, how can you tell if low stomach acid might be your issue? The symptoms overlap heavily with the “too much acid” picture, which is exactly why guessing does not work here. Signs that raise the question include:

- Bloating, belching, and gas shortly after eating.
- Feeling full for a long time after meals.
- Indigestion or a sensation of food “sitting” in your stomach.
- Heartburn/reflux (as explained above).
- Undigested food in your stool.
- Nutrient deficiencies (e.g., iron deficiency anemia, B12 deficiency).
- Brittle nails, hair loss, and fatigue.

Causes of Low Stomach Acid

Hypochlorhydria isn't just bad luck; it often stems from identifiable factors:

- **Aging:** As we get older, our natural production of stomach acid tends to decline.
- **Chronic Stress:** When you're constantly in “fight or flight” mode, your body prioritizes survival functions over digestion. This diverts blood flow away from your gut and suppresses acid production.

- **Nutrient Deficiencies:** Key nutrients like zinc and B vitamins are essential for HCl production. Deficiencies can impair your stomach's ability to produce acid.
- **H. pylori Infection:** This common bacterial infection, which we'll discuss in detail in Chapter 2, is a major cause of low stomach acid. H. pylori neutralizes stomach acid to survive, leading to reduced production.²
- **Medications:** Ironically, the very medications often prescribed for reflux, like proton pump inhibitors (PPIs) and antacids, are designed to reduce stomach acid. While helpful in acute situations, long-term use can perpetuate and worsen hypochlorhydria, creating a vicious cycle.³

Testing for Low Stomach Acid

While a definitive diagnosis of hypochlorhydria often requires clinical testing, there are a few ways to get an indication:

- **The Baking Soda Test (At-Home, Informal):** mix 1/4 teaspoon of baking soda in 4 to 6 ounces of cold water first thing in the morning before eating or drinking anything else, and drink it quickly. The idea is that adequate acid reacts with the bicarbonate to produce carbon dioxide and you belch within two to three minutes. **This is not a validated test**, it has never been shown to correlate with measured gastric acid output, and I include it because you will encounter it, not because I want you making decisions on it.
- **Heidelberg pH Test (Clinical):** a clinical test in which you swallow a capsule that transmits pH readings from your stomach, measuring the ability to acidify and rebound after a challenge. This is the closest thing to a direct measurement, and availability is limited.
- **Blood work for autoimmune gastritis:** parietal cell and intrinsic factor antibodies, gastrin, and pepsinogen I and II with the ratio. **This is the workup I would actually run** if low acid is genuinely on the table, and it is covered in the lab guide in this kit.
- **Trial of Betaine HCl (Under Guidance):** taking betaine HCl with a protein-heavy meal and observing symptoms. The at-home version, where you increase capsules until you feel warmth, is widely described and **is not a validated test**. It can be informative in practice and it should not be treated as a diagnosis, particularly when the same symptoms would be produced by an ulcer. More below.

Dietary Support for Stomach Acid Production

Fortunately, there are simple dietary strategies that can help stimulate your stomach's natural acid production:

- **Bitters:** Think of traditional aperitifs! Bitter foods and herbal tinctures stimulate digestive juices, including stomach acid. Incorporate foods like artichoke, dandelion greens, radicchio, arugula, and bitter melon. You can also find herbal bitter tinctures. Take them about 15 minutes before meals.
- **Apple Cider Vinegar (ACV):** While not providing actual HCl, ACV is acidic and can help lower stomach pH slightly, signaling to your body to produce more of its own acid. Mix 1-2 teaspoons of raw, unfiltered ACV

in a small glass of water and drink 10-15 minutes before meals. Start with a small amount and increase gradually if tolerated.

- **Ginger:** This versatile root is a powerful digestive aid. Fresh ginger tea before or during meals can stimulate stomach acid, calm digestive upset, and even act as a natural prokinetic (helping food move through the digestive tract).

Supplements to Support Stomach Acid (Backed by Research)

For those with confirmed or suspected low stomach acid, certain supplements can provide targeted support.

- **Betaine HCl:** This is a supplement that provides a supplemental source of hydrochloric acid.
- **How it Works:** By adding HCl to your stomach, Betaine HCl helps to lower the stomach's pH to the optimal acidic range. This aids in protein digestion, nutrient absorption, and proper closure of the LES.
- **Proper Dosing:** Dosing Betaine HCl is highly individualized and should ideally be done under the guidance of a healthcare professional. A common approach is to start with one capsule (e.g., 650 mg) with a protein-heavy meal. If you don't feel any burning or warmth, increase by one capsule with each subsequent protein-heavy meal until you feel a slight warmth or burning sensation. At that point, reduce by one capsule, and that's your optimal dose for that meal size.
- **Contraindications:** Do not take Betaine HCl if you have a history of stomach ulcers, gastritis, or are currently taking NSAIDs (like ibuprofen or aspirin), as it could irritate the stomach lining. Always consult your doctor before starting Betaine HCl.
- **Research Note,** stated straight: **betaine HCl does not have trial evidence supporting its use across reflux generally, and I am not going to pretend otherwise in a document you paid for.** Large outcome trials cost tens to hundreds of millions of dollars and that money only exists where a patent can pay it back, so nobody has funded one on an unpatentable compound. That explains the absence; it does not fill it. Where it has a coherent rationale is in someone with demonstrated hypochlorhydria, which is why the workup comes first. That is what I do clinically: demonstrate the deficit, then decide with the patient, then hold it to a defined endpoint rather than running it forever.
- **Digestive Enzymes:** Even with improved stomach acid, sometimes your body might need extra help breaking down fats, carbohydrates, and proteins. Broad-spectrum digestive enzyme supplements contain enzymes like amylase, lipase, and protease, which assist in the breakdown of all macronutrients. Taking these with meals can reduce the digestive burden, especially if food isn't being properly broken down in the stomach. Look for enzyme blends that include HCl or are specifically designed for low stomach acid support.

To close this chapter where I opened it: acid matters, low acid is real and testable, and the claim that most reflux is caused by too little acid is not something the evidence supports. Test before you treat, and do not add acid to a stomach you have not investigated. There is another driver worth ruling out first anyway, because it is common, cheap to test, and genuinely consequential: a bacterial infection.

Chapter 2: The Hidden Invader: H. pylori and Its Role in Reflux

Imagine a tiny, spiral-shaped bacterium that can live comfortably in the acidic environment of your stomach, often for decades, without you even knowing it's there. This unassuming microbe, *Helicobacter pylori*, or simply H. pylori, is far more common than many realize, infecting roughly half the world's population. While often asymptomatic, H. pylori can be a silent saboteur, quietly undermining your digestive health and playing a significant, yet frequently overlooked, role in chronic reflux.

Understanding H. pylori: A Common Bacterial Infection with Big Consequences

H. pylori is unique because it's one of the few bacteria that can survive the harsh, acidic conditions of the stomach. It does this by producing an enzyme called urease, which converts urea into ammonia and carbon dioxide. Ammonia acts as a shield, neutralizing the stomach acid around the bacterium, allowing it to burrow into the protective mucus lining of your stomach.

This colonization leads to several problems that directly contribute to reflux:

- **Reduced Stomach Acid Production:** Over time, H. pylori colonization often leads to chronic inflammation (gastritis) and can directly impair the parietal cells in your stomach that are responsible for producing hydrochloric acid. This further exacerbates the problem of hypochlorhydria (low stomach acid) that we discussed in Chapter 1. As we learned, low stomach acid means a relaxed LES, leading to reflux.
- **Inflammation and Damage:** The constant presence of H. pylori causes chronic inflammation of the stomach lining. This inflammation can not only lead to discomfort but also increases the risk of more serious conditions.
- **Link to Gastritis, Ulcers, and Stomach Cancer:** While many people with H. pylori never develop severe symptoms, it is the primary cause of chronic gastritis, a major cause of peptic ulcers (sores in the lining of the stomach or duodenum), and a significant risk factor for certain types of stomach cancer.⁴ This underscores the importance of identifying and addressing this infection.

It's important to recognize that H. pylori doesn't cause symptoms in everyone, and some individuals may carry the bacterium without ever experiencing problems. However, if you are struggling with chronic reflux, it is a key factor to investigate.

Identifying H. pylori

Since H. pylori can often be silent, specific testing is necessary to confirm its presence. There are several reliable methods:

- **Urea Breath Test:** This is a common and non-invasive test. You drink a special liquid containing a harmless substance (urea). If *H. pylori* is present, its urease enzyme will break down the urea, releasing carbon dioxide that can be detected in your breath. This test is highly accurate for active infection.
- **Stool Antigen Test:** This test detects specific *H. pylori* proteins (antigens) in your stool. It's also highly accurate and non-invasive, excellent for both initial diagnosis and confirming eradication after treatment.
- **Endoscopy with Biopsy:** This is a more invasive procedure performed by a gastroenterologist. A thin, flexible tube with a camera is inserted down your throat into your stomach and duodenum. Biopsies (small tissue samples) can be taken and examined under a microscope to detect the presence of the bacteria and assess any inflammation or damage. This is often done if there are other concerns, such as ulcers or suspicious lesions.
- **Blood Test (Antibody Test):** Blood tests can detect antibodies to *H. pylori*, indicating exposure at some point. However, they cannot differentiate between a current active infection and a past, resolved infection. Therefore, breath or stool tests are generally preferred for diagnosing active infection.

If you have chronic reflux symptoms and haven't been tested for *H. pylori*, discuss these testing options with your healthcare provider.

Strategies for Removal (Integrative Approach)

Once *H. pylori* is identified, eradication is key to long-term healing. The approach to removal can involve conventional medicine, natural remedies, or often, a beneficial combination of both.

- **Conventional Treatment:** standard eradication is a prescription antibiotic regimen, “triple therapy” or “quadruple therapy,” combining two or three antibiotics with a proton pump inhibitor, usually for 10 to 14 days. These are prescription drugs that some physicians use for confirmed *H. pylori* infection. They are not something I prescribe, and they are included here so you know they exist and can raise them with the physician managing that part of your care. I will say plainly that this is the one place in this book where I think the drug route deserves first consideration, because eradication has real outcome evidence behind it and the alternatives do not. Antibiotics do disrupt the microbiome, which is what the probiotic support below is for. Follow your prescriber's instructions precisely, and get a confirmatory test afterward.
- **Natural Support (Backed by Research):** For those seeking a less aggressive approach or looking to support conventional treatment and prevent recurrence, several natural compounds have demonstrated anti-*H. pylori* activity in scientific research. These should be considered under the guidance of a healthcare professional.
- **Mastic Gum:** Derived from the mastic tree, this resin has been used for centuries for digestive complaints.
 - **Research Note:** Numerous studies have shown mastic gum's ability to inhibit and kill *H. pylori*. For example, research published in the *New England Journal of Medicine* in 1998 demonstrated its bactericidal effect.⁵ Subsequent studies have confirmed its efficacy, often showing it can reduce *H. pylori* load and improve symptoms.

- **Manuka Honey:** This unique honey from New Zealand is renowned for its potent antimicrobial properties, attributed to its high concentration of Methylglyoxal (MGO).
 - **Research Note:** Studies have indicated that Manuka honey can inhibit the growth of *H. pylori* and may enhance the effects of antibiotics, making it a valuable adjunct.⁶ Its soothing properties can also benefit the stomach lining.
- **Broccoli Sprouts (Sulforaphane):** Broccoli sprouts are a concentrated source of sulforaphane, a potent compound with significant anti-cancer and anti-bacterial properties.
 - **Research Note:** Research, including human trials, has shown that sulforaphane-rich broccoli sprouts can reduce *H. pylori* colonization and inflammation in the stomach.⁷ They are also powerful antioxidants.
- **Bismuth Subcitrate:** While often considered a conventional treatment, bismuth compounds (like Pepto-Bismol) can be part of natural protocols. Bismuth has direct antimicrobial effects against *H. pylori* and also forms a protective barrier over the stomach lining.
- **Probiotics:** Supporting your gut microbiome is paramount, especially when tackling *H. pylori* (whether with antibiotics or natural remedies).
 - **Research Note:** Specific probiotic strains, particularly *Saccharomyces boulardii* and certain *Lactobacillus* and *Bifidobacterium* species, have been shown to inhibit *H. pylori* growth, reduce side effects from antibiotics, and aid in restoring gut balance.⁸ *S. boulardii* is especially notable for its ability to compete with *H. pylori* and its resistance to antibiotics.
- **Deglycyrrhized Licorice (DGL):** DGL does not directly kill *H. pylori*, but it helps to protect and heal the stomach lining, which is often irritated by the infection. It can enhance mucus production and promote healthy tissue repair.
- **Dietary Considerations during Treatment:** During any *H. pylori* eradication protocol, focus on a whole-foods diet rich in nutrient-dense, easily digestible foods. Avoid processed foods, excessive sugar, alcohol, and anything that triggers your reflux symptoms. Lean proteins, cooked vegetables, and healthy fats should form the cornerstone of your meals.

One honest caveat on all of the natural agents above. Their evidence is mostly laboratory and animal work plus small human studies, and none of them has been shown to match a full eradication regimen. I use several of them, mostly as adjuncts alongside eradication and for the recovery period afterward, and that is what I am recommending them as. Testing for *H. pylori* is cheap and the finding is actionable, which is why it sits this early in the book. Food itself, though, can sometimes be the problem, even when *H. pylori* is gone.

Chapter 3: When Food Fights Back: Sensitivities & Allergies

You've worked on optimizing stomach acid and explored the potential of *H. pylori* as a culprit. But sometimes, despite these efforts, certain foods still seem to trigger your reflux, or leave you feeling generally unwell. This is where the often-complex world of food sensitivities and allergies comes into play. It's not always the obvious "acidic" or "spicy" foods; sometimes, it's seemingly healthy foods that are silently creating inflammation and digestive distress, contributing to your reflux.

Differentiating Between Allergies, Sensitivities, and Intolerances

Before we go further, it's important to understand the distinctions between these terms:

- **Food Allergy (IgE-mediated):** This is an immediate, often severe, immune response to a food protein, mediated by IgE antibodies. Symptoms typically appear rapidly (minutes to a few hours) and can range from hives, swelling, and difficulty breathing to anaphylaxis, a life-threatening reaction. Common allergens include peanuts, tree nuts, milk, eggs, soy, wheat, fish, and shellfish. True food allergies are serious and require strict avoidance.
- **Food Intolerance (Non-immune mediated):** This is a digestive issue, not an immune response. It occurs when your body lacks the enzymes needed to break down a certain food component. The most common example is lactose intolerance, where there's insufficient lactase enzyme to digest the sugar in dairy. Symptoms are typically digestive (gas, bloating, diarrhea) and depend on the amount consumed.
- **Food Sensitivity (Non-IgE Immune-mediated):** This is a delayed, often dose-dependent, immune response that involves different immune pathways (often IgG antibodies, though other mechanisms exist). Symptoms can be diverse, affecting almost any system in the body, and may not appear for hours or even days after consuming the offending food. This delayed reaction makes them incredibly difficult to pinpoint without a systematic approach. Food sensitivities are a major contributor to chronic inflammation and can profoundly impact gut health, making them highly relevant to reflux.

For the purpose of addressing chronic reflux, our primary focus will be on identifying and managing food sensitivities, as they are frequently hidden triggers of inflammation and digestive dysfunction that lead to reflux symptoms.

Common Food Triggers for Reflux

While we're aiming to go beyond the obvious, it's worth acknowledging the widely recognized food triggers that can exacerbate reflux symptoms, often by irritating an already compromised esophagus or stomach, or by relaxing the LES:

- **Acidic Foods:** Citrus fruits (oranges, lemons, grapefruit), tomatoes, tomato products (sauce, paste), vinegar. While these are often blamed, they are more likely secondary irritants if your primary problem is a

weak LES due to low stomach acid or other issues.

- **Spicy Foods:** Chili peppers, hot sauces, and other spicy ingredients can directly irritate the esophageal lining.
- **High-Fat Foods:** Fried foods, greasy meals, high-fat dairy, and large portions of fatty meats. Fat slows down stomach emptying and can relax the LES, allowing contents to reflux.
- **Peppermint and Chocolate:** These contain compounds that can relax the LES.
- **Onions and Garlic:** For many, these can cause bloating and gas, increasing abdominal pressure, which can push stomach contents upwards. They can also be high in FODMAPs (fermentable carbohydrates), linking to SIBO (which we'll cover in Chapter 5).
- **Caffeine:** Coffee, tea, and energy drinks can relax the LES.
- **Alcohol:** Can relax the LES and irritate the esophageal lining.
- **Carbonated Beverages:** The gas from these drinks can increase stomach pressure and cause belching, leading to reflux.

While avoiding these foods can provide relief, it's like putting a band-aid on a deeper wound. True healing often requires looking beneath the surface for less obvious culprits.

Beyond the Obvious: Identifying Deeper Sensitivities

For chronic reflux sufferers, the key often lies in identifying IgG food sensitivities. When you consume a food you're sensitive to, your immune system may mount a low-grade, delayed inflammatory response. This chronic inflammation can:

- **Damage the Gut Lining:** Leading to "leaky gut" or increased intestinal permeability, where larger food particles and toxins can pass through the gut barrier and trigger systemic inflammation.
- **Impair Digestion:** Inflamed digestive tissues don't function optimally.
- **Increase Histamine Release:** A connection we'll explore further in Chapter 5.
- **Contribute to Motility Issues:** Affecting how quickly food moves through your digestive tract.
- **Directly Irritate the Esophagus:** Leading to reflux symptoms even without significant acid.

Common culprits for IgG food sensitivities that drive chronic inflammation and digestive issues include:

- **Gluten:** The protein found in wheat, barley, rye, and some oats. Gluten sensitivity is increasingly recognized as a widespread issue, contributing to a vast array of symptoms beyond celiac disease.
- **Dairy:** Specifically casein (protein) and lactose (sugar). Many people struggle to digest dairy due to enzyme deficiencies or immune reactions to its proteins.
- **Soy:** A common allergen and sensitivity trigger, particularly in processed foods.
- **Corn:** Another pervasive ingredient in processed foods, often a hidden sensitivity.

- Eggs: Both the yolk and white can be problematic for some.

Testing for Food Sensitivities

Identifying food sensitivities can be challenging due to their delayed and varied symptoms. There are two primary approaches:

- The Elimination Diet: the most useful method available to you, because it observes your body's response directly rather than inferring it from a marker. It is not a validated diagnostic and I would not call it a gold standard, but nothing else on this list beats it.
- How it works: For a period (typically 3-4 weeks), you completely eliminate common culprits and any foods you suspect might be triggers. This includes gluten, dairy, soy, corn, eggs, peanuts, and often other inflammatory foods like sugar, alcohol, and processed items. The goal is to allow your immune system to calm down and inflammation to subside.
- Reintroduction Phase: After the elimination period, you systematically reintroduce foods, one at a time, every 2-3 days, while carefully monitoring your symptoms. Reintroduce a small amount of the food and wait. If no symptoms appear, try a slightly larger amount. If symptoms (reflux, bloating, headaches, fatigue, skin issues, joint pain, etc.) return or worsen, that food is likely a trigger for you. Keep a detailed food and symptom journal throughout this process.
- Pros: Highly accurate for identifying your unique triggers. No expensive lab tests.
- Cons: Requires discipline, patience, and careful planning. Can be socially challenging.
- IgG Antibody Testing: These blood tests measure your immune system's IgG antibody response to various foods.
- How it works: A blood sample is sent to a lab, which tests it against a panel of common foods. The results typically show a numerical score or a scale (e.g., low, moderate, high) indicating your IgG reactivity to each food.
- Pros: Provides a list of potential triggers, which can be a helpful starting point, especially if an elimination diet feels overwhelming.
- Cons: IgG reactions indicate exposure and an immune response, but not necessarily a symptomatic one. Some debate exists in the scientific community about their sole diagnostic value, as a high IgG level might simply mean you eat that food frequently. They should always be interpreted in conjunction with symptoms and ideally followed up with an elimination and reintroduction phase.
- Recommendation: If you choose IgG testing, use the results as a guide for your elimination diet, rather than relying solely on them for dietary decisions.

Creating a Personalized Reflux-Friendly Diet Plan

Once you've identified your individual triggers, you can build a sustainable, reflux-friendly diet.

- Phase 1: Elimination (Initial 3-4 Weeks):
- Focus: Remove all identified food sensitivities, common inflammatory foods, and typical reflux triggers (as listed earlier).
- Emphasize: Whole, unprocessed foods.
 - Lean Proteins: Grass-fed meats, poultry, wild-caught fish, eggs (if tolerated).
 - Abundant Vegetables: Non-starchy vegetables, both cooked and raw (depending on tolerance). Focus on variety. Avoid high-FODMAP vegetables if suspecting SIBO.
 - Healthy Fats: Avocado, olive oil, coconut oil, ghee (if dairy-free).
 - Complex Carbohydrates (in moderation): Gluten-free grains (quinoa, brown rice, millet, buckwheat, if tolerated), sweet potatoes, squash.
 - Fruits (in moderation): Lower-acid fruits like berries, melon, pears, bananas.
- Hydration: Drink plenty of filtered water throughout the day, between meals rather than with them to avoid diluting stomach acid. Herbal teas (chamomile, ginger, slippery elm) can be soothing.
- Phase 2: Reintroduction & Sustainable Eating:
- Carefully reintroduce eliminated foods one by one, observing symptoms.
- Once your triggers are identified, your long-term diet becomes one that avoids these specific problematic foods, while emphasizing nutrient density and gut healing.
- Key Principles for Ongoing Reflux Management:
 - Whole, Unprocessed Foods: Minimize packaged foods, artificial ingredients, and added sugars.
 - Mindful Eating: Eat slowly, chew your food thoroughly (at least 20-30 times per bite), and avoid eating on the go or when stressed. This allows your digestive system to prepare and function optimally.
 - Portion Control: Don't overeat. Large meals put pressure on the LES.
 - Timing of Meals: Avoid eating within 2-3 hours of bedtime to allow for proper digestion before lying down.
 - Hydration: Continue to drink plenty of water, but not large amounts with meals.
 - Listen to Your Body: Pay attention to how different foods make you feel. Your body provides valuable clues.

Identifying and eliminating food sensitivities can dramatically reduce inflammation, ease digestive burden, and significantly improve reflux symptoms. It requires detective work and patience, but the relief and improved well-being are often well worth the effort. However, even the most perfect diet can be undermined if the communication highway between your brain and gut is dysfunctional.

Chapter 4: The Mind–Gut Connection: Vagus Nerve Dysfunction & Poor Parasympathetic Tone

We've explored the importance of stomach acid and the impact of unseen infections and food triggers. Now, let's turn our attention to an often-overlooked, yet profoundly influential, player in your digestive health and reflux: the vagus nerve. This remarkable nerve acts as the primary communication highway between your brain and your gut, profoundly influencing everything from stomach acid production to gut motility. When it's not functioning optimally, your entire digestive system, including your esophagus and LES, can suffer, perpetuating reflux.

Understanding the Vagus Nerve: The Highway of Communication

The vagus nerve is the longest cranial nerve in your body, extending from your brainstem down through your neck, chest, and into your abdomen, innervating your heart, lungs, and most of your digestive organs. It's a key component of your autonomic nervous system, which controls involuntary bodily functions. Specifically, the vagus nerve is the main nerve of the parasympathetic nervous system, often referred to as the “rest and digest” system.

When the vagus nerve is properly activated and your parasympathetic system is dominant, your body is in a state conducive to healing and digestion. This includes:

- **Stimulating Stomach Acid Production:** The vagus nerve signals the stomach to produce HCl and digestive enzymes.
- **Promoting Bile and Pancreatic Enzyme Release:** It tells the gallbladder to release bile and the pancreas to release enzymes, essential for breaking down fats, proteins, and carbohydrates.
- **Regulating Gut Motility:** It helps to keep food moving efficiently through your digestive tract, preventing stagnation and bacterial overgrowth.
- **Modulating Inflammation:** The vagus nerve has anti-inflammatory effects throughout the body, including the gut.
- **Strengthening the LES:** A properly functioning vagus nerve helps maintain the tone and strength of the Lower Esophageal Sphincter (LES), preventing reflux.
- **Reducing Stress and Anxiety:** Its activation promotes a sense of calm and relaxation, which directly benefits digestion.

How Stress Impairs Vagus Nerve Function

In modern life, many of us spend far too much time in the “fight or flight” mode, governed by the sympathetic nervous system. This is our body's survival mechanism, designed for short bursts of acute stress. However,

when stress becomes chronic, it keeps the sympathetic nervous system on overdrive, effectively sidelining the vagus nerve and the parasympathetic “rest and digest” system.

When your body perceives stress (whether it’s a real threat or just a looming deadline), it diverts resources away from “non-essential” functions like digestion. This means:

- **Reduced Blood Flow to the Gut:** Less blood means less oxygen and nutrients for digestive organs.
- **Suppressed Stomach Acid Production:** Your stomach doesn’t get the signal to produce adequate acid.
- **Impaired Enzyme Release:** Pancreatic and bile flow are reduced.
- **Altered Gut Motility:** Leading to either too fast (diarrhea) or too slow (constipation) movement.
- **Increased Gut Permeability (“Leaky Gut”):** Chronic stress can directly contribute to the breakdown of the gut lining.
- **Relaxation of the LES:** When the nervous system is in a constant state of alarm, the LES can lose its tone, making it prone to opening inappropriately.

This constant state of sympathetic dominance creates a vicious cycle where stress worsens digestion, and poor digestion, in turn, can contribute to stress and anxiety, ultimately fueling your reflux symptoms.

Symptoms of Vagus Nerve Dysfunction

Vagus nerve dysfunction can manifest in a wide array of symptoms beyond just reflux, offering clues that this nerve might need attention:

- **Digestive Issues:** Chronic reflux, bloating, constipation, diarrhea, slow digestion, feeling full quickly.
- **Mood & Mental Health:** Anxiety, depression, poor stress resilience.
- **Cardiovascular:** Low heart rate variability (HRV), lightheadedness upon standing (orthostatic intolerance).
- **Inflammation:** Chronic pain, widespread inflammation.
- **Immune System:** Frequent infections, slow wound healing.
- **Other:** Difficulty swallowing, chronic fatigue, gag reflex issues.

Lifestyle Modifications to Activate the Vagus Nerve

The good news is that you can actively train and strengthen your vagus nerve. By consciously engaging in practices that stimulate parasympathetic activity, you can significantly improve your digestive function and reduce reflux. These are simple yet powerful tools:

- **Deep Diaphragmatic Breathing (Belly Breathing):** This is one of the most effective ways to stimulate the vagus nerve.⁹

- **How to practice:** Lie down or sit comfortably. Place one hand on your chest and the other on your belly. Inhale slowly and deeply through your nose, allowing your belly to rise (your chest should remain relatively still). Exhale slowly through pursed lips, feeling your belly fall. Make your exhale longer than your inhale. Aim for 5-10 minutes, several times a day, especially before meals.
- **Gargling, Humming, and Singing:** The vagus nerve passes through your vocal cords and the back of your throat.
- **How to practice:** Vigorously gargle with water for 30-60 seconds, several times a day. Hum a tune or sing loudly (even in the shower!) for a few minutes daily. These actions stimulate the vagus nerve.
- **Cold Exposure:** Short, acute exposure to cold can “shock” the vagus nerve into action.
- **How to practice:** Finish your shower with 30-60 seconds of cold water. Splash cold water on your face. This activates a “diving reflex” that stimulates the vagus nerve. Start slowly and gradually increase duration.
- **Mindfulness & Meditation:** Regular meditation practices have been shown to increase vagal tone.
- **How to practice:** Spend 10-20 minutes daily focusing on your breath, observing your thoughts without judgment, or using guided meditations. Apps like Calm or Headspace can be very helpful.
- **Yoga and Tai Chi:** These practices combine mindful movement, deep breathing, and meditation, making them excellent for vagal nerve stimulation and stress reduction.
- **Gentle Exercise:** Regular, moderate physical activity helps regulate the nervous system. Avoid overtraining, which can be a stressor.
- **Social Connection and Laughter:** Positive social interactions and genuine laughter release endorphins and can shift your nervous system into a more relaxed state.
- **Massage:** Gentle massage, particularly around the neck and feet, can stimulate vagal nerve points.
- **Acupuncture:** Can be effective in modulating vagal nerve activity and reducing stress.

Supplements to Support Nervous System Health

While lifestyle changes are primary, certain supplements can offer supportive roles for nervous system balance:

- **Magnesium:** Often called “nature’s tranquilizer,” magnesium is essential for over 300 biochemical reactions in the body, including nerve and muscle function. Many people are deficient. It helps to relax muscles, including those in the digestive tract, and supports a calm nervous system. Look for highly absorbable forms like magnesium glycinate or citrate.
- **L-Theanine:** An amino acid found primarily in green tea, L-Theanine promotes relaxation without causing drowsiness. It can help reduce stress and anxiety, thereby supporting vagal tone.
- **Adaptogens:** These are herbs that help your body adapt to stress, normalizing physiological processes.
- **Ashwagandha:** An Ayurvedic herb known for its calming and stress-reducing properties.

- Rhodiola Rosea: Helps improve mental and physical fatigue, enhancing resilience to stress.
- Holy Basil (Tulsi): Another Ayurvedic herb known for its ability to lower cortisol and promote a sense of calm.

By consciously cultivating a more relaxed state and actively engaging your vagus nerve, you're not just managing stress; you're directly improving the fundamental functioning of your digestive system. This connection between your mind and gut is a powerful key to unlocking your reflux. A calm mind and an active vagus nerve are not the whole answer, though. An imbalance in the gut microbiome can still stall your healing.

Chapter 5: The Overgrown Garden: SIBO (Small Intestinal Bacterial Overgrowth) and Histamine Intolerance

We've covered a lot of ground so far, from stomach acid to the mind-gut connection. Now, let's look at two complex but increasingly recognized conditions that can profoundly impact digestive health and contribute to persistent reflux: Small Intestinal Bacterial Overgrowth (SIBO) and Histamine Intolerance. These issues are often overlooked in conventional approaches, yet they can be significant roadblocks to healing.

Part 1: SIBO - When Bacteria Migrate Upstream

Your digestive tract is a long, winding tube, and different sections are designed to host different populations of microbes. Your large intestine (colon) is teeming with trillions of bacteria, a diverse and beneficial ecosystem. Your small intestine, however, should have relatively few bacteria. When bacteria from the large intestine migrate and overgrow in the small intestine, it's called Small Intestinal Bacterial Overgrowth (SIBO).

- What is SIBO? It's a condition where an excessive amount of bacteria, normally found in the large intestine, colonize the small intestine. These bacteria ferment carbohydrates, producing gases like hydrogen and methane.
- How SIBO Contributes to Reflux:
- Gas Production and Abdominal Pressure: The fermentation of food by overgrown bacteria in the small intestine creates excessive gas (hydrogen and/or methane). This gas builds up, leading to significant bloating and increased intra-abdominal pressure. This pressure acts like a piston, pushing stomach contents upwards and putting constant strain on the Lower Esophageal Sphincter (LES), making it more likely to open inappropriately and cause reflux.¹⁰
- Nutrient Malabsorption: The overgrown bacteria can consume nutrients (especially B vitamins and iron) that you need, leading to deficiencies. They can also damage the small intestinal lining, impairing nutrient absorption.
- Inflammation: The presence of bacteria in the “wrong place” creates chronic inflammation in the small intestine, which can contribute to generalized gut dysfunction and impact motility.
- Impaired Motility: SIBO often exists alongside, or is caused by, impaired gut motility (the coordinated muscular contractions that move food through your digestive tract). If food isn't moving efficiently, bacteria have more time to ferment and proliferate.
- Identifying SIBO:
- The most reliable way to diagnose SIBO is through a Hydrogen/Methane Breath Test. This involves drinking a sugar solution (lactulose or glucose) and then collecting breath samples over a period of 2-3 hours. The gas levels in your breath indicate whether bacteria are fermenting the sugar in your small intestine. Different gas patterns (high hydrogen, high methane, or both) indicate different types of SIBO.

- Addressing SIBO (Integrative Approach): Addressing SIBO requires a multi-faceted approach, often involving diet, targeted antimicrobials, and addressing the underlying root causes.
- Dietary Strategies: The Low-FODMAP Diet:
 - What it is: FODMAPs (Fermentable Oligosaccharides, Disaccharides, Monosaccharides, and Polyols) are types of carbohydrates that are poorly absorbed in the small intestine and readily fermented by bacteria. A low-FODMAP diet temporarily restricts these carbohydrates to “starve” the overgrown bacteria, reducing gas production and symptoms.¹¹
 - How it works: This is typically a temporary elimination diet (2-6 weeks) to reduce symptoms, followed by careful reintroduction to identify triggers. It’s not a long-term solution but a key tool for symptom relief while addressing the root cause.
- Herbal Antimicrobials: several herbs show antimicrobial activity in the laboratory and are used clinically to reduce bacterial overgrowth in the small intestine. Rifaximin is a prescription antibiotic that some physicians use for SIBO. It is not something I prescribe, and it is mentioned here so you know it exists and can raise it with the physician managing that part of your care.
 - Berberine: found in herbs like Goldenseal, Oregon Grape, and Barberry, berberine has broad-spectrum antimicrobial activity in laboratory work against various bacteria, fungi, and parasites. Laboratory activity is not a demonstrated clinical outcome, and it should be read that way.
 - Oregano Oil: an essential oil with antibacterial and antifungal activity in laboratory work. Use an enteric-coated form to prevent irritation, and **cap it at one to two capsules total per day**.
 - Allicin (from Garlic): While whole garlic is high in FODMAPs, concentrated allicin extracts can be effective against specific bacteria, particularly methane-producing organisms.
- Prokinetics: These are agents that stimulate the Migrating Motor Complex (MMC), the “housekeeping wave” of contractions that sweeps undigested food and bacteria through the small intestine between meals. Improving MMC function is essential for preventing SIBO recurrence.
 - Natural Prokinetics: Ginger (fresh or extract), Iberogast (an herbal blend), artichoke extract. These are typically taken before bed or between meals.
- Addressing Root Causes: The most important aspect of long-term SIBO resolution is identifying and addressing the underlying reasons for its development. These often include:
 - Low Stomach Acid: As discussed in Chapter 1, inadequate acid allows bacteria to survive the stomach and enter the small intestine.
 - Vagus Nerve Dysfunction/Poor Motility: A sluggish MMC is a major driver of SIBO.
 - Adhesions from Surgery: Scar tissue can create kinks or blockages in the intestines, impeding flow.
 - Structural Issues: Ileocecal valve dysfunction (the valve between the small and large intestine).
 - Repeated Antibiotic Use: Can disrupt the microbiome and allow resistant strains to flourish.

Part 2: Histamine Intolerance - An Overlooked Trigger

Histamine is a natural compound involved in your immune system, digestion, and nervous system. It plays a role in allergic reactions, signals stomach acid release, and acts as a neurotransmitter. While essential, too much histamine in the body can lead to a wide range of symptoms, including, for some, reflux. This is known as Histamine Intolerance.

- What is Histamine Intolerance? It's not a true allergy, but rather a condition where your body has an excess of histamine, either because it's producing too much, or more commonly, it's not breaking it down efficiently.
- How Histamine Intolerance Relates to Reflux:
- Increased Inflammation: High histamine levels can trigger a systemic inflammatory response, irritating the digestive tract and potentially contributing to a leaky gut.
- Direct Irritation: Histamine can directly irritate the esophageal lining, mimicking or exacerbating reflux symptoms.
- Acid Secretion: While histamine promotes stomach acid secretion (via H2 receptors, which H2 blockers block), an overabundance can lead to dysregulation and discomfort.
- Mast Cell Activation: Histamine is released by mast cells. When these cells are overactive (due to sensitivities, stress, or gut issues), they dump histamine into the system.
- Symptoms of Histamine Intolerance: The symptoms are diverse and can mimic many conditions, which is why it's often missed. Beyond reflux, they can include:
 - Headaches/Migraines
 - Skin Issues: Hives, rashes, flushing, itching.
 - Respiratory: Nasal congestion, sneezing, asthma-like symptoms.
 - Cardiovascular: Palpitations, low blood pressure.
 - Mental Health: Anxiety, panic attacks, insomnia.
 - Digestive: Nausea, abdominal pain, diarrhea.
 - Reflux/Heartburn: Often worsened by high-histamine foods.
- Causes of Histamine Intolerance:
 - DAO Enzyme Deficiency: The primary enzyme responsible for breaking down dietary histamine in the gut is Diamine Oxidase (DAO). If you have insufficient DAO activity (due to genetics, gut issues, certain medications, or nutrient deficiencies like copper or Vitamin C), histamine from food accumulates.
 - Gut Dysbiosis: Some gut bacteria produce histamine, while others break it down. An imbalance can lead to an excess of histamine-producing bacteria. SIBO can also contribute to histamine overload.
 - Leaky Gut: An impaired gut lining can allow more histamine to enter the bloodstream from the gut.
 - Other Conditions: Mast cell activation syndrome, certain medications, and alcohol can also contribute.

- Identifying Histamine Intolerance:
- Elimination Diet (Low-Histamine Diet): This is the most practical diagnostic tool. Eliminate high-histamine foods for 2-4 weeks and observe if your symptoms improve. Then, reintroduce them carefully.
 - High-Histamine Food Examples: Fermented foods (sauerkraut, kombucha, aged cheeses, yogurt), aged meats/fish, processed meats, alcohol, spinach, tomatoes, eggplant, avocado, citrus fruits, chocolate, certain spices.
 - Histamine Releasers: Foods that cause your body to release its own histamine (e.g., strawberries, bananas, chocolate, citrus).
 - DAO Blockers: Foods/substances that inhibit DAO enzyme activity (e.g., alcohol, black tea, green tea, energy drinks).
- DAO Enzyme Levels (Blood Test): Some labs offer a blood test to measure DAO enzyme activity. Low levels can indicate a predisposition to histamine intolerance.
- Dietary Support for Histamine Intolerance:
- Strict Low-Histamine Diet (Temporarily): Focus on fresh, unprocessed, naturally low-histamine foods.
 - Examples: Freshly cooked meat/poultry/fish, fresh vegetables (except the high-histamine ones), gluten-free grains (rice, quinoa), fresh herbs, olive oil.
 - Key is Freshness: Histamine levels increase as food ages, ferments, or is processed. Cook meat and fish immediately after purchase. Avoid leftovers.
- Avoid Histamine Releasers and DAO Blockers.
- Supplements to Support Histamine Breakdown:
- DAO Enzyme: Direct supplementation with a DAO enzyme product before meals can help your body break down dietary histamine and reduce symptoms.
- Vitamin C: A natural antihistamine that helps stabilize mast cells and aids in histamine breakdown.
- Quercetin: A powerful flavonoid that acts as a natural antihistamine and mast cell stabilizer, reducing histamine release.
- Stinging Nettle: Another herb with natural antihistamine properties.
- Probiotics (Carefully Selected): Some probiotic strains can produce histamine, while others help degrade it. Look for specific strains like *Bifidobacterium infantis* or *Lactobacillus plantarum*, which are often considered histamine-friendly or histamine-degrading. Avoid high-histamine producing strains if you suspect histamine intolerance.

Addressing SIBO and histamine intolerance can be complex due to their intricate connections and the wide array of potential triggers. Working with a knowledgeable practitioner is highly recommended for proper testing and a personalized protocol. These internal imbalances are only part of the picture. Something as external and fundamental as your posture has a surprisingly powerful impact of its own.

Chapter 6: The Foundation: The Impact of Poor Posture

We've worked through complex digestive processes, hidden infections, and the mind-gut connection. Now let's bring it back to basics, to something as fundamental as how you hold your body. It might seem too simple, but posture and diaphragm function are direct mechanical inputs to the valve, and this is the chapter people skip on their way to buying something.

How Posture Affects Reflux

Your body is an integrated system, and its structural alignment directly impacts the function of your internal organs. When it comes to reflux, poor posture can contribute in several ways:

- **Increased Abdominal Pressure:**
- **Slouching/Hunching:** When you slouch, particularly while sitting or standing, you compress your abdomen. This literally squeezes your stomach, increasing pressure within the abdominal cavity. This elevated pressure can force stomach contents upwards against the Lower Esophageal Sphincter (LES).
- **Tight Clothing:** Similarly, tight waistbands or belts can put external pressure on your abdomen, mimicking the effects of poor posture.
- **Diaphragmatic Compromise:**
- The diaphragm is a dome-shaped muscle located just below your lungs. It's your primary breathing muscle, but it also plays a critical role in preventing reflux. The esophagus passes through an opening in the diaphragm called the hiatus. The diaphragm provides external support to the LES, helping it stay closed.¹²
- Poor posture (especially hunching forward) can constrict the diaphragm's movement and reduce its ability to properly support the LES. It can also exacerbate or contribute to a hiatal hernia, where part of the stomach pushes up through the diaphragm's opening. While often asymptomatic, a larger hiatal hernia can significantly worsen reflux.
- **Mechanical Impact on the LES:**
- As mentioned, the diaphragm helps cinch the LES. When you're slumped, the natural alignment between the esophagus and the stomach is disrupted. This can reduce the effectiveness of the LES, making it easier for stomach contents to reflux into the esophagus, even if your stomach acid levels are optimal.
- Think of it like a garden hose: if it's kinked or bent, the water flow is restricted or diverted. Similarly, poor posture can create mechanical impediments or weaknesses in your digestive "plumbing."

Common Postural Habits that Worsen Reflux

Consider your daily routine. Do any of these habits sound familiar?

- **Slouching at a Desk:** Hunching forward over a computer or phone for hours, with rounded shoulders and a collapsed chest.
- **Driving Position:** Slumping in your car seat with your stomach compressed against the steering wheel or seatbelt.
- **Eating while Slumped:** Eating meals while hunched over a plate, on the couch, or in bed. This is a double whammy: you're compressing your stomach while it's actively digesting.
- **Lying Down Immediately After Eating:** Gravity is your friend when it comes to keeping stomach contents down. Lying flat removes this advantage and allows stomach acid to pool in the esophagus.
- **Poor Sleeping Position:** Lying completely flat can promote reflux. Sleeping on your right side may also worsen reflux for some due to the anatomy of the stomach and LES.

Practical Posture Modifications

The good news is that posture is something you can consciously improve! Even small changes can make a big difference in reducing the mechanical strain that contributes to reflux.

- **Ergonomics at Work and Home:**
- **Sitting:** Ensure your chair provides good lumbar support. Sit with your feet flat on the floor, knees at a 90-degree angle. Your computer monitor should be at eye level to prevent neck strain and hunching. Use a standing desk if possible, and alternate between sitting and standing.
- **Standing:** Imagine a string pulling you gently from the crown of your head. Keep your shoulders relaxed and back, chest open, and core gently engaged. Avoid locking your knees.
- **Sleeping Position:**
- **Elevate the Head of Your Bed:** This is one of the most effective strategies to use gravity to your advantage. Place 6-8 inch risers or blocks under the bedposts at the head of your bed. Simply using extra pillows often isn't enough, as it can cause neck strain and still leave your torso flat. A wedge pillow designed for reflux can also be helpful.
- **Sleep on Your Left Side:** For many, sleeping on the left side can help reduce reflux. This is thought to be due to the anatomy of the stomach and esophagus, which allows the LES to remain above the level of stomach acid when lying on the left.
- **Mindful Movement Throughout the Day:**
- **Regular Breaks:** If you have a sedentary job, set an alarm to remind yourself to get up, stretch, and walk around every 30-60 minutes.
- **"De-Slump" Breaks:** Every hour or so, practice standing tall, taking a few deep diaphragmatic breaths, and gently stretching your chest open.
- **Stretching and Strengthening Exercises:**

- **Chest Openers:** Stretches that open up the chest and shoulders can counteract the effects of hunching. Examples include doorway stretches or gentle backbends.
- **Core Engagement:** Strengthening your deep core muscles (transverse abdominis) provides better support for your spine and internal organs, reducing abdominal bulging.
- **Diaphragmatic Breathing:** As discussed in Chapter 4, this not only calms the nervous system but also strengthens the diaphragm, improving its role in LES support.

When to Seek Professional Help

If you have persistent postural issues, chronic back or neck pain, or suspect a hiatal hernia, consider seeking professional guidance:

- **Physical Therapy:** A physical therapist can assess your posture, identify muscular imbalances, and provide targeted exercises and stretches.
- **Chiropractic Care:** A chiropractor can address spinal misalignments that may be contributing to poor posture and nerve impingement.
- **Osteopathic Manual Therapy:** Osteopathic practitioners use a hands-on approach to diagnose and treat musculoskeletal imbalances.

Improving your posture is a foundational step in creating an environment conducive to healing your reflux. It's a simple, non-invasive change that can significantly reduce physical stress on your digestive system, allowing the other "keys" to unlocking your reflux to work more effectively. With all these pieces of the puzzle now laid out, it's time to bring it all together into a personalized action plan.

Chapter 7: Your Personalized Action Plan: Bringing It All Together

You've now worked through the factors that contribute to chronic reflux. You understand that it is rarely just "too much acid" but a set of interacting drivers, from the micro-level of bacteria to the macro-level of your physical posture. What that understanding buys you is the ability to investigate in a sensible order rather than trying everything at once.

This chapter is about synthesizing all the knowledge you've gained into a practical, personalized action plan. There's no one-size-fits-all solution for reflux, as each person's underlying "keys" may differ. Your task now is to become the architect of your own healing.

Recap of the "Keys" to Unlock Your Reflux

Let's briefly revisit the critical areas we've covered, which, when addressed, can collectively unlock your reflux:

- **Investigate Stomach Acid Honestly:** the burning is refluxed contents reaching a lining that cannot handle them. Whether insufficient acid is upstream of that in your case is a question to answer with the autoimmune gastritis workup, not to assume. Bitters, ACV, and ginger are low-risk and worth trying. Betaine HCl comes after a demonstrated deficit, under guidance, and never with an ulcer, gastritis, NSAIDs, or alarm features in the picture.
- **Identify & Address *H. pylori*:** This pervasive bacterium can actively suppress stomach acid and cause inflammation. Testing (breath or stool) and targeted protocols (conventional or natural support) are essential.
- **Identify & Remove Food Sensitivities & Allergies:** Beyond the obvious triggers, hidden IgG sensitivities to foods like gluten, dairy, soy, and corn can drive chronic inflammation. The elimination and reintroduction diet is your most powerful tool here.
- **Support Vagus Nerve Function & Improve Parasympathetic Tone:** Chronic stress pushes your body into "fight or flight," hindering digestion. Activating your "rest and digest" system through deep breathing, gargling, cold exposure, mindfulness, and gentle movement is paramount.
- **Address SIBO (Small Intestinal Bacterial Overgrowth) and Histamine Intolerance:** These often-missed conditions can cause gas, pressure, inflammation, and esophageal irritation. Breath testing for SIBO and elimination diets for histamine intolerance are key, followed by targeted protocols.
- **Improve Poor Posture:** Physical alignment significantly impacts abdominal pressure and LES function. Mindful sitting, standing, and sleeping positions, along with specific stretches, can offer surprising relief.

Creating Your Individualized Healing Plan

Now that you have your blueprint, how do you start building?

- Prioritization: Where to Begin?
- Start where the evidence is strongest, which happens to also be where the cost is lowest. **Weight loss if you carry extra, and raising the head of your bed, have the best evidence of anything in this book.** Improving posture and daily diaphragmatic breathing come next, and the highest-value window for the breathing is the half hour after a meal, because that is when reflux events cluster. None of these earns anyone any money, which is roughly why they get mentioned last everywhere else.
- Assess Symptoms: Reflect on which “keys” resonate most with your symptoms.
 - Chronic bloating, gas, and belching after meals might point strongly to low stomach acid or SIBO.
 - Unexplained headaches, skin rashes, or anxiety alongside reflux could suggest histamine intolerance.
 - If you’re constantly stressed, prioritize vagus nerve work.
- Consider Testing: If you haven’t already, discuss H. pylori and SIBO breath testing with your healthcare provider. These objective tests can provide clear direction.
- Start with Diet: The elimination diet for food sensitivities (and potentially the low-FODMAP diet for SIBO or low-histamine diet for histamine intolerance) is foundational. This is often the biggest shift but can yield the most significant results.
- Gradual Introduction of Supplements: Once you have a clearer picture, introduce supplements one at a time, under professional guidance, to address specific imbalances (e.g., Betaine HCl, digestive enzymes, specific probiotics, DAO enzyme, targeted antimicrobials).
- The Importance of Patience and Consistency:
- Healing is rarely linear. There will be good days and challenging days.
- Be patient with your body. It took time for these imbalances to develop, and it will take time for them to heal.
- Consistency is key. Small, daily efforts in diet, lifestyle, and supplement protocols compound over time. Don’t get discouraged by setbacks; simply return to your plan.
- Tracking Progress: Your Personal Data:
 - Symptom Journal: This is invaluable. Keep a daily log of your symptoms (reflux severity, bloating, energy levels, mood, sleep, etc.), the foods you eat, and any supplements or lifestyle practices you implement. This helps you identify patterns, correlate actions with results, and communicate effectively with your healthcare provider.
 - Food Diary: As part of the symptom journal, detail everything you eat and drink. This is essential for identifying food triggers during an elimination and reintroduction phase.

- An Example Action Sequence (Adapt to your needs):
- Phase 1 (Foundation & Assessment):
 - Implement daily deep breathing and start gargling/humming.
 - Begin improving posture throughout the day and elevating the head of your bed.
 - Discuss H. pylori and SIBO testing with your doctor.
 - Start a comprehensive symptom and food journal.
- Phase 2 (Dietary & Initial Support):
 - Begin a targeted elimination diet (e.g., general inflammatory food elimination, then low-FODMAP or low-histamine if tests indicate SIBO/histamine).
 - Incorporate bitters/ACV/ginger before meals.
 - If symptoms suggest, and under guidance, trial Betaine HCl.
- Phase 3 (Targeted Interventions):
 - If H. pylori or SIBO are confirmed, implement specific eradication protocols (conventional or natural) with professional oversight.
 - Introduce targeted supplements based on identified deficiencies or imbalances (e.g., specific probiotics, DAO enzyme, adaptogens).
 - Continue to refine your diet based on reintroduction results.
- Phase 4 (Maintenance & Optimization):
 - Transition to a sustainable, personalized anti-reflux diet.
 - Continue vagus nerve stimulation and good posture habits as part of your daily routine.
 - Address any lingering root causes identified (e.g., chronic stress, nutrient deficiencies).

Long-Term Maintenance: Sustaining Your Healing

Healing reflux isn't just about eliminating symptoms; it's about building a foundation of health that prevents their return.

- **Sustainable Dietary Habits:** Once your triggers are identified, your goal is to incorporate a diet rich in whole, unprocessed foods that nourish your body and avoid your specific sensitivities. This is a lifestyle, not a temporary diet.
- **Stress Management is Non-Negotiable:** Chronic stress is a relentless aggressor against digestion. Continue to prioritize stress reduction techniques daily. Make self-care a non-negotiable part of your routine.
- **Ongoing Gut Health Support:** Consider rotating beneficial probiotics, continuing with digestive support as needed, and ensuring adequate fiber and nutrient intake to maintain a healthy gut microbiome.

- **Listen to Your Body:** Your body is constantly providing feedback. Pay attention to subtle shifts in symptoms. If you notice old patterns emerging, re-evaluate your habits and address potential triggers proactively.

When to Seek Professional Guidance

While this ebook provides a wealth of information, complex health conditions often benefit from personalized, professional guidance. Consider working with:

- **Functional Medicine Practitioner:** These practitioners are uniquely trained to identify and address the root causes of chronic illness, integrating conventional and complementary therapies.
- **Registered Dietitian (RD) with Functional/Integrative Experience:** An RD specializing in gut health can provide tailored dietary guidance, especially for complex elimination diets, SIBO, or histamine intolerance.
- **Gastroenterologist (GI Specialist):** Essential for initial diagnosis, ruling out serious conditions (like Barrett's esophagus or strictures), and performing specific tests (e.g., endoscopy, manometry). They can also help manage conventional treatments when necessary.
- **Chiropractor or Physical Therapist:** If poor posture or musculoskeletal imbalances are significant contributors to your reflux.

Working out what drives your reflux is an act of self-advocacy. By understanding the five drivers and testing them against your own log rather than against anyone's promise, you stop treating symptoms blindly and start working on the thing that is actually failing.

Conclusion

The burning, the constant discomfort, the fear of eating: reflux casts a long shadow. For too long the conversation about it has stopped at acid suppression rather than asking what broke the barrier. You now hold a fuller picture.

You've learned that reflux is rarely a single problem but an interplay of drivers: the barrier, the signal, the terrain, the load, and the capacity. You have also learned which parts of that picture rest on trial evidence and which rest on mechanism and clinical practice, because I marked the difference rather than selling you one grade of certainty for all of it.

By working through these pages you are no longer a passive recipient of symptoms. You have the keys, meaning the understanding of where to look and in what order.

This process requires patience and honest observation. It may involve dietary shifts, mechanical changes, and targeted support. Start with the two measures that have the best evidence and cost nothing, keep a symptom log so you can tell whether anything is actually moving, and give each change enough time to be judged. If four weeks of something has done nothing, stop paying for it.

I will not tell you where this ends, because I have not examined you and I do not know which of the five drivers is yours. What I will tell you is that the people who make the most progress are the ones who changed one thing at a time and wrote it down. Begin there, today.

Working with Root Health

Book a free Root Discovery Session: rootcauseunlocked.com/discovery, or call the Garner, NC office at 919-662-0520.

Consultations and ongoing care are both available wherever you are located. Being outside North Carolina is not a barrier to working with me. Tell me where you are when you reach out and I will lay out exactly how we would work together from there.

If you have been on acid suppression for years, or if you have been handed a protocol with no defined endpoint, bring all of it to the session, including your symptom log. I will tell you honestly which of the five drivers your picture looks like and what I would investigate first. That conversation is free, and it is the fastest way to stop guessing.

If you would rather start on your own, raise the head of your bed tonight, start the log, and read the lab guide in this kit before you buy a single supplement.

Appendix and Resources

- Recommended Reading:
- “Why Stomach Acid Is Good For You” by Jonathan V. Wright, M.D. Listed because it is the origin of the low-acid model and worth reading to understand the argument. Read it alongside Chapter 1 of this book, which explains where that argument does and does not have evidence behind it.
- “The SIBO Made Simple Handbook” by Phoebe Lapine
- “The Wahls Protocol” by Dr. Terry Wahls (for general anti-inflammatory diet principles)
- Helpful Websites/Organizations:
- The Institute for Functional Medicine (IFM): www.ifm.org (Find a practitioner)
- Monash University Low FODMAP Diet: www.monashfodmap.com (Research and App for SIBO diet)
- American Gastroenterological Association (AGA): www.gastro.org (For conventional GI information)
- Glossary of Terms:
- HCl (Hydrochloric Acid): Stomach acid.
- Hypochlorhydria: Low stomach acid.
- LES (Lower Esophageal Sphincter): The muscular valve between the esophagus and stomach.
- H. pylori: Helicobacter pylori, a common stomach bacterium.
- IgG Antibodies: Immune proteins involved in delayed food sensitivities.
- Vagus Nerve: The main nerve of the parasympathetic “rest and digest” system.
- SIBO (Small Intestinal Bacterial Overgrowth): Overgrowth of bacteria in the small intestine.
- FODMAPs: Fermentable Oligosaccharides, Disaccharides, Monosaccharides, and Polyols.
- Histamine Intolerance: Inability to break down histamine efficiently.
- DAO (Diamine Oxidase): The enzyme that breaks down dietary histamine.
- Hiatal Hernia: A condition where part of the stomach pushes through the diaphragm.

References

1. Shahid MS, Ahmed N, Kamal Z, Nathaniel L, Singla B, Singla S, Kumawat S, Batool M, Ekomwereren O, Anika NN, Sahil M. A Systematic Review of Long-Term Use of Proton Pump Inhibitors (PPIs) in Older Adults on Polypharmacy: Do PPIs Deplete Nutrients? *Cureus*. 2025. <https://pubmed.ncbi.nlm.nih.gov/40995253/>
2. Vara-Luiz F, Mendes I, Palma C, Mascarenhas P, Nunes G, Patita M, Fonseca J. Age-Related Decline of Gastric Secretion: Facts and Controversies. *Biomedicines*. 2025. <https://pubmed.ncbi.nlm.nih.gov/40722622/>
3. Targownik LE, Fisher DA, Saini SD. AGA Clinical Practice Update on De-Prescribing of Proton Pump Inhibitors: Expert Review. *Gastroenterology*. 2022. <https://pubmed.ncbi.nlm.nih.gov/35183361/>
4. Casey O, Dobric M, Kelly O, O'Morain CA. The importance of *Helicobacter pylori* eradication: a narrative review. *Frontiers in Gastroenterology (Lausanne)*. 2026. <https://pubmed.ncbi.nlm.nih.gov/41993703/>
5. Huwez FU, Thirlwell D, Cockayne A, Ala'Aldeen DA. Mastic gum kills *Helicobacter pylori*. *New England Journal of Medicine*. 1998. <https://pubmed.ncbi.nlm.nih.gov/9874617/>
6. Abdel-Latif MM, Abouzied MM. Molecular Mechanisms of Natural Honey Against *H. pylori* Infection Via Suppression of NF- κ B and AP-1 Activation in Gastric Epithelial Cells. *Archives of Medical Research*. 2016. <https://pubmed.ncbi.nlm.nih.gov/27751367/>
7. Yanaka A, Fahey JW, Fukumoto A, Nakayama M, Inoue S, Zhang S, Tauchi M, Suzuki H, Hyodo I, Yamamoto M. Dietary sulforaphane-rich broccoli sprouts reduce colonization and attenuate gastritis in *Helicobacter pylori*-infected mice and humans. *Cancer Prevention Research (Philadelphia, Pa.)*. 2009. <https://pubmed.ncbi.nlm.nih.gov/19349290/>
8. Westphal JR, Koch N, Vilchez-Vargas R, Vasapolli R, Malfertheiner P, Schulz C. *Saccharomyces boulardii* as a Probiotic Supplement in *Helicobacter pylori* Eradication Therapy: A Meta-analysis. *Journal of Gastrointestinal and Liver Diseases*. 2026. <https://pubmed.ncbi.nlm.nih.gov/42470702/>
9. Hoshikawa Y, Fitzke H, Sweis R, Fikree A, Saverymuttu S, Kadirkamanathan S, Iwakiri K, Yazaki E, Aziz Q, Sifrim D. Rumination syndrome: Assessment of vagal tone during and after meals and during diaphragmatic breathing. *Neurogastroenterology and Motility*. 2020. <https://pubmed.ncbi.nlm.nih.gov/32383546/>
10. Hu Z, Shan B, Wang Y, Yang J, Lu L, Wang Y, Pang X. Association of gastroesophageal reflux disease with small intestinal bacterial overgrowth: a case-control study. *BMC Gastroenterology*. 2025. <https://pubmed.ncbi.nlm.nih.gov/41131458/>
11. Chuah KH, Loo QY, Loh AJC, Leong JY, Chan WL, Khoo XH, Wong KL, Panirsheeluam S, Natarajan V, Ng AK, Abdul Majid H, Mahadeva S. Clinical Trial: Rifaximin Versus Low FODMAP Diet in Irritable Bowel Syndrome. *Alimentary Pharmacology and Therapeutics*. 2026. <https://pubmed.ncbi.nlm.nih.gov/41098003/>
12. Gu L, Wu F, Cheng J, Shen X. The pathogenesis and surgical treatment progress of gastroesophageal reflux disease. *Frontiers in Surgery*. 2026. <https://pubmed.ncbi.nlm.nih.gov/42266217/>

For educational purposes only. This ebook does not constitute medical advice, diagnosis, or treatment, and reading it does not establish a patient-provider relationship.

Root Health | Dr. Adam Seay, DC | Garner, NC | rootcauseunlocked.com

From Understanding to Doing

Everything up to this point was about understanding: what is happening in your body, why it may be happening, and which directions are worth investigating. That is the half a book can do well.

What follows is the other half, and it is a different kind of document on purpose. Three guides, written to be used rather than read straight through. The first tells you which lab panels answer which question and what the numbers mean, and it includes a request list you can hand to a clinician. The second covers what to eat and, more usefully, how to test a dietary change on yourself so you end up with an answer instead of a habit. The third walks through what has actually been studied, at what dose, and names the products whose trials came back negative.

Each one stands alone, so use them in whatever order matches what you are stuck on. If you are not sure where to begin, begin with the lab guide, because knowing what is driving your symptoms changes what you should do about the other two.

A note on repetition. A few points appear in both halves. That is deliberate. The ones repeated are the ones where getting it wrong causes harm, and they are worth meeting twice.

Read this part first

This guide is educational. It is not medical advice, it does not diagnose anything, and reading it does not make you my patient.

Some reflux symptoms are not reflux. Get medical attention now if you have:

- Chest pain, pressure, or tightness, especially with sweating, nausea, shortness of breath, or pain into the jaw or arm. **Heart attacks are mistaken for heartburn every day. Call 911.**
- Vomiting blood, or stool that is black and tarry
- Food that sticks or will not go down. Complete inability to swallow your own saliva is an emergency.

Book a doctor's appointment, do not self-treat, if you have any of these:

- Difficulty swallowing, or the sensation of food catching
- Unintentional weight loss
- Persistent vomiting
- Anemia, or a known low iron level
- Symptoms that began after age 60
- A family history of esophageal or stomach cancer

Those are the alarm features. Guidelines put patients aged 60 and over presenting with dyspepsia straight to upper endoscopy to exclude organic disease, and note that people at higher risk of malignancy may warrant endoscopy at a younger age (PMID 28631728). This matters more in reflux than in almost anything else I write about, because reflux has a serious downstream complication and because the symptoms are easy to normalize for years.

The honest starting point, and the five drivers

There is no blood test for reflux. Reflux is diagnosed clinically, and confirmed when needed by endoscopy or pH testing, both of which are specialist procedures (PMID 39692638).

That is not a reason to skip testing. It is a reason to test the right thing. Acid is what makes reflux *hurt*. Acid is almost never what makes reflux *happen*. If you only ever measure and suppress the acid, you are treating the last step in a chain and leaving the first four in place, which is exactly why up to half of people remain symptomatic despite adequate acid suppression, a figure that has pushed even the mainstream literature toward describing reflux as a multifactorial condition involving anti-reflux barrier dysfunction rather than a simple acid problem (PMID 42198379).

The five drivers, and what tests each one

Everything in this guide is organized around the same five drivers used in *Unlock Your Reflux*, because the point of testing is to find out which of them is yours.

Driver	What it means	How you find it
1. The barrier	The valve at the top of the stomach is mechanically compromised: sphincter, diaphragm, or hiatal hernia	Endoscopy, symptom pattern, breathing and postural assessment. Not a blood test
2. The signal	The nervous system that runs the valve and reports the pain is dysregulated: vagal tone, stress, sleep, hypervigilance	History, sleep assessment, HRV. Not a blood test
3. The terrain	The microbial environment of the stomach and small intestine is disturbed: H. pylori, overgrowth, dysbiosis	H. pylori breath or stool test, breath testing, stool analysis
4. The load	What is being asked of the system: meal size, meal timing, weight, food reactivity, histamine	Food and symptom diary, structured elimination and reintroduction
5. The capacity	Whether the stomach is doing its job at all: acid, enzymes, emptying	Autoimmune gastritis panel, B12 and iron, gastrin and pepsinogens

Read that table as the map for the rest of this guide. Two of those five drivers, the barrier and the signal, are the ones most often missed, and neither of them has a blood test. That does not make them less real. It makes them the ones you have to look for deliberately.

The blood work in this guide does three jobs inside that map:

- 1. Find a treatable cause underneath the reflux.** H. pylori is the big one, and it sits in the terrain.
- 2. Find the conditions that imitate reflux** and are treated completely differently. Eosinophilic esophagitis and celiac disease belong here.

3. **Find the consequences of the reflux, or of its treatment.** Iron, B12, and magnesium all belong here, and this is the part most often skipped.

The complication you should understand before anything else

Long-standing reflux can change the lining of the lower esophagus into a different type of tissue, called Barrett's esophagus. It is a precancerous change, it results from chronic reflux, and it raises the risk of esophageal adenocarcinoma, a cancer whose incidence has been rising in Western populations (PMID 41230608).

I am not telling you this to frighten you into a scope. Most people with reflux never develop it. I am telling you because it is the reason that "I have had heartburn for fifteen years and I just live with it" is not a neutral choice, and it is the reason the alarm features above are worth taking seriously rather than managing around with antacids.

Driver 1: the barrier, which is mechanical and has no blood test

This is the part of reflux that no lab will ever show you, and for many people it is the whole answer.

What the anti-reflux barrier actually is. Not one valve, but three components working together: the lower esophageal sphincter, the crural diaphragm wrapped around it, and the flap valve formed by the angle at which the esophagus enters the stomach (PMID 40412989).

The diaphragm is half of your sphincter. The crural diaphragm acts as an external sphincter: its contractions raise lower esophageal sphincter pressure with every inspiration and during any rise in abdominal pressure. A sliding hiatus hernia pulls the sphincter up and out of the diaphragm's grip, and it is associated with reflux symptoms, increased acid exposure, and the severity of esophagitis (PMID 15316404). Most reflux episodes are not caused by a permanently weak sphincter at all, but by transient sphincter relaxations, and losing crural support through a hernia significantly compromises the whole barrier (PMID 11003803).

Why this matters more than any supplement in this kit. If your diaphragm is the external sphincter, then how you breathe and how you hold your trunk are not adjunct wellness advice. They are direct inputs to the valve. That is the physiological basis for the breathing work in the supplement guide and for the postural chapter of the ebook, and it is why a chiropractic and structural assessment belongs in a reflux workup rather than being an unrelated appointment.

How the barrier gets assessed.

- **Endoscopy** grades hiatal disruption directly, and there is now a formal endoscopic grading of hiatus integrity that correlates with how well the barrier is holding on prolonged pH recordings (PMID 40412989). If you are having a scope anyway, this is worth asking about by name.
- **Your symptom pattern is informative.** Symptoms tied to bending, lifting, tight waistbands, coughing, straining on the toilet, or a full abdomen point toward pressure and barrier mechanics rather than acid quantity.
- **A breathing and postural assessment.** Whether you breathe into your ribs and belly or into your neck and shoulders, whether your thoracic spine extends, and what happens to your abdomen under load.

One honest note on the emerging piece. A randomized controlled trial of structured inspiratory muscle training in reflux, aimed squarely at the diaphragmatic half of the barrier, has been designed and registered, and it is **published as a protocol with results not yet reported** (PMID 41874149). The rationale is sound and the outcome data are pending. I would rather tell you that than imply the trial already exists.

Driver 2: the signal, where the nervous system sets the volume

The same event can be a minor sensation in one person and disabling pain in another. That difference lives in the nervous system, and it is a genuine physiological finding rather than a way of saying the pain is imaginary.

Stress changes the experience of reflux. In 42 patients undergoing 24-hour pH monitoring, a psychological stressor produced increased subjective symptom ratings, and the authors conclude that the perception of symptoms **in the absence of increased reflux** during stress may explain low response rates to traditional treatment (PMID 16310024). In other words, the acid did not necessarily rise; the volume knob did.

Hypervigilance is measurable and it predicts severity. Esophageal hypervigilance is present across reflux presentations regardless of acid burden or symptom index, and it significantly predicts symptom severity even after controlling for symptom-specific anxiety (PMID 33432708). This is why two people with identical pH studies can have completely different lives.

And the honest counterweight, because it exists. Not every experiment agrees. When acute stress was induced with an IQ test during esophageal acid perfusion in 15 healthy volunteers and 10 reflux patients, it did **not** increase acid perception, even though blood pressure confirmed the stressor worked (PMID 19453516). Short laboratory stress is not the same as months of chronic load, and the honest summary is that chronic stress and hypervigilance have better support here than a single acute stressor does.

The intervention evidence is the encouraging part, and it is in the supplement guide: relaxation training reduced both symptom reports and total esophageal acid exposure compared with an attention-placebo control (PMID 8020690). A mind-body intervention moved an objective acid measurement. That is not a small finding.

Sleep belongs in this driver, and it runs both ways. A systematic review and meta-analysis found a bidirectional relationship between reflux and sleep problems, with reflux associated with poor sleep quality (OR 1.47), sleep disturbance (OR 1.47), and short sleep duration (OR 1.17) (PMID 38646475). Reflux wrecks sleep, and poor sleep worsens reflux, and you can enter that loop from either end.

Sleep apnea deserves its own line. Reflux and sleep apnea are linked, with proposed mechanisms including acid and pepsin irritation of the airway and vagally mediated changes in bronchial tone (PMID 40045407). If you snore, wake unrefreshed, or wake choking, ask for a sleep assessment. Treating the apnea can improve the reflux, and no supplement substitutes for that.

How the signal gets assessed. Not by blood work. By history, by a sleep assessment, by tracking symptoms against stress and sleep rather than against food alone, and by heart rate variability if you want a number to follow. HRV from a wearable is imperfect and it is a reasonable way to watch vagal tone respond to the breathing practice over weeks.

Driver 3: the terrain, and the blood work worth running

Helicobacter pylori

Why it is first. It is common, it is treatable, and it causes real disease. Roughly one in five peptic ulcers is associated with H. pylori infection, and the combination of H. pylori and NSAID use increases bleeding ulcer risk more than sixfold. The test-and-treat strategy is the mainstay of outpatient management: patients under 60 with dyspepsia and no alarm symptoms should be tested and treated if positive (PMID 36791443).

Which test. The two non-invasive options are the urea breath test and the stool antigen test. In a comparison against endoscopy, the ¹³C urea breath test had higher positive and negative predictive values (84.2% and 90.9%) than the stool antigen test (81.3% and 71.4%), with an accuracy of 86.7% versus 76.7%, and the two agreed closely when used to confirm successful eradication (PMID 32635179).

The rule that ruins this test. Both tests are affected by acid suppression and antibiotics. **You generally need to be off PPIs for about two weeks and off antibiotics and bismuth for about four weeks before testing**, or you will get a false negative. Ask your clinician for the exact interval and do not stop a prescription without telling them.

Serology is the weak option. A blood antibody test can stay positive after successful treatment, so it cannot distinguish current from past infection, and it is not the test to use for confirming eradication.

If you are treated, confirm it worked. Retest at least four weeks after finishing therapy. Clarithromycin resistance is rising, and first-line treatment has changed accordingly (PMID 36791443). Eradication is not assumed, it is verified.

Iron studies and a complete blood count

Two reasons, pulling in opposite directions.

Bleeding. An ulcer or an inflamed esophagus can bleed slowly enough that you never see it, and iron deficiency anemia may be the first sign. That is an alarm feature and it needs a doctor.

Absorption. Long-term acid suppression reduces the absorption of several nutrients. Reviews of long-term PPI use list iron alongside magnesium, calcium, and vitamin B12 among the micronutrient deficiencies associated with chronic use (PMID 36791443, PMID 41010960).

Ask for ferritin, serum iron, TIBC, transferrin saturation, and a CBC together.

Vitamin B12 and magnesium, the two the medication takes

B12. Stomach acid is required to liberate B12 from food protein, which is why long-term acid suppression is associated with B12 deficiency (PMID 36791443). If you have been on a PPI for more than a year and nobody has checked, this is the single most reasonable request you can make. B12 deficiency causes fatigue and neurological symptoms that get attributed to almost anything else.

Magnesium. Chronic PPI use is associated with hypomagnesemia, and it is on the standard list of long-term PPI harms (PMID 41010960, PMID 36791443). It can cause cramps, palpitations, and in severe cases arrhythmia. Serum magnesium is an imperfect measure of total body status, and it is still the test you will be able to get. Ask for it if you have been on acid suppression long-term.

The imitators, which must be excluded before anything else

These matter because they are treated completely differently, and because reflux is the default label that gets applied before anyone looks.

Eosinophilic esophagitis

An allergic inflammation of the esophagus that presents with reflux-like symptoms, difficulty swallowing, and food getting stuck. **It is diagnosed only by biopsy at endoscopy**, so no blood test rules it in or out.

Why it deserves a section: esophageal food impaction is the leading complication in people with undiagnosed eosinophilic esophagitis, and in a multicenter study of patients presenting with food impaction, biopsies were taken in only 52% of cases, yet 68.9% of those biopsied received a new diagnosis of the condition (PMID 40971831).

Read that number again. In the people who got the biopsy, most of them had it. **If you have ever had food stick in your esophagus badly enough to need help, ask specifically whether esophageal biopsies were taken.** If they were not, that is a reasonable question to raise.

Celiac disease

Reflux and upper gut symptoms are a recognized presentation. The serology must be drawn while you are still eating gluten, or it is uninterpretable. Removing gluten before testing is the most common way this diagnosis gets missed for years.

Cardiac disease

Not a lab, and the most important item on this page. Reflux and cardiac pain overlap in presentation, and reflux is the more comfortable explanation, which is exactly why it is the dangerous one. New chest pain deserves a cardiac evaluation before it is filed under heartburn, particularly with exertion, and particularly if you have cardiovascular risk factors.

Driver 5: capacity, and how much weight to put on the acid question

Stomach acid, and the honest state of that question

Functional medicine often frames reflux as a problem of **too little** stomach acid rather than too much, and the reasoning is mechanistic: inadequate acid means poor sphincter signaling, slower gastric emptying, and incomplete protein digestion.

Here is where the evidence actually sits, stated plainly so you can judge it.

The strongest published case for replacing lost stomach acid is made for **autoimmune gastritis**, a defined condition in which the parietal cells that make acid are destroyed, producing documented hypochlorhydria and eventually achlorhydria. In that population, the argument is that B12 and iron are routinely replaced while the missing acid is not, that these patients are often prescribed acid suppression for a stomach that is not producing acid, and that re-acidification could improve gastrointestinal symptoms and may theoretically reduce risks tied to hypergastrinemia and N-nitroso compound formation (PMID 38474790).

What that supports and what it does not. It supports acid replacement in someone with **demonstrated** low acid production. It is not evidence that most people with reflux have low stomach acid, and that broader claim does not have trial support behind it.

So the practical position is: test rather than assume. If low acid is genuinely the mechanism in your case, that should be demonstrable, most convincingly through evaluation for autoimmune gastritis, which includes parietal cell and intrinsic factor antibodies, gastrin, pepsinogen I and II with their ratio, plus B12 and iron. A raised gastrin with a low pepsinogen I to II ratio, in someone with B12 or iron deficiency, is a real pattern rather than an inference.

The at-home betaine HCl challenge, where you increase capsules until you feel warmth, is widely described and is not a validated test. It can be informative in practice and it should not be treated as a diagnosis, particularly when the same symptoms would be produced by an ulcer.

The esophageal and gastric microbiome

The terrain thinking that functional medicine has applied to the gut for years has now reached the esophagus, and the mainstream literature is catching up.

A review of the microbiome in esophageal disease describes the picture from the last five years of sequencing work, while being candid that antibiotic use, PPI use, dietary habits, and geography all drive variability and make standardized study difficult (PMID 42132964). A study sampling oral, gastric, and stool microbiomes in patients with reflux esophagitis and Barrett's esophagus found compositional differences between patients and controls, including an association between salivary Tannerellaceae and elevated serum IL-6 (PMID 41486021).

The most useful version of this for you is what it implies about long-term acid suppression. A narrative review from a microbial ecology perspective points out that up to 50% of people remain symptomatic despite adequate acid suppression, describes reflux as multifactorial rather than acid-centric, and examines how suppressing acid alters the microbial ecology of the upper gut (PMID 42198379).

Read that as directional rather than actionable: there is not yet a microbiome test that tells you what to do about your reflux. What it does support is the underlying premise, that acid suppression is not a neutral act and that the environment of the stomach matters beyond its pH.

Small intestinal bacterial overgrowth

A case-control study of 50 patients with GERD confirmed by manometry and 24-hour acid monitoring, against 53 symptom-matched controls, found SIBO positively correlated with GERD, and mean expiratory hydrogen significantly higher in the GERD group. The authors describe a potential connection and a new perspective on the pathogenesis (PMID 41131458).

Read that as an association in 103 people, not a causal chain. Breath testing is reasonable if bloating and distension dominate your picture. It is not where I would start in someone whose main complaint is heartburn.

The rest of the terrain

Thyroid function, fasting glucose and HbA1c, and hs-CRP are all cheap, and they occasionally reframe the picture. Body weight belongs here too, and it is covered in the diet guide, because it has better evidence behind it than any test on this page.

Two notes on how to read those. Ask for **fasting insulin** alongside glucose and HbA1c, because insulin rises years before glucose does and a normal glucose can sit on top of a picture that is already moving. And read **hs-CRP** for what it is: a nonspecific screening marker. It says an inflammatory load exists somewhere, not where it came from, and it confirms nothing on its own. Repeat it when you are well rather than acting on a single reading taken after an illness.

Functional ranges for the markers in this guide

Clinical targets I use at Root Health, not validated diagnostic thresholds, narrower than the standard laboratory range on purpose.

Marker	Units	Functional target	Approximate standard range
Ferritin	ng/mL	50 to 150	25 to 300
Serum iron	ug/dL	65 to 110	40 to 150
TIBC	ug/dL	290 to 370	250 to 400
Transferrin saturation	%	20 to 40	15 to 45
Vitamin B12	pg/mL	500 to 900	200 to 1100
Folate (RBC)	ng/mL	600 to 1500	400 to 2000
Magnesium (serum)	mg/dL	2.0 to 2.5	1.5 to 2.7
Magnesium (RBC)	mg/dL	5.2 to 6.8	4.0 to 7.2
Zinc (plasma)	ug/dL	70 to 110	60 to 120
Vitamin D (25-OH)	ng/mL	40 to 80	20 to 100
hs-CRP	mg/L	under 0.5	under 1.0
Glucose, fasting	mg/dL	75 to 86	70 to 99
HbA1c	%	4.6 to 5.3	4.0 to 5.6
Albumin	g/dL	4.2 to 4.8	3.5 to 5.0
White blood cells	K/uL	5.0 to 7.5	3.5 to 10.5

Reading your results as a pattern

Pattern one: H. pylori positive. The clearest good news in this guide, because it is treatable and verifiable. Treat, then confirm eradication at least four weeks later.

Pattern two: long-term PPI with depleted nutrients. Low B12, low ferritin, low magnesium in someone who has been on acid suppression for years. Both halves need attention: replace what is low, and have a real conversation about whether the medication is still needed at the current dose. The deprescribing section of the supplement guide covers how that is done safely.

Pattern three: everything normal, symptoms persist. Common. It moves the work to the things labs do not measure: weight, meal timing, the position you sleep in, and the breathing and vagal work in Chapter 4 of the ebook. Those are covered in the diet guide, and several of them outperform anything on this lab list.

Pattern four: normal labs, mechanical story. Symptoms on bending, lifting, tight clothing, or a full stomach, worse lying flat, better standing. That is a barrier pattern, and the work is mechanical: bed elevation, meal size, breathing, posture, and a conversation about hiatal anatomy.

Pattern five: normal labs, nervous system story. Symptoms track stress and poor sleep more than they track food, and the intensity is out of proportion to what any scope shows. That is the signal driver, and hypervigilance predicts symptom severity independently of acid burden (PMID 33432708). The work is vagal, behavioral, and sleep-focused, and it is not a lesser diagnosis.

Pattern six: swallowing trouble. Stop reading and get the endoscopy. Difficulty swallowing is not a terrain problem.

The conversation with your doctor

A request list you can bring

- H. pylori testing by urea breath test or stool antigen, after the appropriate washout from acid suppression
- CBC, ferritin, serum iron, TIBC, transferrin saturation
- Vitamin B12, and RBC folate
- Magnesium, if you have been on acid suppression long-term
- Celiac serology, drawn while still eating gluten
- TSH with free T4
- 25-hydroxy vitamin D
- Fasting glucose, HbA1c, and fasting insulin
- hs-CRP
- A discussion of upper endoscopy if you have any alarm feature, are 60 or over with new symptoms, or have had symptoms for many years

Usually through a functional medicine practice, often out of pocket:

- Autoimmune gastritis panel (parietal cell and intrinsic factor antibodies, gastrin, pepsinogen I and II with the ratio), if low stomach acid is genuinely suspected
- Breath testing for bacterial overgrowth, if bloating and distension dominate
- Comprehensive stool analysis, if the terrain picture is broader than reflux alone

Not laboratory tests at all, and for many people the highest-yield part of the workup:

- **A breathing and postural assessment.** Whether you can breathe diaphragmatically, what your thoracic spine does, and what happens to abdominal pressure under load. This assesses Driver 1 directly and no blood test substitutes for it
- **If you are having an endoscopy anyway, ask about the hiatus specifically**, since hiatal integrity can be graded endoscopically and correlates with barrier competence (PMID 40412989)
- **A sleep assessment**, formal if you snore or wake unrefreshed
- **Two weeks of tracking symptoms against stress and sleep**, not only against food

The sentence that changes the appointment

“I have been on acid suppression for X years, and I would like to know whether I still need it and whether it has affected my B12, iron, or magnesium.” That is a specific, reasonable, guideline-aligned request, and it opens the deprescribing

conversation without sounding like you are refusing treatment.

What these labs cannot tell you

- **Whether you have reflux.** That is clinical, and confirmed by scope or pH study when it matters (PMID 39692638).
- **Whether you have Barrett's esophagus.** Only endoscopy answers that (PMID 41230608).
- **Whether a specific food is your trigger.** Only a structured elimination and reintroduction does, and the diet guide covers it.
- **Whether your stomach acid is low.** Not from a routine panel. That takes the specific workup described above.

When to seek clinical review

Emergency: chest pain with sweating, nausea, breathlessness, or pain into the jaw or arm. Vomiting blood or black tarry stools. Inability to swallow saliva.

Soon:

- Difficulty swallowing or food catching, ever
- Unintentional weight loss, persistent vomiting, or anemia
- New symptoms starting at age 60 or over
- Symptoms needing daily treatment for more than a few weeks
- Ten or more years of untreated symptoms, which is a Barrett's screening conversation
- Reflux with cough, hoarseness, or asthma that is not improving

Your next step

Testing tells you what you are dealing with. The companion guides do the rest: **The Reflux Diet Guide** covers what actually has evidence behind it, including which famous trigger foods do not, and **The Reflux Supplement Guide** covers what has been studied and how deprescribing is done safely.

Book a free Root Discovery Session: rootcauseunlocked.com/discovery?utm_source=guide&utm_medium=ebook&utm_campaign=reflux-labs

Consultations and ongoing care are both available wherever you are located. Being outside North Carolina is not a barrier to working with me. Tell me where you are when you reach out and I will lay out exactly how we would work together from there.

Get in touch: rootcauseunlocked.com/contact?utm_source=guide&utm_medium=ebook&utm_campaign=reflux-labs

If you would rather start on your own, start a symptom and meal diary. In reflux it is unusually informative, because the timing between a meal, a position, and a symptom is the diagnosis:

Free symptom tracker: rootcauseunlocked.com/tracker?utm_source=guide&utm_medium=ebook&utm_campaign=reflux-labs

References

All references were verified against live PubMed records on 2026-08-08, including a publication type and corrections check for retractions and errata. Verification detail is documented in `Reflux-Kit-Citation-Ledger.md`.

1. Moayyedi P, Lacy BE, Andrews CN, et al. ACG and CAG Clinical Guideline: Management of Dyspepsia. *Am J Gastroenterol*. 2017;112(7):988-1013. PMID 28631728
2. ASGE Standards of Practice Committee, Desai M, Ruan W, et al. American Society for Gastrointestinal Endoscopy guideline on the diagnosis and management of GERD: summary and recommendations. *Gastrointest Endosc*. 2025;101(2):267-284. PMID 39692638
3. Al Hazzouri A, Attieh P, Azizi T, et al. Barrett's Esophagus. *J Gastroenterol Hepatol*. 2026;41(1):41-53. PMID 41230608
4. McConaghy JR, Decker A, Nair S. Peptic Ulcer Disease and H. pylori Infection: Common Questions and Answers. *Am Fam Physician*. 2023;107(2):165-172. PMID 36791443
5. Alzoubi H, Al-Mnayyis A, Al Rfoa I, et al. The Use of 13C-Urea Breath Test for Non-Invasive Diagnosis of Helicobacter pylori Infection in Comparison to Endoscopy and Stool Antigen Test. *Diagnostics (Basel)*. 2020;10(7):448. PMID 32635179
6. Andrawes M, Andrawes W, Das A, et al. Proton Pump Inhibitors, An Evidence-Based Review of Indications, Efficacy, Harms, and Deprescribing. *Medicina (Kaunas)*. 2025;61(9). PMID 41010960
7. Siegfried Y, Pokryszka J, Hruz P, et al. Wide discrepancy between best practice recommendations and real-life management of suspected eosinophilic esophagitis-associated food bolus impaction. *Dis Esophagus*. 2025;38(5). PMID 40971831
8. Taylor L, McCaddon A, Wolffenbuttel BHR. Creating a Framework for Treating Autoimmune Gastritis, The Case for Replacing Lost Acid. *Nutrients*. 2024;16(5):662. PMID 38474790
9. Hu Z, Shan B, Wang Y, et al. Association of gastroesophageal reflux disease with small intestinal bacterial overgrowth: a case-control study. *BMC Gastroenterol*. 2025;25(1):753. PMID 41131458
10. Kahrilas PJ, Nguyen NT. The Esophagogastric Junction Reconsidered. *Gastrointest Endosc Clin N Am*. 2025;35(3):541-553. PMID 40412989
11. van Herwaarden MA, Samsom M, Smout AJ. The role of hiatus hernia in gastro-oesophageal reflux disease. *Eur J Gastroenterol Hepatol*. 2004;16(9):831-835. PMID 15316404
12. Holloway RH. The anti-reflux barrier and mechanisms of gastro-oesophageal reflux. *Baillieres Best Pract Res Clin Gastroenterol*. 2000;14(5):681-699. PMID 11003803
13. Syropoulos S, Moutzouri M, Grammatopoulou E, et al. Impact of Inspiratory Muscle Training in Individuals with Gastroesophageal Reflux Disease: A Randomized Controlled Trial Protocol. *Methods Protoc*. 2026;9(2). PMID 41874149
14. Wright CE, Ebrecht M, Mitchell R, et al. The effect of psychological stress on symptom severity and perception in patients with gastro-oesophageal reflux. *J Psychosom Res*. 2005;59(6):415-424. PMID 16310024
15. Guadagnoli L, Yadlapati R, Taft T, et al. Esophageal hypervigilance is prevalent across gastroesophageal reflux disease presentations. *Neurogastroenterol Motil*. 2021;33(8):e14081. PMID 33432708
16. Hemmink GJ, Bredenoord AJ, Weusten BL, et al. Does acute psychological stress increase perception of oesophageal acid? *Neurogastroenterol Motil*. 2009;21(10):1055-e86. PMID 19453516
17. McDonald-Haile J, Bradley LA, Bailey MA, et al. Relaxation training reduces symptom reports and acid exposure in patients with gastroesophageal reflux disease. *Gastroenterology*. 1994;107(1):61-69. PMID 8020690

18. Tan X, Wang S, Wu F, et al. Bidirectional correlation between gastroesophageal reflux disease and sleep problems: a systematic review and meta-analysis. *PeerJ*. 2024;12:e17202. PMID 38646475
19. Wang J, Wang P, Lv J, et al. Exploring the silent connection: unveiling the intricate relationship between gastroesophageal reflux disease and sleep apnea syndrome. *Hum Genomics*. 2025;19(1):23. PMID 40045407
20. He P, Stein E. The Microbiome and Esophageal Disease: Where Are We Now? *Curr Gastroenterol Rep*. 2026;28(1). PMID 42132964
21. Yao CC, Tai WC, Liang CM, et al. Alternation of the oral, gastric and stool microbiome in patients with reflux esophagitis and Barrett's esophagus. *J Microbiol Immunol Infect*. 2026;59(4):463-470. PMID 41486021
22. Zaroni A, Facchin S, Mari V, et al. Rethinking Long-Term PPI Therapy in GERD: A Narrative Review from a Microbial Ecology Perspective Beyond Acid Suppression. *Pharmaceuticals (Basel)*. 2026;19(5). PMID 42198379

Dr. Adam Seay, DC · Seay Wellness / Root Health, Garner, NC
rootcauseunlocked.com · (919) 662-0520

Educational content only. Not medical advice. Not a diagnosis. Not a substitute for your clinician. Chest pain is a medical emergency until proven otherwise.

Read this part first

This guide is educational. It is not medical advice, it does not diagnose anything, and reading it does not make you my patient.

Call 911 for chest pain or pressure with sweating, nausea, breathlessness, or pain into the jaw or arm. Heartburn and heart attack are confused constantly, and the comfortable explanation is the dangerous one.

See a doctor before dieting your way through this if you have trouble swallowing, food sticking, unintentional weight loss, vomiting blood, black tarry stools, anemia, or new symptoms after age 60. Those are covered in the companion lab guide and none of them are a diet problem.

If you are pregnant, underweight, diabetic, or have a history of disordered eating, talk with a clinician before restricting food or changing meal timing.

Where food actually sits in the reflux picture

Reflux has five drivers, and they are the spine of this whole kit: **the barrier** (the valve and the diaphragm), **the signal** (vagal tone, stress, sleep), **the terrain** (the microbial environment), **the load** (what you are asking the system to handle), and **the capacity** (whether the stomach is doing its job). The lab guide maps all five.

Food is the fourth. That placement is deliberate and it is the most useful thing in this guide, because most people arrive believing food is the first, spend two years subtracting foods, and never touch the driver that was actually theirs.

Food still matters, and it matters in a specific way: **it changes the load on a barrier that is already compromised**. A large meal distends the stomach and triggers the sphincter relaxations that let reflux happen. A late meal puts that load in your esophagus while you are horizontal. Extra abdominal weight raises the pressure the valve has to hold against. Those are mechanical effects of food, and they are where the evidence is.

What the evidence does not support is the other model, the one where specific foods are chemically toxic to your esophagus and must be eliminated forever.

Most of what you have been told to avoid has never been shown to help.

That sentence will save some readers years. When researchers applied an evidence grading system to the standard lifestyle advice for reflux, reviewing the literature on smoking, alcohol, obesity, weight loss, caffeine and coffee, citrus, chocolate, spicy food, head-of-bed elevation, and late evening meals, they found **no published evidence of efficacy for dietary measures**. What survived the review was weight loss and head-of-bed elevation (PMID 16682569).

A later systematic review reached a compatible place with more nuance: weight loss and smoking cessation should be recommended to people with reflux who are obese or who smoke respectively, and for nocturnal reflux, avoiding late evening meals and elevating the head of the bed are effective. In randomized trials, late evening meals increased time with supine acid exposure compared with early meals by 5.2 percentage points, and head-of-bed elevation decreased supine acid exposure time from 21% to 15% (PMID 25956834).

So the hierarchy is not what the trigger lists imply. It looks like this:

1. **Weight, if you carry extra.** The single most evidenced dietary lever.
2. **When you eat, and how you sleep.** Timing and position beat food choice.
3. **Your own individual triggers**, discovered by testing rather than assumed from a list.
4. **The general pattern of the diet**, which is worth improving for reasons beyond reflux.

Everything below follows that order, because spending your effort in that order is what gets results.

Part one: the two things with the best evidence

Weight

If you are carrying extra weight, particularly around the middle, this is the highest-yield dietary change available to you. Abdominal weight increases pressure on the stomach and the junction between stomach and esophagus, which is mechanical rather than metaphorical. Weight loss improved both acid measurements and symptoms in the evidence review, one of only two measures that did (PMID 16682569).

Two honest notes. First, this is not a reason to attempt something drastic; gradual loss is what the evidence describes and crash approaches make reflux worse in the short term. Second, if this does not apply to you, skip it entirely rather than feeling you have missed the main event. Plenty of people with reflux are not overweight.

Timing and position

Sleep is part of this, in both directions. A systematic review and meta-analysis found a bidirectional relationship between reflux and sleep problems: reflux is associated with poor sleep quality, sleep disturbance, and short sleep duration, and the relationship runs the other way too (PMID 38646475). You can enter that loop from either end, which means fixing the evening meal and fixing the sleep are the same project rather than two.

Stop eating three hours before you lie down. This is the highest-value free change in the entire guide, and in the randomized data late meals measurably increased the time acid sat in the esophagus overnight (PMID 25956834).

Raise the head of the bed. Not extra pillows, which bend you at the waist and can make it worse. Raise the head of the bed frame itself by six to eight inches, or use a wedge under the mattress. Head-of-bed elevation cut supine acid exposure time from 21% to 15% (PMID 25956834).

Sleep on your left side. The left lateral position improved the time esophageal pH stayed above 4 in the evidence review (PMID 16682569). It is anatomy: on the left, the junction sits above the pool of stomach contents rather than in it.

Eat smaller meals. A large meal distends the stomach, and a distended stomach triggers the sphincter relaxations that let reflux happen. Smaller and more frequent beats large and occasional here.

Smoking, with the honest nuance

Smoking cessation is worth doing, and the reflux-specific evidence is narrower than you would expect. In a prospective population cohort of 29,610 people, stopping smoking was associated with improvement in **severe** reflux symptoms only among people of normal body mass index who were using antireflux medication at least weekly, with an odds ratio of 5.67 in that group, and not among overweight individuals or those with minor symptoms (PMID 24322837).

Stop smoking. Just know that if you are overweight, the weight is likely the bigger reflux lever of the two.

Part two: finding your actual triggers

The standard trigger list is coffee, chocolate, citrus, tomato, spicy food, mint, fat, and alcohol. As a general prescription it did not survive evidence review (PMID 16682569). As a list of **candidates to test on yourself**, it is perfectly reasonable, because individual reactions are real even when the population-level evidence is not.

The difference between those two uses is the whole game.

The two-week test, run properly

Days 1 to 7, baseline. Change nothing. Record every meal with a time, and every symptom with a time and a severity out of ten. The gap between the meal and the symptom is the most informative number you will collect.

Look for the pattern before you remove anything. Most people find their diary implicates two or three items, not fifteen, and often it implicates a *timing* or a *portion size* rather than a food.

Days 8 to 21, remove your top three candidates. Only three. Keep everything else stable.

Then reintroduce, one at a time. A normal portion on one day, then two days of watching. A food that reproduces symptoms across two separate trials is a real trigger for you. Everything else goes back in.

Expect a short list. If you finish this process avoiding twelve foods, something has gone wrong with the method rather than with your esophagus.

What is worth knowing about the usual suspects

Alcohol. Relaxes the lower esophageal sphincter and is one of the more commonly reported precipitants. Worth removing during any trial period so it does not confound your results.

Coffee. Frequently blamed, and the population evidence for avoidance is weak (PMID 16682569). Test it. Many people find that the amount and the timing matter more than the coffee itself, particularly coffee on an empty stomach.

Fat and portion size. Large, fatty meals slow gastric emptying and keep the stomach distended for longer. In practice I see portion size matter more often than fat content.

Mint and chocolate. Both are traditionally implicated on sphincter-relaxation grounds. Both belong in the test-it category rather than the avoid-it-forever category.

Carbonation. Frequently reported and easy to test. If fizzy drinks reliably produce symptoms within the hour, you have your answer without a study.

Part two and a half: how you eat, which is the vagus nerve at the table

This section exists because the mind-gut driver from Chapter 4 of the ebook shows up at every meal, and almost nobody works on it.

Digestion is a parasympathetic activity. The vagus nerve drives the sequence: it triggers stomach acid and enzyme release in anticipation of food, it coordinates the muscular activity that moves a meal along, and it governs the rate at which the stomach empties. Eat while stressed, rushed, or working, and you are asking the digestive system to run while the nervous system is prioritizing something else.

The nervous system also sets the volume on the symptom. Esophageal hypervigilance is present across reflux presentations regardless of acid burden and predicts symptom severity (PMID 33432708), and in patients undergoing pH monitoring, a psychological stressor increased symptom ratings, with the authors noting that symptom perception **in the absence of increased reflux** may explain poor responses to standard treatment (PMID 16310024). Not every study agrees, and a short laboratory stressor did not increase acid perception in one experiment (PMID 19453516), so the better-supported version is chronic load rather than a single bad afternoon.

What to actually do, in order of how much it matters:

1. **Take five slow breaths before the first bite.** Longer exhale than inhale. This is the cheapest vagal input available and it takes thirty seconds.
2. **Sit down. No screens, no driving, no standing at the counter.** The posture point is not manners, it is abdominal pressure and diaphragm position.
3. **Chew until the texture is gone.** Chewing is the only part of digestion under conscious control, and it is the step that determines how much work the stomach has to do.
4. **Put the fork down between bites.** The single most reliable way to slow a meal without counting anything.
5. **Stop at comfortably satisfied, not full.** Fullness is stomach distension, and distension is the trigger for the sphincter relaxations that produce reflux.
6. **Do not lie down, and do not exercise hard, for the hour after eating.** Walk instead. A gentle walk after a meal supports emptying without raising abdominal pressure the way crunches or heavy lifting does.

Then breathe again afterward. The highest-value time for the diaphragmatic breathing practice in the supplement guide is the half hour after a meal, because that is when reflux events cluster and when the practice was shown to reduce them.

Part three: the eating pattern

A plant-forward Mediterranean pattern

The most interesting study in this area compared a dietary approach against medication. In a retrospective comparison, patients with laryngopharyngeal reflux treated with alkaline water, a plant-based Mediterranean-style diet, and standard reflux precautions were compared with patients treated with proton pump inhibition and standard precautions. The percentage reduction in the reflux symptom index was significantly greater with the dietary approach, and the authors suggest a plant-based diet and alkaline water should be considered in the treatment of this condition (PMID 28880991).

How to read that honestly. It is retrospective rather than randomized, and it is in laryngopharyngeal reflux, the throat-and-voice presentation, rather than classic heartburn. That is a real limitation. It is also one of the few studies where a dietary approach was measured against the drug and did not lose, which is worth knowing.

What the pattern means practically: vegetables and fruit at most meals, olive oil as the main fat, fish and legumes for protein more often than red and processed meat, whole grains, and far less ultra-processed food.

Fiber, and the reason it belongs here

A pattern built on plants brings fiber, and fiber supports regular bowel function. Straining and constipation raise intra-abdominal pressure, which is the same mechanism by which abdominal weight worsens reflux. This is an unglamorous connection that pays off.

Increase fiber gradually. A sudden jump produces bloating, and a distended abdomen is not what you want.

The histamine question, briefly

Some people with reflux also have histamine-related symptoms, and the aged, fermented, and leftover foods that carry the most histamine overlap with the classic reflux trigger list. If flushing, headache, hives, or palpitations travel with your digestive symptoms, the four-week freshness-based trial in the Allergy Diet Guide is the structured way to test that, and DAO support is covered in the supplement guide. If your symptoms are purely heartburn, this is probably not your thread to pull.

Bacterial overgrowth, if bloating dominates

A case-control study of 50 people with GERD confirmed by manometry and acid monitoring, against 53 symptom-matched controls, found small intestinal bacterial overgrowth positively correlated with GERD and higher mean expiratory hydrogen in the GERD group (PMID 41131458). That is an association in 103 people, not a proven mechanism.

Practically: if bloating and distension are as prominent as the burning, it is worth raising breath testing with a clinician rather than reaching for a low FODMAP diet on your own. Low FODMAP is a demanding, temporary, dietitian-supported protocol, and it is not a reflux diet.

A week that supports the plan

The rhythm is the point: three moderate meals, the last one finished three hours before bed.

Breakfast options. Oats with berries and ground flax. Eggs with sauteed greens. Plain yogurt with fruit if dairy is not on your personal list.

Lunch options. Lentil or vegetable soup with olive oil. Chicken or fish over rice or quinoa with vegetables. A large salad with beans, olive oil, and a protein.

Dinner options, finished early. Baked fish with potatoes and greens. Chicken thighs with roasted vegetables. Turkey and vegetable stir fry over rice.

The rules that matter more than the menu. Moderate portions. Last bite three hours before lying down. Left side, head of bed raised. Alcohol out during any trial period. Water through the day rather than large volumes with meals.

What I deliberately left out, and why

- **The universal trigger list as a prescription.** No published evidence of efficacy for dietary measures in the evidence review (PMID 16682569). Use it as candidates to test, not rules to obey.
- **Baking soda in water as a routine remedy.** It relieves symptoms briefly and it is a large sodium load, it can cause rebound, and in rare cases stomach rupture has been described with large amounts on a full stomach. Alginate products are covered in the supplement guide and are a better answer.
- **Apple cider vinegar as a general reflux treatment.** Widely recommended on the low-acid theory. I could not locate randomized trial evidence for reflux, and if there is an ulcer or erosive esophagitis it can hurt. It is not on this list on the strength of theory alone.
- **Long elimination diets.** Twelve removed foods is a method failure, not a discovery.
- **Alkaline water as a stand-alone answer.** It appeared as one component of a dietary package in a retrospective study (PMID 28880991). That is not the same as the water doing the work.

Telling you what to skip is part of what you paid for.

When to seek professional care

Emergency: chest pain with sweating, nausea, breathlessness, or pain radiating to the jaw or arm. Vomiting blood or black tarry stools. Inability to swallow saliva.

Soon:

- Difficulty swallowing or food sticking, ever
- Unintentional weight loss, persistent vomiting, or anemia
- New symptoms at age 60 or over
- Symptoms requiring daily treatment for more than a few weeks
- Many years of untreated symptoms, which is a Barrett's screening conversation
- Reflux with cough, hoarseness, or worsening asthma

Working with Root Health

This guide is written to be safe and useful for a reader I have never met. A personalized plan starts from your diary, your test results, and your medication list, and sequences changes rather than stacking them.

***Book a free Root Discovery Session: [rootcauseunlocked.com/discovery?](https://rootcauseunlocked.com/discovery?utm_source=guide&utm_medium=ebook&utm_campaign=reflux-diet)
[utm_source=guide&utm_medium=ebook&utm_campaign=reflux-diet](https://rootcauseunlocked.com/discovery?utm_source=guide&utm_medium=ebook&utm_campaign=reflux-diet)***

Consultations and ongoing care are both available wherever you are located. Being outside North Carolina is not a barrier to working with me. Tell me where you are when you reach out and I will lay out exactly how we would work together from there.

***Get in touch: [rootcauseunlocked.com/contact?](https://rootcauseunlocked.com/contact?utm_source=guide&utm_medium=ebook&utm_campaign=reflux-diet)
[utm_source=guide&utm_medium=ebook&utm_campaign=reflux-diet](https://rootcauseunlocked.com/contact?utm_source=guide&utm_medium=ebook&utm_campaign=reflux-diet)***

If you would rather start on your own, the tracker is what turns a food experiment into an answer:

***Free symptom tracker: [rootcauseunlocked.com/tracker?](https://rootcauseunlocked.com/tracker?utm_source=guide&utm_medium=ebook&utm_campaign=reflux-diet)
[utm_source=guide&utm_medium=ebook&utm_campaign=reflux-diet](https://rootcauseunlocked.com/tracker?utm_source=guide&utm_medium=ebook&utm_campaign=reflux-diet)***

Read the companion documents alongside this one. **Understanding Your Reflux Labs** covers H. pylori and the conditions that imitate reflux, and **The Reflux Supplement Guide** covers what has been studied and how to come off acid suppression safely.

References

All references were verified against live PubMed records on 2026-08-08, including a publication type and corrections check for retractions and errata. Verification detail is documented in `Reflux-Kit-Citation-Ledger.md`.

1. Kaltenbach T, Crockett S, Gerson LB. Are lifestyle measures effective in patients with gastroesophageal reflux disease? An evidence-based approach. *Arch Intern Med*. 2006;166(9):965-971. PMID 16682569
2. Ness-Jensen E, Hveem K, El-Serag H, et al. Lifestyle Intervention in Gastroesophageal Reflux Disease. *Clin Gastroenterol Hepatol*. 2016;14(2):175-182. PMID 25956834
3. Ness-Jensen E, Lindam A, Lagergren J, et al. Tobacco smoking cessation and improved gastroesophageal reflux: a prospective population-based cohort study: the HUNT study. *Am J Gastroenterol*. 2014;109(2):171-177. PMID 24322837
4. Zalvan CH, Hu S, Greenberg B, et al. A Comparison of Alkaline Water and Mediterranean Diet vs Proton Pump Inhibition for Treatment of Laryngopharyngeal Reflux. *JAMA Otolaryngol Head Neck Surg*. 2017;143(10):1023-1029. PMID 28880991
5. Hu Z, Shan B, Wang Y, et al. Association of gastroesophageal reflux disease with small intestinal bacterial overgrowth: a case-control study. *BMC Gastroenterol*. 2025;25(1):753. PMID 41131458
6. Guadagnoli L, Yadlapati R, Taft T, et al. Esophageal hypervigilance is prevalent across gastroesophageal reflux disease presentations. *Neurogastroenterol Motil*. 2021;33(8):e14081. PMID 33432708
7. Wright CE, Ebrecht M, Mitchell R, et al. The effect of psychological stress on symptom severity and perception in patients with gastro-oesophageal reflux. *J Psychosom Res*. 2005;59(6):415-424. PMID 16310024
8. Hemmink GJ, Bredenoord AJ, Weusten BL, et al. Does acute psychological stress increase perception of oesophageal acid? *Neurogastroenterol Motil*. 2009;21(10):1055-e86. PMID 19453516
9. Tan X, Wang S, Wu F, et al. Bidirectional correlation between gastroesophageal reflux disease and sleep problems: a systematic review and meta-analysis. *PeerJ*. 2024;12:e17202. PMID 38646475

Dr. Adam Seay, DC · Seay Wellness / Root Health, Garner, NC
rootcauseunlocked.com · (919) 662-0520

Educational content only. Not medical advice. Not a diagnosis. Not a substitute for your clinician.

Read this part first

This guide is educational. It is not medical advice, it does not diagnose anything, and reading it does not make you my patient.

Call 911 for chest pain or pressure with sweating, nausea, breathlessness, or pain into the jaw or arm.

See a doctor before treating yourself if you have difficulty swallowing, food sticking, unintentional weight loss, vomiting blood, black tarry stools, anemia, or new symptoms after age 60.

Do not stop a proton pump inhibitor abruptly on the strength of this document. There is a section below on how deprescribing is actually done, and the short version is that stopping suddenly is the version that tends to fail. If you take a PPI for Barrett's esophagus, a bleeding ulcer, or gastroprotection alongside NSAIDs, aspirin, or a blood thinner, that prescription is doing a specific job and stopping it is a decision for the prescriber, not for a guide.

Supplements in the United States are regulated as food, not as drugs. Quality varies. Look for independent third-party testing.

Why nutrient evidence looks thinner than it is

A large outcome trial costs tens to hundreds of millions of dollars, and that money only exists when a patent can pay it back. You cannot patent magnesium, an herb, or a food. So the evidence hierarchy this guide grades against is not neutral: it systematically over-documents patentable drugs and under-documents nearly everything else. That is measurable, not rhetorical. A Cochrane methodology review found that manufacturer sponsorship of drug and device studies produces more favourable results and conclusions than independent funding, through a bias that standard quality checks do not catch (PMID 28207928).

Two honest consequences follow, and they pull in opposite directions.

First, “no evidence” has three different meanings, and this guide tries to always tell you which one applies. *Tested and failed:* a real trial ran and the answer was no. *Never properly tested:* nobody could afford the trial, so absence of evidence is not evidence of absence. *Tested small and won:* real but modest trials, usually measuring markers rather than outcomes, because markers are what a small budget can afford.

Second, the asymmetry earns nutrients humility, not automatic credit. When isolated nutrients have received large publicly funded trials, the results have been mixed: some genuine wins, some nulls, and some harms. Small nutrient trials also carry their own publication bias toward positive results, just as drug trials tilt toward their sponsor. So this guide applies one rule in both directions: name the funding, name the evidence tier, and let you see both.

Where supplements actually fit in reflux

Reflux has five drivers, and this kit is organized around them: **the barrier** (the valve and the diaphragm), **the signal** (vagal tone, stress, sleep), **the terrain** (the microbial environment), **the load** (meals, weight, timing), and **the capacity** (whether the stomach is doing its job).

Here is the honest placement of the products in this guide: **most of them act on the consequence, not on a driver**. Acid is what makes reflux hurt. It is almost never what makes reflux happen. Soothing and neutralizing has real value when you are in pain, and it does not repair a barrier or retrain a nervous system.

That is not a counsel of despair, because two of the five drivers respond to things you can do yourself, for free, starting today. They come first in this guide for that reason, ahead of anything you can buy.

The two acid-suppressing realities to keep in view. Up to half of people remain symptomatic despite adequate acid suppression, which is itself the argument that reflux is multifactorial rather than acid-centric (PMID 42198379). And the interventions with the best evidence in the whole kit, weight loss and head-of-bed elevation, are dietary and mechanical rather than pharmacological (PMID 16682569).

So the order is: work the barrier, work the signal, fix the terrain if it is disturbed, reduce the load, and use the products below to make all of that survivable while it takes effect.

How to read this guide

The tiers. Tier 1 has randomized human evidence in reflux specifically. Tier 2 has evidence in adjacent conditions such as functional dyspepsia, or in reflux with real limitations. Tier 3 needs a prescriber conversation first.

Track two numbers daily: symptom severity out of ten, and how many doses of any rescue product you took. The second number moves before the first.

One change at a time, and give most of these four weeks.

Foundation 1: retrain the diaphragm, because it is half of your sphincter

This is first because it is free, because it acts on the mechanism rather than the symptom, and because most people have never been told it.

The physiology. The anti-reflux barrier has three parts: the lower esophageal sphincter, the crural diaphragm wrapped around it, and the flap valve at the entry angle (PMID 40412989). The crural diaphragm functions as an external sphincter, raising sphincter pressure with every breath in and during any rise in abdominal pressure (PMID 15316404). Most reflux happens through transient sphincter relaxations rather than a permanently open valve, and crural support is a real part of what holds the line (PMID 11003803).

The trial. In patients with pH-proven upright reflux, diaphragmatic breathing reduced the number of postprandial reflux events by increasing the difference between lower esophageal sphincter and gastric pressure. Esophageal acid exposure in the two hours after a standardized meal fell from 11.8% to 5.2% (PMID 33009052). A 2026 systematic review and meta-analysis found a significant reduction in symptom scores favoring diaphragmatic breathing, while stating plainly that heterogeneity was substantial and the evidence is not yet sufficient for definitive recommendations (PMID 42123139).

A structured inspiratory muscle training trial aimed at this exact mechanism is registered and published as a protocol, with results not yet reported (PMID 41874149). Promising, pending, and I would rather say so.

The practice. Sitting or lying, one hand on the chest and one on the abdomen. Breathe so the lower hand rises and the upper hand stays still. Exhale longer than you inhale. Five to ten minutes, twice daily, and **the highest-value window is the half hour after a meal**, because that is when reflux events cluster.

The structural half, which is my own field. If the diaphragm is half the sphincter, then thoracic mobility, rib position, posture, and how you manage abdominal pressure under load are all inputs to the valve. Bending, lifting badly, straining, tight waistbands, and a slumped seated posture all raise the pressure your barrier has to hold against. That is why the postural chapter of the ebook is not filler, and why a structural assessment belongs in a reflux workup.

Foundation 2: work the nervous system, which sets the volume

Why it belongs in a supplement guide. Because it outperformed most of the products below in the one trial that measured an objective endpoint.

In a controlled study of 20 subjects with documented reflux disease, relaxation training during a stressful task produced significantly lower heart rate and anxiety ratings than an attention-placebo control, and significantly lower reflux symptom ratings **and lower total esophageal acid exposure** (PMID 8020690). A relaxation intervention moved measured acid exposure, not just how people felt about it.

Why it works. Esophageal hypervigilance is present across reflux presentations regardless of acid burden and independently predicts symptom severity (PMID 33432708). Stress increases symptom ratings, and the authors of one pH-monitored study specifically note that symptom perception in the absence of increased reflux may explain poor responses to standard treatment (PMID 16310024). The honest counterweight: a short experimental stressor did not increase acid perception in another study (PMID 19453516), so chronic load has better support than acute stress does.

Sleep is the other half of this driver, and it runs in both directions with reflux: poor sleep quality, sleep disturbance, and short sleep duration are all associated with reflux, bidirectionally (PMID 38646475). If you snore or wake unrefreshed, ask about sleep apnea, which is linked to reflux through airway irritation and vagal pathways (PMID 40045407). Treating apnea can improve reflux, and nothing in this guide substitutes for that.

What to actually do. A daily relaxation practice you will repeat, which matters more than which one. The breathing practice above does double duty here, since it is both a mechanical input to the diaphragm and a vagal one. Protect sleep as a treatment rather than a luxury. And read the eating mechanics section of the diet guide, which applies the same driver at the table.

Then the products: Tier 1, studied in reflux

Alginate

What it is. A seaweed-derived compound that forms a physical raft floating on top of the stomach contents. It is the one item in this guide that acts mechanically rather than chemically, which is why I put it first.

What the research shows. A systematic review and meta-analysis of 14 studies covering 2,095 subjects found alginate-based therapies increased the odds of resolution of reflux symptoms compared with placebo or antacids, with an odds ratio of 4.42 (95% CI 2.45 to 7.97), with moderate heterogeneity. Compared with PPIs or H2 blockers, alginates appeared less effective, though that pooled estimate was not statistically significant (PMID 28375448).

A second systematic review and meta-analysis of randomized trials found alginates may have greater efficacy than placebo or antacids, described the evidence on alginate versus PPI as questionable and suggesting no difference, and found the risk of adverse events no greater than placebo or PPIs (PMID 33275256). A broader meta-analysis of over-the-counter reflux therapies found alginate and antacid combinations superior to placebo in symptomatic improvement, with an absolute benefit increase of 26% (PMID 17229239).

How to read that. Better than placebo and better than plain antacids, with a clean safety profile, and probably not equal to a PPI. That is a genuinely useful place in the toolkit, particularly for breakthrough symptoms and for people trying to reduce acid suppression.

Typical use. After meals and at bedtime, which is when the raft is most useful. It works best taken after eating rather than before.

Cautions. Most products carry a meaningful sodium load, which matters in heart failure or on a sodium-restricted diet. It can bind other medications, so separate it by two hours.

Melatonin

What it is. Better known for sleep, and it also has effects on gastric acid secretion and on the lower esophageal sphincter.

What the research shows. In a randomized double-blind trial, 78 patients with reflux received omeprazole 20 mg daily plus either sublingual melatonin 3 mg or placebo for four weeks. Symptom scores declined significantly more in the melatonin group, quality of life scores were significantly higher, and adverse events were similar between groups with none serious (PMID 37768310).

An earlier comparative study tested a supplement containing melatonin, l-tryptophan, vitamin B6, folic acid, vitamin B12, methionine, and betaine against omeprazole 20 mg in 351 patients. All 176 patients in the supplement group reported complete regression of symptoms after 40 days, compared with 65.7% in the omeprazole group (PMID 16948779).

How to read that second study honestly. A 100% response rate is an extraordinary claim, it was a multi-ingredient formula so no single component can be credited, and it has not been replicated at that magnitude. I include it because it exists and because the melatonin literature in reflux rests partly on it, not because I expect you to get that result.

Typical studied range. 3 mg sublingually at night, as an addition to standard treatment (PMID 37768310).

Cautions. Next-morning grogginess usually means too high a dose. It adds to other sedating medication. Not established in pregnancy or nursing.

Tier 2: evidence in adjacent conditions, or with real limits

Iberogast and similar multi-herb formulas

What it is. A standardized nine-herb liquid preparation used for decades in functional dyspepsia, built around bitter candytuft with chamomile, peppermint, caraway, licorice, lemon balm, celandine, angelica, and milk thistle.

What the research shows. A narrative review describes it as an evidence-based phytomedicine with a favorable safety profile, addressing motility, hypersensitivity, inflammation, and barrier function, and specifically names overlapping dyspepsia and reflux phenotypes as an area for future research (PMID 41319263). In a randomized, double-blinded, placebo-controlled study of 32 patients with functional dyspepsia and bloating, two weeks of the STW 5-II formulation produced significantly lower symptom perception during a gastric gas challenge than placebo, with no difference in abdominal distension (PMID 40684457).

How to read it. Good evidence in functional dyspepsia, which overlaps with reflux without being the same condition. Reasonable if bloating and fullness after meals are as prominent as burning.

Cautions. It contains peppermint, which is traditionally avoided in reflux on sphincter-relaxation grounds, so this is one to test rather than assume. Regulatory authorities have flagged rare liver injury with this class of preparation, so stop and call for jaundice, dark urine, or right-sided abdominal pain. Contains licorice and alcohol in the liquid form.

Curcumin

What it is. The active fraction of turmeric. Poorly absorbed on its own, which is why formulation matters more here than for almost anything else in this guide.

What the research shows. A randomised, double blind controlled trial in Thailand assigned 206 patients with functional dyspepsia to curcumin alone, omeprazole alone, or both, for 28 days, with follow-up to day 56. Symptoms were scored with the Severity of Dyspepsia Assessment. All three groups improved significantly on the pain, non-pain, and satisfaction categories at day 28, and improved further by day 56. There were **no significant differences between the three groups**, no synergistic effect from combining them, and no serious adverse events (PMID 37696679).

Headline version: curcumin performed comparably to omeprazole. That is a striking result for a botanical, and it is the reason this entry exists.

Now the caveat that decides how much weight to give it, and it is the one every summary of this trial leaves out.

There was **no placebo arm**. All three groups got an active treatment, and all three improved. Functional dyspepsia has a large placebo response and a naturally fluctuating course, so a three-arm trial in which everyone improves and nobody differs cannot separate “both treatments work equally well” from “both treatments are tracking the same placebo and regression-to-the-mean effect.” Those two readings are very different and this design does not distinguish them.

Two more things worth knowing. 206 patients enrolled and 151 completed, so roughly a quarter left the study, which is high. And the paper carries a published correction from 2025 (PMID 38866470). A correction is not a retraction and the record is in good standing, but you should hear it from me.

Why it is in Tier 2 rather than Tier 1. The trial was in **functional dyspepsia**, not reflux. Those conditions overlap and are not the same: functional dyspepsia is epigastric pain and fullness, reflux is heartburn and regurgitation. Same reason Iberogast sits in this tier.

Typical studied range. 250 mg four times daily, taken as two capsules per dose, giving 2 g daily for 28 days (PMID 37696679). That is a higher and more frequent dose than most people take, and it is what was tested.

Cautions. Curcumin increases bleeding risk with anticoagulants and antiplatelet drugs, and should be stopped before surgery. It can cause reflux and stomach upset in some people, which matters in this guide specifically. It interacts with several drugs through the same liver enzymes, and it is not established in pregnancy. Plain turmeric powder is not the same as the standardised extract used in the trial.

Zinc L-carnosine

What it is. A zinc compound with an unusual property: it binds preferentially to damaged mucosa and releases zinc locally.

What the research shows. A review of its role in gastrointestinal mucosal disease describes direct mucosal cytoprotective and anti-inflammatory action through antioxidative effects, cytokine modulation, and membrane stabilization, notes that it is not an antacid or acid-suppressing agent, and describes promotion of repair of mucosal injury in humans across several settings (PMID 35659631).

How to read it. Mucosal repair rather than symptom suppression, and the human literature is broader in gastric injury than in reflux specifically. Sensible where the goal is healing an irritated lining rather than blocking acid.

Cautions. Zinc long-term can deplete copper. Do not run high-dose zinc indefinitely without monitoring.

Ginger

What the research shows. In a randomized double-blind crossover study of 11 patients with functional dyspepsia, ginger accelerated gastric emptying, with median half-emptying time of 12.3 minutes versus 16.1 minutes on placebo, and showed a trend toward more antral contractions, while **gastrointestinal symptoms did not differ** (PMID 21218090). A pilot randomized crossover study in 11 healthy volunteers found a standardized ginger and artichoke combination significantly promoted gastric emptying compared with placebo (PMID 26813467).

How to read it. It moves the stomach along, which is a sensible mechanism if slow emptying is part of your picture. It did not change symptoms in the dyspepsia trial, and both studies had 11 participants. Cheap and low risk, with modest expectations.

Cautions. Can increase bleeding risk at higher doses with anticoagulants. Commonly causes heartburn in some people, which is worth noting in a reflux guide.

D-limonene

Widely sold for heartburn. The most-cited source is a review describing it as generally recognized as safe as a flavoring agent, with low toxicity, and noting that because of its gastric acid neutralizing effect and support of normal peristalsis it has been used for relief of heartburn and reflux (PMID 18072821).

Be clear about what that is. It is a safety and applications review describing use, not a randomized trial demonstrating benefit. I could not locate controlled trial evidence in reflux. Low risk, plausible, unproven. If you try it, know which of those three words is doing the work.

Probiotics, mainly as an adjunct to *H. pylori* treatment

This is where the probiotic evidence in this category is strongest, and it is about tolerating the antibiotics rather than treating reflux.

A meta-analysis of *Saccharomyces boulardii* as an adjuvant to standard eradication therapy found it significantly improved eradication rates and reduced total adverse effects (RR 0.49), diarrhea (RR 0.36), abdominal distension (RR 0.49), constipation (RR 0.38), and nausea (RR 0.50), while not reducing abdominal pain, vomiting, or taste disturbance (PMID 40012609). A meta-analysis of eight randomized trials in 1,087 patients found *Lactobacillus reuteri* supplementation tended to increase eradication rates, reduce antibiotic-related side effects, and alleviate gastrointestinal symptoms during treatment (PMID 38846173).

Practical use. If you are going through eradication therapy, this is worth raising with your prescriber. It is not a reflux treatment on its own.

Tier 3: ask your prescriber first

Betaine HCl and the low stomach acid question

The lab guide covers the evidence in full. The summary: the strongest published case for replacing lost stomach acid is made specifically for **autoimmune gastritis**, a defined condition with documented destruction of the acid-producing cells, where the argument is that B12 and iron are routinely replaced while the missing acid is not, and where re-acidification may improve gastrointestinal symptoms (PMID 38474790).

Where that leaves betaine HCl. It has a coherent rationale in someone with demonstrated hypochlorhydria. It does not have trial evidence supporting its use across reflux generally, and I am not going to pretend otherwise in a document you paid for.

Why it is in Tier 3 rather than Tier 1. Because if you have an ulcer, erosive esophagitis, or active gastritis, adding acid can hurt you, and those are exactly the conditions that produce the symptoms leading people to try it. **Do not self-start betaine HCl if you take NSAIDs, aspirin, or steroids, if you have a known ulcer, or if you have alarm features.** If low acid is genuinely suspected, the honest sequence is to demonstrate it first with the workup in the lab guide, then decide with a clinician.

Deglycyrrhizinated licorice

Traditionally used for mucosal soothing. I searched for randomized evidence in reflux and functional dyspepsia and did not find trials that would justify a strong claim, so it sits here as low risk and unproven rather than as a recommendation. The deglycyrrhizinated form exists specifically to remove the compound that raises blood pressure and lowers potassium, so if you use licorice, use that form. Standard licorice is not interchangeable and matters with hypertension, heart disease, and diuretics.

Coming off a PPI, which is what most readers actually came here for

This deserves its own section because it is where the most harm gets done by well-meaning advice, in both directions.

The terrain argument, stated fairly. A narrative review from a microbial ecology perspective notes that up to 50% of people remain symptomatic despite adequate acid suppression, describes reflux as multifactorial rather than acid-centric, and examines how long-term acid suppression alters the microbial ecology of the upper gut, while being explicit that causality is not fully established (PMID 42198379). That is a reasonable argument for not staying on acid suppression by default. It is not an argument for stopping one abruptly, and the distinction matters.

First, the honest evidence on rebound. In a randomized, double-blind, placebo-controlled trial, 120 **healthy volunteers with no reflux** received either 12 weeks of placebo or 8 weeks of esomeprazole 40 mg followed by 4 weeks of placebo. In weeks 9 to 12, significantly more of the PPI group reported dyspepsia, heartburn, or acid regurgitation than the placebo group, around 21% to 22% versus 2% to 7% (PMID 19362552).

Read that carefully: people who never had reflux developed reflux symptoms after stopping a PPI. That is why stopping feels like proof you still need it when it may be proof of nothing.

And the counterweight, because it exists. A systematic review of rebound acid hypersecretion found the studies heterogeneous in design and outcome, with some evidence from uncontrolled trials of increased acid-secreting capacity in *H. pylori*-negative subjects after 8 weeks, and concluded there is **no strong evidence for a clinically relevant increased acid production** after withdrawal (PMID 17229219).

So the symptoms on withdrawal are well demonstrated; whether the underlying acid physiology is truly rebounding is less settled. Either way, the practical implication is the same.

How deprescribing is actually done. A 2025 evidence-based review of PPI indications, harms, and deprescribing describes the standard approaches: tapering the dose, switching to on-demand or intermittent use, and switching to an H2 receptor antagonist, particularly in people with non-erosive reflux disease or functional dyspepsia, with formal tools available to identify candidates for tapering (PMID 41010960).

A structure to bring to your prescriber, not to run alone:

1. **Confirm you are a candidate.** Not if you have Barrett's, a history of bleeding ulcer, or need gastroprotection with NSAIDs or antiplatelet therapy.
2. **Get the foundations in place first**, for at least two to four weeks: the head of the bed raised, three hours between eating and lying down, weight addressed if relevant, alcohol reduced, and the breathing practice started.
3. **Taper rather than stop.** Halve the dose, or move to alternate days, over several weeks.
4. **Have a bridge ready.** This is what alginate is for, and an H2 blocker is the other option your prescriber may use.
5. **Expect two to four weeks of worse symptoms**, and know in advance that this is the documented withdrawal pattern rather than proof of failure (PMID 19362552).
6. **Reassess honestly.** If symptoms genuinely return and stay, the medication was doing a job. That is a legitimate outcome, not a personal failure.

What I deliberately left out, and why

- **Baking soda as a routine remedy.** Fast relief, large sodium load, rebound, and rare reports of serious harm with large amounts. Alginate does the same job better.
- **Apple cider vinegar.** Widely recommended on the low-acid theory. No randomized reflux evidence located, and a real potential to hurt if there is an ulcer or erosive disease.
- **Aloe vera juice.** Some early work in reflux exists, and the products vary enormously and some have a laxative effect from the latex fraction. Not enough to recommend by name.
- **Slippery elm and marshmallow root.** Soothing, harmless, traditional. I could not locate randomized reflux trials, so they are named here honestly rather than sold to you.
- **Digestive enzyme blends for everyone.** Reasonable in demonstrated insufficiency, which is a different diagnosis. Not a general reflux answer.
- **Stopping your PPI cold because a supplement guide said acid is good.** The most common harm in this category.

Telling you what to skip is part of what you paid for.

Realistic expectations

What is realistic. Fewer breakthrough episodes, fewer rescue doses, sleeping through the night, and for some people getting off or down on daily medication with a plan and a prescriber.

What is not realistic. A supplement fixing the mechanics. If the sphincter and the anatomy are the problem, chemistry only manages it.

Timeline. Four weeks for most of these. The mechanical changes, meal timing and bed elevation, often work within days, which is another reason to do them first.

Cost sense. If a product has done nothing after four weeks, stop paying for it. Alginate and a wedge under the mattress cost less than most of the shelf.

When to seek professional care

Emergency: chest pain with sweating, nausea, breathlessness, or pain to the jaw or arm. Vomiting blood or black tarry stools. Inability to swallow saliva.

Soon:

- Difficulty swallowing or food sticking, ever
- Unintentional weight loss, persistent vomiting, or anemia
- New symptoms at 60 or over
- Daily symptoms for more than a few weeks despite treatment
- Many years of untreated reflux, which is a Barrett's screening conversation
- Any plan to come off long-term acid suppression

Stop and call your prescriber for: yellowing of the skin or eyes, dark urine, or right-sided abdominal pain while taking a multi-herb preparation. Also for new black stools, vomiting blood, or worsening pain after starting anything acidic.

Where to find quality versions of these

Nothing here names a brand, on purpose. The evidence supports ingredients and forms, not labels.

Look for third-party testing. NSF, USP Verified, or an equivalent independent seal.

Check the form against the studied one. Sublingual melatonin at 3 mg. Alginate rather than a plain antacid, and check the sodium content. Zinc L-carnosine rather than generic zinc, since the compound is the point.

Read the whole label. Multi-herb digestive formulas frequently include peppermint, which is one of the things you may be trying to avoid.

If you would rather not evaluate all of that yourself, I keep a curated dispensary of professional-grade products at Root Health, screened for the standards above and organized by the categories in this guide.

Root Health dispensary: us.fullscript.com/welcome/drseay

Disclosure: Root Health earns a portion of dispensary purchases. That revenue supports the practice, and it is also why this guide was written to be complete and usable on its own. Nothing here was included, excluded, ranked, or dosed based on what it earns. You can buy any of it anywhere, and the highest-value items in this guide, raising the head of your bed, eating earlier, and the breathing practice, cost nothing at all.

Working with Root Health

Everything here is general education, written to be safe for a reader I have never examined. A personalized plan starts from your history, your test results, and your full medication list.

***Book a free Root Discovery Session: [rootcauseunlocked.com/discovery?](https://rootcauseunlocked.com/discovery?utm_source=guide&utm_medium=ebook&utm_campaign=reflux-supplement)
[utm_source=guide&utm_medium=ebook&utm_campaign=reflux-supplement](https://rootcauseunlocked.com/discovery?utm_source=guide&utm_medium=ebook&utm_campaign=reflux-supplement)***

Consultations and ongoing care are both available wherever you are located. Being outside North Carolina is not a barrier to working with me. Tell me where you are when you reach out and I will lay out exactly how we would work together from there.

***Get in touch: [rootcauseunlocked.com/contact?](https://rootcauseunlocked.com/contact?utm_source=guide&utm_medium=ebook&utm_campaign=reflux-supplement)
[utm_source=guide&utm_medium=ebook&utm_campaign=reflux-supplement](https://rootcauseunlocked.com/contact?utm_source=guide&utm_medium=ebook&utm_campaign=reflux-supplement)***

If you would rather start on your own, track first:

***Free symptom tracker: [rootcauseunlocked.com/tracker?](https://rootcauseunlocked.com/tracker?utm_source=guide&utm_medium=ebook&utm_campaign=reflux-supplement)
[utm_source=guide&utm_medium=ebook&utm_campaign=reflux-supplement](https://rootcauseunlocked.com/tracker?utm_source=guide&utm_medium=ebook&utm_campaign=reflux-supplement)***

Read the companion documents alongside this one. **Understanding Your Reflux Labs** covers H. pylori, the imitators, and the workup for genuinely low stomach acid, and **The Reflux Diet Guide** covers the two measures with the best evidence of anything in this kit.

References

All references were verified against live PubMed records on 2026-08-08, including a publication type and corrections check for retractions and errata. Verification detail is documented in `Reflux-Kit-Citation-Ledger.md`.

1. Kaltenbach T, Crockett S, Gerson LB. Are lifestyle measures effective in patients with gastroesophageal reflux disease? An evidence-based approach. *Arch Intern Med*. 2006;166(9):965-971. PMID 16682569
2. Leiman DA, Riff BP, Morgan S, et al. Alginate therapy is effective treatment for GERD symptoms: a systematic review and meta-analysis. *Dis Esophagus*. 2017;30(5):1-9. PMID 28375448
3. Zhao CX, Wang JW, Gong M. Efficacy and safety of alginate formulations in patients with gastroesophageal reflux disease: a systematic review and meta-analysis of randomized controlled trials. *Eur Rev Med Pharmacol Sci*. 2020;24(22):11845-11857. PMID 33275256
4. Tran T, Lowry AM, El-Serag HB. Meta-analysis: the efficacy of over-the-counter gastro-oesophageal reflux disease therapies. *Aliment Pharmacol Ther*. 2007;25(2):143-153. PMID 17229239
5. Malekpour H, Noori A, Abdi S, et al. Is the Addition of Sublingual Melatonin to Omeprazole Superior to Omeprazole Alone in the Management of Gastroesophageal Reflux Disease Symptoms: A Clinical Trial. *Turk J Gastroenterol*. 2023;34(12):1206-1211. PMID 37768310
6. Pereira Rde S. Regression of gastroesophageal reflux disease symptoms using dietary supplementation with melatonin, vitamins and aminoacids: comparison with omeprazole. *J Pineal Res*. 2006;41(3):195-200. PMID 16948779
7. Halland M, Bharucha AE, Crowell MD, et al. Effects of Diaphragmatic Breathing on the Pathophysiology and Treatment of Upright Gastroesophageal Reflux: A Randomized Controlled Trial. *Am J Gastroenterol*. 2021;116(1):86-94. PMID 33009052
8. Abureesh O, Qaqish F, Abu-Shaban M, et al. Efficacy and Safety of Diaphragmatic Breathing Exercises for Gastroesophageal Reflux Disease: A Systematic Review and Meta-Analysis. *J Clin Med*. 2026;15(9). PMID 42123139
9. Lambiase C, Usai Satta P, Bassotti G, et al. Iberogast in functional dyspepsia: yesterday, today, and tomorrow, a narrative review of a multitarget phytomedicine. *Expert Opin Pharmacother*. 2025;26(17):1867-1876. PMID 41319263
10. Aguilar A, Alcala-Gonzalez L, Barber C, et al. Effect of STW 5-II (Iberogast-N) on Tolerance to Gastric Gas in Patients With Functional Dyspepsia. The IBO-2 Study. *Neurogastroenterol Motil*. 2025;37(11):e70123. PMID 40684457
11. Efthymakis K, Neri M. The role of Zinc L-Carnosine in the prevention and treatment of gastrointestinal mucosal disease in humans: a review. *Clin Res Hepatol Gastroenterol*. 2022;46(7):101954. PMID 35659631
12. Hu ML, Rayner CK, Wu KL, et al. Effect of ginger on gastric motility and symptoms of functional dyspepsia. *World J Gastroenterol*. 2011;17(1):105-110. PMID 21218090
13. Lazzini S, Polinelli W, Riva A, et al. The effect of ginger (*Zingiber officinalis*) and artichoke (*Cynara cardunculus*) extract supplementation on gastric motility: a pilot randomized study in healthy volunteers. *Eur Rev Med Pharmacol Sci*. 2016;20(1):146-149. PMID 26813467
14. Sun J. D-Limonene: safety and clinical applications. *Altern Med Rev*. 2007;12(3):259-264. PMID 18072821
15. Li M, Xie Y. Efficacy and safety of *Saccharomyces boulardii* as an adjuvant therapy for the eradication of *Helicobacter pylori*: a meta-analysis. *Front Cell Infect Microbiol*. 2025;15:1441185. PMID 40012609
16. Li M, Wang X, Dong X, et al. *Lactobacillus reuteri* compared with placebo as an adjuvant in *Helicobacter pylori* eradication therapy: a meta-analysis of randomized controlled trials. *Ther Adv Gastroenterol*. 2024;17:17562848241258021. PMID 38846173

17. Taylor L, McCaddon A, Wolffenbuttel BHR. Creating a Framework for Treating Autoimmune Gastritis, The Case for Replacing Lost Acid. *Nutrients*. 2024;16(5):662. PMID 38474790
18. Reimer C, Sondergaard B, Hilsted L, et al. Proton-pump inhibitor therapy induces acid-related symptoms in healthy volunteers after withdrawal of therapy. *Gastroenterology*. 2009;137(1):80-87. PMID 19362552
19. Hunfeld NG, Geus WP, Kuipers EJ. Systematic review: Rebound acid hypersecretion after therapy with proton pump inhibitors. *Aliment Pharmacol Ther*. 2007;25(1):39-46. PMID 17229219
20. Andrawes M, Andrawes W, Das A, et al. Proton Pump Inhibitors, An Evidence-Based Review of Indications, Efficacy, Harms, and Deprescribing. *Medicina (Kaunas)*. 2025;61(9). PMID 41010960
21. Kahrilas PJ, Nguyen NT. The Esophagogastric Junction Reconsidered. *Gastrointest Endosc Clin N Am*. 2025;35(3):541-553. PMID 40412989
22. van Herwaarden MA, Samsom M, Smout AJ. The role of hiatus hernia in gastro-oesophageal reflux disease. *Eur J Gastroenterol Hepatol*. 2004;16(9):831-835. PMID 15316404
23. Holloway RH. The anti-reflux barrier and mechanisms of gastro-oesophageal reflux. *Baillieres Best Pract Res Clin Gastroenterol*. 2000;14(5):681-699. PMID 11003803
24. Syropoulos S, Moutzouri M, Grammatopoulou E, et al. Impact of Inspiratory Muscle Training in Individuals with Gastroesophageal Reflux Disease: A Randomized Controlled Trial Protocol. *Methods Protoc*. 2026;9(2). PMID 41874149
25. McDonald-Haile J, Bradley LA, Bailey MA, et al. Relaxation training reduces symptom reports and acid exposure in patients with gastroesophageal reflux disease. *Gastroenterology*. 1994;107(1):61-69. PMID 8020690
26. Guadagnoli L, Yadlapati R, Taft T, et al. Esophageal hypervigilance is prevalent across gastroesophageal reflux disease presentations. *Neurogastroenterol Motil*. 2021;33(8):e14081. PMID 33432708
27. Wright CE, Ebrecht M, Mitchell R, et al. The effect of psychological stress on symptom severity and perception in patients with gastro-oesophageal reflux. *J Psychosom Res*. 2005;59(6):415-424. PMID 16310024
28. Hemmink GJ, Bredenoord AJ, Weusten BL, et al. Does acute psychological stress increase perception of oesophageal acid? *Neurogastroenterol Motil*. 2009;21(10):1055-e86. PMID 19453516
29. Tan X, Wang S, Wu F, et al. Bidirectional correlation between gastroesophageal reflux disease and sleep problems: a systematic review and meta-analysis. *PeerJ*. 2024;12:e17202. PMID 38646475
30. Wang J, Wang P, Lv J, et al. Exploring the silent connection: unveiling the intricate relationship between gastroesophageal reflux disease and sleep apnea syndrome. *Hum Genomics*. 2025;19(1):23. PMID 40045407
31. Zaroni A, Facchin S, Mari V, et al. Rethinking Long-Term PPI Therapy in GERD: A Narrative Review from a Microbial Ecology Perspective Beyond Acid Suppression. *Pharmaceuticals (Basel)*. 2026;19(5). PMID 42198379
32. Kongkam P, Khongkha W, Lopimpisuth C, et al. Curcumin and proton pump inhibitors for functional dyspepsia: a randomised, double blind controlled trial. *BMJ Evid Based Med*. 2023;28(6):399-406. PMID 37696679
33. Lundh A, Lexchin J, Mintzes B, et al. Industry sponsorship and research outcome. *Cochrane Database Syst Rev*. 2017;2(2):MR000033. PMID 28207928

Dr. Adam Seay, DC · Seay Wellness / Root Health, Garner, NC
 rootcauseunlocked.com · (919) 662-0520

Educational content only. Not medical advice. Not a diagnosis. Not a substitute for your clinician. Never stop or change a prescribed medication without your prescriber.