



ROOT CAUSE UNLOCKED

# Unlock Your Chemical *Sensitivity*

*A root-cause, terrain-informed roadmap for reducing exposures,  
understanding symptoms, and rebuilding resilience*

Dr. Adam Seay, DC

Root Cause Unlocked, Garner, NC · [rootcauseunlocked.com](http://rootcauseunlocked.com) · (919) 662-0520

# Unlock Your Chemical Sensitivity

*A root-cause, terrain-informed roadmap for reducing exposures, understanding symptoms, and rebuilding resilience*

BEFORE YOU START

# Contents

Important Medical and Legal Disclaimer

---

How to Use This Book

---

Contents

---

## INTRODUCTION

The Terrain Underneath Chemical Sensitivity

---

1. Understanding Multiple Chemical Sensitivity

---

2. Symptoms, Triggers, and Patterns

---

3. What May Be Happening in the Body

---

4. Building a Sound Medical Evaluation

---

5. Environmental and Functional Testing

---

6. Creating a Lower-Exposure Home

---

7. Nutrition for Stability and Resilience

---

8. Supplements, Binders, Sauna, and Detox Claims

---

9. Calming a Sensitized Nervous System

---

10. Work, Healthcare, Travel, and Relationships

---

11. A Personalized 90-Day Recovery Roadmap

---

12. Worksheets and Supplemental Support

---

Closing Perspective

---

Working with Root Health

---

References

---

# Important Medical and Legal Disclaimer

---

This book teaches the approach I use with patients who live with chemical sensitivity. It is intended for education and general wellness support. It does not diagnose Multiple Chemical Sensitivity, Idiopathic Environmental Intolerance, allergy, toxic exposure, mold-related illness, mast cell disease, or any other medical condition. It is not a substitute for examination, diagnosis, or treatment by a qualified healthcare professional, and reading it does not make you my patient.

Symptoms attributed to chemical exposure may also occur with asthma, migraine, heart rhythm disorders, carbon monoxide exposure, medication reactions, endocrine disease, anemia, infection, anxiety and panic, vestibular disorders, occupational poisoning, or other conditions that require medical attention. New, severe, rapidly worsening, or unexplained symptoms should be evaluated promptly. Ruling these out is not a delay tactic. It is the foundation everything else in this book is built on.

Do not stop prescribed medication, begin chelation, use binders, undertake fasting, start high-dose supplements, or begin aggressive sauna or detoxification programs without professional guidance. Pregnancy, breastfeeding, kidney disease, liver disease, cardiovascular disease, diabetes, seizure disorders, eating disorders, medication use, and advanced age can materially change the safety of interventions discussed in this book.

The scientific understanding of Multiple Chemical Sensitivity remains incomplete. I am confident in the terrain-based, nervous-system, and environmental approach I describe here, and I am equally clear about distinguishing established environmental health guidance from emerging hypotheses and patient-reported approaches. No outcome is promised or guaranteed.

# How to Use This Book

---

You do not need to implement every strategy at once. Rapid, sweeping changes can increase cost, anxiety, nutritional restriction, and uncertainty about what is actually helping. I ask patients to use this book in three passes.

Start with Chapters 1 through 4. They explain the condition, the honest uncertainty surrounding it, and why a thoughtful medical evaluation has to happen before symptoms are attributed to chemical sensitivity alone.

After that foundation is in place, Chapters 5 through 9 help you choose a small number of practical, low-risk supports. The 90-day roadmap and worksheets in Chapters 11 and 12 are designed to help you track changes in daily function, not just symptom intensity, and to review your progress with a qualified clinician.

**A guiding principle.** The goal is not to prove that every symptom came from a toxin. The goal is to reduce meaningful exposures, identify treatable conditions, improve stability, and help you regain as much life as possible.

# Contents

---

- Introduction: The Terrain Underneath Chemical Sensitivity
- 1. Understanding Multiple Chemical Sensitivity
- 2. Symptoms, Triggers, and Patterns
- 3. What May Be Happening in the Body
- 4. Building a Sound Medical Evaluation
- 5. Environmental and Functional Testing
- 6. Creating a Lower-Exposure Home
- 7. Nutrition for Stability and Resilience
- 8. Supplements, Binders, Sauna, and Detox Claims
- 9. Calming a Sensitized Nervous System
- 10. Work, Healthcare, Travel, and Relationships
- 11. A Personalized 90-Day Recovery Roadmap
- 12. Worksheets and Supplemental Support
- Working with Root Health
- References

# The Terrain Underneath Chemical Sensitivity

---

Imagine entering a store, medical office, workplace, or family gathering and becoming ill because someone nearby is wearing perfume, a floor has recently been cleaned, or a room contains new furniture. A smell that barely registers to another person may be followed by headache, dizziness, chest tightness, nausea, cognitive difficulty, exhaustion, or a racing heartbeat.

I see this pattern in my practice regularly, and I want to say plainly what I believe about it: the reaction is real, and it is rarely explained by the room alone. In the patients I work with, chemical sensitivity almost always sits on top of a terrain that has already been destabilized. An autonomic nervous system running hot. A gut and immune system carrying more inflammatory load than they can quietly absorb. Mitochondrial and detox capacity that is already stretched thin. Blood sugar and sleep that are not giving the nervous system a chance to reset. And often an initiating exposure or illness that never fully resolved. Reducing exposures matters. Addressing that terrain is what changes how much exposure a person can tolerate before it fires.

I want to be equally direct about the other side of that confidence. The scientific understanding of MCS is genuinely incomplete, and I am not going to pretend otherwise anywhere in this book. Where the evidence is solid, I say so. Where an idea is a reasonable, influential hypothesis and nothing more, I label it that way, every time. That distinction is not hedging. It is what keeps a root-cause approach honest and keeps you safe.

For people living with Multiple Chemical Sensitivity, these reactions can reshape nearly every part of life. Ordinary environments become unpredictable. Work can become difficult or impossible. Social relationships may suffer. Even seeking healthcare can be challenging when clinics use fragranced products, disinfectants, or recently renovated materials. Qualitative research confirms that people who identify as chemically sensitive can experience major occupational and social losses, including isolation and difficulty maintaining employment.<sup>1,2</sup>

Multiple Chemical Sensitivity, commonly abbreviated MCS, is also described as chemical intolerance or Idiopathic Environmental Intolerance, abbreviated IEI. Some clinicians use Toxicant-Induced Loss of Tolerance, or TILT, for a proposed pattern in which a major exposure or accumulated exposures are followed by reactions to much smaller amounts of multiple unrelated substances. These terms overlap, but they are not perfectly interchangeable.

The terminology remains debated because MCS does not have one universally accepted definition, confirmatory laboratory test, biomarker, or proven biological mechanism. Reviews continue to describe uncertainty about its causes, diagnostic boundaries, and treatment.<sup>3,4</sup> A major systematic review initiated in 2025 was designed specifically to reassess the validity, prevalence, measurement tools, mechanisms, and interventions associated with MCS, which shows how many basic questions remain open.<sup>5</sup>

Scientific uncertainty does not mean symptoms are imaginary. Headache, breathlessness, dizziness, cognitive impairment, nausea, pain, fatigue, and palpitations are genuine physical experiences that deserve careful assessment and compassionate care. Patients also deserve protection from unsupported claims, unnecessary laboratory testing, extreme avoidance, and aggressive detoxification programs that can be costly or harmful. I hold both of those things at once in every case I take on.

This book takes a middle path, and I take it deliberately. It does not reduce chemical sensitivity to “all in your head.” It also does not assume that every reaction is caused by stored toxins, liver failure, mold, heavy metals, or one genetic variant. Instead, I offer a structured process for identifying patterns, ruling out treatable conditions, reducing meaningful exposures, supporting physiological stability, and rebuilding function.

**What recovery can mean.** Recovery may mean fewer reactions, shorter flares, clearer thinking, improved sleep, more reliable nutrition, the ability to attend appointments, or a return to work and relationships. Improvement is not all-or-nothing, and I would rather you measure it honestly than wait for it to be total.



# 1. Understanding Multiple Chemical Sensitivity

---

## **A clinical description, not a single proven mechanism**

MCS generally describes recurrent symptoms involving more than one organ system that a person associates with low-level exposure to multiple chemically unrelated substances. Commonly reported triggers include fragrance, cleaning products, smoke, vehicle exhaust, paint, solvents, pesticides, new carpet, building materials, and personal care products. The amounts involved are often below levels expected to cause acute poisoning in most people.

I use this description with patients because it captures the pattern without claiming that one pathway has already been proven. Reviews have repeatedly noted that definitions vary and that consistent physical findings or laboratory abnormalities have not been established across all patients.<sup>3,6</sup>

## **MCS is not the same as acute poisoning**

A high-level exposure to carbon monoxide, chlorine gas, pesticides, solvents, or other hazardous agents can cause recognized toxic injury. That situation requires immediate exposure control and medical care. MCS, by contrast, usually refers to recurrent symptoms attributed to low-level exposures that do not produce acute toxicity in most people.

The distinction matters. A person can have a documented occupational or environmental exposure and later develop persistent sensitivity, but persistent symptoms do not automatically prove that a chemical remains stored in the body. Likewise, a normal routine blood test does not prove that nothing is wrong. Testing has to be matched to the specific exposure, timing, organ system, and clinical question, not ordered reflexively.

## **MCS is not the same as allergy**

Classic allergy involves identifiable immune mechanisms, commonly IgE-mediated reactions to substances such as pollen, animal dander, foods, or insect venom. Fragrances and volatile irritants can worsen rhinitis or asthma, but MCS as a syndrome is not generally explained by IgE-mediated allergy.<sup>7</sup> A person can have both MCS-like symptoms and a separate allergic condition, so I still refer for appropriate allergy or pulmonary evaluation when it is warranted.

## **A diagnosis of exclusion**

Most clinical guidance treats MCS or IEL as a diagnosis of exclusion. This means I evaluate for other conditions that can explain all or part of the symptom pattern before I attribute anything to MCS.<sup>8</sup> This is not dismissal. It is an essential safety step, because treatable diseases and hazardous exposures can otherwise be overlooked.

## Why language matters

People with MCS often report being dismissed, embarrassed, or treated as difficult. Clinicians may feel uncertain because the condition lacks a standard test. Families may become frustrated by changing triggers and complex accommodations. I keep my language neutral because it lowers conflict. I tell patients directly: I believe these symptoms are real, and I want to understand the pattern, without claiming to know the complete mechanism on day one.

**The stance I take.** Validate the person. Investigate dangerous and treatable causes. Reduce obvious irritants. Avoid overpromising. Build function gradually.

## 2. Symptoms, Triggers, and Patterns

---

### Commonly reported symptoms

Symptoms vary substantially from person to person and can change over time. A trigger associated with headache on one day may be associated with nausea or fatigue on another. This variability is exactly why I have every patient keep a structured diary rather than relying on memory alone.

What patients most often report, grouped by system:

- **Neurological and cognitive.** Headache, migraine, dizziness, disequilibrium, light sensitivity, sound sensitivity, difficulty concentrating, slowed thinking, memory problems, word-finding difficulty, tingling, and sleep disturbance.
- **Respiratory and upper airway.** Nasal burning, congestion, throat irritation, coughing, hoarseness, chest tightness, shortness of breath, wheezing, or a sensation that breathing is restricted.
- **Autonomic and cardiovascular.** Racing heart, palpitations, sweating, tremulousness, temperature changes, lightheadedness, blood pressure changes, or a feeling of impending collapse.
- **Gastrointestinal.** Nausea, abdominal discomfort, reflux, bloating, altered bowel habits, or appetite changes.
- **Skin and eyes.** Burning eyes, tearing, flushing, itching, rash, hives, or skin discomfort. Visible skin findings should be documented and evaluated rather than assumed to be MCS.
- **Musculoskeletal and whole-body.** Fatigue, weakness, muscle pain, joint pain, heaviness, post-exertional worsening, or a flu-like sensation.
- **Emotional and threat-response.** Fear, irritability, panic, urgency to escape, sadness, or helplessness. These may be consequences, amplifiers, comorbid conditions, or part of the physiological response itself. I do not use them to invalidate the person, and neither should anyone else.

### Common trigger categories

Fragrance is one of the most frequently reported triggers. Perfume, cologne, scented personal care products, air fresheners, laundry products, and fragranced cleaning agents can all become difficult to tolerate. Smoke, vehicle exhaust, cooking emissions, paint, adhesives, new flooring, cabinetry, and furniture can create similar problems because they introduce odors, particles, or volatile compounds into the air.

Other commonly reported triggers include pesticides, herbicides, solvents, fuels, and workplace chemicals. Damp or moldy environments can also worsen symptoms, although they can do so through several different pathways, including allergy, asthma, respiratory irritation, or the stress of being in a building associated with previous illness.

The surrounding context matters as much as the trigger itself. Heat, exertion, poor sleep, fasting, infection, emotional stress, migraine activity, and sensory overload can lower a person's threshold. This is one reason an exposure that seems manageable on one day can provoke a much stronger reaction on another, and it is exactly the terrain I ask patients to track alongside the trigger.

## **The value of a pattern diary**

Record exposure, timing, symptoms, duration, context, and recovery. Also record sleep, meals, hydration, menstrual cycle, infection, medication changes, stress, and exertion. The purpose is not to document every odor. It is to identify repeatable patterns and modifiable cofactors.

A useful entry might read: "Tuesday, 2:00 p.m. Entered recently painted office for 12 minutes. Odor was strong. Headache and throat burning began within five minutes. I had skipped lunch and slept five hours. Symptoms improved after leaving and eating. Headache lasted three hours." That is far more clinically useful to me than "paint made me toxic," because it shows both the trigger and the terrain that let it land as hard as it did.

## **Red flags that require urgent attention**

- Difficulty breathing, blue lips, severe wheezing, throat swelling, or rapidly progressing hives.
- Chest pain, fainting, new irregular heartbeat, one-sided weakness, confusion, or severe sudden headache.
- Possible carbon monoxide exposure, pesticide poisoning, solvent intoxication, or chemical burn.
- Persistent vomiting, inability to keep fluids down, or signs of dehydration.
- Suicidal thoughts, inability to care for basic needs, or extreme food and medication restriction.

**Emergency principle.** Do not assume that a severe episode is "just MCS." Follow emergency guidance for the symptoms and exposure involved.

### 3. What May Be Happening in the Body

---

Several models have been proposed. No single model has been proven to explain every case. I am careful, always, to use words like “may,” “could,” and “has been proposed” here, because a responsible explanation requires it. Multiple processes can also coexist in the same patient.

#### **Direct irritant effects**

Some substances irritate the eyes, nose, throat, lungs, or skin. Indoor pollutants can cause headache, dizziness, fatigue, and mucosal irritation, especially when concentrations are elevated. The Environmental Protection Agency notes that volatile organic compounds are emitted by many household products and building materials and that indoor concentrations can exceed outdoor levels.<sup>9</sup> This supports sensible exposure control, but it does not establish that all low-level odor reactions are toxic injury.

#### **Sensory and central nervous system sensitization**

Repeated symptoms and threatening experiences can increase responsiveness within sensory and threat-processing networks. Odors have unusually direct connections with brain systems involved in memory, emotion, and autonomic response. Once a smell is strongly associated with danger or illness, the body can generate rapid, involuntary symptoms before conscious reasoning occurs. This is one of the terrain factors I spend the most time on with patients, because it is genuinely treatable.

But there is a second half to this model that almost never gets explained to patients, and it is the half with the best objective evidence in this entire chapter. It deserves its own heading.

#### **The sensory nerves themselves, and the evidence that survives blinding**

Most research in this field asks people to react to smells, which means the person always knows something is happening. That is the weakness at the center of the whole literature.

Capsaicin gets around it. Capsaicin is the compound that makes chili peppers hot, and inhaled below the concentration where you can taste or smell it, it still activates a specific sensory nerve receptor in the airway called TRPV1. You cannot tell whether you are receiving it, and you cannot expect your way through it.

When that test was run as a randomized double-blind trial in patients meeting the standard criteria for chemical sensitivity, they coughed significantly more than healthy controls at every dose given, and reported more symptoms.<sup>18</sup> Their airway sensory nerves were measurably more reactive. Nobody was guessing.

The same pattern shows up away from the airway. Using quantitative sensory testing, which measures how the nervous system handles a standardized painful stimulus, patients with chemical sensitivity and no overlapping conditions developed a significantly larger area of spreading sensitivity after a capsaicin injection than matched controls, at every time point measured, and rated ordinary mechanical stimuli as more

intense.<sup>19</sup> That is the textbook signature of central sensitization: the volume control on sensory input is turned up, and it is turned up throughout the body rather than only where the chemical landed.

Two other threads support it. Increased capsaicin cough sensitivity has been found alongside altered nerve growth factor in nasal fluid, which is a marker of nerve remodeling rather than of allergy, and brain imaging during odor exposure has shown different activation patterns in people with odor-associated symptoms than in controls.<sup>20</sup> The current mechanistic account gathers all of this under sensitization of the TRP family of sensory receptors, principally TRPV1 and TRPA1, which are the same receptors that respond to irritants, to cold, and to a long list of the exact volatile compounds people report reacting to.<sup>21</sup>

Two honesty notes, because I would rather you heard them from me. The review making the broadest version of that case argues hard against psychological framing and reads as advocacy in places, so I weight the individual studies above it. And the two studies I described were small, twelve and fifteen patients. This is a real finding that needs a larger trial, not a settled one.

## **Detection, expectation, and the part that is harder to hear**

Now the other half, which I am not going to leave out, because you deserve the whole picture rather than the flattering portion of it.

When researchers take the smell away, the chemical specificity largely goes with it. A systematic review of thirty-seven provocation studies covering 784 people with chemical sensitivity found that reactions tracked whether the participant could tell the active substance from the sham. Exposures at or below the odor threshold generally did not produce consistent responses, and the studies that used a masking agent to conceal the stimulus found no association at all.<sup>10</sup>

That does not mean symptoms were fabricated, and I want to be exact about what it does mean. It means detection, expectation, prior learning, and threat appraisal influence the physiological response, and that identifying **which** substance caused a given reaction is far less reliable than it feels from the inside.

## **Putting the two halves together**

Held side by side, those two bodies of evidence say something coherent, and it is the position I actually work from.

**The hyperreactivity is real and measurable. The attribution to a specific chemical is not.**

Your nervous system genuinely is more reactive, and that shows up on tests you have no way to influence. What does not hold up is the confident identification of this cleaner or that perfume as the culprit, because under masked conditions those identifications do not reproduce.

Most people are relieved rather than insulted by this once it is said carefully, because it validates the part that has been dismissed and lifts an exhausting burden at the same time. You do not have to correctly identify



every offending molecule in your environment. That job was never possible, and failing at it was never a failure of yours.

Practically, it changes three things in how I work. Reducing your overall exposure load is worth doing, and building your life around chasing individual culprits is not. The nervous-system regulation work in Chapter 9 is not a consolation prize offered after the real medicine failed, because it targets the mechanism with the strongest evidence in this chapter. And any trial of any treatment has to be structured so that expectation is not doing the work, which is why I ask people to get stable first and then change one thing at a time, rather than starting eight things in a hopeful week.

## **Autonomic nervous system dysregulation**

Palpitations, tremor, sweating, dizziness, nausea, temperature shifts, and urgent escape behavior are compatible with autonomic activation. This can occur alongside migraine, panic, postural intolerance, pain, respiratory irritation, or chronic stress. Autonomic regulation is a reasonable treatment target in nearly every case I see, even when the initiating cause remains uncertain.

## **Migraine and sensory amplification**

Odor sensitivity is common in migraine. Some people who identify as chemically sensitive may have unrecognized migraine, vestibular migraine, or a related sensory processing disorder. Treating migraine can reduce odor-triggered headache, dizziness, nausea, light sensitivity, and cognitive disruption even if other sensitivities remain, which is why I screen for it early.

## **Inflammation, oxidative stress, and mitochondrial hypotheses**

Researchers have investigated oxidative stress, inflammatory markers, nitric oxide pathways, mitochondrial function, and brain imaging. Findings are not yet consistent enough to serve as diagnostic biomarkers or to prove a universal mechanism.<sup>11</sup> The nitric oxide and peroxynitrite cycle proposed by Martin Pall is an influential hypothesis, and I present it to patients exactly that way: as a model that has shaped how researchers think about the condition, not as an established clinical explanation.<sup>12</sup> I find it a useful framework for explaining why inflammation, oxidative stress, and mitochondrial capacity matter, but I do not treat it as proven, and neither should you.

## **Mast cells and histamine, and why they are not the engine here**

Mast cell activation can cause flushing, hives, wheezing, gastrointestinal symptoms, tachycardia, and lightheadedness. A person with these features may warrant specialist evaluation. Symptom overlap alone does not prove Mast Cell Activation Syndrome. Diagnosis requires appropriate clinical criteria and laboratory interpretation. I do not assume a universal low-histamine diet or a mast-cell supplement protocol for every person with MCS.

I want to go further than that, because histamine is probably the single most common thing chemical sensitivity gets blamed on, and the evidence does not support it as the driver.

Only one published study has actually measured histamine in people with self-reported chemical sensitivity. What it found was that substance P, vasoactive intestinal peptide, and nerve growth factor were elevated in those patients, **and histamine was not**. Exposure to volatile organic compounds then raised all of them in the patients but not in healthy controls, and made the skin more reactive to histamine applied directly.<sup>22</sup>

The order matters. The things that were elevated at baseline are nerve signaling molecules. Histamine was normal until the nerves were provoked. Substance P is a known activator of mast cells, so there is a genuine route running from an irritated sensory nerve to a degranulating mast cell, and that route explains why some people with chemical sensitivity do have real histamine-flavored symptoms.

But it runs nerve first, mast cell second. Histamine is a passenger on this trip, and sometimes an amplifier. It is not the engine. And that is one small study, so I hold even this loosely.

What follows for you is practical. If your picture is genuinely mast cell shaped, with flushing, hives, blood pressure swings and food-timed reactions, that is a different problem which deserves its own workup rather than being managed as a subsection of this one, and I would rather send you down that road properly than hand you a low-histamine food list. If your picture is chemical reactivity, calming the nerve sits upstream of blocking the histamine, and that is where I would spend your effort first.

## **Toxicant-Induced Loss of Tolerance**

TILT proposes a two-stage process: an initiating exposure or series of exposures is followed by triggering reactions to smaller amounts of unrelated substances. This model has helped researchers and patients describe chemical intolerance and has supported development of the Quick Environmental Exposure and Sensitivity Inventory.<sup>13</sup> It remains a proposed framework rather than a universally accepted mechanism, and I use it as a useful lens for organizing a patient's history, not as a confirmed diagnosis.

## **A biopsychosocial model does not mean “psychological only”**

Biology, learning, emotion, behavior, environment, and social conditions continuously interact. A person can have real airway irritation, migraine vulnerability, autonomic activation, prior trauma, and conditioned threat responses at the same time. This broad model is what allows me to offer more treatment options and avoid the false choice between “toxic” and “imaginary.”



## 4. Building a Sound Medical Evaluation

---

### Begin with a detailed history

A high-quality evaluation starts with chronology. What was happening before symptoms began? Was there a specific spill, fire, pesticide application, renovation, workplace event, infection, medication change, or period of extreme stress? Which symptoms appeared first? Which triggers are repeatable? What improves symptoms? Has function progressively narrowed?

Bring a concise one-page summary to appointments. Include the five most important symptoms, the five most repeatable triggers, relevant exposures, medical history, medication and supplement list, and the three outcomes you most want help with. Worksheet 1 in Chapter 12 is that page.

### Conditions to consider

Before I attribute a symptom pattern to chemical sensitivity, I make sure these categories have been genuinely considered, not just mentioned in passing:

- **Respiratory.** Asthma, nonallergic rhinitis, chronic sinus disease, hypersensitivity pneumonitis, vocal cord dysfunction or inducible laryngeal obstruction, chronic obstructive pulmonary disease, and occupational lung disease.
- **Neurological.** Migraine, vestibular migraine, seizure disorder, other headache disorders, neuropathy, persistent postural-perceptual dizziness, and post-concussion syndrome.
- **Cardiovascular and autonomic.** Arrhythmia, structural heart disease presenting as palpitations or dizziness, orthostatic hypotension, postural orthostatic tachycardia syndrome, anemia, and dehydration.
- **Endocrine and metabolic.** Thyroid disease, glucose disorders, electrolyte abnormalities, vitamin deficiencies, and adrenal disease when clinically indicated.
- **Allergic and immunological.** IgE-mediated allergy, chronic urticaria, contact dermatitis, and selected mast-cell disorders.
- **Toxicological, environmental, and occupational.** Carbon monoxide, combustion exposure, pesticide exposure, solvent intoxication, medication reactions or interactions, lead or other metal exposure when history supports it, water damage, workplace ventilation problems, and documented occupational poisoning with its own recognized health surveillance requirements.
- **Infectious.** An active or recent infection that can mimic or amplify the symptom picture.
- **Sleep and mental health.** Sleep apnea, insomnia, anxiety, panic disorder, depression, trauma-related illness, health anxiety, or eating disorder. These may coexist with physical illness and deserve treatment in their own right.

- **Overlapping chronic syndromes.** Fibromyalgia, ME/CFS, irritable bowel syndrome, chronic pain, and other conditions involving sensory or central amplification.

This list is not exhaustive, and it is not a substitute for your clinician's judgment. It is the starting checklist I use so that a genuinely treatable condition is never missed because "chemical sensitivity" was the fastest label available.

## **Testing should answer a question**

Useful tests are selected to confirm or exclude a plausible diagnosis. Examples might include spirometry for asthma symptoms, an electrocardiogram for palpitations, blood counts for anemia, thyroid testing for compatible symptoms, or targeted toxicology testing after a documented exposure. Broad panels without a clear clinical question often generate ambiguous findings and expensive follow-up, and I will not order a panel whose purpose I cannot explain to you first.

## **Use an interdisciplinary team when needed**

Depending on the presentation, useful professionals may include a primary-care clinician, occupational and environmental medicine physician, allergist, pulmonologist, neurologist, cardiologist, dermatologist, physical therapist, dietitian, psychologist, or social worker. Interdisciplinary care is especially important when the condition is affecting housing, nutrition, employment, medication access, or mental health.<sup>8</sup>

**A script for your appointment.** "I experience repeatable multisystem symptoms around certain odors and products. I know the mechanism may be uncertain. I would like help ruling out treatable conditions, documenting functional limitations, and developing a safe management plan."

# 5. Environmental and Functional Testing

---

## What a test can and cannot prove

Environmental chemicals can be measured in blood, urine, air, dust, water, or building materials. The meaning depends on the chemical, specimen, timing, laboratory method, reference population, and reason for testing. Detection does not automatically equal poisoning, tissue injury, or causation of symptoms, and I tell every patient this before we order anything.

Urinary measurements often reflect recent exposure and excretion. A higher value may identify an exposure source worth reducing. It does not necessarily prove that the substance is stored in tissues or that removing it will resolve symptoms. A low urinary value does not exclude prior exposure either, because many compounds are cleared or redistributed over time.

## Conventional testing

Conventional laboratories are valuable when matched to a clinical question. Blood lead levels, carbon monoxide testing, cholinesterase testing after certain pesticide exposures, kidney and liver function, blood counts, thyroid tests, and pulmonary testing are examples. The correct specimen and collection window matter. Consultation with poison control, occupational medicine, or a medical toxicologist may be appropriate after a significant exposure.

## Specialty environmental panels

Commercial panels may test urine for mycotoxins, phthalate metabolites, pesticides, volatile organic compound metabolites, bisphenols, parabens, metals, or per- and polyfluoroalkyl substances. These tests can generate extensive data, but clinical interpretation is genuinely difficult, and I will not order one without a specific reason. Before I order a specialty test, I ask whether the result will change a specific clinical decision. A useful test has a clearly stated purpose, an analytically reliable method, and a reference range that can be interpreted in the context of ordinary population exposure as well as disease.

It also matters what the sample actually measures. Some tests reflect recent exposure, while others measure excretion rather than stored body burden or tissue injury. Diet, hydration, occupation, smoking, medications, supplements, and sample timing can all influence a result. Finally, I consider what treatment would actually follow if the result comes back abnormal. The value of a test depends partly on whether the proposed response is evidence-informed, proportionate, and safe.

## Mycotoxin testing

A urine mycotoxin result does not by itself diagnose illness caused by a building. Mycotoxins can enter through food as well as inhalation, and the relationship between urinary levels, building exposure, and

symptoms is complex. When dampness or visible mold is present, my first priority is always to correct moisture and building damage, not to chase a lab number. Medical evaluation should focus on recognized conditions such as allergy, asthma, hypersensitivity pneumonitis, infection in susceptible patients, and irritant symptoms.

## Heavy-metal testing

Heavy-metal testing should be based on exposure history and performed with validated specimens. I do not use “provoked” urine testing, which administers a chelating agent before the urine sample is collected. Provoked testing can produce elevated results that are not comparable with standard reference ranges and can lead directly to unnecessary chelation. Chelation carries real risks and is appropriate only for specific, documented poisonings under qualified medical supervision. I do not use chelation as a general detoxification strategy, and I would encourage you to be very cautious of any practitioner who does.

## Genetic testing

Variants in genes such as MTHFR, GST, CYP, or SOD are common. A variant rarely proves that a person cannot detoxify, that toxins are accumulating, or that a specific supplement is required. Genetic information may contribute to research or selected clinical decisions, but symptoms, examination, standard laboratory data, diet, medication, and exposure history generally matter more than an isolated single-nucleotide polymorphism.

## Organic acids and “detox pathway” testing

Organic acids testing can measure urinary metabolites, but commercial interpretations frequently extend beyond what has been validated for diagnosis or treatment of MCS. I do not use results like these alone to diagnose mitochondrial failure, neurotransmitter imbalance, yeast overgrowth, or impaired detoxification in an MCS patient. They are one data point read in the context of the whole clinical picture, never the whole picture themselves.

**A safeguard before you spend.** Before you commit to an expensive panel, ask the ordering clinician to state in writing what each possible result would mean and how it would change your care. If that question cannot be answered before the sample is collected, the test is not ready to be ordered. Worksheet 4 in Chapter 12 walks through the rest of that decision.

## 6. Creating a Lower-Exposure Home

---

Environmental improvement should be practical, prioritized, and proportional. My goal for patients is a healthier home, not a perfectly chemical-free home, which is neither realistic nor necessary. EPA guidance emphasizes three strategies: source control, ventilation with clean outdoor air, and filtration as a supplement.<sup>9,14</sup>

### Step 1: Address high-priority safety hazards

Begin with hazards that can cause serious illness in anyone. Every home should have functioning smoke and carbon monoxide alarms, and radon testing should be completed according to local guidance. Active leaks and chronic moisture deserve prompt attention. When feasible, indoor humidity is generally best kept below 60 percent and often within a range of approximately 30 to 50 percent.<sup>15</sup>

Combustion appliances should be vented correctly, and a range hood that exhausts outdoors is preferable when available. Cleaning products should never be mixed, especially bleach with ammonia or acids. Suspected asbestos, a gas leak, extensive mold damage, or other hazardous material should be evaluated by appropriately licensed professionals rather than handled through improvised do-it-yourself methods.

### Step 2: Remove obvious sources

Source control is usually more effective than trying to filter pollution after it has entered the room. Remove or isolate air fresheners, scented candles, plug-ins, fragranced laundry products, open solvent containers, unused pesticides, smoke, and strongly off-gassing items. Store fuels, paints, and chemicals outside occupied living areas when safe and permitted. EPA notes that products and building materials can emit volatile organic compounds during use and storage.<sup>16</sup>

### Step 3: Simplify cleaning

Choose fragrance-free products with the fewest ingredients that accomplish the task. “Unscented” can mean that a masking fragrance has been added, so “fragrance-free” is generally the more useful label. Introduce one new product at a time. Adequate ventilation and label instructions remain essential even for products marketed as natural.

Vinegar, baking soda, soap, and hydrogen peroxide may be useful for selected tasks, but they are not interchangeable and should not be mixed casually. Vinegar can damage stone and some surfaces. Hydrogen peroxide can irritate skin and eyes. Natural substances can still cause reactions.

## **Step 4: Ventilate strategically**

Ventilation lowers indoor pollutant concentration when outdoor air is cleaner. Open windows or use exhaust fans during painting, cleaning, cooking, or installation of new products. Outdoor smoke, pollen, ozone, extreme heat, or high humidity may make window ventilation inappropriate, so timing matters.

## **Step 5: Use filtration appropriately**

HEPA filtration is designed primarily for particles, including dust, smoke particles, pollen, and some airborne biological material. It does not efficiently remove gases by itself. Activated carbon or another sorbent medium is needed for some gases and odors, and performance depends heavily on the amount and quality of the medium. EPA advises sizing a portable air cleaner to the room and treating filtration as a supplement to source control and ventilation.<sup>14</sup>

Avoid devices that intentionally generate ozone. Ozone is a lung irritant and should not be used as an occupied-space air-cleaning strategy.

## **Step 6: Make renovations carefully**

When renovation is necessary, correct moisture problems before beginning cosmetic work. Ask for lower-emitting materials, obtain samples before making large purchases, and notice how you respond to those samples in a controlled setting.

New items can sometimes be unpacked in a garage or protected outdoor area when that can be done safely. Painting, flooring, cabinetry, and adhesive work are easier to manage when the sensitive person can remain elsewhere during the most intense period of odor and curing. Ventilation and adequate curing time matter.

Avoid changing many materials at once unless safety requires it. A slower approach is less disruptive, makes reactions easier to interpret, and reduces the risk of spending large amounts of money without knowing which change was helpful.

## **A low-cost first week**

During the first week, I focus patients on a few changes with a high likelihood of benefit. Remove air fresheners, scented candles, dryer sheets, and unnecessary sprays. Move paints, gasoline, pesticides, and solvents out of occupied living areas, and wash bedding and frequently worn clothing with a tolerated fragrance-free detergent.

Check the home for moisture, blocked vents, combustion concerns, and expired alarm batteries. Rather than attempting to transform the entire house immediately, create one calm sleeping and recovery area first. A dependable room can provide meaningful relief while the rest of the home is assessed gradually.



## 7. Nutrition for Stability and Resilience

---

There is no scientifically established MCS diet, and I say that to every patient up front. Nutrition should first prevent malnutrition, stabilize energy, support bowel function, and reduce unnecessary restriction. Food fear can become one of the most damaging consequences of chronic sensitivity, and I actively guard against creating it.

### **Build a reliable food foundation**

A reliable food foundation is more helpful than an elaborate detoxification diet, in my experience, without exception. Eat regular meals and avoid prolonged fasting unless it has been medically recommended. Include a tolerated source of protein at each meal, along with a varied selection of fruits, vegetables, whole grains or other tolerated starches, and healthy fats.

Hydration should be adequate but not excessive. Food should be stored safely, and leftovers should be refrigerated promptly. When laboratory testing or clinical evaluation identifies a genuine nutrient deficiency, I correct it starting with food and, when appropriate, carefully supervised supplementation.

### **Organic food: useful option, not a requirement**

Organic choices can reduce exposure to some pesticide residues, but a 100 percent organic diet is not required for health or recovery. Cost, access, nutritional adequacy, and individual tolerance matter. Washing produce, peeling when appropriate, varying food choices, and following food-safety guidance are reasonable strategies when organic options are limited.

### **Low-histamine diets**

A low-histamine trial may be considered when symptoms such as flushing, hives, nasal symptoms, headaches, tachycardia, or gastrointestinal symptoms are closely linked to aged or fermented foods. I keep it time-limited and structured, followed by reintroduction whenever possible. Long-term broad restriction can impair nutrition and quality of life, and I watch for that closely.

### **Elimination diets**

Elimination diets should test a specific hypothesis. Remove a limited number of suspected foods for a defined period, usually with dietitian support, and then reintroduce them systematically unless there is a history of severe allergy. Symptoms that fluctuate naturally can create false conclusions when many foods are removed at once.

## **Sulfur foods and cruciferous vegetables**

Cruciferous vegetables and allium vegetables provide valuable nutrients and bioactive compounds. They can be part of a balanced diet when tolerated. Claims that they “open detox pathways” or rapidly clear stored toxicants are oversimplified, and I do not make them. Food should nourish the person rather than become a treatment test performed at every meal.

## **Bowel regularity**

Constipation can worsen bloating, nausea, and discomfort and can complicate use of supplements or binders. Adequate fluid, fiber, movement, and evaluation of medications or pelvic-floor problems can help. Sudden or severe constipation, blood in stool, weight loss, or persistent abdominal pain requires medical evaluation.

**A nutrition warning I take seriously.** Seek prompt help if the diet is shrinking, weight is falling, meals provoke fear, or fewer than ten to fifteen foods are being eaten reliably. Early dietitian support can prevent severe restriction, and I would rather bring one in early than clean up after months of narrowing.



## 8. Supplements, Binders, Sauna, and Detox Claims

---

People with MCS are often unusually cautious about medications and supplements, sometimes because they have experienced prior reactions. I respect that caution completely. It also means I approach anything more complex slowly and only when there is a clear purpose. This is the chapter where I am most careful to separate what I am confident about from what I am not, because it is also the chapter where patients are most vulnerable to overpromising claims from elsewhere.

### A safer introduction method

Introduce supplements cautiously and for a clearly defined reason. Review the product and intended goal with a qualified clinician or pharmacist, especially when prescription medications, kidney or liver disease, pregnancy, allergies, or multiple sensitivities are involved. Excipients and inactive ingredients can matter as much as the active ingredient for some patients, and in this population they matter more often than people expect.

Change only one variable at a time. When a trial is reasonable, begin with the lowest appropriate dose, define what improvement would look like, and record both benefits and adverse effects over a predetermined period. Stop the product and seek guidance if a significant reaction occurs.

### The commonly promoted supplements, read honestly

These are the products patients most often arrive already taking or already being sold. Here is what each one actually is, and what the evidence does and does not support in MCS specifically.

- **N-acetylcysteine.** A precursor for glutathione, with approved mucolytic uses. The evidence does not establish it as a treatment for MCS. It can cause gastrointestinal symptoms, interact with some medications, and rarely trigger bronchospasm.
- **Glutathione.** A biologically important antioxidant. Oral and liposomal products vary considerably. Clinical evidence for MCS outcomes is insufficient.
- **Vitamin C.** An essential nutrient and antioxidant. High doses may cause diarrhea or increase kidney-stone risk in susceptible people.
- **Magnesium.** May help a documented deficiency, constipation, migraine, or muscle symptoms depending on the formulation. Excess can cause diarrhea and is unsafe in significant kidney failure.
- **Quercetin.** Has laboratory and limited clinical interest for allergic and inflammatory pathways. It is not an established MCS therapy and may interact with medications.
- **Alpha-lipoic acid.** Used in selected neuropathy settings and studied as an antioxidant. It can lower blood glucose, and it should not be used as unsupervised metal chelation.

- **Milk thistle.** Studied in some liver conditions, but the evidence for MCS is lacking. Product quality and drug interactions matter.
- **Methylated B vitamins.** Appropriate when a deficiency or another indication exists. Common MTHFR variants do not automatically justify high doses. Excess vitamin B6 can cause neuropathy.
- **Coenzyme Q10.** May be considered for selected conditions, but the evidence for MCS is insufficient and interactions are possible.

Read that list honestly and a pattern jumps out: not one of these has outcome evidence for MCS as a condition. Nobody has run those trials, and nobody is likely to fund them for compounds that cannot be patented. That is a real gap and I am not going to paper over it. Here is what I do instead of guessing. I treat what is actually documented. If testing shows a nutrient deficiency, I correct it, starting with food. If a coexisting condition is identified, whether that is migraine, anemia, thyroid disease, constipation, or a sleep disorder, I treat that condition on its own merits with the tools that do have evidence behind them. Anything beyond that gets trialed one product at a time, at the lowest appropriate dose, against an endpoint we agreed on in advance, with a written stopping rule. A supplement that cannot name what it is supposed to change, and by when, does not get started.

## Where I source what I do use

I hold no physical inventory. I keep a screened dispensary through Fullscript, which ships directly to you ([us.fullscript.com/welcome/drseay](https://us.fullscript.com/welcome/drseay)). Products from any reputable retailer work equally well. Nothing in this chapter is a recommendation to buy anything; it is here so that when something is genuinely indicated for you, sourcing it is not one more obstacle.

*Root Health earns a portion of dispensary purchases.*

## Binders

Activated charcoal, clay products, modified citrus pectin, and algae are often marketed as intestinal toxin binders. Evidence supporting routine binder use for MCS is limited, and I do not use binders as a routine daily wellness product. Activated charcoal can bind medications and nutrients and can cause constipation or aspiration. Clay and algae products may contain contaminants. Binders are not harmless daily wellness products, and where a binder is genuinely indicated it belongs 1 to 2 hours away from food, other supplements, and medication.

## Sauna and sweating

Sauna may provide relaxation, warmth, and a sense of well-being for some people. Small studies in other contexts have detected certain chemicals in sweat, but this does not prove that sauna removes a clinically meaningful toxic burden or treats MCS, and I do not present it that way. Heat can worsen dehydration, orthostatic intolerance, migraine, cardiovascular symptoms, and medication effects.

Anyone using sauna should begin conservatively, avoid it during acute illness or dehydration, replace fluids appropriately, and obtain medical clearance when cardiovascular, kidney, neurological, pregnancy, or autonomic concerns are present. There is not sufficient evidence to recommend taking binders before sauna, and I do not recommend that combination.

## **Chelation**

I want this section to be unambiguous. Chelation is a medical treatment for specific documented poisonings, not a general detoxification strategy, and I do not use it as one. Chelating drugs can damage kidneys, disturb minerals, and cause serious adverse events. I do not use chelation based on hair testing, provoked urine testing, or nonspecific symptoms, and I would urge you to decline it from anyone who recommends it on that basis alone. A confident, root-cause approach to MCS does not require chelation, provoked testing, or aggressive detoxification protocols as a first-line strategy. It requires patience, sequencing, and a nervous system that is actually being supported rather than shocked.

## **When you feel worse**

Some people feel worse when they start something new, and the wellness world has a ready explanation for it: worsening proves the protocol is working. I do not accept that framing, because a framework in which feeling worse is always evidence of progress cannot be tested and cannot be wrong. Worsening is information. Sometimes it settles within a few days. Sometimes the dose is too high, sometimes the groundwork was skipped, and sometimes the product is simply not right for you. Significant worsening is a reason to stop and reassess with me, not a milestone to push through.

## **The nocebo and iatrogenesis problem**

When every symptom is interpreted as evidence of accumulating poison, a person can become increasingly afraid of food, medication, buildings, and other people. Unproven testing and treatment can unintentionally reinforce disability. Earlier reviews have warned that promises of unproven therapies may perpetuate illness and financial strain.<sup>17</sup> A good plan reduces danger without teaching the nervous system that all exposure is catastrophic, and protecting against that is one of my central jobs in this work.

## 9. Calming a Sensitized Nervous System

---

Nervous-system support is not an accusation that symptoms are imagined. The brain, autonomic nervous system, endocrine system, immune signaling, breathing pattern, and muscles all participate in every physical reaction. Improving regulation reduces symptom intensity regardless of the original trigger, and this is where I put a meaningful share of my clinical attention, because it is genuinely actionable terrain.

### **Pacing before pushing**

Repeated cycles of overexertion and collapse can destabilize sleep, migraine, pain, and autonomic symptoms. Pacing means balancing activity and recovery so that function can expand without repeated major setbacks. Track what you can do reliably, not what you can do once on a good day.

### **Breathing and grounding**

Slow breathing with a comfortable, slightly longer exhalation may reduce hyperventilation and sympathetic activation. Avoid forceful breath holds or aggressive breathing exercises that provoke dizziness. Pair breathing with grounding: feel the feet, name objects in the room, orient to exits and fresh air, and remind yourself that symptoms can be intense without automatically signaling tissue damage.

### **Sleep and circadian stability**

Sleep becomes more stable when the body receives consistent timing cues. A regular wake time and morning light exposure, when tolerated, can help regulate circadian rhythm. Limiting late caffeine and alcohol may also improve sleep quality.

The sleeping area should feel predictable, comfortable, and low in odor. Pain, reflux, nasal obstruction, restless legs, and possible sleep apnea deserve treatment because each can fragment sleep and lower resilience the following day. I discourage patients from turning the bedroom into a constantly changing laboratory of devices, filters, supplements, and new products. Stability itself is therapeutic.

### **Migraine-informed care**

Regular meals, hydration, sleep consistency, light management, appropriate exercise, and migraine treatment can reduce sensory sensitivity. Preventive and acute migraine medications exist and help a great many people. They are not something I prescribe, and I include them here so that you know the option exists and can raise it with the physician managing that part of your care. Medication sensitivity concerns can often be addressed with low starting doses, careful formulation selection, and one change at a time.

## Psychological and behavioral support

Therapy can help with grief, trauma, panic, isolation, conflict, and the fear created by unpredictable symptoms. Cognitive behavioral therapy, acceptance-based approaches, trauma-informed care, and exposure-informed rehabilitation can be useful when delivered respectfully. The purpose is never to argue that exposures are harmless. It is to reduce avoidable suffering and improve choice.

## Neural retraining programs

Commercial neural retraining programs use repetition, attention, imagery, movement, and cognitive techniques to alter threat responses. Some participants report benefit, but high-quality evidence specific to MCS remains limited. I present these programs as optional self-regulation tools, not guaranteed cures. They should never require a person to enter genuinely unsafe environments or deny symptoms.

## Graded re-engagement

Total avoidance may be necessary during an acute hazardous exposure or severe destabilization. Over the long term, however, extreme avoidance can shrink life, increase vigilance, and make future participation harder. Re-engagement should be individualized and collaborative. Begin with environments that are low-risk, meaningful, and controllable. Establish an exit plan. Measure recovery time. Increase duration or complexity gradually.

**A functional goal.** “Attend my grandchild’s school event for twenty minutes with a planned seat near an exit” is far more useful to me than “never react to fragrance again.”

# 10. Work, Healthcare, Travel, and Relationships

---

## Workplace support

The social and occupational burden of MCS can be substantial. Systematic qualitative research describes job disruption, restricted participation, and social losses.<sup>1</sup> A workplace plan should focus on functional barriers and practical modifications rather than debating the mechanism.

At work, a practical request is usually more successful than a demand for an environment that no organization can guarantee. Fragrance-reduction policies, advance notice of painting or pest control, and relocation away from printers, cleaning storage, or high-traffic areas can meaningfully reduce exposure.

Remote or hybrid work, flexible scheduling, improved ventilation, and access to a private workspace can also help. Written communication is useful because it creates a clear record of the employee's functional limitations and the accommodations being requested. Occupational health, human resources, a union representative, or an accommodation specialist can assist when those resources are available.

## Workplace request template

I coach patients to keep workplace requests specific, written, and tied to a functional limitation rather than a demand for a scent-free building no employer can fully deliver. Asking to begin an interactive accommodation process is often the phrase that opens the right door. A short template that works well:

*"I have a documented medical condition that causes significant symptoms, including [headache, respiratory irritation, cognitive difficulty, or your specific symptoms], when exposed to [fragrance, recent renovation odors, or your specific triggers]. I am requesting an interactive accommodation process and the following reasonable accommodations: (1) advance notice before painting, pest control, or deep cleaning near my workspace; (2) relocation away from [printers, cleaning storage, high-traffic areas]; (3) [a fragrance-reduction request for my immediate work area, remote or hybrid scheduling on specific days, or another specific accommodation]. I am glad to provide medical documentation and to discuss which of these are feasible given the workplace."*

Send the request in writing to the appropriate channel, whether that is human resources, an occupational health office, or a designated accommodation coordinator, and keep a copy. A specific, documented request is far more likely to result in a workable accommodation than a verbal, general one.

## Healthcare visits

Healthcare visits are easier to navigate when the office is contacted in advance. Ask about fragrance practices, recent renovation, and whether waiting outside or in a vehicle is possible. The first appointment of



the day may reduce exposure to accumulated cleaning products or other patients' fragrances, although this cannot be guaranteed.

Bring a concise medical summary, current medication list, emergency information, water, and a tolerated mask when appropriate. Staff may be able to avoid room sprays or nonessential scented products, but disinfectants may still be required for infection control. Environmental concerns should never delay urgently needed emergency care.

## **Travel**

Travel adds uncertainty from hotel cleaners, vehicle fragrances, fuel, and unfamiliar food. Start with short trips. Confirm cancellation policies. Ask lodging staff not to use air fresheners and, when possible, to avoid same-day deep cleaning. Bring a familiar pillowcase, tolerated toiletries, simple food, medication, and a written plan for flares.

## **Family and social relationships**

Loved ones may not understand why a product that was tolerated last month is a problem today. Explain the functional effect without demanding that they adopt the same beliefs about mechanism. Make requests specific and time-limited.

**A script for family.** "I am not asking you to understand every mechanism. I am asking for one practical change: please avoid perfume and scented laundry products when we meet indoors. That gives me the best chance of staying and participating."

## **Isolation and mental health**

Isolation can become both a consequence and a driver of worsening health. Maintain connection through outdoor visits, video calls, fragrance-reduced gatherings, faith communities, support groups, or adapted hobbies. I steer patients toward support groups that validate experience without escalating fear, discouraging medical care, or promoting costly cures.

# 11. A Personalized 90-Day Recovery Roadmap

---

This roadmap is not a detox protocol. It is a staged process designed to reduce immediate risks, improve medical clarity, strengthen basic physiology, and expand function. Move more slowly when symptoms are severe.

## **Days 1 to 14: Stabilize**

During the first two weeks, the priority is stabilization. Address emergency symptoms and obvious hazardous exposures first, then create one lower-exposure place for sleep and recovery. Remove air fresheners, smoke, open solvents, and unnecessary fragranced products from that area.

Regular meals, hydration, sleep timing, and bowel regularity provide a basic physiological foundation. Prepare a one-page medical history and begin the exposure and symptom diary. Resist the urge to introduce multiple supplements, devices, restrictive diets, and laboratory tests at the same time.

## **Days 15 to 30: Evaluate**

From days 15 through 30, shift toward evaluation. Complete a primary medical assessment and any targeted referrals that are clinically indicated. Review medications, supplements, occupational history, housing conditions, and relevant exposures.

Use the diary to identify the three most repeatable triggers and the three strongest cofactors, such as poor sleep, fasting, migraine, heat, or stress. Choose no more than two environmental changes to test. At the same time, establish a baseline for function by recording work hours, walking tolerance, social contact, food variety, sleep quality, and recovery time after a reaction.

## **Days 31 to 60: Build capacity**

Between days 31 and 60, the focus turns to building capacity. Treat identifiable conditions such as asthma, migraine, anemia, sleep disorders, rhinitis, reflux, or anxiety. Maintain a nutritionally adequate diet and, when safe, begin restoring foods that were removed without clear benefit.

Add gentle movement or rehabilitation at a tolerable level and practice one nervous-system regulation method consistently. Implement one practical workplace or household accommodation. Supplements should be trialed one at a time and only when there is a clear reason to use them.

## **Days 61 to 90: Expand life**

From days 61 through 90, begin expanding life. Choose one meaningful activity that has been lost and break it into smaller, controllable steps. Environmental planning, pacing, and a clear exit strategy can make re-engagement feel safer and more predictable.



Repeat the activity at a tolerable level before increasing the duration or complexity. Review every intervention honestly. Keep the changes that improved function, and discontinue those that added cost, fear, or complexity without a measurable benefit. Finish the period with a maintenance plan and a list of early signs that indicate the need to slow down or seek help.

## **Measure more than symptoms**

Symptom scores matter, but function can improve before symptoms disappear. Track hours out of bed, food variety, sleep continuity, time spent with others, work participation, walking or exercise capacity, number of urgent escapes, and recovery time after a flare. Your baseline is your own log from day one, not somebody else's number.

## 12. Worksheets and Supplemental Support

---

### Worksheet 1: One-page clinical summary

Bring this to every appointment.

**Main symptoms:**

---

**Most repeatable triggers:**

---

**Date and circumstances of onset:**

---

**Known occupational or environmental exposures:**

---

**Medical diagnoses already evaluated:**

---

**Current medications and supplements:**

---

**Three goals for care:**

---

### Worksheet 2: Exposure and symptom diary

Keep one row per event. Track date and time, place or product, duration, symptoms, onset, context and cofactors, and recovery. Alongside those, record sleep, meals, hydration, stress, and exertion for the day, because the cofactor column is usually where the answer is hiding.

### Worksheet 3: Home priority audit

Check every item that applies to your home right now, then work down the list in order.

- ☐ Smoke or vaping indoors
- ☐ Carbon monoxide alarms absent or expired

- ☐ Gas or combustion concerns
- ☐ Active leak or moisture damage
- ☐ Visible mold or persistent musty odor
- ☐ Air fresheners or scented plug-ins
- ☐ Open paint, solvent, fuel, or pesticide storage
- ☐ Poor kitchen or bathroom exhaust
- ☐ Recent renovation or new furnishings
- ☐ Strongly fragranced laundry products
- ☐ Ozone-generating air cleaner
- ☐ Bedroom lacks a reliable recovery space

### Worksheet 4: Test decision guide

Answer these before any specialty panel is ordered.

1. What exact question will this test answer?
2. How will each possible result change care?
3. Does it measure exposure, body burden, effect, or excretion?
4. Is the specimen and collection timing appropriate?
5. Is the laboratory method validated?
6. Could the result create unnecessary treatment?
7. What is the total financial cost, including follow-up?

### Worksheet 5: Function tracker

Record each area at baseline, then again at 30, 60, and 90 days: sleep quality, food variety, time outside the home, work or household activity, social connection, movement tolerance, flare recovery time, and confidence managing symptoms. Compare each entry against your own baseline, never against anyone else's.

### Worksheet 6: Flare plan

**My early warning signs:**

---

**My safest immediate actions:**

---

**Symptoms that require urgent care:**

---

**People to contact:**

---

**Medication or rescue plan prescribed by my clinician:**

---

## **Choosing supportive communities**

Look for groups that support practical accommodations, medical evaluation, nutrition, mental health, and gradual restoration of function. Be cautious when a group insists that one test proves the cause, discourages conventional care, promotes fear of nearly all foods or medications, or sells a single expensive cure.

## **Questions to ask a clinician or program**

- What evidence supports this recommendation specifically for MCS?
- What are the risks, contraindications, and alternatives?
- How will we know whether it is working?
- What is the stopping rule if I worsen?
- How does this plan protect nutrition, finances, relationships, and function?
- Are you willing to coordinate with my other clinicians?

I welcome all six of those questions. Any clinician worth your time will.

# Closing Perspective

---

People living with chemical sensitivity deserve more than dismissal. They also deserve more than frightening explanations and unproven protocols. The path I take with patients combines compassion, careful diagnosis, sensible environmental health, nourishment, symptom treatment, nervous-system support, and a gradual return to meaningful life.

You are not required to solve every scientific controversy before taking helpful action. Start with safety. Reduce obvious irritants. Seek a thoughtful medical evaluation. Protect nutrition and connection. Measure function. Change one variable at a time. Recovery is built through many small, credible steps, and that is exactly how I build it with every patient who sits across from me.

# Working with Root Health

---

Everything in this book is general education, written to be safe for a reader I have never examined. A personalized plan starts from your history, your exposures, your labs, and your medication list.

***Book a free Root Discovery Session: [rootcauseunlocked.com/discovery?](https://rootcauseunlocked.com/discovery?utm_source=ebook&utm_medium=ebook&utm_campaign=mcs)  
[utm\\_source=ebook&utm\\_medium=ebook&utm\\_campaign=mcs](https://rootcauseunlocked.com/discovery?utm_source=ebook&utm_medium=ebook&utm_campaign=mcs)***

Consultations and ongoing care are both available wherever you are located. Being outside North Carolina is not a barrier to working with me. Tell me where you are when you reach out and I will lay out exactly how we would work together from there.

***Get in touch: [rootcauseunlocked.com/contact?](https://rootcauseunlocked.com/contact?utm_source=ebook&utm_medium=ebook&utm_campaign=mcs)  
[utm\\_source=ebook&utm\\_medium=ebook&utm\\_campaign=mcs](https://rootcauseunlocked.com/contact?utm_source=ebook&utm_medium=ebook&utm_campaign=mcs)***

**If you would rather start on your own:**

***Free tracker: [rootcauseunlocked.com/tracker?](https://rootcauseunlocked.com/tracker?utm_source=ebook&utm_medium=ebook&utm_campaign=mcs)  
[utm\\_source=ebook&utm\\_medium=ebook&utm\\_campaign=mcs](https://rootcauseunlocked.com/tracker?utm_source=ebook&utm_medium=ebook&utm_campaign=mcs)***

*Dr. Adam Seay, DC · Seay Wellness / Root Health, Garner, NC  
[rootcauseunlocked.com](https://rootcauseunlocked.com) · (919) 662-0520*

*Educational content only. Not medical advice. Not a diagnosis. Never stop, skip, or change a prescribed medication based on this book. If a reaction involves difficulty breathing, chest pain, fainting, confusion, or a suspected poisoning, seek emergency care now.*

# References

---

1. Driesen L, Patton R, John M. The impact of multiple chemical sensitivity on people's social and occupational functioning: a systematic review of qualitative research studies. *Journal of Psychosomatic Research*. 2020;132:109964. doi:10.1016/j.jpsychores.2020.109964
2. Gibson PR, Elms ANM, Ruding LA. Perceived treatment efficacy for conventional and alternative therapies reported by persons with multiple chemical sensitivity. *Environmental Health Perspectives*. 2003;111(12):1498-1504. doi:10.1289/ehp.5936
3. Rossi S, Pitidis A. Multiple Chemical Sensitivity: review of the state of the art in epidemiology, diagnosis, and future perspectives. *Journal of Occupational and Environmental Medicine*. 2018;60(2):138-146. doi:10.1097/JOM.0000000000001215
4. Zucco GM, Doty RL. Multiple Chemical Sensitivity. *Brain Sciences*. 2022;12(1):46. doi:10.3390/brainsci12010046
5. Hempel S, Zhang D, Robinson KA, et al. Multiple chemical sensitivity validity, prevalence, tools and interventions: systematic review protocol. *BMJ Open*. 2025;15:e088136. doi:10.1136/bmjopen-2024-088136
6. Graveling RA, Pilkington A, George JPK, Butler MP, Tannahill SN. A review of multiple chemical sensitivity. *Occupational and Environmental Medicine*. 1999;56(2):73-85. doi:10.1136/oem.56.2.73
7. Binkley KE. Multiple Chemical Sensitivity/Idiopathic Environmental Intolerance: a practical approach to diagnosis and management. *Journal of Allergy and Clinical Immunology: In Practice*. 2023. doi:10.1016/j.jaip.2023.08.039
8. Schwenk M. Multiple chemical sensitivity: a guide for dermatologists on how to manage affected individuals. *Journal der Deutschen Dermatologischen Gesellschaft*. 2020;18. PMID 32026633
9. U.S. Environmental Protection Agency. Indoor Air Quality. Updated 2026. Accessed July 22, 2026.
10. Das-Munshi J, Rubin GJ, Wessely S. Multiple chemical sensitivities: a systematic review of provocation studies. *Journal of Allergy and Clinical Immunology*. 2006;118(6):1257-1264. doi:10.1016/j.jaci.2006.07.046
11. Multiple Chemical Sensitivity: a clinical perspective. *Brain Sciences*. 2024;14(12):1261. PMID 39766460 (PMCID PMC11674335)
12. Pall ML. Nitric oxide synthase partial uncoupling as a key switching mechanism for the NO/ONOO cycle. *Medical Hypotheses*. 2007;69(4):821-825. doi:10.1016/j.mehy.2007.01.070
13. Palmer RF, Jaén CR, Walker TJ, Miller CS. Validation of a brief screening instrument for chemical intolerance in a large U.S. national sample. *International Journal of Environmental Research and Public Health*. 2021;18(16):8714. doi:10.3390/ijerph18168714
14. U.S. Environmental Protection Agency. Guide to Air Cleaners in the Home. Updated 2026. Accessed July 22, 2026.
15. U.S. Environmental Protection Agency. What can I do to improve indoor air quality in my home? Updated June 4, 2026. Accessed July 22, 2026.
16. U.S. Environmental Protection Agency. Volatile Organic Compounds' Impact on Indoor Air Quality. Updated 2026. Accessed July 22, 2026.
17. Bornschein S, Hausteiner C, Zilker T, Förstl H. Psychiatric and somatic disorders and multiple chemical sensitivity. *Psychotherapy and Psychosomatics*. 2002;71(4):191-199. doi:10.1159/000063645



18. Ternesten-Hasséus E, Bende M, Millqvist E. Increased capsaicin cough sensitivity in patients with multiple chemical sensitivity. *Journal of Occupational and Environmental Medicine*. 2002;44(11):1012-1017. doi:10.1097/00043764-200211000-00006. PMID 12448352
19. Tran MT, Arendt-Nielsen L, Kupers R, Elberling J. Multiple chemical sensitivity: on the scent of central sensitization. *International Journal of Hygiene and Environmental Health*. 2013;216(2):202-210. doi:10.1016/j.ijheh.2012.02.010. PMID 22487274
20. Millqvist E. Mechanisms of increased airway sensitivity to occupational chemicals and odors. *Current Opinion in Allergy and Clinical Immunology*. 2008;8(2):135-139. doi:10.1097/ACI.0b013e3282f647ec. PMID 18317021
21. Molot J, Sears M, Anisman H. Multiple chemical sensitivity: it's time to catch up to the science. *Neuroscience and Biobehavioral Reviews*. 2023;151:105227. doi:10.1016/j.neubiorev.2023.105227. PMID 37172924
22. Kimata H. Effect of exposure to volatile organic compounds on plasma levels of neuropeptides, nerve growth factor and histamine in patients with self-reported multiple chemical sensitivity. *International Journal of Hygiene and Environmental Health*. 2004;207(2):159-163. doi:10.1078/1438-4639-00262. PMID 15031958

## Additional background reading

1. Damiani G, Alessandrini M, Caccamo D, et al. Italian expert consensus on clinical and therapeutic management of Multiple Chemical Sensitivity. *International Journal of Environmental Research and Public Health*. 2021;18(21):11294. doi:10.3390/ijerph182111294
2. Miller CS. The compelling anomaly of chemical intolerance. *Annals of the New York Academy of Sciences*. 2001;933:1-23. PMID 12000012
3. Cullen MR. The worker with multiple chemical sensitivities: an overview. *Occupational Medicine*. 1987;2(4):655-661. PMID 3328594