



ROOT CAUSE UNLOCKED

Unlock Your *Hashimoto's*

A Complete Functional Medicine Guide to Calming the Autoimmune Attack, Addressing the Root Triggers, and Reclaiming Your Thyroid Health

Dr. Adam Seay, DC

Root Cause Unlocked, Garner, NC · rootcauseunlocked.com · (919) 662-0520

Unlock Your Hashimoto's

*A Complete Functional Medicine Guide to Calming the Autoimmune Attack,
Addressing the Root Triggers, and Reclaiming Your Thyroid Health*

BEFORE YOU START

Contents

INTRODUCTION

The Medication Is Not Enough

KEY 1

What Hashimoto's Actually Is

KEY 2

The Complete Testing Framework

KEY 3

The Gut-Thyroid Connection

KEY 4

Lyme Disease and the Thyroid

KEY 5

Drainage First

KEY 6

The Environmental Drivers, and Which Ones Actually Hold Up

KEY 7

The Anti-Inflammatory Diet for Hashimoto's

KEY 8

Strategic Supplementation for Hashimoto's

KEY 9

Your Next Steps with Root Health

References

The Medication Is Not Enough

There is a conversation that happens in functional medicine practices every week, repeated in thousands of variations but always carrying the same essential frustration. A patient sits down and explains that they have been on thyroid medication for two, five, sometimes ten years. Their TSH is controlled. Their endocrinologist is satisfied. But they are exhausted in a way that sleep does not touch. They cannot lose weight despite genuine effort. Their hair continues to thin. Their brain fog is worse than ever. Their mood is unpredictable. They feel nothing like themselves.

The medication is doing exactly what it was prescribed to do. It is replacing the thyroid hormone that the gland is no longer producing adequately. What it is not doing, and was never designed to do, is stop the immune system from attacking the thyroid. In Hashimoto's thyroiditis, the root problem is the immune system, not the thyroid. Treating the thyroid in isolation while leaving the immune attack unaddressed addresses the hormone deficiency but not the process driving it.

Hashimoto's thyroiditis is the most common autoimmune disease in the United States and the primary cause of hypothyroidism in developed countries.¹ Despite its prevalence, the majority of patients who carry the diagnosis have never been told what drives the autoimmune attack, what maintains it, or what specifically can be done to reduce it. They have been given a medication and told their labs are managed. The progressive destruction of their thyroid tissue continues.

This guide takes a different position. The autoimmune attack in Hashimoto's is not random and it is not irreversible in all patients. It has specific, identifiable, and addressable triggers. Gut permeability allows antigen traffic that activates and sustains the immune response. Gluten cross-reactivity produces antibodies that attack both intestinal tissue and thyroid tissue through molecular mimicry. A common foodborne pathogen called *Yersinia enterocolitica* produces an outer membrane protein associated with cross-reactive immunity against the TSH receptor. *Helicobacter pylori* gastric colonization drives both molecular mimicry and secondary gut permeability that amplifies the autoimmune signal. Gut dysbiosis creates an LPS-driven inflammatory environment that sustains Th17 polarization. Lyme disease drives the same immune polarization that characterizes Hashimoto's lymphocytic thyroid infiltration. Nutrient deficiencies impair the enzyme systems that govern thyroid hormone production and conversion.

Every one of these drivers is testable. Every one is addressable. In my clinical experience, the patients who recover their energy and cognitive function are the ones whose care addresses these upstream factors rather than only the downstream hormone deficiency.

One correction to make before you read another page, because it governs everything that follows.

You have almost certainly been told that your antibodies matter and that lowering them is the goal. Antibody titers are the most-measured, most-discussed, and least outcome-validated number in this condition. Several interventions genuinely lower them. What nobody has demonstrated is that lowering them makes

you feel better or changes where your disease goes. They are still worth watching. They are just not a scoreboard, and you should be suspicious of anyone, including me, whose success metric is a falling TPO number. Throughout this book I use antibodies to establish that the process is autoimmune and to monitor it. I do not use them as the outcome, and neither should you. The outcomes worth tracking are energy, cognition, sleep, hair, cold tolerance, and how you feel by mid-afternoon.

What you are buying here, stated plainly. The mechanisms in this book are drawn from immunology and from association studies, and they are largely sound as mechanisms. What is thin, almost everywhere in this field, is outcome evidence: trials showing that acting on a given driver changes how a Hashimoto's patient feels or where the disease goes. Selenium and vitamin D have antibody data of varying quality and essentially no quality-of-life data. The infectious triggers have association data and, in the case of H. pylori, eradication data. The rest is reasoned clinical practice held to an endpoint. That is what happens when nobody can patent the intervention: the trials never get funded. So read this as a framework for investigation rather than as proven protocol, put your money in the order the evidence sits, and hold everything you start to a defined endpoint.

This guide walks you through eight keys. Read it as a clinical framework, not as a supplement shopping list. The goal is understanding, and understanding is what makes every subsequent clinical decision more effective.

What Hashimoto's Actually Is

The distinction between hypothyroidism and Hashimoto's thyroiditis matters clinically, because it separates a treatment plan focused only on hormone replacement from one that also addresses the immune process driving the destruction. Understanding what Hashimoto's actually is, at the mechanistic level, is the foundation on which everything else in this guide rests.

Hypothyroidism vs. Hashimoto's

Hypothyroidism is a state: insufficient thyroid hormone production. Hashimoto's thyroiditis is a cause: an autoimmune disease in which the immune system generates antibodies against thyroid tissue, progressively destroying the gland's functional capacity. Most cases of hypothyroidism in the United States are caused by Hashimoto's, but the reverse is not true. Not all Hashimoto's patients are hypothyroid, at least not at diagnosis. Early in the disease process, the inflammatory destruction can actually cause transient hyperthyroid episodes as stored thyroid hormone leaks from damaged tissue into circulation. Over time, as destruction progresses, the gland loses functional mass and hypothyroidism becomes established.

This distinction matters because it determines what you investigate. For a physician focused on hypothyroidism, a controlled TSH is the endpoint. For a clinician addressing Hashimoto's, the controlled TSH is where the work starts rather than where it stops, because hormone replacement addresses the hormone deficiency and does nothing to the immune process, the gut, the nutrient status, or the load. That is the reason people can be perfectly medicated and still feel unwell, and it is why this book exists. Note what I am not saying: I am not saying a persistently high antibody number means treatment has failed. Antibodies establish that the process is autoimmune and are worth monitoring; they have not been shown to track how you feel or where you are heading.

The Autoimmune Mechanism

The immune system maintains a critical distinction between self and non-self. Tolerance to self-antigens is established during immune development through a process of eliminating or suppressing immune cells that react to the body's own proteins. When this tolerance breaks down, the immune system generates antibodies and cytotoxic cells that attack the body's own tissues. This is the definition of autoimmunity.

In Hashimoto's, the primary antibody targets are thyroid peroxidase (TPO), the enzyme essential for thyroid hormone synthesis, and thyroglobulin, the protein matrix in which thyroid hormones are stored. When the immune system produces TPO antibodies (TPOAb) and thyroglobulin antibodies (TgAb), it recruits lymphocytes into the thyroid gland in a process called lymphocytic infiltration. These lymphocytes release cytokines that drive chronic inflammation within the gland. Over years, the inflammatory destruction reduces the number of functioning follicular cells, and thyroid hormone output declines.

This process can be active for years or decades before TSH rises above the conventional laboratory cutoff for hypothyroidism. Patients with significant antibody elevations and progressive thyroid destruction may have entirely normal TSH results during this period. This is why antibody testing is not optional in a thyroid workup and why a normal TSH does not rule out active Hashimoto's disease.

The Thyroid-Immune Bidirectional Loop

One of the less-discussed aspects of Hashimoto's is that the destruction of thyroid tissue itself perpetuates the autoimmune response. As thyroid cells are damaged and lysed by lymphocytic attack, their contents, including TPO and thyroglobulin, are released into circulation. This ongoing antigen exposure continuously stimulates the immune system to produce more antibodies. The process is self-reinforcing. This is the mechanism that explains why, without addressing the upstream triggers that initiated the immune response, the autoimmune attack tends to progress over time rather than spontaneously resolving.

Addressing Hashimoto's clinically means two things simultaneously: reducing the upstream drivers that are activating and sustaining the immune response, and supporting the nutritional and metabolic conditions the immune system depends on. Neither alone is sufficient. What that combination is aimed at is how you function, not a titer. Judge it there.

The Complete Testing Framework

The standard thyroid test, a TSH drawn at an annual physical, is the beginning of a workup for Hashimoto's, not the end. Understanding what each marker reveals, and which markers are routinely omitted that should not be, is essential for any patient who wants to understand the full picture of their autoimmune thyroid status.

The Core Thyroid Panel

TSH, or thyroid-stimulating hormone, is produced by the pituitary gland and reflects the brain's assessment of circulating thyroid hormone levels. It is a valuable screening marker, but its limitations are significant in the context of Hashimoto's. TSH can remain in the conventional normal range (0.4 to 4.5 mIU/L) for years while active autoimmune destruction is progressing. The functional target I work toward, 1.0 to 2.5 mIU/L, is considerably tighter and more clinically meaningful for patients with active Hashimoto's. It is the same target printed in the lab interpretation guide in this kit, and it is a clinical target used to guide investigation rather than a validated diagnostic threshold.

Free T4 is the primary hormone produced by the thyroid gland. It is inactive until converted to the active T3 form. Free T3 is the active thyroid hormone that binds to receptors on virtually every cell in the body. Low Free T3, even with normal TSH and normal Free T4, produces the full clinical picture of hypothyroidism at the cellular level and is among the most common explanations for persistent symptoms in medicated Hashimoto's patients. The conversion of T4 to T3 depends on selenium, zinc, adequate liver function, and low inflammatory burden, all of which are commonly compromised in Hashimoto's.

Reverse T3 is an inactive metabolite of T4 that competes with active T3 for receptor binding. When the body is under chronic inflammatory stress, as it is in active Hashimoto's, T4 is preferentially converted to Reverse T3 rather than active T3. Elevated Reverse T3 produces cellular hypothyroidism even when circulating hormone levels appear adequate. The Free T3 to Reverse T3 ratio is one of the most useful single numbers in a Hashimoto's workup.

TPO antibodies and thyroglobulin antibodies are the markers that establish the process is autoimmune rather than something else. That is diagnostically useful, and positive antibodies alongside a normal TSH also predict a higher likelihood of developing hypothyroidism later, which justifies monitoring. What they are not is a treatment target. Several interventions lower them, and none has been shown to make anyone feel better by doing so, so I do not chase them toward undetectable and I would treat any protocol that judges itself on your falling TPO number with suspicion: it has chosen the outcome that is easier to move, not the one you came for. The lab guide in this kit takes this apart in full.

The Extended Hashimoto's Workup

Beyond the core thyroid panel, a complete Hashimoto's workup at Root Health includes several additional assessments that address the upstream drivers of the autoimmune process.

Test	What It Reveals	Clinical Significance
Vibrant Hormone Zoomer	Full thyroid panel (TSH, Free T3, Free T4, Reverse T3, TPOAb, TgAb), adrenal cortisol rhythm, comprehensive sex hormones	Single draw captures thyroid conversion, autoimmune activity, and adrenal status.
Vibrant Wheat Zoomer	Tight junction antibodies (occludin/zonulin), gluten and wheat peptide reactivity across 26 antigens	Suggestive rather than diagnostic for gut permeability and gluten reactivity. Does not replace celiac serology.
LabCorp H. pylori Stool Antigen or Urea Breath Test	Active H. pylori gastric colonization. Stool antigen or urea breath test are preferred over serology for detection of active infection.	Inexpensive, high-yield first-line screen. Eradication studies show measurable TPO antibody reduction. Order before the expensive panels.
LabCorp Yersinia Serology (IgG/IgA/IgM)	Antibodies to Yersinia enterocolitica, a foodborne pathogen associated with thyroid autoimmunity in cross-sectional research.	Consider in non-responders after initial gut and dietary intervention. Not on any standard thyroid or tick-borne panel. Order through the office.
Vibrant Tickborne 2.0 PCR	Borrelia burgdorferi (41 antigens), Babesia, Bartonella, Ehrlichia, Anaplasma	For patients in tick-endemic areas or with non-resolving antibodies after gut and dietary intervention. The most expensive panel on this list, which is why it comes last.
LabCorp EBV Panel (VCA IgG/IgM, EA IgG, EBNA IgG)	Distinguishes past resolved EBV infection, active reactivation (EA positive + VCA IgG positive + EBNA positive), and recent primary infection (VCA IgM positive)	EBV is a plausible driver rather than a certainty, and there is no validated antiviral protocol to act on a positive with. Add when fatigue, lymph node history, or immune dysregulation are prominent.
Vibrant Gut Zoomer Complete	Microbiome composition, pathogen detection, permeability markers, secretory IgA, short-chain fatty acids, Blastocystis hominis	Essential when gut symptoms are present, when antibodies are not responding, or to characterize dysbiosis before antimicrobial intervention.
Vibrant Micronutrient Panel	Selenium, zinc, iodine status, B vitamins, vitamin D, magnesium, iron	Nutrient deficiencies impair T4-to-T3 conversion. Ferritin, B12, folate, zinc, and vitamin D can also be run individually for a fraction of the cost, and often should be.

The order of investigation should be guided by the clinical picture. For all confirmed Hashimoto's patients, the Hormone Zoomer and H. pylori stool antigen are the standard first-line tests, providing the highest yield at the lowest cost. Yersinia serology is added for patients who are not responding to initial gut and dietary

intervention. Wheat Zoomer is added when gluten reactivity is suspected. Tickborne 2.0 PCR is added for patients with tick exposure history or antibodies not responding after gut and infectious workup.

The Gut–Thyroid Connection

The relationship between gut health and thyroid autoimmunity is mechanistic and specific, and it is addressable. Understanding the gut-thyroid pathways in Hashimoto's changes the entire treatment framework from one focused only on the thyroid to one that treats the upstream drivers where the autoimmune process actually begins.

Pathway One: Intestinal Permeability

The gut barrier, when functioning correctly, allows properly digested nutrients to enter circulation while blocking undigested proteins, bacterial endotoxins, and pathogenic organisms. When tight junction proteins are disrupted, this barrier becomes permeable, and substances that should remain in the gut lumen gain access to systemic circulation. The immune system encounters these foreign antigens and mounts a response.

In Hashimoto's, the specific consequence of this increased antigen exposure is activation of the autoimmune cascade through a process called bystander activation, in which immune cells activated against gut-derived antigens generate cross-reactive responses against thyroid tissue.² The persistent antigen traffic through a leaky gut maintains a chronic state of immune activation that keeps thyroid antibody production elevated even when other triggers are addressed.

This is the reasoning behind adding gut barrier repair when dietary change alone has not helped. One caution on how far to push that reasoning: a 2025 study found intestinal permeability to be a modifiable contributor in Hashimoto's with causality unresolved.² “Modifiable contributor with causality unresolved” is a fair reason to work on the gut. It is not a promise that fixing the gut resolves the disease, and read any permeability panel as a pattern rather than as proof from a single marker.

Pathway Two: Gluten and Molecular Mimicry

Gliadin, the immunogenic fraction of gluten, is proposed to share structural epitopes with thyroid tissue, so that in patients carrying HLA-DQ2 or HLA-DQ8 the immune response to gliadin generates antibodies that cross-react with thyroid antigens. That is molecular mimicry, it is a coherent immunological hypothesis, and the effect of gluten-free diets in non-celiac Hashimoto's has been examined in a systematic review and meta-analysis.³

Here is where I part company with most thyroid content, though I am not dismissing gluten. The standard advice is: you have Hashimoto's, go gluten free. I think that advice has the order of operations backwards, and the diet guide in this kit lays out why in full, including a head-to-head trial and a Mendelian randomization analysis. The short version is that in celiac disease the pathogenic role of gluten is firmly

established, while in Hashimoto's gluten is better viewed as a **candidate modifier rather than a proven universal trigger**.

So here is the sequence I actually use.

Get tested for celiac disease first, while you are still eating gluten. Hashimoto's raises your celiac risk measurably. Celiac is a real diagnosis with real testing and a treatment that genuinely works, and that testing depends on you currently eating gluten. Go gluten free first and you lose the ability to get a clean answer without a deliberate gluten challenge later. This is the single most valuable action in the whole dietary section, and skipping it is the most common expensive mistake I see.

If celiac testing is positive, gluten free is the treatment. Not a wellness experiment. A treatment, and strict is the right word there, because cross-contamination from shared surfaces, wheat starch in processed foods, and soy sauce all count.

If celiac testing is negative and you still want to try it, that is reasonable. Run it as a time-limited experiment with a defined endpoint, eight to twelve weeks, judged on how you feel rather than on a repeat antibody panel.

The Wheat Zoomer panel tests reactivity to 26 gluten and wheat peptides beyond the standard celiac antigens, which can be informative when celiac serology is negative and the question is still open. Read it as suggestive rather than diagnostic. It does not replace celiac serology and it does not establish that gluten is driving your thyroid disease.

Pathway Three: Yersinia Enterocolitica

Yersinia enterocolitica is a gram-negative foodborne bacterium transmitted through contaminated pork, unpasteurized dairy, and undercooked meats. It is not a tick-borne organism, and no standard thyroid panel covers it. Its association with Hashimoto's thyroiditis is one of the more studied infectious connections in thyroid autoimmunity research, with multiple cross-sectional studies documenting significantly elevated *Yersinia* antibodies in Hashimoto's patient populations compared to healthy controls.

The proposed mechanism involves a low molecular weight envelope protein (LP) produced by *Yersinia* that shares structural homology with the human TSH receptor.⁴ The hypothesis is that antibodies generated against this protein during *Yersinia* infection may cross-react with TSH receptor epitopes through molecular mimicry. This mechanism is biologically plausible and supported by the cross-sectional association data. A prospective cohort study (Efthymiou et al., 2011) did not find *Yersinia* seropositivity predictive of de novo thyroid autoimmunity development, which means the causal relationship has not been confirmed and the association may reflect shared genetic susceptibility.⁵ The clinical position I take at Root Health is that *Yersinia* is worth ruling out in patients who are not responding to dietary and gut barrier intervention, because identified *Yersinia* burden is addressable and the test is inexpensive.

Testing is a standalone LabCorp order for *Yersinia enterocolitica* IgG, IgA, and IgM serology. Elevated IgG indicates past or chronic exposure. A raised IgA suggests active mucosal infection, and a raised IgM

indicates recent or acute infection. The infection can be subclinical, producing no gastrointestinal symptoms while maintaining an antigenic burden that keeps thyroid antibodies elevated.

Clinical Note: *Elevated TPO antibodies that do not normalize despite dietary intervention and gut repair are a strong clinical signal that an upstream infectious trigger is maintaining the immune activation. H. pylori and Yersinia are the two most cost-effective investigations at that point. Do not stop investigating at the gut barrier.*

Pathway Four: H. pylori and Gastric Immune Activation

Helicobacter pylori is a gram-negative bacterium that colonizes the gastric mucosa in approximately 44 percent of the global population, making it one of the most prevalent chronic bacterial infections in the world. Its connection to Hashimoto's thyroiditis is supported by multiple cross-sectional studies showing significantly elevated H. pylori seroprevalence in Hashimoto's patients compared to healthy controls, and by eradication studies demonstrating measurable reductions in TPO antibody levels following successful treatment.

The mechanisms are two-pronged. First, the H. pylori CagA protein (cytotoxin-associated gene A) shares structural similarities with thyroid antigens, generating cross-reactive antibodies through molecular mimicry.⁶ Second, H. pylori colonization produces chronic gastric inflammation that progressively impairs stomach acid production. Adequate gastric acid is required for protein denaturation in the stomach. When acid production is insufficient, intact food proteins reach the small intestine without proper denaturation, increasing the antigen load that reaches the gut-associated lymphoid tissue and promoting the permeability-driven immune activation described in Pathway One.

The clinical significance of H. pylori in Hashimoto's management is substantial because eradication data shows actual outcome benefit. Unlike the Yersinia association, which has cross-sectional support but limited prospective outcome data, H. pylori eradication studies have shown direct antibody reduction in Hashimoto's patients following successful treatment. This makes H. pylori one of the highest-yield investigations in Hashimoto's patients who are not responding to dietary and gut barrier intervention. Testing is inexpensive: a LabCorp stool antigen or urea breath test detects active infection with high sensitivity. Serology is less preferred because it cannot distinguish active infection from past exposure.

There is a second reason to care about H. pylori here, and it has nothing to do with autoimmunity. Levothyroxine needs stomach acid to dissolve before it can be absorbed, and H. pylori gastritis lowers acid production. In a New England Journal of Medicine study, patients with H. pylori-related gastritis, atrophic gastritis, or both required 22 to 34 percent more thyroxine than a reference group, and eradicating the organism largely reversed the effect.²³ So in someone whose TSH will not settle despite dose increases, H. pylori may be driving the autoimmune process and blocking the treatment for it at the same time. The lab guide in this kit covers what to do about that.

Pathway Five: Gut Dysbiosis and the LPS-Autoimmunity Loop

Gut dysbiosis, an imbalance in the microbial ecosystem characterized by reduced diversity, depletion of beneficial short-chain fatty acid producers, and relative overgrowth of pro-inflammatory gram-negative species, creates a systemic inflammatory environment that sustains Hashimoto's autoimmunity through a mechanism distinct from gut permeability itself. The two pathways interact, but understanding them separately improves the precision of clinical intervention.

The primary mechanism is LPS endotoxemia. Lipopolysaccharide, the outer membrane component of gram-negative bacteria, ranks among the most potent activators of the innate immune system via Toll-like receptor 4 signaling. In a healthy microbiome with adequate gut barrier function, LPS is contained within the intestinal lumen. When dysbiosis increases the gram-negative bacterial load and gut permeability allows LPS translocation into systemic circulation, TLR-4 activation drives a pro-inflammatory cytokine cascade that sustains the Th17-dominant immune environment characteristic of Hashimoto's.⁷ The autoimmune thyroid attack is, at its base, a Th17-mediated process. Persistent LPS signaling from a dysbiotic gut is a consistent upstream driver of that polarization.

Short-chain fatty acids, particularly butyrate produced by Firmicutes species including *Faecalibacterium prausnitzii* and *Akkermansia muciniphila*, are the primary energy source for colonocytes and are essential for maintaining tight junction integrity and regulatory T-cell function. Treg cells, which suppress autoimmune reactivity, are butyrate-dependent. Dysbiosis reduces SCFA production, which simultaneously impairs gut barrier function and reduces the Treg activity that would normally suppress the autoimmune response. This makes microbiome restoration a distinct intervention from gut barrier repair, addressing a different layer of the same problem.

One specific organism within the dysbiosis picture warrants individual attention: *Blastocystis hominis*, a parasitic protozoan found at significantly elevated rates in Hashimoto's patients compared to healthy controls in multiple studies. *Blastocystis* disrupts tight junction proteins, elevates zonulin, and generates antigen traffic through the gut barrier that sustains autoimmune activation. It does not produce prominent gastrointestinal symptoms in most carriers, making it easy to miss without targeted testing. *Blastocystis* is detected on the Gut Zoomer Complete, so no additional panel is required. Its presence should be treated as a contributing driver of gut permeability rather than incidental colonization in the context of active Hashimoto's.

The Gut Zoomer Complete is the appropriate assessment for dysbiosis characterization in Hashimoto's patients. It provides microbiome composition, short-chain fatty acid status, pathogen detection including *H. pylori*, *Blastocystis*, and other parasitic organisms, permeability markers, and secretory IgA in a single panel. The combination of dysbiosis profiling with permeability assessment in a single draw makes it the most clinically efficient gut workup for the Hashimoto's patient who is not responding to initial dietary intervention.

Pathway Six: Epstein-Barr Virus Reactivation

Epstein-Barr Virus (EBV) is the most significant viral trigger in Hashimoto's thyroiditis and the most consistently underinvestigated in clinical practice. Over 90 percent of adults carry EBV as a latent infection following primary exposure, typically in adolescence. The distinction that matters for Hashimoto's is reactivation rather than past exposure: the shift from latent viral dormancy to active viral replication, which generates the immune activation that drives thyroid autoimmunity through multiple overlapping mechanisms.⁸

The molecular mimicry evidence for EBV is more experimentally grounded than for most bacterial triggers. The EBV nuclear antigen EBNA-1 shares structural epitopes with thyroid autoantigens, and cross-reactive antibodies against EBNA-1 and TPO have been demonstrated in patient populations. EBV can also directly infect thyroid follicular cells, establishing a reservoir of viral replication within the thyroid gland itself that generates ongoing immune activation at the site of autoimmune attack. This direct thyroid tropism distinguishes EBV from most other infectious triggers, which operate through systemic immune mechanisms rather than local thyroid infection.

EBV reactivation drives the same Th17 polarization pathway shared by Lyme disease and active Hashimoto's. EBV-driven immune activation suppresses regulatory T-cell function, promotes IL-17 production, and creates the immunological environment in which thyroid autoimmunity escalates. In patients with both EBV reactivation and elevated thyroid antibodies, the viral reactivation is maintaining the immune polarization that makes antibody reduction difficult regardless of dietary and gut interventions.

Testing distinguishes four clinically distinct states. A positive VCA IgM indicates acute or recent primary infection. When VCA IgG is positive but EBNA IgG is negative, the picture is recent primary infection with incomplete seroconversion. Active reactivation, the state most clinically relevant to Hashimoto's, shows as VCA IgG, EBNA IgG, and early antigen (EA) IgG all positive. Past resolved infection with no current reactivation looks the same, except that EA is negative. The full four-marker panel is available through LabCorp and is one of the less expensive items on the testing list. EBV testing is worth considering for any Hashimoto's patient with prominent fatigue, lymph node involvement history, a history of mononucleosis, or a picture that has not moved after the gut, dietary, and bacterial workup is done.

Now the part that most content on this subject leaves out. Everything above is mechanistic and associational. A 2025 review integrates the immunological and virological evidence for EBV across thyroid disease including Hashimoto's, and what it is not is a trial showing that treating a virus changes thyroid antibodies or changes how a patient feels. **There is no validated antiviral protocol for Hashimoto's, and anyone selling you one is ahead of the evidence.** That includes me if I ever describe one to you as established.

So here is what I actually do with a confirmed reactivation pattern, stated at its real evidentiary weight. Prescription antivirals are a drug class that some physicians use for specific viral indications. They are not something I prescribe, and they are mentioned here so you know they exist and can raise them with the physician managing that part of your care. On my side, I support the immune system generally rather than

claiming to treat the virus: sleep, load reduction, correcting documented nutrient deficiencies, and the general immune support tools discussed in Key 8. Those are cheap, low-risk, and defensible on their own terms whatever the virus is doing. What I do not do is build an expensive open-ended antiviral supplement protocol around a marker, because the evidence for that does not exist and the cost is real.

Other Infectious Considerations

Several additional infectious agents have documented or proposed associations with Hashimoto's that are clinically relevant in specific patient contexts.

Hepatitis C Virus: well-established bidirectional relationship. HCV can trigger thyroid autoimmunity through direct thyroid infection and molecular mimicry. Interferon-alpha treatment for HCV is a known thyroid autoimmunity trigger. Relevant for any patient with HCV history or prior interferon treatment. Standard HCV antibody testing applies.

Post-COVID Thyroiditis: SARS-CoV-2 infects thyroid follicular cells via ACE2 receptors, confirmed in autopsy studies.⁹ Post-COVID Hashimoto's-pattern thyroiditis is increasingly documented. For any patient whose antibody escalation followed COVID infection, this context affects treatment framing. No specific additional test; antibody trajectory tracking applies.

Klebsiella pneumoniae and Mycoplasma pneumoniae: cross-sectional association data exist for both. Klebsiella shares thyroid antigen homology through molecular mimicry. Mycoplasma produces immune activation via cold agglutinin mechanisms. Consider in patients with respiratory illness history preceding Hashimoto's onset. LabCorp Mycoplasma IgG/IgM available if clinically indicated.

Parvovirus B19: associated with multiple autoimmune conditions including thyroid autoimmunity. IgG/IgM available through LabCorp. Consider in patients with joint symptoms alongside thyroid antibody elevation.

Lyme Disease and the Thyroid

The biological overlap between Lyme disease and Hashimoto's thyroiditis is clinically important for a specific subset of patients: those in tick-endemic areas whose antibodies are not responding to gut and dietary intervention. The immunological mechanisms are plausible and the shared Th17 pathway is well-characterized. This section presents that evidence accurately, including where it is strong and where it remains at the level of working hypothesis.

The Borrelia-Thyroid Shared Immune Environment

A working hypothesis in functional immunology proposes that *Borrelia burgdorferi* outer surface proteins, particularly OspA and OspC, may share structural similarity with human thyroid tissue antigens. If confirmed, antibodies generated against these proteins during *Borrelia* infection could cross-react with thyroid proteins through molecular mimicry. This hypothesis is currently based on in silico sequence homology analysis and has not been confirmed through experimental demonstration of cross-reactive antibodies in patients or through animal models of Lyme-induced thyroiditis. It should be understood as a plausible but unvalidated mechanism, not as established molecular biology.

What is well-established is that chronic Lyme infection and Hashimoto's share a common immune polarization. *Borrelia* drives a sustained Th17-dominant immune environment, characterized by elevated IL-17, IL-21, and IL-6, with suppressed regulatory T-cell activity. This is the same immune environment that drives lymphocytic thyroid infiltration in Hashimoto's. The two conditions share an immune polarization pattern, which means that active Lyme disease creates and maintains conditions that are broadly permissive to autoimmune thyroid activity, regardless of whether direct molecular mimicry is operating.

The practical clinical implication is that in patients with concurrent Lyme and Hashimoto's, reducing the Lyme burden reduces the Th17 inflammatory environment that sustains the thyroid autoimmunity. This is a mechanistically coherent therapeutic target. The limitation is that no study has directly demonstrated TPO antibody reduction specifically attributable to Lyme treatment, so the claim should be framed as a rational clinical inference from shared immunology rather than a proven treatment outcome.

Symptom Overlap and Missed Diagnoses

Lyme disease and Hashimoto's share a significant symptom overlap. Fatigue, brain fog, cold intolerance, weight changes, hair thinning, anxiety, cognitive impairment, mood instability, and joint pain are features of both conditions. In patients presenting with this symptom cluster, each condition can mask the other. A provider who finds the Hashimoto's stops investigating and misses the Lyme, and one who finds the Lyme misses the Hashimoto's the same way. Patients end up being treated for half of their clinical picture.

The gut represents a third layer of interaction. *Borrelia* infection suppresses secretory IgA production, the gut's primary immune defense against pathogenic colonization. This suppression allows dysbiotic organisms to establish in the gut, promoting intestinal permeability that contributes to autoimmune thyroid activation through the mechanisms described in Key 3. Gut repair in Lyme-Hashimoto's patients is not optional because the Lyme infection is actively degrading the gut barrier that needs to be restored for the thyroid autoimmunity to quiet.

Who Should Be Tested

Tick-borne co-infection testing is indicated for Hashimoto's patients who meet any of the following criteria: have lived in or spent significant time in tick-endemic areas, have a history of a tick bite regardless of whether a bull's-eye rash was observed, have antibodies that are not responding to dietary intervention, gut repair, and *H. pylori* and *Yersinia* workup, have significant neurological or psychiatric symptoms alongside thyroid symptoms, or have been previously treated for Lyme disease but continue to have symptoms.

The Vibrant Tickborne 2.0 PCR panel tests *Borrelia burgdorferi* across 41 antigens, more than the standard Western Blot, and it also covers *Babesia*, *Bartonella*, *Ehrlichia*, and *Anaplasma*, co-infections that most standard panels miss. It is available through rootcauseunlocked.com/order-labs. It is the most expensive panel discussed in this book, which is why it belongs as a secondary investigation after the cheaper gut and infectious workup is done.

Drainage First

Every treatment intervention for Hashimoto's, whether dietary, supplemental, or targeted at an infectious trigger, mobilizes waste products, inflammatory mediators, and cellular debris that must be processed and eliminated through the body's drainage systems. The lymphatic system, the liver, and the kidneys bear the primary burden of this clearance. When drainage capacity is insufficient for the load being generated, mobilized toxins recirculate, inflammatory burden increases, and the autoimmune process can actually flare rather than improve.

This is a leading explanation for why patients who begin aggressive Hashimoto's protocols, particularly antimicrobial protocols for *H. pylori*, *Yersinia*, or Lyme, experience a significant worsening of symptoms in the first weeks. The antimicrobial intervention is working. But the drainage capacity cannot handle the microbial debris load. The result is a Herxheimer reaction that the patient interprets as the treatment making them worse.

At Root Health, I apply the drainage-first sequence to every Hashimoto's case regardless of severity or protocol complexity. The drainage triad runs for a minimum of two weeks before any targeted intervention is introduced, and it is maintained throughout the full protocol.

The Drainage Triad for Hashimoto's

Product	Clinical Role	Dosing
Lymph-Tone II (Energetix)	Extracellular matrix and lymphatic drainage for ECM congestion and xenobiotic accumulation. The appropriate selection for most Hashimoto's patients with dietary toxin burden and compensation history. Taken as drops in warm water.	Per label direction
Methylated B12, liposomal or sublingual	Methylation support for Phase 2 liver detoxification. Low methylation capacity impairs clearance of endotoxins, hormones, and the inflammatory byproducts generated by autoimmune activity.	Per label, twice daily
ReHydration (Energetix)	Electrolyte and mineral foundation for cellular hydration. Required for efficient intracellular waste transport into the lymphatic system.	Per label direction

All three are available through my Fullscript dispensary, wherever you are located.

For patients with significant chronicity, greater than three years of active symptoms, or confirmed Lyme co-infection, a longer drainage phase of three to four weeks before introducing antimicrobial support produces

substantially better clinical outcomes. The investment of time in drainage preparation pays dividends in the form of protocols that are tolerated rather than abandoned.

Lymph Formula Selection

Lymph-Tone I fits acute inflammation, a recent exposure, or an active infection phase.

Most Hashimoto's cases belong on Lymph-Tone II, the formula for extracellular matrix congestion and xenobiotic accumulation.

Lymph-Tone III is for longstanding degeneration, heavy metal burden, or symptoms present more than 3 years.

Never use more than one Lymph-Tone formula simultaneously. All are taken as drops in warm water.

The Environmental Drivers, and Which Ones Actually Hold Up

If you have spent any time reading about Hashimoto's online, you have met the halogen argument. It goes like this. Fluoride, bromide, and chlorine sit near iodine on the periodic table, they are chemically similar enough to take iodine's place, and modern life soaks you in all three. Filter your water, throw out your toothpaste, and your thyroid recovers.

Parts of that are true. The parts that are true are not the parts usually emphasized, and the difference matters, because the version circulating sends people after their toothpaste while the exposure with the best evidence behind it goes unmentioned. Here is what I can defend.

A real mechanism, with a name

Your thyroid pulls iodide out of the bloodstream using a transporter called the sodium-iodide symporter, usually written NIS. It is the front door for every iodine atom your thyroid will ever use. Several environmental anions are shaped enough like iodide to be carried through that same door, and while they occupy it, iodide is not moving.¹³

This is established transporter biology rather than a fringe idea, and medicine once used the same mechanism deliberately: potassium perchlorate was given as a drug precisely because it shuts down thyroidal iodide uptake.

What the mechanism does not tell you is whether the amounts in your actual life are enough to matter. That is a separate question, and it has to be answered compound by compound.

Perchlorate, the one with the strongest case

Perchlorate is the most potent NIS inhibitor of the group, and the thyroid transporter has a higher affinity for it than for iodide itself.¹³ It reaches people through contaminated drinking water and through produce grown with perchlorate-containing irrigation water and fertilizer.

A 2025 systematic review and meta-analysis pooled the human studies on NIS inhibitors and found perchlorate exposure positively associated with TSH across the literature.¹⁴ The effect size is small at population level. It is also the only halogen-type exposure in this chapter with pooled human evidence pointing consistently in one direction. This is the exposure I would prioritize, and it is the one almost nobody talks about.

Bromide, where the mechanism is confirmed and the dose is the question

Bromide does compete at NIS. That is not in dispute. The European Food Safety Authority's 2025 assessment of bromide in food and feed identifies the thyroid as a target organ and states that bromide competes with NIS-mediated iodide uptake.¹⁵

The dose is where the popular version overreaches. In the human data that assessment reviewed, thyroid hormone concentrations shifted at intakes around 9 mg per kilogram of body weight per day, and even at that level the values stayed inside normal limits.¹⁵ The tolerable daily intake the committee derived is 0.4 mg per kilogram per day, and it was set conservatively, with a hundredfold uncertainty factor, to protect neurodevelopment rather than because ordinary diets were producing thyroid disease.

So: a confirmed mechanism, at doses well above what a normal diet delivers. Worth reducing where it is easy. Not the thing that gave you Hashimoto's.

The bromide sources worth knowing are brominated flame retardants in furniture and electronics dust, methyl bromide fumigation residues on some conventionally grown produce, and, until the FDA revoked its authorization in 2024, brominated vegetable oil in citrus-flavored soft drinks. On the flame retardants specifically, a 2024 review of the human literature found the results genuinely heterogeneous, with studies split between positive, negative, and null associations with TSH and T4.¹⁶ I am not going to tell you that finding is stronger than it is.

Fluoride, where I part company with most of what you have read

This is the one where being honest costs me the better story.

The best available dose-response meta-analysis, published in 2024, found that TSH did not change across the lower range of water fluoride exposure and began to rise only above roughly 2.5 mg/L.¹⁷ Community water fluoridation in the United States targets 0.7 mg/L, which sits well below the point where that curve turns. The studies driving the finding were also mostly cross-sectional, mostly conducted in populations with naturally high groundwater fluoride, and mostly in children.

A separate 2024 systematic review looked at the same question, found seven studies that met inclusion criteria, and rated overall confidence in the evidence as low for every outcome, with methodological problems and risk of bias in all seven.¹⁸

Put those together and the evidence position is this: fluoride at the levels found in some natural groundwater does appear to affect thyroid function, and at the levels in American municipal tap water it has not been shown to.

My own position is not the same as where that evidence stops, and I would rather tell you that plainly than let you work it out later. I think limiting fluoride, chlorine, and bromine matters for thyroid health and for general health, and I say so to patients. That is a clinical judgment about total halogen load, and it is not something I can hand you a trial for. Weigh it as what it is. What the published literature currently supports and what I

would do in your position are two different statements, and you are getting both rather than one dressed up as the other.

Practically: if you are on well water, test it, because what is in it is knowable and cheap to find out. On treated municipal water, filtration is still worth doing, and the action list at the end of this chapter says how far I would take it.

Chlorine, where the evidence is absent and I reduce it anyway

Chlorine is the weakest link in the periodic-table argument. I could not find human evidence tying chlorinated drinking water at treatment concentrations to thyroid dysfunction or to thyroid autoimmunity, and I looked. The claim travels because it completes a tidy pattern, not because someone measured it.

So I will not tell you there is a study behind this one. There is not. I still reduce chlorine exposure, for the same reason given in the fluoride section: it belongs to the total halogen load, the reduction is cheap and carries no downside, and a carbon filter that costs very little handles it. That is my judgment rather than a finding, and now you can tell the difference between the two, which is more than most of what you will read on this subject offers you.

The part the halogen story gets backwards

Every halogen argument rests on the same premise: these compounds cause a relative iodine shortage, therefore your thyroid is starved, therefore you should be worried about getting more iodine. In an iodine-deficient population that premise holds. The review literature is consistent that the harms from NIS inhibitors concentrate in iodine-deficient people, in pregnancy, and in children.¹³

But Hashimoto's is not an iodine-deficiency disease, so the premise does not transfer cleanly, and an elevated halogen reading is not by itself a reason to start taking iodine.

I want to be careful here, because my position on iodine is not the one you will find in most Hashimoto's material. I do use iodine clinically in this condition, including kelp, and the diet guide in this kit lays out the reasoning in full: the antibody rise that iodine can produce appears from population data to be a phase rather than a permanent state, thyroid function is minimally disrupted through it, and selenium is the other half of the reaction and goes alongside.²¹²² What I do not do is start iodine because someone's perchlorate came back high. Halogen exposure is an argument for reducing the exposure. Iodine is a separate decision, made deliberately, with selenium, and with monitoring.

Beyond the halogens

Two other exposures have better human data than chlorine and get a fraction of the attention.

Mercury. In an analysis of US women in the 2007 to 2008 NHANES survey, women in the highest blood mercury group had 2.24 times the odds of thyroglobulin antibody positivity, with a confidence interval of 1.22

to 4.12.¹⁹ The same relationship did not appear for thyroid peroxidase antibodies. This is a cross-sectional association in one survey, not a demonstration that mercury causes thyroid autoimmunity, and I want to be precise about that. It is still a finding about thyroid autoantibodies, which is more directly relevant to Hashimoto's than most of the halogen literature.

Bisphenols. A 2025 systematic review and meta-analysis of bisphenol exposure and the hypothalamic-pituitary-thyroid axis found associations with TSH and free T4 in adults, and reported that adult women were more affected than adult men.²⁰ The effect sizes are small and the direction differs by sex, so this is a signal rather than a settled story.

What you are buying in this chapter, stated plainly

Most of this is mechanistic and observational. There is no randomized trial in which reducing an environmental exposure improved Hashimoto's, and I do not expect one, because nobody funds trials on interventions that cannot be patented. What exists is transporter biology that is well characterized, human association studies of varying quality, and regulatory assessments that took the mechanism seriously enough to set intake limits.

I treat that as reason to reduce exposure where reduction is cheap and low-risk, not as reason to reorganize your life.

What I actually do

I do order an environmental exposure panel in Hashimoto's, and I want to be precise about what it does and does not give you.

I use the Vibrant Wellness Environmental Toxins panel, which measures 38 environmental chemicals and includes perchlorate. A urinary panel documents recent exposure. It is suggestive rather than diagnostic for a disease process, and no exposure panel can tell you that a chemical caused your thyroid autoimmunity. Nobody has run a trial showing that testing changes outcomes in Hashimoto's, and I am not going to imply one exists. I order it anyway, because knowing whether perchlorate is actually elevated in you changes how hard I push on water filtration and how closely I watch your iodine status, and because a number in front of someone moves them to act in a way that general advice does not. Read that as my clinical practice rather than as guideline-concordant care.

One limit worth knowing before you pay for it: the panel covers perchlorate, and it does not measure bromide or fluoride. For those two, source control is the only strategy, because there is no test in this workup that will find them for you. The lab guide in this kit covers how I read the panel when it comes back.

Alongside the panel, in this order:

Get iodine status right before anything else. Not by reflex in either direction. Iodine is the most contested question in this condition and the diet guide works through it properly, including where I differ from the standard advice. What matters here is that iodine is decided on its own merits, with selenium alongside and

thyroid function monitored, rather than raised because a halogen result frightened you or dropped because a headline did.

Filter your drinking water. This is the item where I am more aggressive than the evidence strictly requires, and the fluoride section says why. Reverse osmosis is what I suggest when the budget allows, because it removes perchlorate and handles fluoride in the same step. A carbon filter costs far less, reduces chlorine, and is a real improvement over nothing, so it is the answer when reverse osmosis is not realistic. If you are on well water, test it first. Filtration equipment is available from any reputable retailer, and it is not something I sell or profit from.

Reduce the easy bromide sources. Skip the citrus soft drinks, wash and peel conventionally grown produce where practical, and vacuum and dust more often if you have older upholstered furniture, since flame retardants concentrate in household dust.

Handle food storage. Glass rather than plastic for hot food and for anything acidic, and never heat food in plastic. This is the practical end of the bisphenol literature and it costs nothing.

Bring mercury up if your history warrants it. Frequent high-mercury fish consumption, or an occupational exposure, is worth raising with the clinician managing your care. Chelation is not something I do, and not something I would recommend on the strength of the evidence above.

Then let the drainage work from Key 5 carry the load. Reducing what comes in and supporting the systems that clear what is already there are the same project approached from two ends, and the drainage protocol is the half with the clearer clinical rationale.

If you take one thing from this chapter, make it the ranking. Perchlorate is the exposure with the real evidence behind it and the one nobody names, and filtering your water is the single action that addresses it along with the chlorine and fluoride I reduce on judgment rather than on trial data.

The Anti-Inflammatory Diet for Hashimoto's

Diet in Hashimoto's is a targeted clinical intervention, one that addresses specific mechanisms driving the autoimmune process, rather than a general health recommendation. The dietary changes most consistently associated with measurable antibody reduction in clinical research are not general anti-inflammatory eating patterns, though those matter. They are specific to the molecular interactions between dietary components and the immune and gut pathways that sustain the autoimmune attack.

The Removals, in the order I would actually do them

Gluten, handled the way Key 3 describes rather than the way the internet describes. Gliadin is proposed to trigger zonulin release and to generate cross-reactive antibodies against thyroid tissue through molecular mimicry. Both are coherent mechanisms and neither is settled as a driver of your disease. **Test for celiac first, while still eating gluten.** If that is positive, strict elimination is the treatment and strict means labels on every packaged product, asking about preparation when eating out, and treating shared cooking surfaces as a real exposure. If it is negative, an eight to twelve week trial judged on how you feel is reasonable, and a permanent restriction adopted on faith is not.

A note on what the diet evidence actually showed. In the one head-to-head comparison, a Mediterranean pattern outperformed a gluten-free diet on oxidative stress markers, while the gluten-free arm did not move them. No diet in that comparison changed thyroid function or thyroid antibodies. The diet guide in this kit covers it properly. I still think diet matters here. I just think the Mediterranean pattern is the better default and the celiac test is the better first move.

The second removal worth making is conventional dairy, specifically the casein A1 protein present in most commercially available cow's milk products. A1 casein is cleaved during digestion into beta-casomorphin-7, a peptide with opioid receptor activity that has been shown to impair gut barrier integrity and promote intestinal inflammation.¹⁰ In patients with Hashimoto's who are already managing a compromised gut barrier, continuing to consume A1 dairy maintains one of the upstream barrier-disrupting inputs that allows ongoing antigen traffic into systemic circulation. Some patients tolerate A2 dairy, from certain cow breeds, goats, and sheep, without the same permeability consequences.

Refined sugar and high-glycemic carbohydrates create an intestinal environment that favors the dysbiotic organisms most likely to maintain gut permeability and sustain the LPS-driven inflammatory signaling that activates the immune response. For Hashimoto's patients with concurrent *H. pylori* or *Yersinia* colonization, high sugar intake provides a substrate that supports bacterial survival in the gut ecosystem. Removal of refined carbohydrates is both a gut barrier intervention and an anti-pathogenic dietary strategy.

Seed oils, particularly canola, soybean, corn, sunflower, and safflower oils, promote the arachidonic acid cascade and generate pro-inflammatory prostaglandins that amplify the cytokine environment in which

lymphocytic thyroid infiltration occurs. The omega-6 to omega-3 ratio in a standard Western diet is approximately 15 to 20:1. The ratio associated with low autoimmune reactivity is closer to 4:1. Replacing seed oils with olive oil, avocado oil, coconut oil, and fat from grass-fed animal sources shifts this ratio and reduces the inflammatory signaling that drives autoimmune activity at the thyroid.

What to Prioritize

Wild-caught fatty fish, particularly salmon, mackerel, and sardines, provide EPA and DHA that directly resolve inflammatory signaling through specialized pro-resolving lipid mediators. These are not merely anti-inflammatory in a general sense. They actively promote immune resolution. Consuming them three to four times per week during the active treatment phase provides a consistent substrate for inflammatory resolution that cannot be replicated with supplementation alone at the same clinical impact.

Selenium-rich foods deserve particular attention in Hashimoto's. Selenium is essential for the selenoprotein thioredoxin reductase, which regulates the redox environment of the thyroid gland and protects thyroid cells from the hydrogen peroxide generated during hormone synthesis. Brazil nuts are by far the most concentrated food source, with a single nut providing approximately 70 to 90 micrograms. Two to three Brazil nuts per day provides therapeutic selenium intake for most patients. The evidence for selenium in reducing TPO antibody levels is among the more consistent in Hashimoto's nutritional research, though the magnitude of antibody reduction varies across trials.¹¹

Cruciferous vegetables, including broccoli, cauliflower, Brussels sprouts, and cabbage, require individualized guidance in Hashimoto's. Raw cruciferous vegetables contain glucosinolates that can be converted to goitrogens, compounds that interfere with iodine uptake by the thyroid. For patients with adequate iodine status, this is not a significant concern in moderate amounts. Cooking deactivates the majority of the goitrogenic compounds and makes cruciferous vegetables safe for most Hashimoto's patients in standard dietary quantities.

Strategic Supplementation for Hashimoto's

The supplement protocol for Hashimoto's follows the same sequencing principle as every protocol at Root Health: drainage must be established before targeted interventions are introduced, and targeted interventions are selected based on lab findings rather than applied universally. The products named here are the practitioner-grade formulations I use in clinical practice. The framework below explains the clinical reasoning so that you understand what each phase is accomplishing and why the sequence matters.

Phase One: Drainage (Weeks 1 through 2, continued throughout)

A lymphatic botanical, methylated B12, and an electrolyte and trace mineral base form the drainage foundation as described in Key 5. No targeted thyroid or immune interventions are introduced until the drainage triad has been running for a minimum of two weeks. For patients with significant Lyme co-infection or heavy metal burden, three to four weeks of drainage preparation is the clinical standard.

Phase Two: Gut Barrier Repair (Weeks 3 through 6)

Gut barrier repair is the second priority because restoring the physical barrier that prevents ongoing antigen traffic into systemic circulation is prerequisite to meaningful autoimmune modulation. L-glutamine, ideally in a powder that also supplies glycine and proline, provides the amino acid substrate for tight junction protein synthesis and enterocyte regeneration. Bovine colostrum supports the gut-associated lymphoid tissue function that regulates the immune response to gut microbial inhabitants. Bioavailable curcumin provides anti-inflammatory support for the gut environment that accompanies most Hashimoto's cases.

For patients with confirmed celiac disease, strict gluten elimination runs concurrently with gut barrier repair from day one, because there the removal is the treatment. For patients with reactivity on the Wheat Zoomer but negative celiac serology, I run the gluten trial as a time-limited experiment alongside the barrier work and we decide honestly at the end of it.

Phase Three: Thyroid and Immune Support (Weeks 4 through 8)

A thyroid support formula supplying the nutritional cofactors of thyroid hormone production, chiefly selenium, zinc, iron, tyrosine, and vitamin A, is the primary matrix support here. I prefer it over an iodine-focused formula as the default matrix support, because iodine deserves to be a deliberate decision at a known dose rather than an incidental ingredient in a blend. Iodine has a place in this condition. Where I do use iodine, and I do, it goes in knowingly and with selenium alongside. The diet guide covers that reasoning.

Methylated B12 continues from the drainage phase through the full protocol. Beyond its methylation and drainage role, B12 supports neurological function, mood stability, and energy metabolism, all of which are commonly compromised in Hashimoto's patients due to the metabolic burden of chronic autoimmunity.

Vitamin D3 with K2 as MK-7 addresses the vitamin D deficiency that is nearly universal in Hashimoto's patients. Vitamin D functions as an immune modulator, promoting Treg cell activity that suppresses autoimmune reactivity, and correlates inversely with TPO antibody levels in multiple clinical studies.¹² The target is 40 to 80 ng/mL, the same functional range I use across every program, with ongoing monitoring for calcium metabolism in patients requiring extended high-dose supplementation to reach that level.

Phase Four: Targeted Antimicrobial Support (Weeks 6 through 12, based on lab findings)

When EBV reactivation is present on the four-marker panel, general immune support is introduced after drainage and gut barrier repair are established. To restate what Key 3 said, because it matters more than the ingredient list: there is no validated antiviral protocol for Hashimoto's. Lysine is proposed to compete with arginine in viral replication and monolaurin has antiviral activity in laboratory work; both are laboratory findings rather than demonstrated clinical outcomes in this condition, and they should be read that way. Zinc, vitamin C, and adaptogens are general immune support with a reasonable safety profile. EBV is never eradicated in anyone, and the honest framing is that this phase supports the host rather than treating the virus. Hold it to a defined window and judge it on how you feel, not on a titer.

When *H. pylori* is identified, support is introduced after drainage and gut barrier repair are established. Eradication itself is a prescription antibiotic regimen that some physicians use for confirmed *H. pylori* infection. It is not something I prescribe, and it is included here so you know it exists and can raise it with the physician managing that part of your care. This is the one infectious trigger in the book where eradication has outcome evidence behind it, so it is worth having that conversation rather than substituting botanicals for it. The drainage phase matters here because gastric die-off generates an endotoxin load that has to clear.

When *Yersinia* serology is elevated, berberine has shown activity against *Yersinia enterocolitica* in vitro. In vitro activity is a laboratory finding rather than a demonstrated clinical outcome, and it should be read that way. Alongside it, a broad-spectrum botanical antimicrobial provides general support for the wider dysbiotic environment that typically accompanies *Yersinia* colonization. Both sit at the level of reasoned clinical practice, not proven treatment, and both get an endpoint.

When Lyme co-infection is confirmed, the antimicrobial protocol is substantially more involved and requires individualized clinical management based on the specific organisms identified, their burden levels, and the patient's drainage and detoxification capacity. The complete Lyme antimicrobial protocol is addressed in the companion ebook *Unlock Your Lyme*. The sequencing principle is the same: drainage first, then barrier support, then targeted antimicrobial intervention with drainage maintained throughout.

Phase	Products	Duration	Goal
1. Drainage	Lymphatic botanical blend, methylated B12, electrolyte and trace mineral base	Weeks 1 to 2 minimum; continue throughout	Open lymphatic, liver Phase 2, cellular hydration
2. Gut Barrier	L-glutamine with glycine and proline, colostrum, bioavailable curcumin	Weeks 3 to 6	Restore barrier, reduce antigen traffic, repair GALT
3. Thyroid/Immune	Thyroid cofactor formula, methylated B12 (continuing), vitamin D3 with K2, selenium (food or supplemental)	Weeks 4 to 8	Support hormone conversion, modulate immune response, optimize D3
4. Targeted support, by finding	EBV: general immune support (zinc, vitamin C, adaptogens); no validated antiviral protocol exists. H. pylori: eradication with your prescriber. Yersinia: berberine plus broad-spectrum botanical support, on laboratory activity rather than trial evidence. Lyme: see Unlock Your Lyme.	Weeks 6 to 12, based on lab findings	Address identified infectious drivers with drainage and barrier in place

Where to get these

Every item in this protocol is a single ingredient or a standardized botanical, available from any reputable retailer. Look for independent third-party testing (NSF, USP Verified, or equivalent) and match the form named here rather than the name on the front of the bottle.

If you would rather not evaluate that yourself, I keep a curated dispensary of professional-grade products at Root Health, screened for those standards and organized by the phases above.

Root Health dispensary: us.fullscript.com/welcome/drseay

Disclosure: Root Health earns a portion of dispensary purchases. Nothing here was included, excluded, ranked, or dosed based on what it earns, and you can buy any of it anywhere.

Your Next Steps with Root Health

The framework in this guide is the clinical picture that most Hashimoto's patients never receive. It identifies the upstream drivers, provides a testing sequence that captures the full autoimmune picture, and lays out a treatment approach organized by mechanistic priority rather than symptom suppression. What it cannot do is replace individualized clinical care. Your specific antibody pattern, your gut findings, your H. pylori and Yersinia status, your nutrient deficiencies, and your history all determine how this framework applies to your case.

Start with the Right Testing

Start cheap and specific, not broad and expensive. For most people that means celiac serology drawn while still eating gluten, a full thyroid panel including free T3, free T4, reverse T3, and both antibodies, and ferritin, B12, folate, zinc, and vitamin D. Low ferritin in particular is extraordinarily common in this population and is the single most frequent reason someone with a good TSH still feels terrible. Add a LabCorp H. pylori stool antigen or urea breath test, which is inexpensive and the one infectious finding with eradication evidence behind it.

The Vibrant Hormone Zoomer captures the thyroid, adrenal, and sex hormone picture in a single draw if you would rather do it that way. Add the Wheat Zoomer when the gluten question is still open after celiac serology. Yersinia serology comes in when the picture has not moved after the gut and dietary work. The Tickborne 2.0 PCR goes last, and only with a real exposure history, because it is the most expensive item on the list. All panels except H. pylori are available at rootcauseunlocked.com/order-labs.

If you want the working half of this program

This book is the reasoning. The parts you actually work from are separate documents, and they exist: **The Hashimoto's Root Cause Kit** collects the lab interpretation guide, the diet guide and the supplement guide in one place. It is the difference between understanding why a panel matters and having the page that tells you how to read yours when it comes back.

If you would rather have everything together, the Ultimate edition bundles this book with all three. Both are at rootcauseunlocked.com/shop.

Neither is required. Everything in this book stands on its own, and the free conversation below is open to you whether you buy anything or not.

Book a Root Discovery Session

The Root Discovery Session is a free clinical conversation with me, and it is a focused clinical intake rather than a sales call. Your symptom history, prior testing, and current picture are reviewed to identify the most targeted path forward. If you are uncertain which workup is appropriate for your case, or if you want clinical interpretation of prior results before spending money on more testing, this is the right starting point.

Book at rootcauseunlocked.com/discovery, or call the Garner, NC office at (919) 662-0520.

Consultations and ongoing care are both available wherever you are located. Being outside North Carolina is not a barrier to working with me. Tell me where you are when you reach out and I will lay out exactly how we would work together from there.

If you would rather start on your own, get the celiac test done while you are still eating gluten, and read the lab guide in this kit before you order anything else.

A Personal Note

If you have been on thyroid medication for years and you still do not feel well, that is information. It means the immune attack is continuing. It means the upstream drivers have not been addressed. It does not mean the medication is wrong or that nothing can be done. It means the right investigation has not happened yet.

That investigation is exactly what Root Health is built for.

Dr. Adam Seay, DC · Root Health, Garner NC · rootcauseunlocked.com

References

1. Ralli et al., "Hashimoto's thyroiditis: An update on pathogenic mechanisms, diagnostic protocols, therapeutic strategies, and potential malignant transformation," *Autoimmunity Reviews*, 2020. <https://pubmed.ncbi.nlm.nih.gov/32805423/>
2. Cayres et al., "Detection of Alterations in the Gut Microbiota and Intestinal Permeability in Patients With Hashimoto Thyroiditis," *Frontiers in Immunology*, 2021. <https://pubmed.ncbi.nlm.nih.gov/33746942/>
3. Araújo et al., "Effects of Gluten-Free Diet in Non-Celiac Hashimoto's Thyroiditis: A Systematic Review and Meta-Analysis," *Nutrients*, 2025. <https://pubmed.ncbi.nlm.nih.gov/41228508/>
4. Wang et al., "Identification of outer membrane porin f protein of *Yersinia enterocolitica* recognized by antithyrotropin receptor antibodies in Graves' disease and determination of its epitope using mass spectrometry and bioinformatics tools," *The Journal of Clinical Endocrinology and Metabolism*, 2010. <https://pubmed.ncbi.nlm.nih.gov/20484489/>
5. Effraïmidis et al., "No causal relationship between *Yersinia enterocolitica* infection and autoimmune thyroid disease: evidence from a prospective study," *Clinical and Experimental Immunology*, 2011. <https://pubmed.ncbi.nlm.nih.gov/21488870/>
6. Shi et al., "Associations of *Helicobacter pylori* infection and cytotoxin-associated gene A status with autoimmune thyroid diseases: a meta-analysis," *Thyroid*, 2013. <https://pubmed.ncbi.nlm.nih.gov/23544831/>
7. Flower et al., "Investigating the Link between the Intestinal Microbiome and Th17/Treg Dysregulation in Hashimoto's Thyroiditis, and the Therapeutic Potential of Vitamin D," *Journal of the American Nutrition Association*, 2026. <https://pubmed.ncbi.nlm.nih.gov/41920977/>
8. Mertowska et al., "Does Epstein-Barr virus and intracellular Toll-like receptors affect the course of Hashimoto disease? Findings from studies on newly-diagnosed patients," *Polish Archives of Internal Medicine*, 2024. <https://pubmed.ncbi.nlm.nih.gov/39508448/>
9. Macedo et al., "Detection of SARS-CoV-2 infection in thyroid follicular cells from a COVID-19 autopsy series," *European Thyroid Journal*, 2022. <https://pubmed.ncbi.nlm.nih.gov/35900859/>
10. Robinson et al., "Effects of Different Cow-Milk Beta-Caseins on the Gut-Brain Axis: A Narrative Review of Preclinical, Animal, and Human Studies," *Nutrition Reviews*, 2025. <https://pubmed.ncbi.nlm.nih.gov/39024213/>
11. Huwiler et al., "Selenium Supplementation in Patients with Hashimoto Thyroiditis: A Systematic Review and Meta-Analysis of Randomized Clinical Trials," *Thyroid*, 2024. <https://pubmed.ncbi.nlm.nih.gov/38243784/>
12. Sun et al., "Vitamin D deficiency in Hashimoto's thyroiditis: mechanisms, immune modulation, and therapeutic implications," *Frontiers in Endocrinology*, 2025. <https://pubmed.ncbi.nlm.nih.gov/40822954/>
13. Serrano-Nascimento and Nunes, "Perchlorate, nitrate, and thiocyanate: Environmental relevant NIS-inhibitors pollutants and their impact on thyroid function and human health," *Frontiers in Endocrinology*, 2022. <https://pubmed.ncbi.nlm.nih.gov/36339434/>
14. Jang et al., "Associations between exposure to sodium/iodide symporter inhibitors and markers of thyroid function: A systematic review and meta-analysis," *Chemosphere*, 2025. <https://pubmed.ncbi.nlm.nih.gov/39798720/>
15. EFSA Scientific Committee, "Risks to human and animal health from the presence of bromide in food and feed," *EFSA Journal*, 2025. <https://pubmed.ncbi.nlm.nih.gov/39877303/>
16. Yeshoua et al., "A Review of the Association between Exposure to Flame Retardants and Thyroid Function," *Biomedicines*, 2024. <https://pubmed.ncbi.nlm.nih.gov/38927574/>

17. Iamandii et al., "Does fluoride exposure affect thyroid function? A systematic review and dose-response meta-analysis," *Environmental Research*, 2024. <https://pubmed.ncbi.nlm.nih.gov/38029816/>
18. Ferreira et al., "Is there any association between fluoride exposure and thyroid function modulation? A systematic review," *PLOS ONE*, 2024. <https://pubmed.ncbi.nlm.nih.gov/38593166/>
19. Gallagher and Meliker, "Mercury and thyroid autoantibodies in U.S. women, NHANES 2007-2008," *Environment International*, 2012. <https://pubmed.ncbi.nlm.nih.gov/22280926/>
20. Sultan et al., "Association of Bisphenol Exposure and Serum Hypothalamic-Pituitary-Thyroid Axis Hormone Levels in Adults and Pregnant Women: A Systematic Review and Meta-Analysis," *Toxics*, 2025. <https://pubmed.ncbi.nlm.nih.gov/41150534/>
21. Atapattu et al., "Thyroid Autoimmunity During Universal Salt Iodisation: Possible Short-Term Modulation with Longer-Term Stability," *Nutrients*, 2024. <https://pubmed.ncbi.nlm.nih.gov/39770919/>
22. Negro, "Selenium and thyroid autoimmunity," *Biologics*, 2008. <https://pubmed.ncbi.nlm.nih.gov/19707359/>
23. Centanni et al., "Thyroxine in goiter, *Helicobacter pylori* infection, and chronic gastritis," *The New England Journal of Medicine*, 2006. <https://pubmed.ncbi.nlm.nih.gov/16641395/>

For educational purposes only. This ebook does not constitute medical advice, diagnosis, or treatment, and reading it does not establish a patient-provider relationship.

Root Health | Dr. Adam Seay, DC | Garner, NC | rootcauseunlocked.com