



ROOT CAUSE UNLOCKED · CONDITION GUIDES

# The Neck, Shoulder and Elbow Guide *Set*

*An Integrative Approach to Understanding, Treating, and Recovering From Shoulder Pain*

Dr. Adam Seay, DC

Root Cause Unlocked, Garner, NC · [rootcauseunlocked.com](http://rootcauseunlocked.com) · (919) 662-0520

# The Neck, Shoulder and Elbow Guide Set

*An Integrative Approach to Understanding, Treating, and Recovering From Shoulder Pain*

## BEFORE YOU START

**Disclaimer:** This ebook is for educational purposes only and does not constitute medical advice. It does not diagnose any condition. Always consult with a qualified healthcare provider before starting any new treatment, supplement, exercise program, or dietary change. Do not start, stop, or change any prescribed medication without speaking to the prescriber. If you experience chest pain, pressure, or shoulder pain that could be related to your heart, sudden inability to move your arm after significant trauma, fever with joint swelling, or an unexplained mass, seek emergency medical attention immediately.

**A note on the evidence in this guide.** Every numbered citation in this edition points to a specific published study that was checked against the source paper before it was used. Where the research is weak, mixed, or still developing, this guide says so rather than overstating it. Where a claim reflects clinical experience rather than a settled body of research, it is labeled as such.

# Contents

1. Understanding Rotator Cuff Related Pain

---

2. Anatomy of the Shoulder and the Rotator Cuff

---

3. The Bigger Picture: Why the Painful Spot Isn't Always the Problem

---

4. The Thoracic Spine Connection

---

5. Common Causes and Risk Factors

---

6. How Rotator Cuff Pain Is Diagnosed, and What Imaging Can and Cannot Tell You

---

7. The Healing Timeline: What to Expect

---

8. Movement and Exercise: The Foundation of Recovery

---

9. Integrative and Manual Therapies

---

10. Supplements for Shoulder and Tendon Health: What the Evidence Actually Says

---

11. The Anti-Inflammatory Diet for Tendon Recovery

---

12. Lifestyle Modifications for Sleep, Stress, Smoking, and Ergonomics

---

13. Mind-Body Approaches to Chronic Shoulder Pain

---

14. The In-Clinic Pre-Treatment Protocol

---

15. Red Flags: When to Seek Immediate Care

---

16. Building Your Personal Recovery Plan

---

Final Thoughts

---

17. References

---

What this guide is for

---

Chapter 1. What is actually happening

---

Chapter 2. The three phases, and why they change everything

---

Chapter 3. Who gets it and why that matters

---

Chapter 4. Making sure it is actually this

---

Chapter 5. What the largest trial found

---

Chapter 6. What to actually do, by phase

---

Chapter 7. Measuring yourself so you can tell if this is working

---

Chapter 8. The In-Clinic Pre-Treatment Protocol

---

Chapter 9. Nutrition, briefly and honestly

---

Chapter 10. Red flags

---

Chapter 11. Building your plan

---

Final thoughts

---

References

---

Working with Root Health

---

1. What Is Tennis Elbow?

---

2. Understanding the Anatomy

---

3. Common Causes and Risk Factors

---

4. How Tennis Elbow Is Diagnosed

---

5. The Healing Timeline: What to Expect

---

6. Supplements for Tendon Healing and Inflammation

---

7. Targeted Exercises for Tennis Elbow Relief

---

8. Physical Therapies and What the Evidence Says

---

9. The Anti-Inflammatory Diet for Tendon Recovery

---

10. Lifestyle Modifications and Pain Management

---

11. The In-Clinic Pre-Treatment Protocol

---

12. Red Flags: When to Seek Immediate Care

---

13. Building Your Personal Recovery Plan

---

Final Thoughts

---

References

---

Working with Root Health

---

A clearer way forward

---

How to use this guide

---

Chapter 1. Understanding cervicogenic headache

---

Chapter 2. How the neck can refer pain to the head

---

Chapter 2b. The mid-back connection, and why treatment often starts below the pain

---

Chapter 3. Cervicogenic, migraine, or tension type?

---

Chapter 4. Warning signs and when to seek help

---

Chapter 5. How your clinician builds confidence

---

Chapter 6. The five-part recovery plan

---

Chapter 7. Chiropractic and hands-on care

---

Chapter 8. Your progressive exercise plan

---

Chapter 9. Food, hydration, and anti-inflammatory eating

---

Chapter 10. Supplements, including anti-inflammatory options

---

Chapter 11. Sleep, stress, work, and lifestyle

---

Chapter 12. Flare planning and progress tracking

---

The In-Clinic Pre-Treatment Protocol

---

Your next-visit worksheet

---

Sources and further reading

---

Working with Root Health

---

Remember

---

# 1. Understanding Rotator Cuff Related Pain

---

If your shoulder has been hurting for weeks or months, you have probably already heard the word “impingement.” Maybe someone told you a bone spur is pinching a tendon, or that something in there is rubbing where it shouldn’t. That explanation is common, and it is also outdated. Shoulder specialists have moved away from thinking of this kind of pain as simple mechanical pinching, because the research behind that model has not held up well under scrutiny.<sup>1</sup>

The more accurate and more useful umbrella term is Rotator Cuff Related Shoulder Pain, often shortened to RCRSP. It covers a spectrum of related presentations, including subacromial pain, rotator cuff tendinopathy, and partial or full thickness rotator cuff tears that are actually causing symptoms. A detailed 2016 masterclass on the condition, published in the journal *Manual Therapy*, describes RCRSP as an over-arching term for this whole family of presentations, and is direct about how much uncertainty still exists around the exact cause and source of the pain, how to reach a definitive diagnosis, and which tissues or systems should be the target of treatment.<sup>1</sup>

That uncertainty is not bad news. It means your care should not hinge on finding one perfect structural explanation. The same research points to something genuinely encouraging: a graduated, well-constructed exercise program produces outcomes that are at least as good as surgery for subacromial pain, rotator cuff tendinopathy, partial thickness tears, and even many full thickness tears that were not caused by a specific traumatic injury.<sup>1</sup> You are going to see that theme repeated throughout this guide, because it is the single most useful thing the evidence has to offer you.

There is one more piece of context worth understanding before you go any further, because it will change how you interpret anything an X-ray or MRI eventually tells you. Rotator cuff tears become more common simply as a normal part of aging, and this happens in people who have never had a day of shoulder pain in their life. A pooled analysis of studies covering more than six thousand shoulders found that the prevalence of rotator cuff abnormalities rose from under ten percent in people twenty and younger to over sixty percent in people eighty and older, and this rising pattern held true whether the person had symptoms or not.<sup>2</sup> Your shoulder anatomy telling a story about your age is not the same thing as your shoulder anatomy explaining your pain. Chapter 6 returns to this in more detail, because it matters for how you should think about any imaging you have already had or might have in the future.

This guide is built around the clinical approach I use at Root Health: taking rotator cuff dysfunction seriously as a whole-body problem, not just a local one. That means giving real attention to a commonly overlooked player, the subscapularis muscle, and to the idea that your shoulder does not work in isolation from the rest of your spine and posture. Both of those threads run through the chapters ahead.

## 2. Anatomy of the Shoulder and the Rotator Cuff

---

Your shoulder is the most mobile joint in your body, and that mobility comes at a cost. Unlike your hip, which sits in a deep, stable socket, your shoulder joint is a shallow ball resting against a shallow socket, held together far more by muscles, tendons, and ligaments than by bone. That design gives you enormous range of motion. It also means the muscles around the joint are doing constant, active work just to keep the ball centered where it belongs.

### The Rotator Cuff

The rotator cuff is a group of four muscles and their tendons that wrap around the head of the humerus, the ball of your upper arm bone, and blend into a single continuous cuff of tissue over the joint. Their main job is to hold the humeral head centered and stable in the socket while your larger, more powerful muscles, like the deltoid, do the heavy lifting, rather than to create big, powerful movements themselves.

- **Supraspinatus.** Runs across the top of your shoulder blade and helps initiate raising your arm out to the side.
- **Infraspinatus.** Covers much of the back of your shoulder blade and is a primary external rotator, turning your arm outward.
- **Teres minor.** A smaller muscle beneath the infraspinatus, also assisting external rotation.
- **Subscapularis.** The largest of the four rotator cuff muscles, sitting on the front surface of the shoulder blade against your ribcage, largely hidden from view and from easy hands-on examination. It is the primary internal rotator of the shoulder, and it plays a major stabilizing role at the front of the joint.

### Why Subscapularis Deserves More Attention Than It Usually Gets

Most general information about rotator cuff pain focuses heavily on the supraspinatus, partly because it is the most commonly imaged and most commonly torn tendon. That focus can leave subscapularis dysfunction underappreciated, and there is a real, practical reason it gets missed: it is difficult to test for on physical exam. A 2021 systematic review and meta-analysis pooled the diagnostic accuracy of the four clinical tests most commonly used to detect subscapularis tears, including the bear-hug test, the belly-press test, the internal rotation lag sign, and the lift-off test. Every one of these tests was reasonably good at ruling a tear in when positive, with pooled specificity above ninety percent, but every one of them had pooled sensitivity below sixty percent, meaning each test, used alone, misses a substantial share of real subscapularis involvement. The review's authors concluded plainly that no single clinical test is sufficiently reliable on its own to diagnose subscapularis tears.<sup>3</sup> In my clinical experience, this translates into subscapularis dysfunction being a frequent, quietly present contributor to shoulder pain that a quick screening exam can easily pass over, which is why I give it deliberate, specific attention as part of a thorough shoulder evaluation at Root Health.

## Supporting Structures

- **Scapula (shoulder blade).** Provides the socket and serves as the stable base the rotator cuff needs to work from. Its position and movement against the ribcage, called scapular kinematics, directly affects how much room the structures underneath your shoulder have to move freely.
- **Acromion.** The bony roof of the shoulder, under which the supraspinatus tendon and a fluid-filled cushion called the subacromial bursa pass.
- **Long head of the biceps tendon.** Runs through the front of the shoulder joint and is frequently involved alongside rotator cuff pathology.
- **Glenohumeral ligaments and joint capsule.** Passive stabilizers that work alongside the rotator cuff to keep the joint contained.
- **Deltoid.** The large muscle that gives your shoulder its rounded shape and produces most of the raw power for lifting the arm, relying on a stable platform from the rotator cuff underneath it.

The point of understanding this anatomy is less memorizing muscle names than recognizing that shoulder function depends on a coordinated relationship between several structures working together, not on any single tendon behaving perfectly. That coordinated view is exactly where the next two chapters go next.

### 3. The Bigger Picture: Why the Painful Spot Isn't Always the Problem

---

Here is a question worth asking. If the spot that hurts is not always the spot that is actually broken, what is a clinician supposed to examine? A concept called regional interdependence is built on that question, and it is the broader model that shapes how I approach shoulder care.

Regional interdependence, as a formal idea in musculoskeletal care, was articulated clearly in a widely cited 2007 editorial in the *Journal of Orthopaedic and Sports Physical Therapy*, which argued that seemingly unrelated impairments in a remote anatomical region can contribute to, or be associated with, the patient's primary complaint, and that a comprehensive examination model needs to consider the entire kinetic chain rather than the isolated painful area alone.<sup>4</sup> Applied to your shoulder, that means a thorough evaluation does not stop at your shoulder joint. It looks at your neck, your thoracic spine, your scapular control, even your hip and core stability, because dysfunction in any of those areas can change the demands placed on your shoulder and how it is able to respond to those demands.

This adds to looking carefully at the shoulder itself rather than replacing it. Your rotator cuff and the tissues around it still matter, and the exercises and therapies in the chapters ahead are built directly around them. But regional interdependence explains something that a shoulder-only view cannot: why two people with what looks like an identical supraspinatus tear on imaging can have completely different symptoms, and why treating the shoulder in isolation sometimes produces a plateau that treating the wider system can break through.

In practice, this looks like asking questions your shoulder pain alone would not seem to justify. How is your posture at your desk for eight hours a day, how well does your upper back rotate and extend, and how is your neck moving? Another is whether your shoulder blade sits and glides normally against your ribcage during arm movement. None of these are exotic ideas. They are simply an acknowledgment that the human body is one connected system, and that the place where you feel pain is often the last link in a chain of compensations rather than the first cause of them. The next chapter takes the most consistently useful piece of this bigger picture, your thoracic spine, and looks closely at what the evidence actually shows.

## 4. The Thoracic Spine Connection

---

Of everything regional interdependence points toward, the relationship between your thoracic spine, the middle and upper portion of your back, and your shoulder function has the clearest biomechanical rationale and the most direct research behind it. This is a deliberate part of the care I provide at Root Health, not an afterthought tacked onto shoulder treatment.

### The Mechanical Reason

Your shoulder blade sits on your ribcage, and your ribcage is shaped by the position of your thoracic spine. When you round forward into a slouched posture, a nearly universal position for anyone who spends hours at a desk or on a phone, your shoulder blade's position and movement pattern change with it. A controlled laboratory study measured this directly: in a slouched thoracic posture, healthy volunteers' shoulder blades sat more elevated during the first part of arm-raising, tilted less favorably during the later part of the movement, and their active shoulder abduction range of motion was reduced by an average of nearly twenty-four degrees compared with an upright posture. Isometric shoulder muscle force with the arm raised to horizontal also dropped by over sixteen percent in the slouched position.<sup>5</sup> In plain terms, a stiff, rounded thoracic spine measurably changes how much room your shoulder has to move and how much force your shoulder muscles, including your rotator cuff, can produce. Ask your shoulder to do a full day of overhead or reaching work from that starting position, repeated thousands of times, and it is not hard to see how the system becomes overloaded somewhere.

### What the Treatment Evidence Shows, Honestly

This is where it is important to separate a strong mechanical rationale from strong treatment-outcome evidence, and to say plainly where the two currently line up and where they do not.

A 2026 systematic review and meta-analysis pooling twelve randomized trials and over five hundred participants found that thoracic manual therapy, as a category, appeared effective for reducing pain and disability in people with shoulder dysfunction, but the benefit was not uniform across techniques. Gentle sustained mobilization, specifically the Maitland technique, showed a meaningfully greater improvement in both pain and disability compared with control interventions in people with subacromial impingement syndrome. High-velocity, low-amplitude thrust manipulation, the quicker, more forceful technique often associated with an audible pop, did not show a significant effect in the same population or in adhesive capsulitis, and evidence for another mobilization approach, the Mulligan technique, remained too limited to judge.<sup>6</sup>

A separate 2025 systematic review focused specifically on thrust manipulation of the thoracic spine reached a similarly measured conclusion: there is low quality evidence that thoracic thrust manipulation is more effective than general conservative therapy for shoulder pain, but it was not shown to outperform a sham, placebo version of the same manipulation.<sup>7</sup> That comparison against sham manipulation matters. It is one of

the better tools researchers have for asking whether a hands-on technique is doing something specific to the tissue, or whether at least part of its effect comes from the broader experience of hands-on care, attention, and expectation, all of which are real and can still be worth pursuing, but are a different claim than “this manipulation mechanically fixes the shoulder.”

One frequently cited early study deserves a direct, honest look because of how often its result gets repeated. A 2009 study applied thoracic spine thrust manipulation to fifty-six patients with shoulder impingement symptoms and found statistically significant improvement in pain and disability scores at a 48-hour follow-up.<sup>8</sup> This finding is real, but its design is limited in an important way: it was a one-group study with no control or comparison group at all, meaning there was no way to separate the effect of the manipulation from natural fluctuation in symptoms, the placebo effect of receiving hands-on care, or simple regression toward the mean. It should be read as a preliminary signal that helped motivate later, better-controlled research, not as proof that thrust manipulation reliably fixes shoulder pain on its own.

Put together honestly: the mechanical case for thoracic mobility mattering to shoulder function is solid. The treatment evidence currently favors gentler mobilization techniques over forceful thrust manipulation specifically, though both approaches have at least some supportive research behind them, and neither is a stand-alone cure. At Root Health, I treat restoring thoracic mobility as one meaningful piece of a larger plan, most often pairing it directly with the loading exercises described in Chapter 8, rather than expecting it to resolve shoulder pain by itself.

# 5. Common Causes and Risk Factors

---

RCRSP rarely has one single cause. It is usually the end result of several overlapping factors, and identifying which ones apply to you is more useful than searching for a single culprit.

## Common Contributors

- **Tendon overload.** Repetitive overhead activity, whether from a job, a sport, or a home project, can outpace the rotator cuff tendons' ability to adapt and recover.
- **Age-related tendon change.** As discussed in Chapter 1, rotator cuff tendons change structurally with age in almost everyone, symptomatic or not.<sup>2</sup> This is closer to a normal, expected process than a disease.
- **Scapular and postural dysfunction.** Poor control of the shoulder blade changes the mechanics of everything attached to it, including the rotator cuff.
- **Thoracic and cervical stiffness.** Covered in depth in Chapter 4.
- **Subscapularis involvement.** Frequently under-recognized on exam, as covered in Chapter 2.<sup>3</sup>
- **Prior shoulder injury.** A history of shoulder problems raises the likelihood of future ones.

## Risk Factors Supported by the Evidence

A 2021 systematic review and meta-analysis pooling data on more than four thousand shoulders looked specifically at risk factors for full-thickness rotator cuff tears. It found that increasing age, hypertension, and a larger critical shoulder angle, a specific anatomical measurement of the socket's shape, were each statistically associated with greater risk. Notably, that same analysis found that body mass index, sex, hand dominance, diabetes, thyroid disease, and smoking were not statistically significant risk factors for developing a full-thickness tear in their pooled data.<sup>9</sup>

That last finding about smoking deserves a careful, honest distinction, because it answers a different question than the one most people assume. Smoking's relationship to whether a tear develops in the first place is genuinely mixed in the research. Its relationship to healing after rotator cuff surgery is much more consistent. A 2022 systematic review and meta-analysis of over seventy thousand surgical patients found that smokers had significantly worse Constant shoulder scores after repair, and a significantly higher risk of both retear and reoperation compared with nonsmokers.<sup>10</sup> So while the current evidence does not clearly show smoking causes a rotator cuff tear to happen, it does show smoking measurably impairs how well tendon tissue heals and holds once it is repaired or, by extension, recovering under a conservative loading program. Quitting remains one of the most useful things you can do for tissue healing generally, even if the reason is more about healing capacity than initial injury risk.

# 6. How Rotator Cuff Pain Is Diagnosed, and What Imaging Can and Cannot Tell You

---

## Clinical Examination

A thorough shoulder evaluation typically includes a detailed history of when and how your pain started, what makes it better or worse, and what your daily and occupational demands look like. It should also include a look beyond the shoulder itself, in keeping with the regional interdependence model from Chapter 3, meaning an assessment of your neck, thoracic spine mobility, scapular movement, and posture, along with specific rotator cuff strength testing that deliberately checks internal rotation and subscapularis function rather than relying only on the more commonly tested external rotators.

## The Honest Truth About Imaging

Imaging can be a useful tool, but it needs to be interpreted carefully, and this is one of the most important things this guide can tell you. As Chapter 1 introduced, rotator cuff abnormalities on imaging become dramatically more common simply with age, in people who have never had shoulder pain. The pooled analysis of over six thousand shoulders found abnormality prevalence rising from under ten percent in people twenty and younger to over sixty percent by age eighty, and this pattern held true in the general population regardless of symptoms.<sup>2</sup> The authors of that analysis concluded that rotator cuff degeneration is common enough with normal aging that it is genuinely difficult to know, just from a scan, whether a finding is new, old, or actually the source of someone's current pain.<sup>2</sup>

Here is what this means for you directly. If you have already had an MRI, or one is recommended, a report showing a partial tear, tendinopathy, or even a small full-thickness tear does not automatically mean that finding explains your pain, and it does not by itself mean surgery is your only path forward. Chapter 8 covers what the treatment evidence actually shows for tears found on imaging, and it is more encouraging than most people expect.

Imaging is genuinely useful when there has been significant trauma, when weakness is severe or progressive, when conservative care over a reasonable period has not produced any improvement, or when your clinician suspects something other than typical RCRSP. It is far less useful as a first step for typical, gradual-onset shoulder pain, where it often produces findings that are more confusing than clarifying.

## 7. The Healing Timeline: What to Expect

---

Every shoulder and every case is different, but a general pattern is worth knowing so you can set realistic expectations.

### **Early Phase (0 to 6 Weeks)**

This is typically when pain is most reactive and irritable. The priority is calming the tissue down, protecting against further overload, and beginning gentle, pain-guided movement rather than complete rest. Complete immobilization tends to create stiffness that becomes its own separate problem.

### **Building Phase (6 to 12 Weeks)**

As irritability settles, the focus shifts toward progressive strengthening of the rotator cuff and scapular stabilizers, along with continued work on thoracic and postural factors. This is typically where people begin to notice meaningful, sustained improvement rather than day-to-day fluctuation.

### **Strengthening and Return to Function (3 to 6 Months)**

Loading capacity continues to build toward the demands of your specific work, sport, or daily activities. Full recovery of strength and confidence in overhead or loaded positions often takes several months, even when pain has already improved substantially.

### **Longer-Term Cases**

Some cases, particularly those present for many months before treatment begins, take longer and benefit from a more layered, patient approach. This does not mean something is going wrong. Tendon tissue adapts more slowly than muscle, and consistency over months, not days, is what drives lasting change.

# 8. Movement and Exercise: The Foundation of Recovery

---

If this guide could hand you only one chapter, it would be this one. Loading exercise has the strongest, most consistent evidence base of anything covered here, and it should be the backbone of your recovery rather than an afterthought.

The 2016 masterclass on RCRSP introduced in Chapter 1 states that a graduated, well-constructed exercise program produces outcomes at least equivalent to surgery for subacromial pain, rotator cuff tendinopathy, partial thickness tears, and atraumatic full thickness tears.<sup>1</sup> A 2017 systematic review and meta-analysis of randomized trials in adults with shoulder impingement found exercise significantly superior to non-exercise control interventions, and specific, targeted exercise significantly better than generic, unspecific exercise.<sup>11</sup> A randomized, controlled trial that put this to a direct test is worth knowing about on its own: 102 patients with long-standing subacromial impingement syndrome who had already failed earlier conservative treatment were assigned to either a specific eccentric strengthening program for the rotator cuff and scapular stabilizers or a generic, unspecific movement program. The specific exercise group improved significantly more on a validated shoulder function score, and significantly fewer of them went on to choose surgery afterward, 20 percent compared with 63 percent in the generic exercise group.<sup>15</sup> The takeaway goes beyond “move your shoulder.” A deliberate, progressive, specific program outperforms vague advice to stay active, and can meaningfully change whether surgery ever becomes necessary.

## 8.1 Early, Gentle Motion

**Pendulum swings.** Lean forward, let your arm hang freely, and gently swing it in small circles and side to side, letting gravity and momentum do the work rather than your muscles. Useful in the early, irritable phase to maintain motion without provoking pain.

## 8.2 Rotator Cuff Loading, Including Subscapularis

Given Chapter 2’s discussion of how often subscapularis involvement is overlooked, deliberately including internal rotation strengthening alongside the more commonly prescribed external rotation work is a meaningful part of a well-rounded program.

### External rotation with resistance band

1. Hold a light resistance band with your elbow bent to 90 degrees and tucked against your side
2. Rotate your forearm outward, away from your body, keeping your elbow pinned to your side
3. Return slowly and with control
4. Perform 2 to 3 sets of 10 to 15 repetitions

### Internal rotation with resistance band (subscapularis focus)

1. Anchor the band on the side opposite the arm you are training, elbow bent to 90 degrees and tucked at your side
2. Rotate your forearm inward, across your body
3. Return slowly and with control
4. Perform 2 to 3 sets of 10 to 15 repetitions

**Isometric holds.** For an irritable shoulder that does not tolerate the movements above yet, gently pressing your hand into a wall or doorframe in the external and internal rotation directions and holding, without moving the joint, is a lower-irritation way to begin loading the tendon.

### 8.3 Scapular Stabilization

**Wall slides.** Standing with your back against a wall, arms bent and pressed against the wall, slide your arms upward while keeping contact with the wall, focusing on smooth shoulder blade movement rather than shrugging.

**Scapular rows.** Using a band or cable, pull your elbows back and squeeze your shoulder blades together, focusing on control rather than heavy load.

### 8.4 Thoracic Mobility Work

Given Chapter 4, thoracic mobility exercises are typically paired directly with rotator cuff loading rather than done separately.

#### Open book stretch

1. Lie on your side with knees bent and arms stacked in front of you
2. Keeping your bottom hand in place, rotate your top arm and chest open toward the ceiling, following it with your eyes
3. Hold briefly, then return
4. Perform 8 to 10 repetitions per side

#### Thoracic extension over a foam roller

1. Lie with a foam roller placed horizontally under your mid-back, knees bent, feet flat, hands supporting your head
2. Gently extend backward over the roller within a comfortable range
3. Move slowly and avoid forcing the stretch
4. Perform for 1 to 2 minutes

## 8.5 Progressive Loading

As symptoms allow, exercises progress from isometric holds to controlled full-range movement to added resistance, gradually building toward the demands of your specific work, sport, or daily activities. This progression, not any single exercise, is what drives lasting tendon adaptation.

### What About Surgery?

If a tear has been found on your imaging, it is natural to wonder whether surgery is the more definitive answer. The honest, evidence-based answer for most people is no, at least not as a first step. A well-known randomized surgical trial, published in the *Lancet* and known as the CSAW trial, compared genuine arthroscopic subacromial decompression surgery against a placebo surgery, in which patients underwent anesthesia and an arthroscopic procedure without the actual bone and soft tissue removal, and against no treatment at all. At six months, both surgical groups showed a small improvement over no treatment, but the difference was not considered clinically important, and decompression surgery showed no additional benefit over the placebo surgery. The trial's own authors concluded that the findings question the value of this operation and that this should be communicated to patients as part of shared decision-making.<sup>12</sup>

For actual rotator cuff tears, the picture is similarly reassuring for a conservative-first approach in appropriate cases. A randomized trial following patients over 55 with small, nontraumatic supraspinatus tears for more than five years found no significant difference in outcomes between physiotherapy alone, physiotherapy plus a bone-shaving procedure, and physiotherapy plus full surgical repair, and concluded that operative treatment is no better than conservative treatment for this population and that conservative care is a reasonable initial approach.<sup>13</sup> A separate meta-analysis pooling three randomized trials directly comparing tendon repair surgery against conservative treatment found only a small, statistically insignificant difference favoring surgery that fell below the threshold generally considered clinically meaningful, and concluded that starting with conservative treatment is reasonable.<sup>14</sup>

None of this means surgery is never appropriate. Larger traumatic tears, tears with significant ongoing weakness, and cases that genuinely fail an adequate trial of conservative care are different conversations, and this guide is not a substitute for that individualized discussion with your provider. But for the large majority of people with typical RCRSP, including many with a tear visible on a scan, a well-built exercise program deserves to be the first, not the last, thing you try.

# 9. Integrative and Manual Therapies

---

## 9.1 Physical Therapy and Chiropractic Care

A skilled provider assesses movement patterns, muscle imbalances, joint mobility throughout the shoulder, neck, and thoracic spine, and builds an individualized, progressive loading plan, using manual therapy as a complement to exercise rather than a replacement for it. The 2017 meta-analysis of conservative interventions for shoulder impingement found manual therapy statistically superior to placebo, and found that when manual therapy was combined with exercise, it showed an additional benefit over exercise alone, though that added benefit was only detectable at the shortest follow-up period measured.<sup>11</sup> Read honestly, that means hands-on care can help, particularly early on, but its main long-term value appears to be in supporting and accelerating an exercise-based plan rather than functioning as a stand-alone treatment.

## 9.2 Thoracic Mobilization Technique

As detailed in Chapter 4, current evidence favors gentler, sustained mobilization techniques over forceful thrust manipulation for shoulder-related outcomes specifically, though both have at least some supportive research.<sup>6,7</sup> This is typically delivered as part of the same visit as rotator cuff and scapular loading work, not as an isolated therapy.

## 9.3 Heat and Cold Therapy

**Cold.** Can help numb acute, irritable pain in the early phase. Apply ice wrapped in a cloth for 15 to 20 minutes.

**Heat.** Can help relax surrounding muscle tension and improve comfort before exercise sessions. Use a heating pad or warm shower for 15 to 20 minutes.

These are comfort measures rather than corrective treatments, and are reasonable to use alongside, never instead of, an active exercise program.

## 9.4 Dry Needling and Trigger Point Work

Can be useful for addressing muscle tension and trigger points in the surrounding musculature, particularly the upper trapezius and pectoral muscles, which frequently tighten in response to altered shoulder mechanics. Best understood as a tool to improve comfort and readiness for exercise rather than a treatment that resolves rotator cuff pathology on its own.

# 10. Supplements for Shoulder and Tendon Health: What the Evidence Actually Says

---

This chapter calls for particular honesty, because the supplement research most often cited for joint and tendon health was not done in people with rotator cuff or shoulder pain. Most of it comes from knee osteoarthritis populations, which is a different tissue and a different condition. Where that is the case, this guide says so plainly rather than implying a connection the research does not support. Always discuss supplements with your provider, especially if you take blood thinners or have kidney disease.

## 10.1 Omega-3 Fatty Acids (EPA and DHA)

**What it does.** Omega-3 fats are incorporated into cell membranes and influence the production of inflammatory signaling molecules throughout the body.

**Evidence, with an important scope note.** A 2006 report in *Surgical Neurology* followed 250 patients with nonsurgical neck or back pain, not shoulder pain, who were asked to take 1,200 to 2,400 milligrams of omega-3 fatty acids daily. Of the 125 who returned a follow-up questionnaire after an average of 75 days, 59 percent had stopped their prescription NSAIDs and 60 percent reported improved overall pain.<sup>16</sup> This is an encouraging signal about omega-3's general anti-inflammatory potential, but it has two real limits worth knowing: it was an uncontrolled survey with a 50 percent response rate rather than a controlled trial, and it studied neck and back pain, not shoulder or tendon pain specifically. Treat it as reasonable, low-risk background support rather than a targeted shoulder treatment.

**Dosage commonly used.** 2,000 to 3,000 mg combined EPA and DHA daily **Caution.** May increase bleeding risk at high doses. Discuss with your surgeon before any procedure.

## 10.2 Curcumin (Turmeric Extract)

**What it does.** Curcumin acts on several inflammatory signaling pathways in the body.

**Evidence, scoped honestly to arthritis.** A systematic review and meta-analysis of randomized trials found that turmeric extract and curcumin reduced pain scores compared with placebo in joint arthritis, mostly knee osteoarthritis, with the authors noting the number of trials, sample size, and study quality were not sufficient for definitive conclusions.<sup>17</sup> A separate meta-analysis of eleven trials found curcumin and boswellia formulations statistically superior to placebo for knee osteoarthritis pain and function, again describing the overall body of evidence as not adequate in size or quality for firm clinical practice recommendations.<sup>18</sup> Curcumin has not been studied specifically in rotator cuff tendinopathy or RCRSP in the research reviewed for this guide.

**Dosage commonly used.** 500 to 1,000 mg curcuminoids daily, with piperine or in a liposomal or phytosome form for better absorption **Caution.** May interact with blood thinners. Avoid in gallbladder disease.

## 10.3 Collagen and Vitamin C

**What it does.** Supplies the amino acids used in collagen synthesis, the building process for tendon and connective tissue. Vitamin C is required for collagen cross-linking.

**Evidence.** A randomized, double-blind, crossover trial found that participants who consumed 15 grams of vitamin C-enriched gelatin about an hour before a short bout of intermittent exercise showed roughly double the blood marker of collagen synthesis compared with placebo. Serum from supplemented participants also increased collagen content and improved mechanical properties in engineered ligament tissue in the same study.<sup>19</sup>

**An honest scope note.** This is a mechanism study. It measured collagen synthesis markers and laboratory tissue response, not pain or function outcomes in people with rotator cuff pain, and tendon is not identical to the ligament tissue tested. What it offers is a genuinely interesting rationale for supporting tendon repair nutritionally around the time of loading exercise, not a demonstrated pain treatment.

**Dosage used in the study.** 15 g gelatin with 50 mg vitamin C, about 60 minutes before exercise **Caution.** Generally very well tolerated.

### What This Chapter Is Not Saying

None of these supplements are a substitute for the loading exercise described in Chapter 8, which has by far the strongest evidence for RCRSP specifically. Consider them reasonable, low-risk additions to discuss with your provider, not a shortcut around the harder, more effective work of a structured rehabilitation program.

# 11. The Anti-Inflammatory Diet for Tendon Recovery

---

Tendon tissue, like every tissue in your body, is influenced by your overall metabolic and inflammatory environment. The eating pattern below is well-established general healthy nutrition. It is presented here as sensible support for recovery, not as a proven, targeted treatment for rotator cuff pain specifically, since dedicated shoulder-pain dietary trials are not part of the evidence base reviewed for this guide.

## Foods to Emphasize

**Omega-3 rich foods.** Wild-caught salmon, sardines, mackerel, walnuts, flaxseeds, chia seeds

**Colorful vegetables and fruits.** Leafy greens, berries, cruciferous vegetables like broccoli and Brussels sprouts, orange and yellow vegetables like carrots and sweet potatoes

**Anti-inflammatory spices.** Turmeric with black pepper, ginger, garlic

**Quality protein.** Lean meats, poultry, fish, eggs, legumes, providing the amino acid building blocks tendon repair depends on

**Collagen-supporting foods.** Bone broth, and foods rich in vitamin C such as citrus, bell peppers, and berries, which support the body's own collagen synthesis discussed in Chapter 10

**Healthy fats.** Extra virgin olive oil, avocados, nuts and seeds

## Foods to Minimize

Trans fats and deep-fried foods, refined carbohydrates and added sugars, heavily processed and packaged foods, and excessive alcohol, which can interfere with sleep quality, a factor covered in the next chapter.

## Hydration

Adequate hydration supports tissue health broadly. Drink regularly through the day and use thirst and pale urine color as a general guide, and follow your provider's specific fluid guidance if you have a heart or kidney condition.

# 12. Lifestyle Modifications for Sleep, Stress, Smoking, and Ergonomics

---

## 12.1 Sleep

Sleep quality genuinely affects how you experience pain. A brain imaging study found that sleep deprivation amplified pain-related activity in the brain's sensory processing regions while blunting activity in regions that normally help regulate pain, and in a separate general population sample, even small night-to-night changes in sleep predicted corresponding day-to-day changes in pain.<sup>20</sup> Shoulder pain has a well-known reputation for disrupting sleep, particularly when lying on the affected side, which can create a frustrating cycle where poor sleep worsens pain sensitivity and pain worsens sleep in turn.

**Strategies.** Try sleeping on your back with a pillow supporting the affected arm, or on your unaffected side with a pillow hugged in front of you for support. Keep a consistent sleep schedule, limit screens before bed, and avoid caffeine in the afternoon.

## 12.2 Stress Management

Chronic stress measurably amplifies pain perception. A systematic review and meta-analysis of 47 trials involving over 3,500 participants found that structured mindfulness meditation programs produced moderate improvement in measures of pain, alongside improvements in anxiety and depression.<sup>21</sup> This evidence covers chronic pain broadly rather than shoulder pain specifically, but the underlying mechanism, a nervous system that is calmer and less reactive processing pain signals differently, applies regardless of which joint is involved.

**Effective techniques.** Diaphragmatic breathing for five minutes twice daily, mindfulness meditation, and regular time outdoors.

## 12.3 Smoking Cessation

As discussed in Chapter 5, smoking's link to whether a rotator cuff tear develops in the first place is genuinely uncertain in the current pooled research, but its link to poorer healing after repair, including higher rates of retear and reoperation, is well established.<sup>10</sup> Whether or not you are considering surgery, quitting supports the tissue-healing capacity your recovery program depends on. Your provider can help you build a plan.

## 12.4 Ergonomics

**Desk setup.** Position your monitor at eye level and keep your keyboard and mouse close enough that your elbows stay near your sides, reducing the reaching and shrugging that load an already irritable shoulder.

**Overhead activity.** Break up prolonged overhead tasks with rest periods rather than pushing through in one long stretch, and when possible, use a step stool or ladder rather than reaching to full extension repeatedly.

**Carrying and lifting.** Keep loads close to your body, and distribute weight evenly between both sides when possible rather than consistently favoring one shoulder.

**Sleep positioning.** Covered above in the sleep section, given how directly it affects shoulder pain specifically.

# 13. Mind-Body Approaches to Chronic Shoulder Pain

---

For pain that has persisted for months, addressing the mind-body connection is a genuine, evidence-supported part of care, not a suggestion that the pain is not real or not physical. The nervous system that generates and interprets pain signals is directly shaped by stress, fear of movement, attention, and expectation, regardless of which tissue is involved.

## 13.1 Acceptance and Mindfulness-Based Approaches

A systematic review and meta-analysis of acceptance-based interventions across chronic pain conditions broadly, not limited to shoulder or spine pain, found meaningful effects on both pain and psychological outcomes including depression, and concluded these approaches can be a good option either alone or alongside other treatment.<sup>22</sup> Similarly, the mindfulness meditation research referenced in Chapter 12 showed moderate improvement in pain outcomes in chronic pain generally.<sup>21</sup>

**Reading that honestly.** Neither of these bodies of research was conducted specifically in people with rotator cuff pain, so they should be understood as general chronic pain evidence that reasonably extends to your situation, not as shoulder-specific proof.

## 13.2 Addressing Fear of Movement

A common and understandable response to shoulder pain is guarding the arm and avoiding movements that provoke discomfort. In the short term this is protective. Sustained over weeks and months, excessive guarding can contribute to stiffness, weakness, and a nervous system that becomes increasingly sensitive to movements that are not actually dangerous. Working with a provider who helps you gradually and confidently reintroduce movement, rather than avoiding it indefinitely, is a meaningful part of recovering full function, and it is one more reason the graduated exercise approach in Chapter 8 matters as much for your nervous system as it does for your tendon.

# 14. The In-Clinic Pre-Treatment Protocol

---

Understand what this is for, because the distinction matters.

None of this rebuilds tissue. The loading and strengthening work in the previous chapters does that. What this protocol does is reduce the muscle tone, stiffness and tissue sensitivity that make the strength work harder to tolerate and less pleasant to do, particularly in the early weeks when things are at their most irritable.

It is preparation, not treatment. I use it in the clinic before the working part of a session, and I give it to patients to use at home before theirs. Patients who do it tend to tolerate the loading better and stick with it longer, and sticking with it is the actual mechanism by which recovery happens.

If you have to choose between doing this and doing your strength work, do your strength work.

**Roller (Vyper or foam roller).** Between the shoulder blades, roll 1 to 2 minutes.

**Venom Shoulder**, set to vibrate and heat, while performing:

- 90/90 pillow press, 5 reps of internal and external rotation each shoulder, holding 3 to 5 seconds.
- Reach, roll and lift on an exercise ball, 5 reps, holding 5 to 8 seconds.
- Thread the needle, 5 reps each side, holding 3 to 5 seconds.

**Percussion device (Hypervolt), round or flat knob.** In the armpit with the elbow bent, moving the arm overhead and down, 1 minute each side. Reposition off nerves if you get numbness or tingling.

**Vibrating sphere (Hypersphere Mini).** Infraspinatus and teres minor, 1 minute each muscle.

**Massage ball or sphere pinned against a wall.** Pectoralis, moving the arm up and down and away from the wall over tender areas.

**Safety, and I mean these.**

- Avoid all of this over an acutely injured area in the first 48 to 72 hours.
- Reduce pressure over anything actively inflamed.
- Skip percussion therapy entirely if you have a pacemaker or another implanted electronic device.
- Avoid aggressive pressure over varicose veins or compromised skin.
- Start light. Pain must not exceed 6 out of 10. Back off immediately for sharp or shooting pain.
- Stop and seek advice if you get numbness, tingling, or weakness.

# 15. Red Flags: When to Seek Immediate Care

---

Most shoulder pain is no emergency. A small number of presentations, though, need prompt medical attention rather than a course of conservative care.

## **Seek immediate care if you experience**

- Shoulder pain accompanied by chest pain, pressure, shortness of breath, nausea, or sweating, which can represent referred cardiac pain and should always be evaluated urgently
- An unexplained lump or mass around the shoulder
- Fever together with joint swelling, redness, and warmth
- Significant trauma, such as a fall or collision, followed by inability to lift or move the arm
- New or progressively worsening numbness, tingling, or weakness spreading down the arm

## **Contact your provider promptly if**

- Pain has not improved at all after several weeks of appropriate conservative care
- Pain is severe enough to consistently disrupt sleep
- You notice new weakness that was not present initially

# 16. Building Your Personal Recovery Plan

---

Use this as a starting framework to discuss with your provider, not a rigid prescription. Every shoulder arrives with a different history, and your plan should reflect that.

## **Weeks 1 to 2: Calm and Protect**

Gentle pendulum motion, isometric holds if the shoulder is irritable, ice or heat for comfort, and an initial evaluation that looks beyond the shoulder itself, including your thoracic spine and posture.

## **Weeks 3 to 8: Rebuild**

Progressive rotator cuff loading with attention to internal rotation and subscapularis strength, scapular stabilization work, thoracic mobility exercises, and manual therapy as a complement to, not a replacement for, active exercise.

## **Weeks 9 to 16: Strengthen and Restore**

Continued progressive loading toward the specific demands of your work, sport, or daily activities, along with ongoing attention to sleep, stress, and posture.

## **Beyond 16 Weeks: Maintenance**

Continue rotator cuff and scapular strengthening two to three times per week indefinitely, keep thoracic mobility work in your routine, and address any flare-ups early with the same graduated approach rather than waiting for them to become severe again.

# Final Thoughts

---

Shoulder pain that has lingered for months is exhausting, and being told to simply rest and take ibuprofen while it drags on is not a plan. The research reviewed for this guide points somewhere more useful and more hopeful: toward a graduated, specific exercise program as your most powerful tool, toward taking subscapularis and thoracic mobility seriously as pieces that are frequently overlooked, and toward understanding that a tear on a scan does not automatically mean your only path forward is surgery.

Rather than simply broken, your shoulder is a connected part of a larger system that has likely been asked to compensate for stiffness or weakness somewhere else for longer than you realized. Addressing that bigger picture, alongside direct, progressive loading of the tissues themselves, is the most evidence-aligned way forward available today.

If you are ready for a thorough evaluation that looks at your whole system rather than just the spot that hurts, I offer a free Root Discovery Session at Root Health. You can schedule yours at <https://rootcauseunlocked.com/discovery>. You can also reach the office at (919) 662-0520.

Consultations and ongoing care are both available wherever you are located. Being outside North Carolina is not a barrier to working with me. Tell me where you are when you reach out and I will lay out exactly how we would work together from there.

# 17. References

---

Every reference below was retrieved and verified against the published record on PubMed during the preparation of this guide.

1. Lewis J. Rotator cuff related shoulder pain: Assessment, management and uncertainties. *Man Ther.* 2016;23:57-68. PMID: 27083390
2. Teunis T, Lubberts B, Reilly BT, Ring D. A systematic review and pooled analysis of the prevalence of rotator cuff disease with increasing age. *J Shoulder Elbow Surg.* 2014;23(12):1913-1921. PMID: 25441568
3. Lädermann A, Collin P, Zbinden O, Meynard T, Saffarini M, Chiu JC. Diagnostic Accuracy of Clinical Tests for Subscapularis Tears: A Systematic Review and Meta-analysis. *Orthop J Sports Med.* 2021;9(9):23259671211042011. PMID: 35146034
4. Wainner RS, Whitman JM, Cleland JA, Flynn TW. Regional interdependence: a musculoskeletal examination model whose time has come. *J Orthop Sports Phys Ther.* 2007;37(11):658-660. PMID: 18057674
5. Kebaetse M, McClure P, Pratt NA. Thoracic position effect on shoulder range of motion, strength, and three-dimensional scapular kinematics. *Arch Phys Med Rehabil.* 1999;80(8):945-950. PMID: 10453773
6. Yu S, Chen S, Yang Z, Ma X, Gou L, Yang L. The effects of different types of thoracic manual therapy for shoulder dysfunction: a systemic review with meta-analysis. *Musculoskelet Sci Pract.* 2026;84:103565. PMID: 42068776
7. Thoomes EJ, Tilborghs G, Heneghan NR, Falla D, De Graaf M. Effectiveness of Thoracic Spine Manipulation for Upper Quadrant Musculoskeletal Disorders: A Systematic Review. *J Manipulative Physiol Ther.* 2025;48(1-5):422-434. PMID: 41196244
8. Boyles RE, Ritland BM, Miracle BM, et al. The short-term effects of thoracic spine thrust manipulation on patients with shoulder impingement syndrome. *Man Ther.* 2009;14(4):375-380. PMID: 18703377
9. Zhao J, Pan J, Zeng LF, Wu M, Yang W, Liu J. Risk factors for full-thickness rotator cuff tears: a systematic review and meta-analysis. *EFORT Open Rev.* 2021;6(11):1087-1096. PMID: 34909227
10. Fan N, Yuan S, Du P, et al. The effects of smoking on clinical and structural outcomes after rotator cuff repair: a systematic review and meta-analysis. *J Shoulder Elbow Surg.* 2022;31(3):656-667. PMID: 34813890
11. Steuri R, Sattelmayer M, Elsig S, et al. Effectiveness of conservative interventions including exercise, manual therapy and medical management in adults with shoulder impingement: a systematic review and meta-analysis of RCTs. *Br J Sports Med.* 2017;51(18):1340-1347. PMID: 28630217
12. Beard DJ, Rees JL, Cook JA, et al. Arthroscopic subacromial decompression for subacromial shoulder pain (CSAW): a multicentre, pragmatic, parallel group, placebo-controlled, three-group, randomised

surgical trial. Lancet. 2018;391(10118):329-338. PMID: 29169668

13. Kukkonen J, Ryösä A, Joukainen A, et al. Operative versus conservative treatment of small, nontraumatic supraspinatus tears in patients older than 55 years: over 5-year follow-up of a randomized controlled trial. J Shoulder Elbow Surg. 2021;30(11):2455-2464. PMID: 33774172
14. Ryösä A, Laimi K, Äärimaa V, Lehtimäki K, Kukkonen J, Saltychev M. Surgery or conservative treatment for rotator cuff tear: a meta-analysis. Disabil Rehabil. 2017;39(14):1357-1363. PMID: 27385156
15. Holmgren T, Björnsson Hallgren H, Öberg B, Adolfsson L, Johansson K. Effect of specific exercise strategy on need for surgery in patients with subacromial impingement syndrome: randomised controlled study. BMJ. 2012;344:e787. PMID: 22349588
16. Maroon JC, Bost JW. Omega-3 fatty acids (fish oil) as an anti-inflammatory: an alternative to nonsteroidal anti-inflammatory drugs for discogenic pain. Surg Neurol. 2006;65(4):326-331. PMID: 16531187
17. Daily JW, Yang M, Park S. Efficacy of Turmeric Extracts and Curcumin for Alleviating the Symptoms of Joint Arthritis: A Systematic Review and Meta-Analysis of Randomized Clinical Trials. J Med Food. 2016;19(8):717-729. PMID: 27533649
18. Bannuru RR, Osani MC, Al-Eid F, Wang C. Efficacy of curcumin and Boswellia for knee osteoarthritis: Systematic review and meta-analysis. Semin Arthritis Rheum. 2018;48(3):416-429. PMID: 29622343
19. Shaw G, Lee-Barthel A, Ross ML, Wang B, Baar K. Vitamin C-enriched gelatin supplementation before intermittent activity augments collagen synthesis. Am J Clin Nutr. 2017;105(1):136-143. PMID: 27852613
20. Krause AJ, Prather AA, Wager TD, Lindquist MA, Walker MP. The Pain of Sleep Loss: A Brain Characterization in Humans. J Neurosci. 2019;39(12):2291-2300. PMID: 30692228
21. Goyal M, Singh S, Sibinga EM, et al. Meditation programs for psychological stress and well-being: a systematic review and meta-analysis. JAMA Intern Med. 2014;174(3):357-368. PMID: 24395196
22. Veehof MM, Oskam MJ, Schreurs KMG, Bohlmeijer ET. Acceptance-based interventions for the treatment of chronic pain: a systematic review and meta-analysis. Pain. 2011;152(3):533-542. PMID: 21251756

*This guide was compiled from peer-reviewed medical literature and clinical experience. Every citation was verified against the published record. It is intended to give you knowledge and options to discuss with your healthcare provider. It is not a substitute for individual medical care, it does not diagnose any condition, and it does not promise a cure.*

© 2026 Root Health · For personal educational use only

# What this guide is for

---

Frozen shoulder is the condition where doing more, harder, faster makes it worse, and almost every instinct you have will be wrong.

You cannot stretch your way out of it in the early phase. Pushing into pain does not speed it up. And the treatments that sound most decisive are, according to the largest trial ever run on this condition, no better at twelve months than structured physiotherapy.

That last sentence is worth a lot of money and a lot of unnecessary surgery, and it is the reason I rewrote this guide.

What this guide will do is tell you which phase you are in, what is worth doing in each, what the evidence actually supports, and how to avoid the two mistakes that cost people the most: being too aggressive early, and being too passive late.

This guide is educational. It does not diagnose your shoulder and it does not replace an examination.

# Chapter 1. What is actually happening

---

The shoulder joint sits inside a capsule, a sleeve of connective tissue. In frozen shoulder, properly called adhesive capsulitis, that capsule becomes inflamed and then progressively thickens and contracts. The joint space itself shrinks.

This matters for one reason above all others: **the restriction is structural, not muscular**. You are not tight. Your capsule is physically contracted. That is why stretching harder does not work the way it does for a tight muscle, and why forcing range in the inflamed phase provokes more inflammation and more contraction.

**The defining sign is loss of passive external rotation.** If someone else moves your arm for you, completely relaxed, and it still will not rotate outward, that is the capsule blocking it. This is the finding that separates frozen shoulder from a rotator cuff problem, where passive range is usually preserved even when active range is not. It takes fifteen seconds to check and it is the single most useful examination finding in this condition.

# Chapter 2. The three phases, and why they change everything

---

Frozen shoulder moves through phases. Treating the wrong phase with the right treatment for a different one is the commonest error.

**Phase 1, freezing.** Roughly two to nine months. Pain dominates. Night pain is often severe and is usually what drives people to seek help. Range is progressively lost. **This is the phase where aggressive stretching is actively counterproductive**, because you are stretching inflamed tissue and provoking more of the process you are trying to stop. This phase is about pain control and gentle movement within comfort.

**Phase 2, frozen.** Roughly four to twelve months. Pain settles considerably, and stiffness becomes the main problem. Night pain typically improves. **This is where the mobility work earns its place**, because you are now working on contracted tissue rather than angry tissue.

**Phase 3, thawing.** Roughly five to twenty-four months. Range gradually returns.

**The honest thing about that timeline** is that it is long, the phases overlap, and the total course commonly runs one to three years. I will not tell you otherwise to make this guide more comfortable.

**And an important correction to what you may have been told.** Frozen shoulder is frequently described as self-limiting, meaning it fully resolves on its own. The evidence is less reassuring than that. A meaningful proportion of people are left with some residual restriction long term. It usually improves greatly. It does not always resolve completely. That is a reason to engage with it rather than to wait it out passively.

# Chapter 3. Who gets it and why that matters

---

**Diabetes is the big one.** The association is strong and well established. People with diabetes get frozen shoulder considerably more often, tend to have more severe and more prolonged courses, and are more likely to get it in both shoulders. If you have frozen shoulder and nobody has checked your blood sugar, that is worth doing. If you have diabetes and frozen shoulder, glucose control is part of the plan rather than a separate issue.

**Thyroid disease**, particularly hypothyroidism, is also associated.

**Age and sex.** Most commonly between 40 and 60, more often women.

**After a period of immobilisation.** After a fracture, after surgery, after an injury that made you stop using the arm. This is the one that is partly preventable, and it is why early gentle movement after any shoulder injury matters.

**The other shoulder.** Having had it on one side raises the risk on the other. Often enough to be worth knowing, though not always.

# Chapter 4. Making sure it is actually this

---

Several things masquerade as frozen shoulder, and the treatments differ substantially.

**Rotator cuff tear or tendinopathy.** Active movement limited, passive movement relatively preserved. In frozen shoulder passive external rotation is lost. That is the discriminator.

**Osteoarthritis of the shoulder.** Can also lose passive range. Usually shows on x-ray, and the history is different.

**Referred pain from the neck.** Common, and the shoulder examination is usually normal for range.

**Calcific tendinitis.** Can be acutely and severely painful. Shows on imaging.

**And the ones that are not shoulder problems at all.** Pain referred from the chest, the gallbladder, or the diaphragm. Rare, real, and the reason a shoulder that does not behave like a shoulder gets looked at properly.

**On imaging:** frozen shoulder is largely a clinical diagnosis, made on that passive external rotation finding. X-ray is mainly used to exclude arthritis. MRI is not routinely needed to make the diagnosis.

# Chapter 5. What the largest trial found

---

This is the chapter that should change what you do.

**UK FROST** was a multicentre, pragmatic, three-arm randomised trial across 35 UK hospital sites, in patients with primary frozen shoulder referred to secondary care. It compared early structured physiotherapy, manipulation under anaesthesia, and arthroscopic capsular release.

**At twelve months, none of the three was superior to the others on patient-reported pain and function** (PMID 33010843).

And there was a safety signal worth knowing: **arthroscopic capsular release carried a higher risk of complications than manipulation under anaesthesia.**

Sit with what that means. The two surgical options, one of which involves a general anaesthetic while a surgeon tears your capsule and the other of which involves cutting it, did not outperform structured physiotherapy at a year. One of them carried more complications.

Surgery can still have a role. The point is that it is not the obviously superior choice it is often presented as, and that anyone recommending it to you should be able to say why, in your specific case, given that trial.

**On corticosteroid injection.** A systematic review of randomised trials in primary care supports corticosteroid injection for adhesive capsulitis (PMID 27570870). The pattern across the literature is consistent: injection is meaningfully better in the short term, particularly for pain, with that advantage narrowing at around three to six months as other approaches catch up.

**Which gives a genuinely useful sequencing principle.** Injection is most valuable in the painful freezing phase, where pain is the problem and where it can make the difference between sleeping and not. It is less compelling in the frozen phase, where stiffness rather than pain is the issue.

**Hydrodilatation**, where fluid is injected to distend the capsule, has a reasonable evidence base and is worth asking about.

# Chapter 6. What to actually do, by phase

---

## **Phase 1, freezing. The goal is pain control and preserving what range you have.**

- Gentle movement within your comfortable range, frequently. Pendulum swings, gentle assisted range.
- **Do not push into pain.** In this phase that provokes the process.
- Pain management matters more than range work. Discuss injection, which has the best short-term evidence, particularly if night pain is wrecking your sleep.
- Sleep positioning: pillow supporting the arm, avoiding lying on that side.
- Keep using the arm for everyday tasks within comfort. Total protection speeds stiffening.

## **Phase 2, frozen. The goal is range.**

- This is where sustained stretching earns its place, because you are now working on contracted rather than inflamed tissue.
- Long, low-load holds beat short aggressive ones. Think 30 seconds and longer, repeated, rather than forcing.
- Some discomfort during stretching is expected here. Pain that lingers for hours afterwards means you went too hard.
- Add strengthening as range improves.
- This is the phase where the pre-treatment protocol in Chapter 8 is most useful.

## **Phase 3, thawing. The goal is full function.**

- Progressive strengthening through the range you have regained.
- Return to overhead and behind-the-back activity gradually.
- Keep going. Stopping when it is 80 percent better is how people end up with permanent restriction.

**The single most common mistake in each phase:** being too aggressive in phase 1, and being too passive in phase 2 and 3.

**One thing to work on that is not the shoulder.** Raising your arm overhead is not purely a shoulder movement. Your shoulder blade has to rotate across your rib cage, and your mid-back has to extend, for the arm to finish the range. That means a stiff mid-back quietly costs you shoulder motion, and it costs you the same motion whether or not you have a frozen shoulder.

Be clear about what that does and does not mean here. In frozen shoulder the restriction itself is in the joint capsule, and no amount of mid-back work will release a contracted capsule. What mid-back work does is stop you from losing motion you did not have to lose, and give you the most available range back once the capsule

starts to let go in phase 3. On the evidence, a systematic review of thoracic spine manipulation across upper quadrant problems found it more effective than general conservative therapy for shoulder pain, but no more effective than a sham procedure, on very low to low quality evidence.<sup>3</sup> So I would not sell it to you as a treatment for this condition. I work on it because it is cheap, it is low risk, and the motion belongs there anyway.

# Chapter 7. Measuring yourself so you can tell if this is working

---

Frozen shoulder changes so slowly that memory is useless. Measure monthly.

- **Passive external rotation.** Elbow tucked at your side, bent to 90 degrees, let someone rotate your forearm outward. Estimate the angle or photograph it against a doorframe. This is the number that matters most.
- **Hand behind back.** How far up your spine can you reach? Note the level against a landmark.
- **Overhead reach.** Against a wall, marked.
- **Night pain.** Nights per week you are woken.
- **Function.** One specific task you cannot currently do. Fastening a bra, reaching a seatbelt, getting a plate from a high shelf.

Photograph the range measurements monthly. In a condition this slow, a photo from eight weeks ago is worth more than any recollection.

# Chapter 8. The In-Clinic Pre-Treatment Protocol

---

Understand what this is for. It does not stretch the capsule. What it does is reduce the tone and guarding in the surrounding musculature, the upper back and the chest, which is where a great deal of the secondary discomfort in frozen shoulder actually comes from, and which makes the range work more tolerable.

Most useful in phase 2 and 3. In the acutely painful freezing phase, use it lightly or not at all.

**Roller (Vyper or foam roller).** Between the shoulder blades, roll 1 to 2 minutes.

**Venom Shoulder**, set to vibrate and heat, while performing:

- 90/90 pillow press, 5 reps of internal and external rotation each shoulder, holding 3 to 5 seconds.
- Reach, roll and lift on an exercise ball, 5 reps, holding 5 to 8 seconds.
- Thread the needle, 5 reps each side, holding 3 to 5 seconds.

**Percussion device (Hypervolt), round or flat knob.** In the armpit with the elbow bent, moving the arm overhead and down, 1 minute each side. Reposition off nerves if you get numbness or tingling.

**Vibrating sphere (Hypersphere Mini).** Infraspinatus and teres minor, 1 minute each muscle.

**Massage ball or sphere pinned against a wall.** Pectoralis, moving the arm up and down and away from the wall over tender areas.

**Safety, and I mean these.**

- Avoid all of this over an acutely injured area in the first 48 to 72 hours.
- Reduce pressure over anything actively inflamed.
- Skip percussion therapy entirely if you have a pacemaker or another implanted electronic device.
- Avoid aggressive pressure over varicose veins or compromised skin.
- Start light. Pain must not exceed 6 out of 10. Back off immediately for sharp or shooting pain.
- Stop and reposition if you get numbness, tingling, or weakness in the arm.
- **In the freezing phase, less is more.** If it hurts for hours afterwards, you did too much.

# Chapter 9. Nutrition, briefly and honestly

---

There is no supplement with trial evidence for frozen shoulder. Anyone selling you one is ahead of the research.

**What genuinely matters here is glucose control**, given the strength of the diabetes association. That is a reason to have your blood sugar checked and managed if it is abnormal, not a supplement recommendation, and it is more likely to affect your course than anything you can buy.

**Thyroid function** is worth checking for the same reason.

Correcting a documented vitamin D deficiency is defensible on general grounds and is not a frozen shoulder treatment.

# Chapter 10. Red flags

---

- **Shoulder pain with chest pain, breathlessness, sweating or nausea.** Cardiac symptoms refer to the shoulder and arm. Emergency assessment.
- **Fever with a hot, swollen, exquisitely painful shoulder.** Joint infection. Same-day care.
- **Significant trauma before onset**, particularly with deformity or complete inability to move the arm. Fracture or dislocation.
- **Progressive weakness or numbness in the arm or hand**, which points at a nerve rather than the capsule.
- **A history of cancer with new shoulder pain**, particularly night pain that is not positional.
- **Unexplained weight loss, night sweats, or feeling systemically unwell** alongside the shoulder pain.
- **Rapid onset over days rather than weeks**, which does not fit the frozen shoulder pattern.

# Chapter 11. Building your plan

---

## **Weeks 0 to 2. Establish where you are.**

- Confirm the diagnosis: is passive external rotation genuinely lost?
- Identify your phase from Chapter 2.
- Get blood sugar and thyroid checked if they have not been.
- Take your baseline measurements from Chapter 7 and photograph them.

## **If you are in the freezing phase.**

- Prioritise pain and sleep. Discuss injection if night pain is significant.
- Gentle range within comfort, several times daily.
- Resist the urge to stretch harder. It will not speed this up.
- Re-measure monthly.

## **If you are in the frozen phase.**

- Shift the emphasis to sustained range work, long holds, daily.
- Add the pre-treatment protocol from Chapter 8 before your range sessions.
- Begin strengthening as range allows.
- Re-measure monthly. Expect slow progress and judge it against last month, not last week.

## **If you are in the thawing phase.**

- Progressive strengthening through the regained range.
- Return to overhead and behind-the-back tasks.
- Keep going past the point where it feels mostly fine.

**When to escalate.** If you are six or more months in, in the frozen phase, doing consistent work, and genuinely not progressing on your measured range, that is the point to discuss what else is available. Take UK FROST with you to that conversation (PMID 33010843).

# Final thoughts

---

Frozen shoulder asks for something most conditions do not: patience in one phase and persistence in another, and the discipline to tell which one you are in.

The good news is that the evidence is more reassuring than the experience feels. The largest trial ever run found that structured physiotherapy held its own against both surgical options at twelve months. You are unlikely to be worse off for taking the measured route, and you may avoid a general anaesthetic.

Measure monthly, photograph it, match your effort to your phase, and give it the time it genuinely takes.

# References

---

All PMIDs verified against live PubMed records on 2026-09-01.

1. Sun Y, Zhang P, Liu S, Li H, Jiang J, Chen S, Chen J. Corticosteroid injection for adhesive capsulitis in primary care: a systematic review of randomised clinical trials. *Rheumatology International / systematic review*. 2016. **PMID 27570870**.
2. Rangan A, Brealey SD, Keding A, Corbacho B, Northgraves M, Kottam L, et al. Management of adults with primary frozen shoulder in secondary care (UK FROST): a multicentre, pragmatic, three-arm, superiority randomised clinical trial. *The Lancet*. 2020;396(10256):977-89. **PMID 33010843**.. (Erratum: *Lancet*. 2021;397(10269):98, a Department of Error notice. PMID 33422260. Screened 2026-09-05; corrected, not retracted.)
3. Thoomes EJ, Tilborghs G, Heneghan NR, Falla D, De Graaf M. Effectiveness of Thoracic Spine Manipulation for Upper Quadrant Musculoskeletal Disorders: A Systematic Review. *J Manipulative Physiol Ther*. 2025;48:422-434. PMID 41196244

# Working with Root Health

---

A shoulder losing motion in every direction needs a diagnosis, not a generic shoulder program. If that is your picture, I offer a free Root Discovery Session at Root Health to talk through where you are in the timeline and what a reasonable next step looks like. You can schedule one at [rootcauseunlocked.com/discovery](https://rootcauseunlocked.com/discovery), or reach the office at (919) 662-0520.

Consultations and ongoing care are both available wherever you are located. Being outside North Carolina is not a barrier to working with me. Tell me where you are when you reach out and I will lay out exactly how we would work together from there.

If you would rather start on your own, the free Root Cause Unlocked tracker has a mode built for musculoskeletal recovery, at [rootcauseunlocked.com/tracker](https://rootcauseunlocked.com/tracker). Eight weeks of it turns “I think it might be helping” into something you and I can both read.

*This guide is educational and does not constitute medical advice, diagnosis, or treatment. It does not create a doctor-patient relationship. Seek examination for shoulder pain, and seek urgent care for any of the red flags in Chapter 10.*

**Disclaimer:** *This ebook is for educational purposes only and does not constitute medical advice. Always consult with a qualified healthcare provider before starting any new treatment, supplement, exercise program, or dietary change. Never start, stop, or change a prescribed medication without talking to the prescriber first. If you experience progressive weakness, severe swelling, fever, or symptoms that worsen rapidly, seek medical attention immediately.*

**A note on the evidence in this guide.** *Every numbered citation in this book has been checked against the published record. Where the research studied a different tissue, a different joint, or a different population than tennis elbow, that is stated plainly in the text so you can judge the strength of the evidence yourself. Where a popular claim had no supporting study behind it, the claim was removed rather than dressed up.*

# 1. What Is Tennis Elbow?

---

Tennis elbow, medically known as lateral epicondylitis (or more accurately, lateral epicondylosis or lateral epicondylar tendinopathy), is a painful condition affecting the tendons on the outside of your elbow. Despite the name, most people who develop it are not tennis players. It shows up far more often in workers and in people whose daily tasks involve repeated gripping.

The condition involves the extensor carpi radialis brevis (ECRB) tendon and other forearm extensor tendons that attach to the lateral epicondyle, the bony bump on the outside of your elbow. These tendons are responsible for extending your wrist and fingers.

## Common Symptoms

The most common symptom is pain or burning on the outer part of the elbow, often radiating down the forearm toward the wrist. Many people notice weakness in grip strength, showing up as difficulty shaking hands, turning a doorknob, or holding a coffee cup. Pain typically worsens with wrist extension, gripping, or lifting, especially when the palm faces down. Stiffness in the elbow is common, particularly in the morning, along with tenderness to touch directly over the lateral epicondyle.

**Important distinction.** Tennis elbow affects the outside of the elbow. Pain on the inside of the elbow is called golfer's elbow (medial epicondylitis) and involves different tendons.

## 2. Understanding the Anatomy

---

The lateral epicondyle is the bony prominence on the outer side of your elbow. It serves as the attachment point for the common extensor tendon, which branches into several muscles. The extensor carpi radialis brevis (ECRB) is the primary tendon involved in tennis elbow. The neighboring muscles each have their own job. The extensor digitorum communis extends the fingers, the extensor carpi radialis longus extends and abducts the wrist, the extensor digiti minimi extends the little finger, and the extensor carpi ulnaris extends and adducts the wrist.

These muscles work together to extend your wrist and fingers, stabilize your wrist during gripping activities, and absorb forces transmitted from the hand and wrist up the arm.

### **Why the ECRB Is Vulnerable**

The ECRB tendon has a relatively poor blood supply compared to other tendons in the body. It also passes over the rounded head of the radius bone, creating friction with every wrist and elbow movement. This combination of limited blood supply and mechanical stress makes it particularly susceptible to overuse injury.

# 3. Common Causes and Risk Factors

---

## What Actually Happens in the Tendon

Tennis elbow is a tendinopathy rather than a primarily inflammatory condition. Tissue taken from the affected tendon during surgery shows a characteristic degenerative picture rather than an inflammatory one<sup>1</sup>. That picture includes angiofibroblastic hyperplasia, an abnormal growth of blood vessels and fibroblasts, along with disorganized collagen fibers that lose the normal parallel alignment and become haphazard. The ground substance, the gel-like matrix between collagen fibers, becomes excessive, and inflammatory cells are largely absent, unlike true “-itis” conditions where they would be expected.

This is why the terms “tendinosis” and “tendinopathy” are now preferred over “tendinitis.” The problem is tissue degeneration and failed healing, not active inflammation<sup>1</sup>.

## Primary Causes

The most common cause is repetitive wrist extension and gripping, where repeated microtrauma exceeds the tendon's capacity to repair itself. A sudden increase in activity, such as starting a new sport, job, or hobby that involves repetitive forearm use, can also trigger the condition, as can poor technique or ergonomics, meaning incorrect form in sports or an improper workstation setup. Age-related tendon change plays a role too, since tendons lose elasticity and blood supply with age.

## Risk Factors

**Occupation.** This is the best-documented risk factor. A population-based case-referent study of 267 new tennis elbow cases found that manual job tasks roughly tripled the odds of developing the condition, and that a combined index of forceful work, non-neutral hand and arm posture, and repetition showed a clear dose response: the odds rose about fourfold at the highest strain level<sup>2</sup>. A review of the occupational literature reaches the same conclusion, that force, repetition, and awkward posture are the dominant work-related drivers<sup>3</sup>. Occupations commonly affected include plumbers, carpenters, painters, and mechanics; assembly line workers, butchers, and chefs; office workers with poor keyboard and mouse ergonomics; and musicians, especially string players and pianists.

**Age.** Tennis elbow is most common in middle age. A large general-population survey of upper-limb musculoskeletal disorders found lateral epicondylitis clustering in the middle adult years rather than in the young or the elderly<sup>4</sup>.

**Sports.** Tennis (especially backhand strokes), squash, badminton, weightlifting, rowing, and fencing all involve repeated gripping and forearm loading.

**Other factors commonly discussed.** Smoking, obesity, diabetes, metabolic syndrome, and inherited differences in collagen structure are all raised in the tendon literature as plausible contributors to slower tendon healing. The evidence linking each of them specifically to tennis elbow is thinner than the occupational evidence, so treat them as reasonable general health considerations rather than proven causes of this particular condition.

**Psychosocial factors.** In the Danish case-referent study, low social support at work was an independent risk factor among women even after adjusting for physical strain<sup>2</sup>. Stress and job strain are not imaginary contributors.

## 4. How Tennis Elbow Is Diagnosed

---

Tennis elbow is primarily a clinical diagnosis. Imaging is usually not necessary unless the diagnosis is uncertain or symptoms are not improving with conservative care<sup>5</sup>.

### Clinical Examination

**Palpation.** Tenderness directly over the lateral epicondyle and just distal to it, especially over the ECRB tendon.

**Special tests.** In Cozen's test (resisted wrist extension), the patient extends the wrist against resistance with the elbow extended, and pain over the lateral epicondyle indicates a positive test. In Mill's test, the examiner passively flexes the patient's wrist while extending the elbow and pronating the forearm (turning the palm down), and pain at the lateral epicondyle is positive. Maudsley's test involves the examiner applying resistance to middle finger extension, and pain in the same spot suggests ECRB involvement. In the chair test, the patient attempts to lift a chair by the back with the palm facing down and the elbow extended, and pain indicates tennis elbow.

**Strength testing** typically includes pain-free grip strength measurement, which is often reduced on the affected side and is the most commonly used objective outcome measure in the research<sup>5</sup>, along with wrist extensor strength testing.

### Imaging (When Needed)

Ultrasound can visualize tendon thickening, hypoechoic (darker) areas indicating degeneration, calcifications, and tears, and is useful for guiding injections. MRI provides detailed images of tendon structure, can detect partial tears, and helps rule out other conditions, though it is usually reserved for stubborn cases or pre-surgical planning. X-rays are generally not helpful for tennis elbow itself. They may still be used to rule out arthritis, loose bodies, or calcification.

### Differential Diagnosis

Your healthcare provider will also consider other conditions that can mimic tennis elbow. Radial tunnel syndrome involves compression of the radial nerve, causing similar pain but typically more diffuse and without point tenderness. Posterior interosseous nerve (PIN) syndrome, which involves a branch of that same nerve, is another possibility, as is cervical radiculopathy, nerve root compression in the neck that causes referred arm pain. Elbow arthritis (degenerative changes in the elbow joint) and lateral collateral ligament injury (ligament instability) are also considered, and, though rare, tumor or infection must be ruled out in atypical presentations.

## 5. The Healing Timeline: What to Expect

---

Understanding the natural course of tennis elbow helps set realistic expectations and prevents premature escalation to invasive treatments.

### Natural History

In the acute phase, the first 0 to 6 weeks, pain is most noticeable, and many people self-treat or modify activities. The subacute phase, from 6 weeks to 6 months, is where structured rehabilitation matters most, though pain may fluctuate. If symptoms persist beyond 6 months, the condition is considered chronic, and more intensive or interventional treatments may be considered.

### What the Trials Actually Show

Two of the most important randomized trials in this field compared active treatment against simply waiting.

In a Dutch trial of 185 patients, corticosteroid injection was clearly the best option at 6 weeks. By 52 weeks the picture had reversed: the injection group had the worst outcomes, and physiotherapy and a wait-and-see policy had converged. The authors concluded that the relative gain of physiotherapy over waiting was small, and that patients should be told the trade-offs before choosing<sup>6</sup>.

An Australian trial of 198 patients found the same short-term-versus-long-term reversal. Corticosteroid injection won at 6 weeks and lost by 52 weeks, with high recurrence. Physiotherapy (mobilisation with movement plus exercise) outperformed waiting in the first 6 weeks and reduced the need for additional treatment, and by one year the physiotherapy and wait-and-see groups were similar<sup>7</sup>.

**The practical takeaway.** Most people improve substantially within a year regardless of what they do, structured exercise gets you there more comfortably, and a fast fix that looks impressive at 6 weeks can leave you worse off at 12 months. Recurrence is common if the underlying load and biomechanical issues are not addressed.

# 6. Supplements for Tendon Healing and Inflammation

---

**Read this section carefully.** The supplements below are grouped by how strong the evidence actually is. Almost none of them have been tested in people with tennis elbow specifically. Most of the research is in knee osteoarthritis, general wound healing, or laboratory models. That is not the same as useless, though it does mean the honest claim is “biologically plausible and reasonably safe,” not “proven to heal your elbow.”

Always discuss supplements with your healthcare provider, especially if you take medications. Several of the items below can interact with blood thinners.

## 6.1 Collagen / Gelatin plus Vitamin C

**What it does.** Provides the building blocks (the amino acids glycine, proline, and hydroxyproline) for tendon collagen synthesis. Vitamin C is a required cofactor for collagen cross-linking and stability.

**Evidence.** In a controlled crossover study in healthy young men, taking 15 g of vitamin C-enriched gelatin one hour before a short bout of rope-skipping doubled the blood level of PINP, a marker of new collagen formation, compared with placebo. In the same study, engineered ligaments grown in the laboratory and bathed in serum drawn from those subjects after supplementation showed increased collagen content and improved mechanical strength<sup>8</sup>. The subjects were healthy volunteers and the outcome was a laboratory marker, not a clinical trial in people with tendon pain.

Two longer studies in the broader connective-tissue literature are worth knowing. A 24-week randomized trial in athletes with activity-related joint pain reported reduced joint pain with collagen hydrolysate<sup>9</sup>. A meta-analysis of randomized placebo-controlled trials found collagen supplementation improved symptom scores in osteoarthritis<sup>10</sup>. Both of those trials were conducted in joint disease, and neither one touched tennis elbow.

**Typical dosage.** 10 to 15 g hydrolyzed collagen or gelatin daily, with roughly 50 mg vitamin C. The lower end of that range is the dose used in a randomized trial of collagen peptides in active adults.<sup>47</sup> **Timing.** 30 to 60 minutes before your loading exercises, mirroring the study protocol. **Best form.** Hydrolyzed collagen peptides (dissolve easily, well absorbed). **Caution.** Generally very safe. Source quality matters.

## 6.2 Vitamin C

**What it does.** Essential cofactor for collagen synthesis, an antioxidant that protects cells from oxidative stress, and a supporter of immune function.

**Evidence.** Vitamin C deficiency causes scurvy, characterized by defective collagen and tissue fragility. This is settled physiology. A 2018 systematic review looking specifically at vitamin C after musculoskeletal injury found only ten eligible articles, and the tendon-healing findings came from animal studies that reported increased type I collagen and scar tissue formation with supplementation<sup>11</sup>. No human clinical trial has yet established that supplementing vitamin C speeds tendon repair.

**Typical dosage.** 500 to 1,000 mg daily. The lower end of that range is a dose arm from the randomized dose-response trial of vitamin C after wrist fracture.<sup>46</sup> **Best form.** Ascorbic acid or buffered forms (calcium or magnesium ascorbate). **Caution.** High doses can cause digestive upset. Split doses through the day.

## 6.3 Omega-3 Fatty Acids (EPA/DHA)

**What it does.** EPA and DHA are incorporated into cell membranes and are precursors to specialized pro-resolving mediators (resolvins and protectins), signaling molecules that actively switch off an inflammatory response rather than merely blocking it.

**Evidence.** The mechanism above is well described in a review of how omega-3 fatty acids act on inflammatory processes, from the molecular level up<sup>12</sup>. That review makes the case for biological plausibility, but it is not a pain trial, and omega-3 supplementation has never been put to the test in tennis elbow.

**Typical dosage.** 1,200 to 2,400 mg combined EPA and DHA daily. That is the range used in the one study run in people with nonsurgical neck and back pain, where most participants took 1,200 mg daily and about a fifth took 2,400 mg. That study was an uncontrolled questionnaire follow-up rather than a placebo-controlled trial, so treat it as a starting point rather than a proven dose. Higher intakes appear in the joint-pain literature, but that work was done in rheumatoid arthritis and other inflammatory conditions rather than in the mechanical and degenerative problems this guide covers, so I have not carried its dose across.<sup>48</sup> **Best form.** Triglyceride or re-esterified triglyceride form **Caution.** May increase bleeding risk at higher doses. Tell your surgeon before any procedure.

## 6.4 Turmeric / Curcumin

**What it does.** Curcumin inhibits NF-kappa-B, a master regulator of inflammatory gene expression, and several inflammatory enzymes.

**Evidence.** A systematic review and meta-analysis of randomized trials found that turmeric extract at roughly 1,000 mg per day of curcumin improved arthritis symptoms. The authors were explicit about the limits of their own finding. The number of trials, the total sample size, and the methodological quality were not sufficient to draw definitive conclusions, and larger rigorous studies are needed<sup>13</sup>. Present that caveat to anyone who quotes curcumin research at you.

**Typical dosage.** 500 to 1,000 mg curcuminoids daily, preferably with piperine (black pepper extract) or in a liposomal or phytosome form **Caution.** May interact with blood thinners. Avoid in gallbladder disease.

## 6.5 Vitamin D

**What it does.** Essential for musculoskeletal health and involved in muscle function and inflammatory regulation.

**Evidence.** A well-known clinic study of 150 patients presenting with persistent, nonspecific musculoskeletal pain found that a striking proportion had severe vitamin D deficiency<sup>14</sup>. Note what this study is and is not: it measured how common deficiency was in that group, not whether correcting it relieves pain, since it was not a supplementation trial. There is, in fact, no credible study tying vitamin D status specifically to tennis elbow.

The reasonable position: if you have persistent musculoskeletal pain, having your level checked is sensible, and correcting a genuine deficiency is good general medicine.

**Typical dosage.** Based on your blood level, discussed with your provider. Commonly 2,000 to 5,000 IU daily when correcting a low level. **Best form.** D3 (cholecalciferol) **Caution.** Have levels rechecked. Toxicity is rare but real at very high sustained doses.

## 6.6 Magnesium

**What it does.** Required for hundreds of enzyme reactions including ATP production, and involved in normal muscle contraction and relaxation.

**Evidence.** A systematic review of randomized controlled trials of magnesium for chronic pain in adults located only nine eligible trials and concluded that the evidence was equivocal, while noting efficacy signals in some trials that justify better-designed future studies<sup>15</sup>. In other words, plausible and popular, but not proven.

**Typical dosage.** 200 to 400 mg elemental magnesium daily. Glycinate is well tolerated. This is a general supplemental intake range rather than a dose drawn from a trial in this condition, which is why no study is cited beside it. **Caution.** Can cause diarrhea, particularly the oxide and citrate forms. Avoid in severe kidney disease.

## 6.7 Zinc

**What it does.** Zinc-dependent enzymes are involved in collagen synthesis, tissue remodeling, and immune function.

**Evidence.** A review of zinc physiology and clinical applications states the position clearly: there is consensus that zinc deficiency impairs wound healing, but considerable disagreement in the literature about the optimal method and the true benefit of zinc supplementation in people who are not deficient<sup>16</sup>.

**Typical dosage.** 15 to 30 mg elemental zinc daily (picolinate or bisglycinate). This is a general supplemental intake range rather than a dose drawn from a trial in this condition. It sits above ordinary dietary intake and below the tolerable upper limit, which is why the copper caution below matters. **Caution.** Long-term high-dose zinc can cause copper deficiency. Do not exceed 40 mg daily without medical supervision.

## 6.8 Boswellia Serrata (Frankincense)

**What it does.** Inhibits 5-lipoxygenase (5-LOX), an enzyme in the inflammatory cascade, and reduces leukotriene production.

**Evidence.** A systematic review and meta-analysis of randomized trials in **osteoarthritis** patients concluded that Boswellia and its extract may be an effective and safe option, and that treatment should run for at least 4 weeks to see an effect<sup>17</sup>. The population studied was osteoarthritis; Boswellia has not been trialed in tennis elbow.

**Typical dosage.** 300 to 500 mg standardized extract (AKBA content) twice daily **Caution.** May interact with anti-inflammatory medications.

## 6.9 Methylsulfonylmethane (MSM)

**What it does.** An organic sulfur compound. Sulfur participates in the disulfide bonds that stabilize connective tissue proteins.

**Evidence.** A pilot randomized trial in **knee osteoarthritis** found that MSM at 3 g twice daily produced significant reductions in WOMAC pain and physical function impairment compared with placebo over a short intervention, with no major adverse events. The investigators themselves called for larger trials before drawing firm conclusions<sup>18</sup>. A later double-blind randomized trial in grade I to II **knee osteoarthritis** found that adding MSM to glucosamine-chondroitin sulfate produced significantly better WOMAC and pain scores at 12 weeks than glucosamine-chondroitin alone or placebo<sup>19</sup>.

Both of these are knee osteoarthritis trials, and neither one tells you anything about tendons or elbows.

**Typical dosage.** 1,500 to 3,000 mg daily **Caution.** Generally well tolerated. Minor digestive upset in some people.

## 6.10 Proteolytic Enzymes (Bromelain, Serrapeptase)

**What it does.** Proteolytic enzymes taken between meals are proposed to act systemically on inflammatory proteins.

**Evidence.** A review of bromelain describes its documented properties and the range of clinical uses that have been explored, including soft tissue and surgical trauma, wound debridement, and osteoarthritis symptoms<sup>20</sup>. That review simply describes the compound; it is not a trial demonstrating benefit in tendinopathy.

Serrapeptase is widely sold and has a long history of use in Europe and Asia, but there are no controlled trials supporting it for tennis elbow. If you use it, use it knowing that.

**Typical dosage.** Bromelain 500 to 1,000 mg (2,000 to 4,000 GDU) daily between meals; serrapeptase 60,000 to 120,000 SPU daily on an empty stomach **Caution.** May interact with blood thinners. Take away from meals for a systemic rather than digestive effect.

## 6.11 Hyaluronic Acid

**What it does.** Hyaluronic acid is a genuine component of the extracellular matrix around tendons, contributing to gliding and lubrication.

**Evidence, stated honestly.** A literature review of hyaluronic acid in tendon lesions found encouraging preclinical results, in the laboratory and in animals, showing that HA improves tendon gliding, reduces adhesions, improves architectural organization, and limits inflammation. The clinical results were described as limited but encouraging short-term effects on pain and function, and the authors concluded that controlled randomized studies are still needed<sup>21</sup>.

Two things follow. First, that evidence concerns hyaluronic acid **injected** by a clinician, not an oral supplement. Second, nothing shows that oral hyaluronic acid capsules reach an injured tendon or do it any good. This guide does not recommend oral hyaluronic acid for tennis elbow. It is listed here so that you recognize the marketing claim when you meet it and know exactly how thin the support is.

# 7. Targeted Exercises for Tennis Elbow Relief

---

**Critical rule.** Exercises for tennis elbow should be performed slowly and with control. Mild discomfort (2 to 3 out of 10) during exercise is acceptable, but sharp pain, or increased pain lasting more than 24 hours after exercise, means you need to reduce intensity or volume.

## 7.1 Isometric Wrist Extension

**Best for.** Early-stage rehabilitation when even movement causes pain

### How to perform

1. Sit with your forearm supported on a table, wrist hanging over the edge, palm facing down
2. Place your unaffected hand on top of the back of your affected hand
3. Try to raise that hand while resisting with your other hand
4. Hold the contraction for 30 to 45 seconds
5. Perform 3 to 5 repetitions, 2 to 3 times daily

**Why it works.** Isometric contractions load the tendon without joint movement, which many people tolerate when movement hurts. The best-known study of this effect was performed in athletes with **patellar tendinopathy**, where a single bout of isometric contractions produced immediate and sustained pain reduction and reduced cortical inhibition of the quadriceps<sup>22</sup>. That study was in the knee, not the elbow. The principle has been widely adopted in upper-limb rehabilitation, but the direct elbow evidence is weaker than the popularity of the technique suggests.

## 7.2 Eccentric Wrist Extension

**Best for.** The cornerstone of tennis elbow rehabilitation

### How to perform

1. Sit with your forearm supported, palm facing down, holding a light weight (1 to 3 lbs) or a resistance band
2. Use your unaffected hand to lift the weight up (concentric phase)
3. Slowly lower the weight over a count of 3 to 4 seconds (eccentric phase)
4. Perform 3 sets of 10 to 15 repetitions
5. Progress the weight gradually as tolerated

**Why it works.** Eccentric loading, lengthening the muscle-tendon unit under tension, stimulates collagen remodeling and promotes proper fiber alignment. The landmark trial establishing heavy slow eccentric loading was conducted in **Achilles tendinopathy**, where the eccentric group returned to running while the control group did not<sup>23</sup>. That is Achilles evidence extrapolated to the elbow, and this guide states that plainly rather than pretending otherwise.

For the elbow itself, a systematic review of twelve studies of eccentric exercise in lateral epicondylitis found that every group whose program included eccentric exercise reported decreased pain and improved function and grip strength from baseline,

with seven studies reporting improvements for treatment packages that included eccentric work<sup>24</sup>. Most of those programs combined eccentrics with other treatments, so eccentric loading is best described as a well-supported component of care rather than a proven stand-alone cure.

**Progression.** Start with 1 lb and increase by 0.5 to 1 lb weekly as long as pain remains acceptable.

### 7.3 Tyler Twist (FlexBar Exercise)

**Best for.** Moderate to advanced rehabilitation, and one of the few tennis elbow exercises with a dedicated trial

#### How to perform

1. Hold a flexible rubber bar vertically in front of you
2. Grip the top of the bar with the affected arm, wrist extended
3. Grip the bottom with the unaffected arm, wrist flexed
4. Slowly rotate the bar by flexing the unaffected wrist while resisting with the affected wrist
5. Hold the fully twisted position for 2 seconds
6. Slowly return to the starting position
7. Perform 3 sets of 10 to 15 repetitions

**Why it works.** The movement isolates an eccentric load on the wrist extensors. In a prospective randomized trial, adding this isolated wrist extensor eccentric exercise to standard physical therapy for chronic lateral epicondylitis produced greater improvement than standard therapy alone<sup>25</sup>. This is one of the more directly applicable pieces of evidence in the whole guide.

### 7.4 Wrist Flexor Stretching

**Best for.** Reducing tension in the opposing muscle group and improving forearm flexibility

#### How to perform

1. Extend your affected arm straight out with the palm facing up
2. Use your other hand to gently bend the wrist down, palm toward the floor
3. You should feel a stretch along the flexor side of the forearm
4. Hold for 30 seconds, repeat 3 times
5. Do this 2 to 3 times daily

**Why it works.** This is general rehabilitation practice rather than a specifically studied intervention. Stretching the flexors is comfortable for most people, improves range of motion, and is low risk.

### 7.5 Forearm Pronation and Supination

**Best for.** Restoring full forearm rotation

### How to perform

1. Sit with your elbow bent at 90 degrees, tucked at your side
2. Hold a light hammer or small dumbbell vertically
3. Slowly rotate your forearm so the palm faces up (supination), then down (pronation)
4. Perform 2 to 3 sets of 10 to 15 repetitions
5. Keep the movement slow and controlled

**Why it works.** The forearm extensors are active during rotation as well as wrist extension. Restoring comfortable, controlled rotation is a normal part of returning to work and sport. This is standard practice, not a separately proven treatment.

## 7.6 Scapular Stabilization

**Best for.** Addressing the wider kinetic chain

### Scapular retraction

1. Sit or stand with good posture
2. Squeeze your shoulder blades together and down, as if tucking them into your back pockets
3. Hold for 5 seconds, repeat 10 times

### Wall angels

1. Stand with your back against a wall, arms in a “W” position
2. Slowly slide your arms up into a “Y” position, keeping contact with the wall
3. Return to “W”
4. Perform 8 to 10 repetitions

**Why it is included, and what the evidence really shows.** A controlled study comparing people with lateral epicondylalgia to healthy controls found that the involved side had significantly lower middle trapezius, serratus anterior, and lower trapezius strength, lower endurance, and reduced serratus anterior thickness change. The authors were careful to note that some of the differences between the involved and uninvolved arms were small and of potentially limited clinical relevance<sup>26</sup>.

Read that precisely. It shows an **association** between scapular muscle performance and lateral epicondylalgia. It does not show that scapular exercises improve tennis elbow outcomes. No trial has demonstrated that. These exercises are included here as sensible general conditioning that costs nothing, carries essentially no risk, and addresses a measurable deficit, not as an evidence-backed treatment for elbow pain.

## 7.7 Grip Strengthening

**Best for.** Restoring functional grip strength once pain has decreased

### How to perform

1. Squeeze a soft stress ball or therapy putty

2. Hold for 5 seconds, release
3. Perform 10 to 15 repetitions
4. Progress to firmer resistance as tolerated

**Why it works.** Pain-free grip strength is the standard functional measure used to track recovery in the research<sup>5</sup>. Rebuilding it is both the goal and the yardstick. Add this late, not early.

# 8. Physical Therapies and What the Evidence Says

---

## 8.1 Physical Therapy

Physical therapy is the usual first-line treatment for tennis elbow. A skilled therapist will assess your specific movement patterns and biomechanics, prescribe a progressive loading program, and use manual therapy to improve tissue mobility. They will address contributing factors including posture and ergonomics, educate you on activity modification and technique, and progress you through a structured return-to-activity program.

**Evidence.** A clinical review of physiotherapy management of lateral epicondylalgia summarizes exercise and manual therapy as the mainstays of conservative care and pain-free grip strength as the key outcome measure<sup>5</sup>. A systematic review of 58 randomized controlled trials of non-surgical treatments concluded that physical therapy and exercise-based approaches are appropriate first-line care, while noting how variable the trial quality is across this literature<sup>27</sup>.

For manual therapy specifically, a systematic review and meta-analysis of joint mobilization found a modest pooled effect for mobilization with movement on pain ratings, with a mean effect size of 0.43 (95% CI 0.15 to 0.71)<sup>28</sup>. Modest and real is the accurate description, not dramatic.

The strongest exercise evidence is covered in Chapter 7: eccentric loading<sup>24</sup> and the wrist extensor eccentric program tested by Tyler and colleagues<sup>25</sup>.

## 8.2 Extracorporeal Shockwave Therapy (ESWT)

ESWT delivers acoustic shockwaves to the affected tendon.

**Evidence, and it is genuinely mixed.** A large randomized trial of 114 patients with chronic lateral epicondylitis found ESWT without local anesthesia significantly better than sham on pain and function<sup>29</sup>. A double-blind randomized controlled trial published three years earlier found no benefit over placebo<sup>30</sup>. A systematic review and meta-analysis pooling 13 randomized trials and 1,035 patients found better pain scores and grip strength with ESWT, while stating explicitly that the limited quality and quantity of the included studies means more high-quality trials are needed<sup>31</sup>.

**What to expect.** Typically 3 to 5 sessions spaced 1 to 2 weeks apart. Mild discomfort during treatment is normal. Improvement may take 4 to 8 weeks to appear.

## 8.3 Low-Level Laser Therapy (LLLT) / Photobiomodulation

LLLT uses specific wavelengths of light with the aim of stimulating cellular activity in the tissue.

**Evidence.** A systematic review and meta-analysis specific to lateral elbow tendinopathy pooled 13 randomized placebo-controlled trials covering 730 patients and reported benefit, while also detecting publication bias in a negative direction and noting that many of the included patients had already failed other treatments<sup>32</sup>. A broader systematic review across tendinopathies found conflicting results: of 25 controlled trials, 12 showed positive effects and 13 were inconclusive or negative. The authors' key observation was that the positive trials clustered around a particular dosage window, suggesting that dose and technique matter a great deal<sup>33</sup>.

**Practical reading.** LLLT may help if it is delivered at the right dose by someone who knows the protocol. It is not a reliable stand-alone treatment.

**What to expect.** Painless treatment, 10 to 15 minutes per session, 2 to 3 times per week for 4 to 6 weeks.

## 8.4 Acupuncture and Dry Needling

Acupuncture involves inserting thin needles at specific points. Dry needling targets myofascial trigger points in the forearm muscles.

**Evidence.** A systematic review of acupuncture for lateral epicondyle pain found that all included studies suggested a benefit, with five of six showing acupuncture superior to a control treatment, and concluded that there is strong evidence for **short-term** relief<sup>34</sup>. The review did not establish long-term benefit, and short-term is the honest claim.

**What to expect.** 6 to 10 sessions, typically weekly.

## 8.5 Massage Therapy and Myofascial Release

Therapeutic massage addresses muscle tension and tissue restriction in the forearm extensors.

**Evidence.** This is the weakest section in the chapter, and it should be. A pilot randomized study of augmented soft tissue mobilization versus natural history in lateral epicondylitis found that both groups improved, and the study was explicitly designed and reported as a feasibility pilot, too small to detect a difference between the treatments. The authors calculated that a proper trial would need 116 participants and called for one<sup>35</sup>. That trial has not been done.

Massage is comfortable, low risk, and may help you tolerate your loading program. There is no good evidence that it changes the course of the condition.

**Focus areas.** Forearm extensor muscles, trigger points in the forearm, scapular and shoulder musculature, and neck muscles if referred pain is a factor.

## 8.6 Bracing and Orthotics

**Counterforce bracing.** A strap worn just below the elbow, intended to reduce tension at the ECRB tendon origin.

**Evidence.** A prospective, randomized, double-blinded, **placebo-controlled** trial compared a counterforce brace against a placebo brace in acute tennis elbow, following patients out to a mean of three years. Both braces improved pain and function. The counterforce brace produced significantly greater reduction in pain frequency and severity in the short term (2 to 12 weeks) and better overall elbow function at 26 weeks compared with the placebo brace<sup>36</sup>. This is a properly controlled trial with a modest, real, mostly short-term benefit. That is a fair reason to use a brace during aggravating activities and not a reason to expect it to cure anything.

**Wrist splinting.** Immobilizing the wrist (not the elbow) in a neutral position reduces strain on the extensor tendons during provoking tasks.

**What to expect.** Braces are most helpful during aggravating activities. They should not be worn around the clock, since prolonged immobilization leads to stiffness and weakness.

## 8.7 Platelet-Rich Plasma (PRP) Injections

PRP involves drawing your blood, concentrating the platelets, and injecting the concentrate into the affected tendon.

**Evidence, and this is a genuinely contested area.** A double-blind randomized controlled trial comparing PRP with corticosteroid injection found PRP better at one year<sup>37</sup>, and the same cohort followed to two years continued to favor PRP, with success defined as at least a 25% reduction in pain or disability score without reintervention<sup>38</sup>.

Against that, a randomized, double-blind, **placebo-controlled** trial comparing PRP, glucocorticoid, and saline injection found no significant difference between PRP and saline<sup>39</sup>. The distinction matters: the first two trials compared PRP against a steroid injection, which we know does poorly in the long run. The third compared PRP against salt water, and PRP did not win.

**Considerations.** PRP is generally considered only when conservative care has failed for 6 months or more. It is not first-line treatment, insurance usually does not cover it, and the evidence does not currently support paying for it as an early option.

## 8.8 Corticosteroid Injections

Corticosteroids are potent anti-inflammatory agents injected into the painful area.

**Evidence.** A systematic review of randomized controlled trials of injection therapies for tendinopathy found that despite the effectiveness of corticosteroid injections in the short term, the long-term picture is unfavorable for lateral epicondylalgia, and that non-corticosteroid injections may be preferable for longer-term treatment. The authors also cautioned that findings should not be generalized across different tendinopathy sites<sup>40</sup>. Note that this is a systematic review, not a single study, though it is often described as one.

The individual randomized trials tell the same story from the patient's side: excellent relief at 6 weeks, the worst outcomes of any group at 52 weeks, and high recurrence<sup>6,7</sup>.

**The verdict.** A corticosteroid injection may offer meaningful short-term relief for severe pain, and there are situations where that is the right call. It should be used cautiously, not repeatedly, and not as a substitute for a loading program. Discuss the 6-week versus 12-month trade-off explicitly with your provider before agreeing to one.

## 8.9 Surgery

Surgery for tennis elbow involves releasing or debriding the degenerated tendon tissue. It is considered only after 6 to 12 months of genuinely failed conservative treatment.

**Evidence.** A systematic review of surgical treatment for lateral epicondylitis found the literature too thin and too heterogeneous to support a meta-analysis. No technique appeared superior by any measure, and the authors concluded that until randomized controlled trials are performed it is reasonable to defer to the individual surgeon's experience<sup>41</sup>.

**A caution worth taking seriously.** Orthopedic surgery has a documented history of procedures that looked highly successful in uncontrolled series and then failed when tested against a sham operation. The clearest example is arthroscopic surgery for knee osteoarthritis: in a controlled trial where patients were randomly assigned to debridement, lavage, or a placebo procedure, outcomes after the real operations were no better than after the placebo<sup>42</sup>. That trial was about the knee, not the elbow, and it does not prove anything about elbow surgery. It is a general lesson about why "most of my patients got better after the operation" is not the same as "the operation works," and it is a fair question to raise with any surgeon proposing an elective tendon procedure.

**Considerations.** Surgery is rarely necessary. Exhaust conservative options first.

# 9. The Anti-Inflammatory Diet for Tendon Recovery

---

While tennis elbow is a degenerative tendinopathy rather than a true inflammatory condition, nutrition supports tissue repair and collagen synthesis in general. The recommendations in this chapter are ordinary good nutrition applied to a healing tissue. They are not presented as a treatment for tennis elbow, and no diet has been tested for this condition.

## 9.1 Foods to Emphasize

**Collagen-supporting foods.** Bone broth (rich in glycine and proline), chicken and fish skin, egg whites, citrus fruits and berries for vitamin C

**Omega-3 rich foods.** Wild-caught salmon, sardines, mackerel, anchovies, walnuts, flaxseeds, chia seeds. A reasonable goal is 2 to 3 servings of fatty fish per week.

**Colorful vegetables and fruits.** Leafy greens (spinach, kale, Swiss chard), berries, cruciferous vegetables (broccoli, cauliflower), orange and yellow vegetables (carrots, sweet potatoes). Aim for 8 to 10 servings of vegetables and fruits daily.

**Quality protein.** Grass-fed meats, pasture-raised poultry, wild fish, eggs, legumes if tolerated. The meta-analysis and meta-regression across 49 studies and 1,863 participants found that the benefit of added protein plateaus at roughly 1.6 grams per kilogram of body weight daily, which is about 0.7 grams per pound, or around 130 grams for a 180 pound adult.<sup>45</sup> That is the number worth hitting, and eating past it has not been shown to add anything. Most people rehabbing an injury are well under it, which makes this the cheapest correction in this chapter.

**Anti-inflammatory spices.** Turmeric (with black pepper), ginger, garlic, cinnamon

**Healthy fats.** Extra virgin olive oil, avocados, nuts and seeds

**Zinc-rich foods.** Oysters, beef, pumpkin seeds, lentils

## 9.2 Foods to Minimize

**Pro-inflammatory fats.** Trans fats (partially hydrogenated oils), excessive omega-6 vegetable oils (corn, soybean, safflower), deep-fried foods

**Refined carbohydrates and sugars.** White bread, pastries, sugary cereals, soda and sweetened beverages, candy and desserts. Beyond the general metabolic effects, high sugar intake promotes advanced glycation end-products (AGEs), which cross-link collagen.

**Processed foods.** Deli meats, hot dogs, bacon, fast food, packaged snacks

**Excessive alcohol.** Depletes B-vitamins and zinc, impairs protein synthesis, and interferes with sleep, which matters for repair

**AGE-rich preparations.** Heavily charred or high-temperature-grilled meats. AGEs cross-link collagen fibers, making connective tissue stiffer.

## 9.3 Hydration

Adequate hydration supports nutrient delivery and waste removal.

**A simple target.** Half your body weight in pounds, taken as ounces of water daily. For a 160-pound person that is about 80 ounces, roughly 10 cups.

**Tips.** Start the day with 16 to 20 oz of water, drink before meals, limit caffeine and alcohol, add electrolytes if you sweat heavily.

## 9.4 Sample Anti-Inflammatory Day

**Breakfast.** A smoothie with spinach, blueberries, collagen powder, flaxseed, ginger, almond butter, and unsweetened almond milk. Or oatmeal with walnuts, chia seeds, cinnamon, berries, and a scoop of collagen.

**Mid-morning snack.** Apple with almond butter, or a handful of pumpkin seeds

**Lunch.** A large salad with mixed greens, grilled salmon, avocado, cherry tomatoes, olive oil and lemon dressing. Side of bone broth.

**Afternoon snack.** Carrot and celery sticks with hummus, or hard-boiled eggs

**Dinner.** Grass-fed beef or chicken stew with a bone broth base, turmeric, and garlic. Steamed broccoli and sweet potato. Side salad with olive oil.

**Evening (optional).** Ginger or turmeric tea, and a small square of dark chocolate at 70% cacao or higher

# 10. Lifestyle Modifications and Pain Management

---

## 10.1 Activity Modification

This is the single most important change you can make, and it is supported by the strongest risk-factor evidence in the whole guide. Force, repetition, and awkward wrist and forearm posture are the documented drivers of this condition<sup>2,3</sup>. Reducing them is not a soft recommendation.

**Workplace ergonomics.** For keyboard and mouse use, choose a mouse that keeps your wrist in a neutral position. A vertical mouse or trackball helps many people, and positioning the keyboard so your elbows are near 90 degrees and your wrists are straight also helps. Take micro-breaks: every 20 to 30 minutes, stop typing and mousing for 30 seconds and stretch your wrists and forearms. Voice-to-text software can reduce repetitive typing where practical, and tool modification, using tools with larger grips, padded handles, or spring-assisted mechanisms, reduces grip force.

**Sports modification.** Tennis players should check racquet grip size, since too small a grip increases strain, consider a more flexible, lighter racquet, and have a professional assess their backhand technique. Weightlifters should avoid the exercises that aggravate symptoms, such as wrist curls, reverse wrist curls, and heavy gripping, and lifting straps can reduce grip demand. All athletes should return gradually, with a common guideline of increasing volume or intensity by no more than about 10% per week.

**Household activities.** Use two hands to lift heavy pots and pans, use a jar opener to reduce grip force, carry grocery bags on your forearm rather than gripping the handles, and use a backpack instead of a briefcase.

## 10.2 Sleep Optimization

Helpful strategies include maintaining a consistent sleep schedule, keeping your room cool, around 65 to 68 degrees F (18 to 20 C), and limiting screens for an hour before bed. Avoid caffeine after 2 PM, and sleep with the affected arm supported on a pillow to reduce strain.

**Evidence.** A review of the relationship between sleep and pain concluded that the association runs in both directions, and that sleep disturbance appears to be at least as strong a predictor of subsequent pain as pain is of subsequent sleep disturbance<sup>43</sup>. Sleep is not a nice-to-have in a recovery plan.

## 10.3 Stress Management

Chronic stress is a plausible contributor to slower recovery, and in the occupational data on tennis elbow, low workplace social support was an independent risk factor among women<sup>2</sup>.

**Techniques worth trying** include diaphragmatic breathing for 5 minutes twice daily, progressive muscle relaxation, meaning systematically tensing and releasing muscle groups, and gentle yoga or tai chi. Mindfulness meditation has better evidence behind it than most: a randomized clinical trial in 342 adults with **chronic low back pain** found that mindfulness-based stress reduction improved back pain and functional limitation more than usual care at 26 weeks, and performed comparably to cognitive behavioral therapy<sup>44</sup>. That trial studied low back pain, not tennis elbow. It supports mindfulness as a general chronic pain tool, not as an elbow treatment.

## 10.4 Smoking Cessation

Smoking reduces peripheral blood flow and impairs oxygen delivery to tissue, and tendon tissue already has a limited blood supply. Stopping smoking is sound general advice for anyone healing a musculoskeletal injury and for many other reasons besides.

You may see a specific figure quoted for smoking and tendon surgery outcomes. I am not going to give you one, because I could not verify a source for it that I would be willing to stand behind. The general point above rests on physiology rather than on a trial, and that is the honest strength of it.

## 10.5 Weight Management

Maintaining a healthy weight supports overall metabolic health. The link to tennis elbow specifically is weaker than for weight-bearing joints, so treat this as general health advice rather than targeted treatment.

# 11. The In-Clinic Pre-Treatment Protocol

---

Understand what this is for, because the distinction matters.

None of this rebuilds tissue. The loading and strengthening work in the previous chapters does that. What this protocol does is reduce the muscle tone, stiffness and tissue sensitivity that make the strength work harder to tolerate and less pleasant to do, particularly in the early weeks when things are at their most irritable.

It is preparation, not treatment. I use it in the clinic before the working part of a session, and I give it to patients to use at home before theirs. Patients who do it tend to tolerate the loading better and stick with it longer, and sticking with it is the actual mechanism by which recovery happens.

If you have to choose between doing this and doing your strength work, do your strength work.

## **Percussion massager.**

- Triceps, round ball or dampener, level 2 to 3, circles, 60 to 90 seconds. Avoid the elbow joint.
- Biceps and brachialis, round ball, level 2, 45 to 60 seconds. Avoid the bicipital groove.
- Forearm flexor mass, dampener or fork, level 1 to 2, linear strokes, 45 to 60 seconds. Light pressure near the medial epicondyle.
- Forearm extensor mass, dampener, level 1 to 2, 45 to 60 seconds.
- Brachioradialis, round ball, level 1 or 2, 30 to 45 seconds.

## **Massage ball or vibrating sphere.**

- Lateral epicondyle region, gentle circles, 60 to 90 seconds, over the extensor tendon origin. Monitor for nerve irritation and stay off the bone.
- Medial epicondyle region, light, 60 to 90 seconds.
- Lateral and medial intermuscular septum, gentle circles along the septum, a 60 to 90 second pass.

**Firm trigger point tool**, linear strokes. Triceps, then biceps and anterior compartment, 45 to 60 seconds each. Forearm flexors, forearm extensors and brachioradialis, 30 to 45 seconds each.

**Additional cautions for the elbow.** The ulnar nerve runs superficially at the inside of the elbow, which is the funny bone. No direct pressure on either epicondyle. Stop immediately for tingling or numbness.

## **Safety, and I mean these.**

- Avoid all of this over an acutely injured area in the first 48 to 72 hours.
- Reduce pressure over anything actively inflamed.
- Skip percussion therapy entirely if you have a pacemaker or another implanted electronic device.
- Avoid aggressive pressure over varicose veins or compromised skin.
- Start light. Pain must not exceed 6 out of 10. Back off immediately for sharp or shooting pain.

- Stop and seek advice if you get numbness, tingling, or weakness.
- If you are on a blood thinner, use lighter pressure throughout and check with your prescriber first.

## 12. Red Flags: When to Seek Immediate Care

---

Tennis elbow is rarely an emergency, but certain symptoms warrant prompt medical evaluation.

### **Seek immediate care if you experience**

- Severe swelling, redness, and warmth around the elbow (possible infection)
- Fever with elbow pain
- Sudden, severe pain after a specific injury (possible tendon rupture)
- Inability to extend your wrist or fingers (possible tendon rupture or nerve injury)
- Numbness or tingling in the hand (possible nerve involvement)
- Progressive weakness in the hand or forearm
- A visible deformity or gap in the tendon area

### **Contact your healthcare provider promptly if**

- Symptoms worsen after 2 to 3 weeks of self-care
- Pain is severe enough to prevent sleep despite rest
- You develop new symptoms such as numbness, tingling, or weakness
- Symptoms persist beyond 6 to 8 weeks without improvement
- You have a history of significant trauma to the elbow
- You have rheumatoid arthritis or another inflammatory condition

# 13. Building Your Personal Recovery Plan

---

Use this framework as a starting point for a conversation with your provider, not as a prescription.

## Week 1 to 2: Calm and Protect (Acute Phase)

If you choose to use supplements during this phase, collagen plus vitamin C has the most direct mechanistic support for tendon tissue<sup>8</sup>. For movement, rest from aggravating activities and stick to gentle isometric holds and wrist flexor stretches. Therapies at this stage include icing after activity for 15 to 20 minutes, considering a counterforce brace during provoking work tasks<sup>36</sup>, and arranging an initial physical therapy evaluation. On the diet side, begin anti-inflammatory eating and increase hydration, and for lifestyle, modify ergonomics, prioritize sleep, and reduce stress load.

## Week 3 to 6: Rebuild (Subacute Phase)

Reassess supplements during this phase, adding or dropping them based on how you actually feel and what your provider advises. Movement progresses to eccentric wrist extension, continued isometrics, and added forearm rotation work. Therapies include regular physical therapy, considering acupuncture for short-term pain relief<sup>34</sup>, and adding scapular conditioning. Diet moves to the full anti-inflammatory pattern, and lifestyle means implementing the activity modifications properly rather than partially.

## Week 7 to 12: Strengthen and Restore

Movement progresses to the FlexBar Tyler Twist<sup>25</sup>, with grip strengthening added, resistance increased gradually, and a return to modified activity. Therapies at this stage are maintenance physical therapy visits, and considering shockwave therapy if progress has stalled<sup>29,31</sup>. Diet becomes anti-inflammatory eating as a habit rather than a protocol, and lifestyle means a gradual return to sport and full work duties.

## Beyond 12 Weeks: Maintenance and Prevention

Continue eccentric exercises 2 to 3 times per week and maintain the ergonomic changes, since this is where recurrence is prevented. Warm up before demanding activities and address flare-ups early with your established toolkit. Consider injection therapy or surgery only if conservative care has genuinely failed over 6 to 12 months, and only after discussing the long-term trade-offs<sup>40,41</sup>.

# Final Thoughts

---

Tennis elbow can be frustratingly persistent. The encouraging news from the randomized trials is that most people improve substantially within a year, and that the improvement happens with conservative care<sup>6,7</sup>. Tendons remodel in response to controlled, progressive loading, and that loading, combined with genuinely reducing the forces that caused the problem, is the core of recovery.

The key is patience and consistency. Tendons are slow to respond because they have a limited blood supply. Pushing too hard too soon often leads to setbacks, and complete rest leads to deconditioning. The working zone is gradual, progressive loading that challenges the tissue without overwhelming it.

Healing is rarely linear. You may have good weeks and frustrating weeks. The goal is steady progress over months, not days. Stay consistent with your exercises and nutrition, and work closely with your healthcare team.

No book can promise you an outcome. What this one can do is tell you which strategies have real evidence behind them, which are plausible but unproven, and which are being sold to you on a claim that does not hold up.

# References

---

Every reference below was individually retrieved and verified in PubMed. PMIDs are given so you can check any of them yourself.

1. Nirschl RP. Elbow tendinosis/tennis elbow. *Clin Sports Med.* 1992;11(4):851-870. PMID: 1423702
2. Haahr JP, Andersen JH. Physical and psychosocial risk factors for lateral epicondylitis: a population based case-referent study. *Occup Environ Med.* 2003;60(5):322-329. PMID: 12709516
3. Shiri R, Viikari-Juntura E. Lateral and medial epicondylitis: role of occupational factors. *Best Pract Res Clin Rheumatol.* 2011;25(1):43-57. PMID: 21663849
4. Walker-Bone K, Palmer KT, Reading I, Coggon D, Cooper C. Prevalence and impact of musculoskeletal disorders of the upper limb in the general population. *Arthritis Rheum.* 2004;51(4):642-651. PMID: 15334439
5. Bisset LM, Vicenzino B. Physiotherapy management of lateral epicondylalgia. *J Physiother.* 2015;61(4):174-181. PMID: 26361816
6. Smidt N, van der Windt DA, Assendelft WJ, Devillé WL, Korthals-de Bos IB, Bouter LM. Corticosteroid injections, physiotherapy, or a wait-and-see policy for lateral epicondylitis: a randomised controlled trial. *Lancet.* 2002;359(9307):657-662. PMID: 11879861
7. Bisset L, Beller E, Jull G, Brooks P, Darnell R, Vicenzino B. Mobilisation with movement and exercise, corticosteroid injection, or wait and see for tennis elbow: randomised trial. *BMJ.* 2006;333(7575):939. PMID: 17012266
8. Shaw G, Lee-Barthel A, Ross ML, Wang B, Baar K. Vitamin C-enriched gelatin supplementation before intermittent activity augments collagen synthesis. *Am J Clin Nutr.* 2017;105(1):136-143. PMID: 27852613
9. Clark KL, Sebastianelli W, Flechsenhar KR, et al. 24-Week study on the use of collagen hydrolysate as a dietary supplement in athletes with activity-related joint pain. *Curr Med Res Opin.* 2008;24(5):1485-1496. PMID: 18416885
10. García-Coronado JM, Martínez-Olvera L, Elizondo-Omaña RE, et al. Effect of collagen supplementation on osteoarthritis symptoms: a meta-analysis of randomized placebo-controlled trials. *Int Orthop.* 2019;43(3):531-538. PMID: 30368550
11. DePhillipo NN, Aman ZS, Kennedy MI, Begley JP, Moatshe G, LaPrade RF. Efficacy of vitamin C supplementation on collagen synthesis and oxidative stress after musculoskeletal injuries: a systematic review. *Orthop J Sports Med.* 2018;6(10):2325967118804544. PMID: 30386805
12. Calder PC. Omega-3 fatty acids and inflammatory processes: from molecules to man. *Biochem Soc Trans.* 2017;45(5):1105-1115. PMID: 28900017
13. Daily JW, Yang M, Park S. Efficacy of turmeric extracts and curcumin for alleviating the symptoms of joint arthritis: a systematic review and meta-analysis of randomized clinical trials. *J Med Food.* 2016;19(8):717-729. PMID: 27533649
14. Plotnikoff GA, Quigley JM. Prevalence of severe hypovitaminosis D in patients with persistent, nonspecific musculoskeletal pain. *Mayo Clin Proc.* 2003;78(12):1463-1470. PMID: 14661675
15. Park R, Ho AM, Pickering G, Arendt-Nielsen L, Mohiuddin M, Gilron I. Efficacy and safety of magnesium for the management of chronic pain in adults: a systematic review. *Anesth Analg.* 2020;131(3):764-775. PMID: 32049671
16. Kogan S, Sood A, Garnick MS. Zinc and wound healing: a review of zinc physiology and clinical applications. *Wounds.* 2017;29(4):102-106. PMID: 28448263
17. Yu G, Xiang W, Zhang T, Zeng L, Yang K, Li J. Effectiveness of Boswellia and Boswellia extract for osteoarthritis patients: a systematic review and meta-analysis. *BMC Complement Med Ther.* 2020;20(1):225. PMID: 32680575

18. Kim LS, Axelrod LJ, Howard P, Buratovich N, Waters RF. Efficacy of methylsulfonylmethane (MSM) in osteoarthritis pain of the knee: a pilot clinical trial. *Osteoarthritis Cartilage*. 2006;14(3):286-294. PMID: 16309928
19. Lubis AMT, Siagian C, Wonggokusuma E, Marsetyo AF, Setyohadi B. Comparison of glucosamine-chondroitin sulfate with and without methylsulfonylmethane in grade I-II knee osteoarthritis: a double blind randomized controlled trial. *Acta Med Indones*. 2017;49(2):105-111. PMID: 28790224
20. Pavan R, Jain S, Shraddha, Kumar A. Properties and therapeutic application of bromelain: a review. *Biotechnol Res Int*. 2012;2012:976203. PMID: 23304525
21. Kaux JF, Samson A, Crielaard JM. Hyaluronic acid and tendon lesions. *Muscles Ligaments Tendons J*. 2016;5(4):264-269. PMID: 26958533
22. Rio E, Kidgell D, Purdam C, et al. Isometric exercise induces analgesia and reduces inhibition in patellar tendinopathy. *Br J Sports Med*. 2015;49(19):1277-1283. PMID: 25979840
23. Alfredson H, Pietilä T, Jonsson P, Lorentzon R. Heavy-load eccentric calf muscle training for the treatment of chronic Achilles tendinosis. *Am J Sports Med*. 1998;26(3):360-366. PMID: 9617396
24. Cullinane FL, Boocock MG, Trevelyan FC. Is eccentric exercise an effective treatment for lateral epicondylitis? A systematic review. *Clin Rehabil*. 2014;28(1):3-19. PMID: 23881334
25. Tyler TF, Thomas GC, Nicholas SJ, McHugh MP. Addition of isolated wrist extensor eccentric exercise to standard treatment for chronic lateral epicondylitis: a prospective randomized trial. *J Shoulder Elbow Surg*. 2010;19(6):917-922. PMID: 20579907
26. Day JM, Bush H, Nitz AJ, Uhl TL. Scapular muscle performance in individuals with lateral epicondylalgia. *J Orthop Sports Phys Ther*. 2015;45(5):414-424. PMID: 25579691
27. Sims SE, Miller K, Elfar JC, Hammert WC. Non-surgical treatment of lateral epicondylitis: a systematic review of randomized controlled trials. *Hand (N Y)*. 2014;9(4):419-446. PMID: 25414603
28. Lucado AM, Dale RB, Vincent J, Day JM. Do joint mobilizations assist in the recovery of lateral elbow tendinopathy? A systematic review and meta-analysis. *J Hand Ther*. 2019;32(2):262-276. PMID: 29705077
29. Pettrone FA, McCall BR. Extracorporeal shock wave therapy without local anesthesia for chronic lateral epicondylitis. *J Bone Joint Surg Am*. 2005;87(6):1297-1304. PMID: 15930540
30. Speed CA, Nichols D, Richards C, et al. Extracorporeal shock wave therapy for lateral epicondylitis: a double blind randomised controlled trial. *J Orthop Res*. 2002;20(5):895-898. PMID: 12382950
31. Yao G, Chen J, Duan Y, Chen X. Efficacy of extracorporeal shock wave therapy for lateral epicondylitis: a systematic review and meta-analysis. *Biomed Res Int*. 2020;2020:2064781. PMID: 32309425
32. Bjordal JM, Lopes-Martins RA, Joensen J, et al. A systematic review with procedural assessments and meta-analysis of low level laser therapy in lateral elbow tendinopathy (tennis elbow). *BMC Musculoskelet Disord*. 2008;9:75. PMID: 18510742
33. Tumilty S, Munn J, McDonough S, Hurley DA, Basford JR, Baxter GD. Low level laser treatment of tendinopathy: a systematic review with meta-analysis. *Photomed Laser Surg*. 2010;28(1):3-16. PMID: 19708800
34. Trinh KV, Phillips SD, Ho E, Damsma K. Acupuncture for the alleviation of lateral epicondyle pain: a systematic review. *Rheumatology (Oxford)*. 2004;43(9):1085-1090. PMID: 15213328
35. Blanchette MA, Normand MC. Augmented soft tissue mobilization vs natural history in the treatment of lateral epicondylitis: a pilot study. *J Manipulative Physiol Ther*. 2011;34(2):123-130. PMID: 21334545
36. Krosiak M, Pirapakaran K, Murrell GAC. Counterforce bracing of lateral epicondylitis: a prospective, randomized, double-blinded, placebo-controlled clinical trial. *J Shoulder Elbow Surg*. 2019;28(2):288-295. PMID: 30658774

37. Peerbooms JC, Sluimer J, Bruijn DJ, Gosens T. Positive effect of an autologous platelet concentrate in lateral epicondylitis in a double-blind randomized controlled trial: platelet-rich plasma versus corticosteroid injection with a 1-year follow-up. *Am J Sports Med.* 2010;38(2):255-262. PMID: 20448192
38. Gosens T, Peerbooms JC, van Laar W, den Ouden BL. Ongoing positive effect of platelet-rich plasma versus corticosteroid injection in lateral epicondylitis: a double-blind randomized controlled trial with 2-year follow-up. *Am J Sports Med.* 2011;39(6):1200-1208. PMID: 21422467
39. Krogh TP, Fredberg U, Stengaard-Pedersen K, Christensen R, Jensen P, Ellingsen T. Treatment of lateral epicondylitis with platelet-rich plasma, glucocorticoid, or saline: a randomized, double-blind, placebo-controlled trial. *Am J Sports Med.* 2013;41(3):625-635. PMID: 23328738
40. Coombes BK, Bisset L, Vicenzino B. Efficacy and safety of corticosteroid injections and other injections for management of tendinopathy: a systematic review of randomised controlled trials. *Lancet.* 2010;376(9754):1751-1767. PMID: 20970844
41. Lo MY, Safran MR. Surgical treatment of lateral epicondylitis: a systematic review. *Clin Orthop Relat Res.* 2007;463:98-106. PMID: 17632419
42. Moseley JB, O'Malley K, Petersen NJ, et al. A controlled trial of arthroscopic surgery for osteoarthritis of the knee. *N Engl J Med.* 2002;347(2):81-88. PMID: 12110735
43. Finan PH, Goodin BR, Smith MT. The association of sleep and pain: an update and a path forward. *J Pain.* 2013;14(12):1539-1552. PMID: 24290442
44. Cherkin DC, Sherman KJ, Balderson BH, et al. Effect of mindfulness-based stress reduction vs cognitive behavioral therapy or usual care on back pain and functional limitations in adults with chronic low back pain: a randomized clinical trial. *JAMA.* 2016;315(12):1240-1249. PMID: 27002445
45. Morton RW, Murphy KT, McKellar SR, et al. A systematic review, meta-analysis and meta-regression of the effect of protein supplementation on resistance training-induced gains in muscle mass and strength in healthy adults. *Br J Sports Med.* 2018;52(6):376-384. PMID: 28698222 (Correction: *Br J Sports Med.* 2020;54(19):e7. PMID 32943392. The correction adds a competing interest, not a change to the data: one author served on the advisory board of a sports supplement manufacturer while the paper was written. Corrected, not retracted; the breakpoint used here stands.)
46. Zollinger PE, Tuinebreijer WE, Breederveld RS, Kreis RW. Can vitamin C prevent complex regional pain syndrome in patients with wrist fractures? A randomized, controlled, multicenter dose-response study. *J Bone Joint Surg Am.* 2007;89(7):1424-1431. PMID: 17606778
47. Clifford T, Ventress M, Allerton DM, et al. Collagen peptides supplementation improves function, pain, and physical and mental outcomes in active adults. *J Int Soc Sports Nutr.* 2023;20(1):2243252. PMID: 37551682
48. Maroon JC, Bost JW. Omega-3 fatty acids (fish oil) as an anti-inflammatory: an alternative to nonsteroidal anti-inflammatory drugs for discogenic pain. *Surg Neurol.* 2006;65(4):326-331. PMID: 16531187

# Working with Root Health

---

If your elbow has not responded to loading, or you are being offered an injection and want the one-year evidence before you decide, I offer a free Root Discovery Session at Root Health to talk through your options and what a reasonable next step looks like. You can schedule one at [rootcauseunlocked.com/discovery](https://rootcauseunlocked.com/discovery), or reach the office at (919) 662-0520.

Consultations and ongoing care are both available wherever you are located. Being outside North Carolina is not a barrier to working with me. Tell me where you are when you reach out and I will lay out exactly how we would work together from there.

If you would rather start on your own, the free Root Cause Unlocked tracker has a mode built for musculoskeletal recovery, at [rootcauseunlocked.com/tracker](https://rootcauseunlocked.com/tracker). Eight weeks of it turns “I think it might be helping” into something you and I can both read.

*This guide was compiled from peer-reviewed medical literature and clinical best practices. Every citation has been individually verified against the PubMed record. It is intended to give you accurate information and options to discuss with your healthcare provider. It is not a substitute for individual medical care.*

© 2026 Root Health · For personal educational use only

# A clearer way forward

---

You deserve a clearer way to understand the patterns behind persistent symptoms. This guide explains how neck-related headaches are evaluated and how an individualized recovery plan may support you.

Headaches linked to the neck can be frustrating because the pain is felt in the head while an important contributor may be sitting in the cervical spine, several inches away from where you hurt. That mismatch is why these headaches are so often misread, self-treated for years, or dismissed as stress. This guide will help you understand the pattern, recognize the warning signs that need urgent attention, and take an active part in a practical recovery plan.

This ebook is educational. It does not diagnose a headache disorder or replace individualized care. Seek appropriate medical evaluation for a new, severe, changing, or unexplained headache. Use this guide to prepare questions, track progress, and work with your healthcare team. It is not a substitute for an individualized evaluation.

# How to use this guide

---

The first six chapters give you the full picture: what a cervicogenic headache is, why the neck can produce pain in the head, how it is told apart from migraine and tension-type headache, which symptoms demand urgent evaluation, how a clinician builds diagnostic confidence, and what a complete recovery plan contains. Read those in order, because each one sets up the next.

The later chapters are working chapters. Chiropractic and hands-on care, progressive exercise, nutrition, supplements, and daily lifestyle are all meant to be worked through with your clinician rather than adopted wholesale on your own. The final chapter gives you a flare plan and a way to track progress, and the book closes with a worksheet you can bring to your next visit.

The goal throughout is a small plan you can actually repeat, measure, and adjust, rather than a complicated protocol.

# Chapter 1. Understanding cervicogenic headache

---

The headache is felt in the head, but the contributing disorder is in the neck. That single sentence carries the whole idea, and everything else in this book follows from it.

Cervicogenic headache is classified as a secondary headache, which means the headache is attributed to another disorder rather than being a primary disease of its own. In this case the other disorder is a problem of the cervical spine or the soft tissues of the neck.<sup>1</sup> Neck pain often accompanies it, and most people who have it will tell you their neck feels involved. Neck pain alone, however, does not establish the diagnosis. Plenty of people have sore necks and headaches at the same time without one causing the other.

Certain patterns raise suspicion that the neck is genuinely involved. The pain often begins in the upper neck or the back of the head and travels forward, rather than starting behind the eyes and staying there. The familiar headache can frequently be reproduced or aggravated by specific neck movements or by holding a sustained position, which is a clue that mechanical input from the neck is feeding the symptom. Neck movement is often reduced in a way that relates meaningfully to the headache pattern rather than being an incidental finding. The headache pattern tends to improve as the neck disorder and neck function improve, and that response over time is one of the most useful pieces of evidence you and your clinician will ever collect. Pain is also frequently one-sided, although that feature is not unique to cervicogenic headache and should never be used on its own to make the call.

There is an important distinction to hold onto here. Tight muscles, imperfect posture, and age-related findings on imaging do not automatically prove that your neck is causing your headache. Those findings are extremely common in people with no headache at all. What matters is whether your history, your examination, your response to movement, and your progress over time all fit together into one coherent story.

The diagnosis can be genuinely challenging, and it is worth knowing why. Migraine, tension-type headache, occipital neuralgia, jaw disorders, medication-overuse headache, and several other conditions can produce overlapping symptoms. A person can also carry more than one headache type at the same time, which is more common than most patients expect. Good care does not force every headache into a neck-only explanation, and a clinician who never considers an alternative is not being thorough.

# Chapter 2. How the neck can refer pain to the head

---

Shared nerve pathways help explain why a problem in the cervical spine may be experienced as pain in the head. Sensory information arriving from the upper cervical region and from parts of the head converges within related pathways in the nervous system. Because of that overlap, irritation or dysfunction in cervical structures can be perceived by the brain as pain in the head or face. Your nervous system is not making a mistake so much as working with an anatomical arrangement that blurs the line between neck and head.

Several contributors commonly feed that pattern, and each one changes something specific about how your neck behaves. Joint mobility is one: movement limits or mechanical sensitivity in the cervical joints may reproduce your familiar headache pattern when those segments are loaded or tested. Motor control is another, because poor coordination of the small stabilizing muscles increases the effort required for ordinary neck movements that should feel automatic. Muscle endurance matters in a different way. When capacity is low, the demands of driving, computer work, and other sustained positions become harder to tolerate even though no single moment is strenuous.

The problem is rarely confined to the neck itself. Upper-back and shoulder-blade function play a real role, because limited thoracic movement or reduced scapular capacity pushes more demand onto the cervical spine to accomplish the same tasks. Finally, there is recovery load. Poor sleep, ongoing stress, inactivity, and inconsistent pacing all reduce your tolerance and slow the adaptation that exercise is supposed to produce, which is why a plan built only on hands-on care and stretches often stalls.

Whole-person care needs a clear purpose to be worth your time. Every part of your plan should connect to something concrete, whether an examination finding, a daily activity you want to improve, or a treatment response that can actually be measured. If a recommendation cannot be tied to one of those three, it is fair to ask why it is there.

## Chapter 2b. The mid-back connection, and why treatment often starts below the pain

---

Your mid-back, the thoracic spine, is the twelve vertebrae your ribs attach to, and it is designed to be the most mobile part of your spine. Your neck, by design, is comparatively stable. When the mid-back stiffens, which happens to nearly everyone who works at a desk, the neck ends up doing motion that was never its job. The result is a neck that aches by the end of the day without any single injury to explain it, and a headache that follows.

This is why a clinician may treat your mid-back when your head is what hurts, and why it is not a mistake or a distraction when they do.

**The evidence here is genuinely decent.** A meta-analysis of randomized controlled trials found that treating the thoracic spine in people with neck pain produced immediate improvements in pain, neck mobility, and function compared with sham or alternative treatments.<sup>15</sup> A broader systematic review across upper quadrant problems, meaning neck and shoulder together, is more cautious and grades the evidence carefully.<sup>16</sup> Read together, those two say the principle is sound, the neck evidence is the stronger half, and neither one makes thoracic work a cure on its own. It is one part of a program, and the strengthening in Chapter 6 is still the part that holds the result.

# Chapter 3. Cervicogenic, migraine, or tension type?

---

The correct label matters because the most useful treatment plan differs depending on which pattern you are dealing with. Treating a migraine as a neck problem, or a neck problem as though it were a migraine, wastes months.

A cervicogenic pattern typically shows a mechanical relationship with the neck. Movement is restricted, and appropriate cervical provocation reproduces the familiar pain rather than producing some new and unrelated ache. When that picture holds together, condition-directed care makes sense. Careful cervical assessment, manual care when it is appropriate, and active rehabilitation form the center of the plan.

Migraine looks different. It may involve throbbing pain, sensitivity to physical activity, nausea, sensitivity to light or sound, or aura. When those features dominate, migraine-specific acute treatment, preventive treatment, and lifestyle strategies may be what you actually need, and a neck-focused plan alone will underperform.

Tension-type headache tends to feel pressing or tightening rather than throbbing, is commonly felt on both sides, and may come with tenderness of the muscles around the skull. The distinction carries a practical consequence that is easy to miss: the acute and preventive evidence that supports treatment for tension-type headache cannot automatically be transferred to cervicogenic headache. Different condition, different evidence base.

It is also entirely possible to have more than one type. If your features change noticeably between episodes, or if two recognizable patterns keep showing up, that deserves a name rather than being ignored. In that situation the most useful approach is to track the patterns separately and coordinate one treatment plan that addresses both.

One point deserves its own paragraph, because it causes real confusion. Ibuprofen has clinical evidence for acute episodic tension-type headache.<sup>14</sup> If your headache eases after you take ibuprofen, that tells you the medicine helped that episode. It does not, by itself, prove that systemic inflammation caused the headache, it does not confirm cervicogenic headache, and it does not establish that an anti-inflammatory supplement will produce the same effect. A response to a medicine is information, not a diagnosis.

On medication safety: use over-the-counter and prescribed medicines only as directed. Discuss frequent headache medicine use, side effects, kidney or stomach concerns, bleeding risk, and possible drug interactions with an appropriate clinician or pharmacist. Do not change a prescribed treatment without talking to the clinician who prescribed it.

# Chapter 4. Warning signs and when to seek help

---

A safe headache plan starts by recognizing the patterns that need urgent or specialty evaluation. Most headaches are not dangerous. A small number are, and those are worth being able to identify quickly. The lists in this chapter are deliberately short and scannable so that you can check them in the moment.

## Seek immediate care

Call emergency services for any of the following:

- A sudden headache that is the worst you have ever experienced
- New weakness or numbness
- Trouble speaking
- Fainting
- Seizure
- Severe confusion
- New double vision
- Severe loss of balance
- A new drooping eyelid occurring with an unusual headache
- A severe headache after a significant injury

## Contact a medical clinician promptly

These do not require an emergency room. They should not wait for a routine appointment weeks away, either:

- A new headache that is steadily worsening or is substantially different from your usual pattern
- A headache that is new and begins after age 50
- Headache with fever, stiff neck, unexplained weight loss, immune suppression, or a history of cancer
- A new headache during pregnancy or after delivery
- A painful red eye, a major change in vision, or symptoms triggered by coughing, exertion, or a change in position
- New or unusual head or neck pain accompanied by marked dizziness, coordination problems, or other neurologic symptoms
- Frequent use of acute headache medicine, or a headache burden that keeps increasing

One specific condition deserves a direct mention here. Cervical artery dissection, which is a tear in the wall of an artery in the neck, can begin with headache or neck pain before any more obvious neurologic findings appear. The early symptoms are

not always specific, which is exactly what makes them easy to underestimate. That is why a new or unusual presentation deserves careful assessment rather than a wait-and-see approach.<sup>6</sup> Chapter 7 returns to this topic in the context of hands-on care, because the two subjects are related and both deserve honest handling.

# Chapter 5. How your clinician builds confidence

---

Diagnosis here is a process rather than a single test, built on pattern recognition, physical examination, safety screening, and reassessment over time, and each stage either strengthens the working explanation or forces a rethink.

The history comes first, and it carries more diagnostic weight than most people expect. Your clinician should ask about onset, frequency, duration, location, triggers, easing factors, and associated symptoms, because the shape of the pattern is often more revealing than any single finding. Previous injury, illness, imaging, treatment, medication use, and supplement use all matter, both for diagnosis and for safety. Expect specific questions about migraine features, tension-type features, nerve pain, the jaw, sleep, vision, and balance, since those are the conditions most likely to mimic or accompany a neck-driven headache. The conversation should also cover the practical side of your life, including work, driving, exercise, caregiving, and whatever else the headache has taken from you, because those activities become the targets your plan is measured against.

The examination then tests the working idea. Neck and upper-back movement are assessed, including whether testing reproduces your familiar headache rather than simply producing discomfort. Neurologic and vascular screening are performed when indicated. Neck motor control, endurance, shoulder-blade function, and overall load tolerance are examined, since those are the things a rehabilitation plan will actually change. Depending on what your history suggests, screening of the jaw, shoulder, visual system, or vestibular system may be added.

Imaging has a specific job, and it is a narrower job than most patients assume. Routine imaging is generally not needed for an uncomplicated presentation that has been appropriately evaluated, because the findings it turns up are so often incidental. Imaging becomes genuinely useful when warning signs, trauma, progressive neurologic findings, or an unexpected treatment response changes the clinical question that needs answering.<sup>2</sup>

# Chapter 6. The five-part recovery plan

---

The strongest plan combines condition-directed care with active rehabilitation and whole-person support. Each of the five parts below depends on the ones before it, which is why the order is not arbitrary.

It begins with safety and diagnostic clarity. Before anything else, your clinician confirms that your presentation is appropriate for conservative care and establishes goals that can actually be measured. Second comes chiropractic and manual care, where manipulation, mobilization, specific movement techniques, or soft-tissue work are selected according to your examination findings, your preferences, and your individual risk profile. Third is progressive exercise, which builds neck control, upper-back mobility, shoulder-blade capacity, and general conditioning, and which is what most often carries the long-term result. Fourth are the recovery foundations of sleep, food quality, hydration, daily activity, stress regulation, and work tolerance, all of which determine how much your system can absorb. Fifth, and easy to skip, is reassessment and independence. You keep what is producing meaningful benefit, modify what is not, and taper passive care as your own self-management improves.

It helps to know what progress actually looks like, because it rarely arrives as a single dramatic moment. You may notice fewer headache days or shorter episodes. Average intensity may fall, or you may find yourself reaching for acute medication less often. Driving, work, sleep, exercise, or family participation may become easier before the headache itself changes much. Neck movement usually becomes more comfortable and exercise better tolerated. Perhaps most importantly, you gain confidence managing early symptoms without swinging into an all-or-nothing response every time something flares.

# Chapter 7. Chiropractic and hands-on care

---

Manual care can be genuinely useful, but it should serve a measurable plan rather than exist as an open-ended routine.

Current evidence supports offering spinal manipulation to appropriately screened adults with cervicogenic headache. Mobilization, specific movement techniques, and soft-tissue care may also be used, and manual care combined with exercise is the pairing that appears most often in the research.<sup>2,3,4,5</sup> No form of hands-on care should be presented to you as a cure, and none of it removes the need for the active side of the plan.

Manipulation, when the patient is appropriately selected, may improve pain or movement. What you should expect alongside it is informed consent, safety screening, respect for your preference, and a measurable response that gets checked rather than assumed. Mobilization is a gentler option that does not use a thrust, and it is often chosen when mobility is the target, when symptoms are highly irritable, when you prefer it, or when the goal is simply to prepare you for exercise. With mobilization the things to track are your movement and whether your familiar symptom pattern responds. Soft-tissue care has a narrower role: it may reduce a specific barrier to motion or to exercise tolerance, and it works best as an adjunct rather than as a stand-alone explanation for why you hurt. Education is the fourth piece and the most underrated one, because reducing fear and building confidence with safe movement and loading changes what you are willing to do. Good education connects advice to activities that matter to you rather than to the pursuit of perfect posture.

Hands-on care should earn its place in your plan. If it does not improve a relevant symptom, a movement, a function, or your ability to exercise, the approach should be modified rather than continued automatically because it is on the schedule.

## What you should know about neck manipulation and artery dissection

You are entitled to a straight answer on this, so here is one. Neck manipulation has been associated with cervical artery dissection, but research has not shown that it causes it. A tear in a neck artery usually causes neck pain and headache on its own, so people often seek treatment for pain that is already being caused by a tear in progress. One large study found the same association with primary care visits as with chiropractic visits, which supports that explanation, and a review of the research found no evidence of causation.<sup>11,12</sup> Even so, national stroke guidance advises that you be told about this association before neck manipulation, which is why it appears here.<sup>13</sup>

The practical response to that information is disclosure, not fear. Before any neck manipulation, tell your clinician if you take blood thinners, if you have vascular disease, a connective tissue disorder, or inflammatory arthritis, if you have had recent trauma, if you are pregnant, if you have a history of cancer or a current infection, or if you have developed any new neurologic symptoms. Any one of those may change what is appropriate for you, and some of them will move the plan toward gentler options or toward medical evaluation first. Review the urgent warning signs in Chapter 4 as well, and seek assessment promptly for any new or unusual presentation rather than assuming it is the same headache you have always had.

# Chapter 8. Your progressive exercise plan

---

Start small, use good technique, and progress one variable at a time. Exercise is where most of the durable change in a cervicogenic headache plan comes from, and it is also where most people go wrong by doing too much in the first week.

Before you begin, understand that the examples below are for clinician-guided education, not a personal prescription. Stop and seek assessment for a sudden unusual headache, new weakness or numbness, severe dizziness, double vision, trouble speaking, fainting, or loss of balance. Those are the same signals described in Chapter 4, and they apply during exercise just as they do at rest.

## Stage 1: Control and comfortable motion

The first stage is about restoring calm, controlled movement rather than building strength. Order matters here, so work through these as a short sequence.

1. **Small neck nod.** Perform 5 gentle holds of about 5 seconds each. Make a small yes motion without lifting the head, and keep the jaw relaxed throughout.
2. **Comfortable rotation.** Perform 5 repetitions in each direction. Move through a comfortable range and do not force the end position.
3. **Light isometric.** Perform 3 to 5 holds of about 5 seconds each. Press gently into your own hand without any visible head movement.

## Stage 2: Shoulder blade and upper-back capacity

Once stage one is comfortable, the focus shifts behind and below the neck, because the upper back and shoulder blades carry load the neck would otherwise absorb.

1. **Light band row.** Perform 1 to 2 sets of 8. Draw the elbows back without shrugging the shoulders or flaring the ribs.
2. **Upper-back rotation.** Perform 5 repetitions in each direction. Move smoothly through the thoracic region without forcing the neck.
3. **Thoracic extension.** Perform 1 to 2 sets of 6. Use a comfortable support and keep the movement controlled rather than bouncing into range.

## Stage 3: Endurance and return to activity

The third stage is where capacity is rebuilt and normal life comes back. Build walking, cycling, or another tolerated aerobic activity starting from a duration you can genuinely repeat, not the duration you could manage on your best day. Progress resistance exercise for the neck, upper back, shoulders, and the rest of the body, since general conditioning supports the local work. Practice the specific movement demands of your work, your driving, your recreation, or your sport in manageable steps rather than returning to all of it at once. Throughout this stage, increase only one main variable at a time, whether repetitions, resistance, range, or frequency, never several together, or you will not know what caused a flare.

## Judging the dose

Consistency and recovery matter more than aggressive loading, and reading your own response is a skill worth developing. Mild, short-lived muscle effort is usually acceptable when it settles and the next day is not meaningfully worse. A clear headache increase that lasts is a signal to reduce range, resistance, repetitions, or frequency until the response changes. If the same exercise repeatedly produces a flare, pause it and ask your clinician to reassess either the selection or your technique, because something about that movement is not fitting you right now. Any new neurologic symptom or anything unusual means you stop and seek appropriate assessment rather than pushing through.

There is a simple test for whether your starting dose is right. A useful dose is one you can perform with good form and recover from predictably. A heroic session followed by several inactive days almost always builds less capacity than a smaller dose you can repeat, and it teaches you less about your own tolerance.

# Chapter 9. Food, hydration, and anti-inflammatory eating

---

Nutrition supports recovery, but there is no proven cervicogenic headache diet, and any source that offers you one is overselling. A nutrient-dense eating pattern can support general health, steady energy, muscle function, and recovery from exercise. Those benefits are worth pursuing on their own terms, without claiming that a food plan directly corrects the cervical source of a headache.<sup>10</sup>

A practical foundation is not complicated. Include a protein-rich food at meals according to your own needs and preferences, since protein supports the muscle work your rehabilitation plan is asking for. Emphasize vegetables, fruit, legumes, whole grains, or other fiber-rich foods that you tolerate well. Use foods that provide unsaturated fats, including fish, nuts, seeds, avocado, and olive oil where those fit your situation. If skipped meals consistently aggravate a separate headache pattern you already have, eat on a more regular schedule. As for caffeine and alcohol, track them only when your own history suggests a reproducible relationship, rather than eliminating them on principle and losing the information.

It is worth being clear about what anti-inflammatory eating means in this context. The goal is a sustainable pattern, not a restrictive therapeutic diet, that favors minimally processed foods and supports cardiometabolic and musculoskeletal health over years. If you suspect a specific food trigger, track it, test one change at a time, and avoid prolonged elimination without a clear reason and a plan to reintroduce the food.

Hydration is individual, and rigid universal targets do more harm than good. Fluid needs vary with body size, activity level, climate, pregnancy, kidney and heart health, and the medicines you take. Ask for individualized guidance whenever a medical condition affects your fluid balance.

# Chapter 10. Supplements, including anti-inflammatory options

---

Targeted use is a very different thing from a default supplement stack, and the distinction is where most money and most disappointment live.

Start from the evidence boundary. No oral supplement has been proven to treat cervicogenic headache itself. A supplement may still be reasonable for a documented deficiency, for a separate diagnosed condition, or as a carefully selected and monitored adjunctive goal. What it should not be is a hopeful addition with no stated reason.

Magnesium comes up most often, usually because of low dietary intake, a documented deficiency, or a separate evidence-informed migraine plan. It is not a proven cervicogenic headache treatment, and if it is used, kidney function and medication timing should be reviewed first.<sup>7</sup> Vitamin D has a narrower role here: correcting a documented deficiency. High doses can be harmful, and vitamin D is not a routine headache treatment.<sup>8</sup> Omega-3 may be discussed for reasons of overall food quality, low fish intake, or another appropriate nutritional indication. It has no direct efficacy for cervicogenic headache, and bleeding risk, upcoming procedures, and any history of atrial fibrillation should be part of the conversation before you start.<sup>9</sup>

If you and your clinician do decide to trial an adjunct, there is a safer way to go about it. Name the specific reason you are using it, rather than beginning with a vague detox or inflammation claim that cannot be tested. Screen first for pregnancy, kidney or liver disease, bleeding risk, allergy, planned procedures, and interactions with your medications. Introduce one product at a time, because two at once makes both the benefits and the side effects uninterpretable. Agree in advance on a measurable outcome, a review date, and a stop rule with the clinician supervising the trial. And stop and seek advice for any concerning reaction or meaningful worsening rather than waiting for the review date.

This is also the right place to revisit why ibuprofen evidence is different. Ibuprofen has direct trial evidence for acute episodic tension-type headache. That evidence does not establish equivalent effectiveness for fish oil or for any other supplement, and it does not show that every headache is driven by systemic inflammation.

# Chapter 11. Sleep, stress, work, and lifestyle

---

Daily load and daily recovery change how much your neck and nervous system can tolerate. This chapter is not filler around the real treatment. For many people it decides whether the real treatment gets a chance to work.

Sleep is the foundation. Keeping a consistent sleep and wake schedule improves recovery and makes everything else in the plan more effective. Mention snoring, breathing pauses, or severe daytime sleepiness to your clinician, too, because those may point to a sleep-breathing disorder that deserves its own evaluation.

Work and device use are the next lever. Changing position every 30 to 60 minutes, bringing your work closer to you, and supporting your forearms all reduce sustained load on the neck. Notice that the aim is reducing unnecessary strain, not achieving one perfect posture and holding it in fear.

Stress regulation belongs in the plan too. Slower breathing, relaxation practice, mindfulness, counseling, or whatever strategy you actually prefer can reduce arousal and improve coping. This does not mean your pain is imagined, and no clinician should imply otherwise.

Movement across the week is what builds capacity. Short, repeatable activity sessions do more than occasional long ones, and they break the all-or-nothing cycle that keeps many people stuck. Driving deserves specific attention if it has become difficult. Adjust the seat and mirrors, take movement breaks on longer trips, and gradually practice the rotation the task requires so that a meaningful activity comes back safely.

Two ideas are worth stating plainly. First, posture without fear. There is no single perfect posture that must be held all day. A useful ergonomic strategy reduces unnecessary strain, supports the task in front of you, and makes it easy to change position. Capacity and variation are usually far more helpful targets than rigid correction. Second, stress without blame. Stress can change muscle tension, sleep, attention, coping, and symptom sensitivity, and addressing it is one legitimate part of whole-person care. Acknowledging that does not invalidate your pain and does not reduce the problem to psychology.

# Chapter 12. Flare planning and progress tracking

---

A written plan makes early symptoms easier to manage and helps your care team make better decisions, because you arrive with information instead of an impression.

When a flare begins, start by checking for warning signs or a pattern that is substantially different from your usual one, using Chapter 4 as your reference. If nothing there applies, temporarily reduce the specific activity or exercise variable that exceeded your current tolerance, rather than shutting everything down. Continue comfortable movement instead of complete inactivity, unless a clinician has advised otherwise, since prolonged rest usually costs you more than it gains. Use clinician-approved self-care and medication only as directed. Contact your clinician when a flare is severe, unusual, persistent, or repeatedly triggered by the same activity, because that last pattern in particular means something in the plan needs to change.

Tracking progress does not require an app or a spreadsheet. Once a week, write down six things and compare them to the week before so you can see a trend rather than a snapshot: the number of headache days you had, your average intensity on a 0 to 10 scale, the typical duration of an episode, how much acute medicine you used, which activity was most limited, and how many exercise sessions you actually completed. Those six numbers will tell you and your clinician more in four weeks than memory ever will.

Reassessment matters, and it should be scheduled rather than assumed. A conservative care plan should include a formal review, often near the four-week mark. At that review the job is to continue what is producing meaningful improvement, modify what is only partly helping, taper passive care as your independence grows, and genuinely reconsider the diagnosis when progress does not fit what was expected.<sup>2</sup>

# The In-Clinic Pre-Treatment Protocol

---

Understand what this is for, because the distinction matters.

None of this rebuilds tissue. The loading and strengthening work in the previous chapters does that. What this protocol does is reduce the muscle tone, stiffness and tissue sensitivity that make the strength work harder to tolerate and less pleasant to do, particularly in the early weeks when things are at their most irritable.

It is preparation, not treatment. I use it in the clinic before the working part of a session, and I give it to patients to use at home before theirs. Patients who do it tend to tolerate the loading better and stick with it longer, and sticking with it is the actual mechanism by which recovery happens.

If you have to choose between doing this and doing your strength work, do your strength work.

**Vibrating sphere (Hypersphere Mini)** with vibration on, along the upper trapezius, rolling over tender and tight areas, 2 to 3 minutes. Or a **percussion device (Hypervolt)** with your attachment of choice over the same area for the same length of time.

**Neck Trigger Point Tool**, up and down the neck in a 2 to 3 minute pass, moving the handles side to side and up and down over tight tissue.

**Roller (Vyper or foam roller)** between the shoulder blades, with the buttocks lifted and one hand behind the head, rolling 1 to 2 minutes.

## **Safety, and I mean these.**

- Avoid all of this over an acutely injured area in the first 48 to 72 hours.
- Reduce pressure over anything actively inflamed.
- Skip percussion therapy entirely if you have a pacemaker or another implanted electronic device.
- Avoid aggressive pressure over varicose veins or compromised skin.
- Start light. Pain must not exceed 6 out of 10. Back off immediately for sharp or shooting pain.
- Stop and seek advice if you get numbness, tingling, or weakness.

# Your next-visit worksheet

---

Bring focused observations and questions to the conversation. The most productive visits start with you naming what matters most rather than waiting to be asked.

Write down the activity you most want to return to or tolerate better, and then write down the single change that would matter most to you. Two sentences are enough, and they will shape the rest of the appointment.

Then work through the questions that keep a plan honest. Ask what specific findings make your clinician think the headache is coming from your neck, and ask which other headache types or serious causes have already been considered and ruled out. Find out which one or two exercises matter most right now, since a long list rarely gets done. Ask how the two of you will measure whether hands-on care is actually helping. If a supplement has been recommended, ask for the clear reason behind it. Ask what the review date and the stop rule are for each treatment in the plan. And ask when visit frequency should decrease, because a good plan has a direction and an endpoint built into it.

The bottom line is this. The most defensible whole-person plan keeps cervical assessment, appropriate manual care, and progressive exercise at the center. Nutrition, supplements, sleep, stress regulation, and lifestyle all support recovery when each one is connected to a clear rationale and a monitored outcome, and not before.

# Sources and further reading

---

1. Headache Classification Committee of the International Headache Society (IHS). The International Classification of Headache Disorders, 3rd edition. *Cephalalgia*. 2018;38(1):1 to 211. doi:10.1177/0333102417738202. PMID 29368949.
2. Trager RJ, Daniels CJ, Hawk C, et al. Chiropractic Management of Adults with Cervicogenic or Tension-Type Headaches: Development of a Clinical Practice Guideline. *Journal of Integrative and Complementary Medicine*. 2026;32(4):325 to 344. doi:10.1177/27683605251397769. PMID 41685545.
3. Bini P, Hohenschurz-Schmidt D, Masullo V, Pitt D, Draper-Rodi J. The effectiveness of manual and exercise therapy on headache intensity and frequency among patients with cervicogenic headache: a systematic review and meta-analysis. *Chiropractic and Manual Therapies*. 2022;30(1):49. doi:10.1186/s12998-022-00459-9. PMID 36419164.
4. Satpute K, Bedekar N, Hall T. Mulligan manual therapy added to exercise improves headache frequency, intensity and disability more than exercise alone in people with cervicogenic headache: a randomised trial. *Journal of Physiotherapy*. 2024;70(3):224 to 233. doi:10.1016/j.jphys.2024.06.002. PMID 38902195.
5. Demont A, Lafrance S, Gaska C, Kechichian A, Bourmaud A, Desmeules F. Efficacy of physiotherapy interventions for the management of adults with cervicogenic headache: a systematic review and meta-analyses. *PM&R*. 2023;15(5):613 to 628. doi:10.1002/pmrj.12856. PMID 35596553.
6. Yaghi S, Engelter S, Del Brutto VJ, et al. Treatment and Outcomes of Cervical Artery Dissection in Adults: A Scientific Statement From the American Heart Association. *Stroke*. 2024;55(3):e91 to e106. doi:10.1161/STR.0000000000000457. PMID 38299330.
7. National Institutes of Health Office of Dietary Supplements. Magnesium Fact Sheet for Health Professionals. Updated 2026. <https://ods.od.nih.gov/factsheets/Magnesium-HealthProfessional/>
8. National Institutes of Health Office of Dietary Supplements. Vitamin D Fact Sheet for Health Professionals. Updated 2025. <https://ods.od.nih.gov/factsheets/VitaminD-HealthProfessional/>
9. National Institutes of Health Office of Dietary Supplements. Omega-3 Fatty Acids Fact Sheet for Health Professionals. Updated 2025. <https://ods.od.nih.gov/factsheets/Omega3FattyAcids-HealthProfessional/>
10. U.S. Department of Health and Human Services and U.S. Department of Agriculture. Dietary Guidelines for Americans, 2025 to 2030. 10th edition. 2026. <https://odphp.health.gov/our-work/nutrition-physical-activity/dietary-guidelines/current-dietary-guidelines>
11. Cassidy JD, Boyle E, Cote P, et al. Risk of vertebrobasilar stroke and chiropractic care: results of a population-based case-control and case-crossover study. *Spine (Phila Pa 1976)*. 2008;33(4 Suppl):S176-S183. PMID 18204390. (Erratum: *Spine (Phila Pa 1976)*. 2010;35(5):595. Screened 2026-09-05; corrected, not retracted.)
12. Church EW, Sieg EP, Zalatio O, Hussain NS, Glantz M, Harbaugh RE. Systematic Review and Meta-analysis of Chiropractic Care and Cervical Artery Dissection: No Evidence for Causation. *Cureus*. 2016;8(2):e498. PMID 27014532.
13. Biller J, Sacco RL, Albuquerque FC, et al. Cervical arterial dissections and association with cervical manipulative therapy: a statement for healthcare professionals from the American Heart Association/American Stroke Association. *Stroke*. 2014;45(10):3155-3174. PMID 25104849. (Erratum: *Stroke*. 2016;47(11):e261. PMID 27777353. Screened 2026-09-05; corrected, not retracted.)
14. Derry S, et al. Ibuprofen for acute treatment of episodic tension-type headache in adults. *Cochrane Database of Systematic Reviews*. 2015;2015(7):CD011474. PMID 26230487.
15. Pan Y, Chen J, Liu Y, Chen Z, Liu J. The immediate effects of thoracic spine manipulation in patients with neck pain: a meta-analysis of randomized controlled trials. *Front Med (Lausanne)*. 2026;13:1790614. PMID 41987777.

16. Thoomes EJ, Tilborghs G, Heneghan NR, Falla D, De Graaf M. Effectiveness of Thoracic Spine Manipulation for Upper Quadrant Musculoskeletal Disorders: A Systematic Review. *J Manipulative Physiol Ther.* 2025;48(1-5):422-434. PMID 41196244.

One note on the evidence above. Ibuprofen has randomized and systematic-review evidence for acute episodic tension-type headache.<sup>14</sup> That evidence does not speak directly to cervicogenic headache, nor to oral anti-inflammatory supplements.

# Working with Root Health

---

If your headaches start in your neck, and you would like someone to examine the neck rather than treat the headache in isolation, I offer a free Root Discovery Session at Root Health to talk through what may be driving them and what a reasonable next step looks like. You can schedule one at [rootcauseunlocked.com/discovery](https://rootcauseunlocked.com/discovery), or reach the office at (919) 662-0520.

Consultations and ongoing care are both available wherever you are located. Being outside North Carolina is not a barrier to working with me. Tell me where you are when you reach out and I will lay out exactly how we would work together from there.

If you would rather start on your own, the free Root Cause Unlocked tracker has a mode built for musculoskeletal recovery, at [rootcauseunlocked.com/tracker](https://rootcauseunlocked.com/tracker). Eight weeks of it turns “I think it might be helping” into something you and I can both read.

# Remember

---

This guide provides general education. Your diagnosis, treatment options, exercise plan, medication use, and supplement decisions should be individualized with the appropriate healthcare professionals.