



ROOT CAUSE UNLOCKED

Unlock Your *Neuropathy*

A Holistic Guide to Reclaiming Nerve Health

A guide for anyone living with numbness, tingling, or burning in the hands and feet who has been offered medication for the symptom and never an explanation for the cause.

Dr. Adam Seay, DC

Root Cause Unlocked, Garner, NC · rootcauseunlocked.com · (919) 662-0520

Unlock Your Neuropathy

A Holistic Guide to Reclaiming Nerve Health

BEFORE YOU START

Contents

INTRODUCTION

Stepping Beyond the Symptoms

Chapter 1: The Foundation, Understanding Neuropathy Beyond the Symptoms

Chapter 2: Key 1, Unmasking Nutrient Deficiencies

Chapter 3: Key 2, Addressing Chronic Infections and Their Impact

Chapter 4: Key 3, Activating Large Diameter Afferent Pathways

Chapter 5: Key 4, Comprehensive Testing Beyond the Conventional

Chapter 6: Key 5, Diet and Food Sensitivities for Nerve Repair

Chapter 7: The Movement Plan, Restoring Function and Reducing Pain

Chapter 8: Mindset and Monitoring, the Unseen Influencers

CONCLUSION

Your Path to Reclaiming Nerve Health

Part 4: The 90-Day Battle Plan, Meal Plan and Recipes

Your Next Step: A Free Root Discovery Session

References

Stepping Beyond the Symptoms

The persistent tingling, numbness, burning, and sharp pains of neuropathy can feel like an invisible cage, limiting your life and stealing your comfort. For millions worldwide, peripheral neuropathy isn't just a symptom, it's a chronic challenge that profoundly impacts daily activities, sleep, and overall quality of life. Whether you've been diagnosed with diabetic neuropathy, idiopathic neuropathy, or a specific form like post-herpetic neuralgia, the conventional approach often focuses on managing symptoms with medications that may come with their own set of side effects, without truly addressing the underlying cause.

But what if, instead of simply masking the discomfort, we could uncover the root imbalances contributing to your nerve damage and dysfunction? This ebook is your guide to that path, toward understanding, addressing, and potentially reversing the progression of your neuropathy.

Traditional medicine excels at acute care and crisis intervention, but chronic conditions like neuropathy often require a detective's eye and a comprehensive strategy. We believe that true healing comes from identifying and addressing the why behind your symptoms. Your body is an interconnected system, and nerve health is profoundly influenced by everything from your diet and nutrient status to your gut health, immune system, and even your stress levels.

In this guide, we will explore five fundamental "keys" to unlocking your neuropathy. These keys represent critical areas often overlooked in standard care, yet they hold immense power to influence nerve health and regeneration. The first key is unmasking nutrient deficiencies: discovering the vital vitamins and minerals essential for nerve function that may be missing from your diet or not adequately absorbed. The second is addressing chronic infections and their impact, understanding how hidden or persistent infections can trigger inflammation and autoimmune responses that attack your nerves. The third is activating large diameter afferent pathways, learning how to stimulate specific nerve fibers to improve sensory input, modulate pain, and re-educate your nervous system. The fourth is comprehensive testing that goes beyond the conventional, identifying the advanced diagnostics that can reveal deeper insights into inflammation, metabolic imbalances, circulation issues, and more. And the fifth is diet and food sensitivities, fueling nerve repair by harnessing the power of nutrition to reduce inflammation, stabilize blood sugar, and provide the building blocks for nerve regeneration.

This journey requires an open mind and a commitment to self-exploration. By the end of this ebook, you will be equipped with practical strategies, research-backed insights, and a detailed 90-day plan to embark on your path toward reclaiming nerve health and unlocking a life with less pain and greater vitality. It's time to stop just coping and start truly healing.

Chapter 1: The Foundation, Understanding Neuropathy Beyond the Symptoms

To truly unlock your neuropathy, we must first lay a solid foundation of understanding. Neuropathy is a term that describes damage to the peripheral nervous system, the vast network of nerves outside of your brain and spinal cord, rather than a single disease. These nerves are the communication highways that carry messages between your central nervous system and every other part of your body, including your muscles, skin, and internal organs.

When these nerves are damaged, their ability to transmit signals is compromised, leading to a wide array of symptoms, and the type of symptoms you experience depends on which nerves are affected. Sensory nerves transmit sensations like touch, pain, temperature, and vibration; damage to them often leads to numbness, tingling, burning, shooting pains, or an exaggerated sensitivity to touch, and this is the most common type of neuropathy. Motor nerves control muscle movement, and damage to them can result in muscle weakness, cramps, spasms, loss of balance, or difficulty with fine motor skills. Autonomic nerves control involuntary bodily functions like heart rate, blood pressure, digestion, bladder function, and sweating, so autonomic neuropathy can cause symptoms such as dizziness upon standing, digestive issues, bladder control problems, or abnormal sweating.

Defining Neuropathy: A Deeper Dive into Types

While symptoms can overlap, understanding the different classifications of neuropathy can help pinpoint potential causes. Peripheral neuropathy is the most common form, affecting nerves in the extremities, the feet, legs, hands, and arms; it often presents as a “stocking-glove” pattern, meaning symptoms begin in the toes and fingers and spread upwards. Diabetic neuropathy, the most prevalent cause of neuropathy worldwide, results from prolonged high blood sugar levels that damage nerve fibers; it often manifests as peripheral neuropathy but can also affect autonomic nerves. When no clear cause can be identified, it’s labeled idiopathic neuropathy. However, this simply means the underlying cause hasn’t been found yet, and our holistic approach is particularly vital here, as we aim to uncover those hidden factors. Compressive neuropathy is caused by pressure on a nerve, such as carpal tunnel syndrome or sciatica. Inflammatory neuropathy describes conditions like Guillain-Barré syndrome or chronic inflammatory demyelinating polyneuropathy (CIDP), where the immune system mistakenly attacks nerve tissue. And toxin-induced neuropathy is damage from exposure to heavy metals such as lead or mercury, certain industrial chemicals, or even some medications such as chemotherapy drugs.

The Root Cause Approach: Beyond Symptom Management

The conventional medical model often focuses on symptom management, with pain relievers for pain, anti-depressants for burning sensations, or medications to manage blood sugar in diabetics. While these

approaches can offer temporary relief, they rarely address why the nerves are being damaged in the first place. This is where the root cause approach comes in.

Imagine a leaky roof in your house. You could continually mop up the puddles, which is symptom management, but the leak would persist. To truly solve the problem, you need to find the hole in the roof and patch it: that is the root cause approach. Similarly, with neuropathy, understanding the underlying mechanisms of nerve damage is paramount. These mechanisms often involve chronic inflammation, a persistent low-grade inflammatory state that can damage nerve cells and their protective myelin sheaths; oxidative stress, an imbalance between free radicals and antioxidants that can lead to cellular damage, including nerve cells; poor circulation, since nerves require a steady supply of oxygen and nutrients and compromised blood flow can starve nerve tissue; metabolic imbalances, where high blood sugar, insulin resistance, and imbalanced lipid profiles can directly harm nerves; immune dysfunction, where autoimmune attacks on nerve tissue or chronic activation of the immune system by infections can lead to nerve damage; nutrient deficiencies, since nerves have high metabolic demands and require specific nutrients for their structure, function, and repair; and toxic burden, where exposure to environmental toxins or heavy metals can directly poison nerve cells.

By systematically investigating these areas, we can move beyond the “idiopathic” label and empower you with knowledge and actionable strategies. The upcoming chapters work through each of our five “keys,” providing you with the tools to identify potential root causes in your own body and implement a comprehensive plan for healing and restoring nerve health.

Chapter 2: Key 1, Unmasking Nutrient Deficiencies

Our nerves are not static wires; they are living, dynamic tissues with high metabolic demands. They constantly require a precise array of vitamins, minerals, and other micronutrients to build and maintain their structure, transmit signals efficiently, and repair themselves. When even one of these essential building blocks is in short supply, nerve function can falter, leading to the symptoms of neuropathy.

It's a common misconception that a "standard" diet provides all the necessary nutrients. Soil depletion, modern food processing, and individual genetic variations or medical conditions can all contribute to insidious nutrient deficiencies that silently undermine nerve health. This chapter covers what specific nutrients do for your nerves and how to identify and address common deficiencies linked to neuropathy.

The Vital Role of Nutrients for Nerve Health

Think of your nervous system as a complex electrical circuit, with nutrients acting as the insulation, the conductors, the energy source, and the repair crew. They are involved in forming the myelin sheath, the fatty insulation around nerve fibers that allows for rapid signal transmission; in synthesizing the neurotransmitters that allow nerve cells to communicate; in energy production, supplying the vast amount of ATP, or adenosine triphosphate, needed for nerve function; in antioxidant defense, protecting nerves from damaging oxidative stress; in modulating the body's inflammatory response; and in nerve growth and repair, providing the raw materials for nerve regeneration.

Common Deficiencies Linked to Neuropathy

Several key nutrients are repeatedly implicated in the development or worsening of neuropathy. Addressing these is often a critical first step in a holistic healing plan.

B Vitamins: The Nerve Powerhouses

The B-complex vitamins are a family of water-soluble nutrients that are indispensable for nervous system health, and their deficiency can directly lead to nerve damage.

Vitamin B1, or thiamine, is essential for glucose metabolism and energy production, particularly in nerve cells, which rely heavily on glucose, and it is crucial for the synthesis of neurotransmitters and the maintenance of the myelin sheath. Thiamine deficiency, a condition known as beriberi, is a classic cause of neuropathy, often seen in alcoholics or those with malabsorption, producing burning feet, muscle weakness, and impaired sensation.¹ Good dietary sources include whole grains, pork, fish, nuts, seeds, and legumes.

Vitamin B6, or pyridoxine, is involved in over one hundred enzymatic reactions, including neurotransmitter synthesis of serotonin and dopamine and the formation of myelin. Both deficiency and excess of B6 can cause neuropathy: deficiency can lead to sensory neuropathy with numbness and tingling, while high doses,

over 200 milligrams a day taken long term, can cause a toxic neuropathy, so careful monitoring is essential.² Good sources include chickpeas, salmon, tuna, chicken, potatoes, and bananas.

Vitamin B9, or folate and folic acid, is critical for DNA synthesis and repair and for the methylation cycle, which is vital for nerve health and detoxification, and it works synergistically with B12. Folate deficiency can lead to nerve damage, often presenting with symptoms similar to B12 deficiency.³ Good sources include leafy green vegetables, legumes, avocado, and fortified grains.

Vitamin B12, or cobalamin, is absolutely critical for the formation and maintenance of the myelin sheath, for DNA synthesis, and for the health of red blood cells. B12 deficiency is a very common and reversible cause of neuropathy, often presenting as numbness, tingling, weakness, balance issues, and cognitive difficulties, and it can be caused by inadequate dietary intake in vegans and vegetarians, by pernicious anemia (a lack of intrinsic factor), or by malabsorption issues, for example from certain medications like metformin or proton pump inhibitors, or from gut surgery.⁴ Good sources are animal products such as meat, fish, eggs, and dairy.

Vitamin D: Beyond Bone Health

Long known for its role in bone health, vitamin D, specifically D3, is now recognized as a critical neurosteroid with profound effects on nerve and immune function. It acts more like a hormone, influencing gene expression in nerves, reducing inflammation, supporting nerve growth factor production, and modulating immune responses. Deficiency is common in neuropathy patients, and studies suggest vitamin D supplementation can improve nerve function and reduce neuropathic pain, particularly in diabetic neuropathy, thanks to its anti-inflammatory and neuroprotective properties.⁵ Good sources include sunlight exposure, fatty fish such as salmon and mackerel, fortified foods, and supplements.

Magnesium: The Great Relaxer

Magnesium is an essential mineral involved in over three hundred enzymatic reactions, including those behind nerve transmission, muscle contraction, energy production, and pain modulation. It stabilizes nerve membranes, regulates neurotransmitter activity such as glutamate and GABA, and influences cellular energy in the form of ATP; it is a natural muscle relaxant and can help reduce nerve hyperexcitability. Deficiency can lead to muscle cramps, spasms, restless legs, and increased nerve pain sensitivity, and many people are deficient due to modern diets and stress.⁶ Good sources include leafy green vegetables, nuts, seeds, legumes, whole grains, and dark chocolate.

Alpha-Lipoic Acid (ALA): The Master Antioxidant

ALA is a powerful antioxidant naturally produced in the body and found in some foods. It is unique because it is both water soluble and fat soluble, which allows it to work throughout the body. It neutralizes a wide range of free radicals, regenerates other antioxidants such as vitamin C, vitamin E, and glutathione, improves glucose utilization, and supports mitochondrial function. Extensive research, particularly in diabetic neuropathy, shows ALA can significantly improve symptoms like burning, numbness, and pain by enhancing

nerve blood flow, reducing oxidative stress, and promoting nerve regeneration.⁷ It occurs naturally in red meat, organ meats, spinach, broccoli, and yeast, though supplementation is usually needed to reach therapeutic doses.

Acetyl-L-Carnitine (ALCAR): Nerve Repair Agent

An amino acid derivative, acetyl-L-carnitine is central to energy production within cells, especially in mitochondria, and has direct neuroprotective and neuroregenerative properties. It transports fatty acids into mitochondria for energy, promotes nerve growth factor production, supports nerve cell integrity, and has antioxidant and anti-inflammatory effects. Studies suggest ALCAR can help improve nerve conduction velocity, reduce neuropathic pain, and promote nerve fiber regeneration, particularly in chemotherapy-induced and diabetic neuropathies.⁸ It is found in red meat and dairy, and is often used therapeutically as a supplement.

Other Potential Deficiencies

Copper is essential for enzymes involved in nerve function and myelin, and both deficiency, often induced by excess zinc supplementation, and toxicity can cause neuropathy. Zinc is involved in nerve signaling and antioxidant defense, and balance is key. Omega-3 fatty acids, EPA and DHA, found in fatty fish, are powerful anti-inflammatory agents that are crucial for cell membrane integrity, including in nerve cells, and they can reduce nerve inflammation and support overall brain and nerve health.⁹

Testing for Deficiencies

Don't guess, test. Blood tests are crucial to identify specific nutrient deficiencies, and these are worth discussing with your healthcare provider. A complete vitamin B panel should look beyond just B12 to include B1, B6, and folate levels; for B12 specifically, measure both serum B12 and the more sensitive markers methylmalonic acid (MMA) and homocysteine, which rise when B12 is functionally deficient at the cellular level. The 25-hydroxy vitamin D test is the most accurate measure of your vitamin D status. Red blood cell, or RBC, magnesium is more reflective of cellular magnesium stores than serum magnesium. And a trace mineral analysis, whether blood or hair based, can assess levels of copper, zinc, and other important trace minerals.

Food First Approach and Targeted Supplementation

While comprehensive testing is vital, the “food first” principle should always be your foundation. A nutrient-dense, whole-food diet provides nutrients in their most bioavailable forms and with synergistic co-factors. Prioritize a wide variety of colorful vegetables, fruits, healthy fats, quality proteins, and whole grains (more on this in Chapter 6). Once deficiencies are identified through testing, targeted, high-quality supplementation can be transformative. Where possible, choose active forms, for example methylcobalamin or adenosylcobalamin for B12, L-methylfolate for folate, and magnesium glycinate or citrate for magnesium. Work with a knowledgeable practitioner to determine appropriate therapeutic dosages, especially for

nutrients like B6 where excess can be harmful, and opt for reputable brands that offer third-party testing for purity and potency.

Addressing nutrient deficiencies is not just about adding a supplement; it's about providing your nerves with the fundamental resources they need to heal, regenerate, and function optimally. This key lays the groundwork for all other healing strategies.

Chapter 3: Key 2, Addressing Chronic Infections and Their Impact

Imagine your immune system as a vigilant army, constantly patrolling your body to identify and neutralize threats. When it encounters an acute infection, like a cold or a cut, it mounts a swift, decisive attack and then returns to a state of calm. With chronic infections, however, this army remains perpetually on high alert, leading to persistent inflammation and immune dysfunction. This constant state of alert can wreak havoc on your entire system, and your delicate nerves are often caught in the crossfire.

The connection between chronic infections and neuropathy is often overlooked in conventional medicine, yet it represents a significant “key” to unlocking stubborn cases. These infections don’t necessarily directly invade and destroy nerve tissue, though some are capable of it; rather, they trigger systemic inflammation, provoke autoimmune responses, or deplete the body’s resources, all of which can damage nerves.

The Infection-Neuropathy Connection: How Persistent Invaders Damage Nerves

Chronic infections can contribute to neuropathy through several mechanisms. Chronic inflammation is one: the sustained immune response to a persistent pathogen releases inflammatory cytokines and chemical mediators that can directly irritate and damage nerve fibers and their surrounding supportive tissues. An autoimmune response, through what is called molecular mimicry, is another. In some cases, the immune system, trying to fight off an infection, mistakenly identifies healthy nerve tissue as being similar to the invading pathogen, and this molecular mimicry leads the immune system to attack its own nerves, resulting in autoimmune neuropathy. Nutrient depletion also plays a role, since fighting a chronic infection consumes vast amounts of the body’s energy and nutrient reserves, including B vitamins, zinc, and magnesium, exacerbating deficiencies that are already detrimental to nerve health. Some pathogens produce toxins that can be directly neurotoxic, impairing nerve function or causing damage. And chronic inflammation can damage blood vessels, compromising the essential blood supply to nerves.

Common Infections Implicated in Neuropathy

While any persistent infection could theoretically contribute to nerve issues, certain pathogens are more frequently linked to neuropathy.

Lyme disease, caused by *Borrelia burgdorferi* and its co-infections, is a notorious cause of neurological symptoms, including neuropathy.¹⁰ This tick-borne bacterial infection can cause a wide range of peripheral nerve problems, from numbness and tingling to burning pain and weakness, often fluctuating in severity and location, and Lyme can also trigger autoimmune responses.¹¹ Testing requires specialized blood tests, such as Western Blot, PCR, or specific culture tests, that often go beyond standard Lyme screens. Because Lyme so often travels with co-infections, the neurological picture can be genuinely difficult to sort out, which is a strong argument for better diagnostics than standard screens provide.¹²

The herpes viruses are another notable family. Varicella-zoster virus, the virus that causes chickenpox and shingles, can lie dormant in nerve roots for decades after the original chickenpox infection; its reactivation as shingles causes a painful rash and, in many cases, post-herpetic neuralgia, a severe, persistent neuropathic pain in the affected dermatome that continues even after the rash resolves.¹³ Epstein-Barr virus, a very common virus that can cause mononucleosis, is in some individuals linked, through chronic activation, to chronic fatigue, fibromyalgia, and various neurological symptoms, including peripheral neuropathy, possibly via immune dysregulation.¹⁴ Testing for both relies on antibody tests for active or past infection, looking at IgM and IgG.

Hepatitis C virus can cause a painful peripheral neuropathy, often linked to cryoglobulinemia, a condition where abnormal proteins in the blood clump together in cold temperatures, damaging small blood vessels and nerves.¹⁵ Testing involves blood tests for HCV antibodies and viral load.

Small intestinal bacterial overgrowth, or SIBO, and gut dysbiosis more broadly are not “classic” infections of the nerves. Even so, an imbalance in gut bacteria, or an overgrowth of bacteria in the small intestine, can lead to systemic inflammation, nutrient malabsorption (especially of B vitamins), and increased intestinal permeability, commonly known as leaky gut. This allows bacterial toxins and undigested food particles to enter the bloodstream, triggering systemic inflammation and potentially autoimmune responses that can impact nerve health.¹⁶ Testing includes lactulose or glucose breath tests for SIBO and comprehensive stool analysis for gut dysbiosis.

Mycoplasma pneumoniae, an atypical bacterium that commonly causes respiratory infections, can also lead to neurological complications, including neuropathy, often through autoimmune mechanisms where antibodies generated against the bacteria cross-react with nerve components.¹⁷ Testing looks for Mycoplasma antibodies in the blood.

Diagnostic Approaches: Uncovering Hidden Infections

Identifying chronic infections requires a thorough clinical history and often specialized laboratory testing, since standard blood tests might miss these subtle, persistent threats; this is a good reason to work with a functional medicine practitioner experienced in complex chronic infections. Expanded antibody panels look beyond acute infection markers to include IgG antibodies that indicate past exposure or chronic activation, such as EBV early antigen, VZV IgG, or specific Lyme co-infection panels. PCR testing, or polymerase chain reaction testing, can detect the genetic material of pathogens, indicating active infection, and in some cases specialized cultures may be used as well. A comprehensive stool analysis can identify bacterial, fungal, or parasitic overgrowths and imbalances in the gut microbiome. And environmental toxin screening, while not testing for an “infection” as such, matters too, since heavy metals and mold toxins can mimic and exacerbate infection symptoms and compromise the immune system, making the body more susceptible.

Holistic Strategies for Infection Management

Once a chronic infection is identified, the approach has several parts, focusing on supporting the immune system and reducing the pathogen's load without relying solely on long-term antibiotics, which can further disrupt the gut microbiome.

On the dietary side, an anti-inflammatory diet, as discussed in Chapter 6, reduces inflammatory foods such as processed foods, sugar, and refined oils, which is crucial to lower the systemic inflammatory burden and support immune function. Immune-boosting foods rich in vitamin C, zinc, vitamin D, and antioxidants, such as colorful fruits and vegetables, bone broth, garlic, onions, ginger, and turmeric, are worth incorporating regularly. Fermented foods like sauerkraut, kimchi, and kefir, along with prebiotic fibers like onions, garlic, and asparagus, support a healthy microbiome, which is foundational to a robust immune system.

Many natural compounds possess antimicrobial, antiviral, and antiparasitic properties that can help the body fight off chronic pathogens with fewer side effects than pharmaceuticals. Examples include olive leaf extract, oregano oil, berberine, garlic, artemisia, cat's claw, colloidal silver, and specific probiotics. These should always be used under the guidance of a qualified healthcare practitioner, given potential interactions and the importance of appropriate dosing.

Immune system modulation is another lever. Adaptogenic herbs like ashwagandha, rhodiola, and ginseng can help the body adapt to stress and support immune balance. Beta-glucans, found in medicinal mushrooms such as reishi and turkey tail, can prime the immune system to respond more effectively. Vitamin D and zinc, discussed in Chapter 2, remain critical for immune cell function. And glutathione, the body's master antioxidant, is vital for detoxification and immune health, with supplementation of precursors like N-acetyl cysteine (NAC) often beneficial.

Finally, stress reduction and sleep optimization matter more than they might seem to. Chronic stress suppresses the immune system, making the body more vulnerable to persistent infections, so techniques like meditation, yoga, and deep breathing are vital, and adequate, restorative sleep is non-negotiable for immune repair and regulation.

By identifying and strategically addressing chronic infections, you remove a significant burden on your immune system and reduce a potent source of inflammation that may be driving your neuropathy. This “key” is about supporting your body's innate ability to heal and restore balance.

Chapter 4: Key 3, Activating Large Diameter Afferent Pathways

Neuropathy often leads to a disruption in the delicate balance of your nervous system. While small nerve fibers transmit pain and temperature, it's the large diameter afferent pathways that carry crucial information about touch, pressure, vibration, and proprioception, your sense of body position in space. When these larger, myelinated fibers are damaged or their signals are overwhelmed by pain signals from smaller, unmyelinated fibers, the brain receives distorted or incomplete sensory information.

Activating and stimulating these large diameter afferent pathways is a powerful strategy in neuropathy management. This process, often referred to as sensory retraining or gating, helps in several ways at once. According to the Gate Control Theory of Pain, stimulating large afferent fibers can “close the gate” to pain signals from smaller fibers, reducing the perception of pain. Restoring accurate sensory input helps the brain better understand where your body is in space, improving coordination and reducing fall risk. By re-educating the nervous system and improving nerve signaling, numbness and tingling can gradually lessen as sensation returns or improves. And the process promotes neuroplasticity, the brain's ability to reorganize itself by forming new neural connections, which can help compensate for damaged areas.

This chapter explores various methods to activate these vital pathways, from simple at-home techniques to advanced therapeutic devices.

1. Vibration Therapy

Vibration is a powerful stimulus for large diameter afferent fibers, particularly Pacinian corpuscles, which are highly sensitive to vibration. Applying specific frequencies of vibration directly to affected areas, such as the feet or hands, stimulates these mechanoreceptors, sending strong signals up the large afferent pathways to the brain, which can help override pain signals and improve sensory perception. In practice, this can take the form of whole-body vibration platforms used for general stimulation and to improve balance, handheld vibrators applied to the soles of the feet, palms of the hands, or specific numb areas for direct stimulation, or medical-grade tuning forks, for example at 128 Hz, applied to bony prominences to assess and stimulate vibratory sensation. Vibration therapy is reported to reduce neuropathic pain, improve balance, and enhance sensory perception.

2. Tactile Stimulation

Simple touch and texture can re-engage sensory pathways. Light touch, brushing, or exposure to varied textures activates different types of mechanoreceptors, sending diverse sensory information to the brain, which helps to desensitize hypersensitive areas and re-awaken numb ones. This can look like walking barefoot on different textured surfaces such as grass, sand, or textured mats; using a soft sensory brush to gently stroke numb or painful areas; or receiving gentle massage, which can improve circulation and

stimulate nerve endings. Tactile stimulation helps to normalize sensory processing, reduce allodynia (pain from non-painful stimuli), and improve tactile discrimination.

3. Proprioceptive Exercises

These exercises focus on your body's awareness in space. Engaging in activities that challenge balance and coordination forces the brain to rely more heavily on proprioceptive input from joints and muscles, strengthening these pathways. Balance exercises such as standing on one leg, heel-to-toe walking, or using a wobble board or balance disc are one avenue; tai chi or yoga, which inherently improve balance, body awareness, and coordination, are another; and gait training, focused on proper walking mechanics, rounds this out. Together, these improve balance, reduce fall risk, enhance coordination, and strengthen the neural connections responsible for body awareness.

4. Photobiomodulation (Laser and Red Light Therapy)

Photobiomodulation, often referred to as low-level laser therapy or red light therapy, uses specific wavelengths of light to stimulate cellular function and promote healing. While it does not directly activate large diameter afferents in the same mechanical way as vibration, it creates an optimal environment for nerve health and repair, indirectly supporting the function of all nerve fibers, including large afferents.

Its mechanism works on several fronts. Red and near-infrared light penetrates tissues and is absorbed by chromophores, primarily cytochrome c oxidase, within the mitochondria of cells, and this absorption increases the production of adenosine triphosphate (ATP), the primary energy currency of cells, meaning nerve cells have more energy to function optimally and repair themselves. Photobiomodulation also helps modulate inflammatory pathways and reduce oxidative stress by neutralizing free radicals; since chronic inflammation and oxidative stress are major contributors to nerve damage, reducing them creates a healthier environment for nerve function and regeneration. Light therapy can promote vasodilation and angiogenesis, the widening of blood vessels and the formation of new ones, leading to increased blood flow to nerve tissues and ensuring nerves receive a vital supply of oxygen and nutrients. It has been shown to stimulate nerve growth factors, promote axonal sprouting, and support the regeneration of damaged nerve fibers and their myelin sheaths, which can lead to improvements in nerve conduction and overall function. And while it's not a direct activation of large afferents for gating, it can reduce pain by decreasing inflammation, improving nerve health, and potentially influencing nerve cell action potentials and neurotransmitter release.

For neuropathy specifically, the reported benefits include pain reduction (burning, shooting pain), improved sensation (reduced numbness and tingling), enhanced nerve conduction velocity, support for nerve regeneration, and reduced inflammation in nerve tissues. In application, clinic-based high-power therapeutic lasers (Class IV) are often used in clinical settings for deeper penetration and faster treatment times, while home-use red light devices, lower-power LED panels, wraps, and handheld units, are available for home use but require more consistent and longer application times. These devices typically use red light around 600 to 700 nanometers and near-infrared light around 780 to 1,100 nanometers for their ability to penetrate tissue. A systematic review found that low-level laser therapy has a positive effect in controlling diabetic neuropathic

pain,¹⁸ and research suggests that near-infrared laser light can rapidly block pain transmission and, with lower irradiances, offer prolonged pain relief by altering chronic inflammation.¹⁹

5. Electrical Stimulation Devices, Such As the ReBuilder Medical Device

Electrical stimulation has long been used in pain management and rehabilitation. Devices like the ReBuilder Medical Device are designed to deliver specific electrical signals to nerve pathways, aiming to re-educate nerves and improve their function.

The ReBuilder device is designed to generate a proprietary electrical waveform that reportedly mimics the natural electrical signals of healthy nerves, which is distinct from standard TENS (transcutaneous electrical nerve stimulation) units that often use simpler, less varied waveforms. By applying electrodes to specific areas, such as the soles of the feet or the hands, the device sends these signals along the nerve pathways, and the manufacturer states that it uses biofeedback to adjust its output to the individual's electrical parameters, aiming for more personalized and effective stimulation. By delivering precise electrical impulses, the ReBuilder aims to directly stimulate large diameter afferent nerve fibers, which can help “wake up” dormant nerves that have become less functional due to damage, improve nerve conduction by enhancing the speed and efficiency of signal transmission, and modulate pain through gating, since, similar to vibration, the strong, regular signals from the device can activate large afferents and help “close the gate” on pain signals from smaller, damaged nerves. The electrical stimulation can also induce muscle contractions and promote vasodilation, increasing local blood flow and oxygenation to nerve tissues, which is vital for nerve health and repair.

Reported benefits for neuropathy include a reduction in neuropathic pain (burning, aching, shooting pains), a decrease in numbness and tingling, improved sensation and proprioception, increased local blood circulation, relaxation of muscle spasms and reduction of cramps, and the potential for reduced reliance on pain medications. In practice, the ReBuilder device typically involves placing conductive pads on the soles of the feet, or the hands, lower back, or shoulders, or using a footbath with a conductive solution; treatments are usually 30 minutes, once or twice daily, with reported immediate relief that can last for several hours and cumulative improvements with continued use.

There are a few important considerations to state plainly. The ReBuilder Medical Device is FDA registered as a transcutaneous electrical nerve stimulator for pain relief. While anecdotal reports and manufacturer-conducted studies report high success rates, independent, large-scale, peer-reviewed clinical trials specifically on the ReBuilder device for neuropathy are less abundant in databases like PubMed compared to broader categories like TENS or photobiomodulation, so it's important to discuss this option with your healthcare provider to determine if it's appropriate for your specific condition. It is also contraindicated for individuals with certain pacemakers, especially those with defibrillator accessories, and for pregnant women.

Integrating Activation Strategies

The power of activating large diameter afferent pathways lies in consistency and combination. Incorporate a variety of these techniques into your daily routine: for example, combine targeted electrical stimulation with daily balance exercises, or use a handheld vibrator on numb areas while also receiving regular photobiomodulation treatments. By providing consistent, positive sensory input, you can help retrain your nervous system, reduce pain, and improve your overall nerve function. This key is about actively engaging your body's innate capacity for neuroplasticity and healing.

Chapter 5: Key 4, Comprehensive Testing Beyond the Conventional

When facing the complexities of neuropathy, relying solely on standard diagnostic tests can often leave crucial stones unturned. Conventional evaluations typically include nerve conduction studies (NCS) and electromyography (EMG) to assess nerve damage, along with basic blood work to rule out common culprits like overt diabetes or severe B12 deficiency. While these are important starting points, they often fail to uncover the nuanced, underlying imbalances that contribute to the majority of neuropathy cases.

The root cause approach to neuropathy demands a more sophisticated and comprehensive diagnostic strategy. This key involves looking beyond superficial markers to identify hidden inflammation, subtle metabolic dysregulation, circulatory compromise, and other systemic factors that may be silently driving your nerve damage. By truly understanding the physiological landscape of your body, you can create a truly targeted and effective treatment plan.

The Limitations of Standard Tests: Why a Deeper Dive Is Needed

Nerve conduction studies and electromyography assess the function of large, myelinated nerve fibers and muscle electrical activity. While invaluable for diagnosing the presence and type of nerve damage, they may not detect small fiber neuropathy, which involves the tiny nerves responsible for pain and temperature sensation, and more importantly, they tell you what nerve damage exists, but rarely why. Basic blood work, a fasting glucose test or HbA1c, may flag diabetes, and a basic B12 level might catch severe deficiency, but these often miss pre-diabetic states, functional B12 deficiencies, or other critical nutrient imbalances that aren't severe enough to show up on standard ranges but are still detrimental to nerve health. And when standard tests yield no clear cause, patients are often labeled with idiopathic neuropathy, which means the cause is unknown, not that there isn't a cause, and that is precisely where comprehensive testing becomes crucial.

Expanding the Diagnostic Lens: Key Areas for Deeper Investigation

Working with a functional medicine practitioner who specializes in neuropathy means accessing a broader array of diagnostic tools designed to uncover the physiological stressors impacting your nerves.

Inflammation markers are the silent nerve attackers, since chronic, low-grade inflammation is a major driver of nerve damage even without a diagnosed autoimmune condition, so identifying and quantifying this inflammation is paramount.²⁰ High-sensitivity C-reactive protein, or hs-CRP, is a sensitive but entirely nonspecific screening marker for systemic inflammation: an elevated result says an inflammatory load exists somewhere, not where it is coming from, and it confirms nothing on its own. Read it as a prompt to keep looking, not as an answer. Erythrocyte sedimentation rate is another general marker of inflammation, and less specific still. Homocysteine belongs in a different category, and it is one of the most consequential

numbers on this panel rather than a piece of trivia to wave off. It is a methylation-cycle marker. It rises when B12, folate, or B6 run short, and with thyroid or kidney trouble, and elevated levels are associated with increased inflammation, oxidative stress, and vascular changes that work against nerve health.²¹ Rather than chasing the number with blind supplementation, find which input is short (B12, folate, B6, thyroid, or kidney function), correct that one, and let homocysteine confirm the fix on retest. Fibrinogen, a protein involved in blood clotting and inflammation, can indicate chronic inflammation and increased risk of vascular issues affecting nerve blood supply when elevated. And more advanced cytokine testing, for markers like IL-6 and TNF-alpha, can measure specific inflammatory cytokines that directly contribute to nerve pain and damage, offering a deeper insight into immune system activation.

Blood sugar regulation matters well beyond a diabetes diagnosis, since many neuropathy cases are linked to blood sugar dysregulation even if a person isn't officially diagnosed with diabetes, and pre-diabetes or insulin resistance can cause nerve damage over time.²² Measuring fasting glucose and insulin together provides insight into how much insulin your body needs to keep blood sugar stable, and high fasting insulin indicates insulin resistance. Hemoglobin A1c gives an average blood sugar level over the past two to three months; while it's a standard test, looking at optimal ranges, below 5.4 percent, rather than just "non-diabetic" ranges, is important. The oral glucose tolerance test with insulin measures blood glucose and insulin levels at intervals after a glucose drink, and it can uncover subtle dysregulation that fasting tests miss. It is the most informative test that is realistically orderable for this question, which is not the same thing as a gold standard, and no single one of these numbers confirms insulin resistance on its own. And for some patients, wearing a continuous glucose monitor for a few days or weeks can provide invaluable real-time data on how different foods, stress, and activities impact blood sugar, revealing spikes and crashes that contribute to nerve stress.

Circulation assessment recognizes that nerves, like all tissues, depend on a robust blood supply for oxygen and nutrients, and compromised circulation is a direct pathway to nerve damage. A physical exam can reveal signs of poor circulation, and non-invasive tests like the ankle-brachial index can screen for peripheral artery disease, which often coexists with neuropathy. Blood tests for endothelial function markers like asymmetric dimethylarginine (ADMA) or endothelin-1 can provide insight into the health of your blood vessel lining, which is critical for healthy blood flow. And the inflammatory markers already discussed, high hs-CRP and homocysteine, also negatively impact vascular health.

Autoimmune markers matter because neuropathy can be an autoimmune condition where the immune system mistakenly attacks nerve components, and specific testing can help identify these cases.²³ Antinuclear antibodies (ANA) offer a general screen for autoimmune activity. Beyond that, specific autoantibody panels look at ganglioside antibodies, such as anti-GM1, anti-GD1a, and anti-GQ1b, which are associated with various autoimmune neuropathies like Guillain-Barré syndrome or multifocal motor neuropathy; MAG antibodies (myelin-associated glycoprotein antibodies), linked to demyelinating neuropathies; SSA and SSB antibodies associated with Sjögren's syndrome, which can cause significant small fiber neuropathy; and thyroid antibodies (TPOAb, TgAb), since autoimmune thyroid conditions like Hashimoto's can be associated with neuropathy.

Heavy metal toxicity testing addresses a hidden poison, since exposure to heavy metals can be profoundly neurotoxic, causing direct damage to nerve cells and their myelin sheaths. Hair mineral analysis can reflect long-term exposure to certain heavy metals. A urine heavy metal challenge test, which involves collecting urine before and after administering a chelating agent that mobilizes metals from tissues, provides a comprehensive assessment of body burden, while a blood heavy metal test is useful for recent or acute exposure. A study of metal industry workers with chronic occupational exposure to lead, cadmium, and arsenic found clear, measurable impairment of peripheral nerve function, most pronounced in the small sensory fibers that carry pain and temperature signals.²⁴

The Organic Acids Test, or OAT, offers metabolic insights by providing a snapshot of various pathways in the body, including mitochondrial function, neurotransmitter metabolism, and nutrient needs. It identifies imbalances in gut flora, such as yeast and bacterial overgrowth, nutrient deficiencies in B vitamins and antioxidants, and markers of oxidative stress that can impact nerve health; for example, high markers of certain organic acids can indicate issues with energy production or detoxification crucial for nerves. It helps reveal functional deficiencies that might not show up on simple blood tests and provides insight into the efficiency of various metabolic processes.

Comprehensive stool testing captures the gut-nerve connection, since your gut microbiome has a profound influence on your overall health, including your nervous system, and dysbiosis, an imbalance of gut bacteria, can lead to systemic inflammation, nutrient malabsorption, and the production of neurotoxic compounds. It analyzes the balance of beneficial and pathogenic bacteria, yeast, parasites, markers of inflammation, and digestive function, and it can identify issues like SIBO, candidiasis, or parasitic infections that contribute to systemic inflammation and nerve symptoms.¹⁶

The Value of Comprehensive Testing

Investing in these advanced diagnostic tests moves you from guessing to knowing. It allows your healthcare team to develop a highly personalized treatment plan that targets the precise mechanisms contributing to your neuropathy, rather than a generic approach. This key empowers you to peel back the layers of your condition and address the root causes, paving the way for more effective and lasting relief.

Chapter 6: Key 5, Diet and Food Sensitivities for Nerve Repair

Imagine your body as a high-performance vehicle. You wouldn't fill a sports car with low-grade fuel and expect it to run optimally. Similarly, the food you consume profoundly impacts every cell, tissue, and organ, especially your delicate nervous system. Diet is not merely about calories; it's about providing the specific nutrients for repair, reducing inflammation, stabilizing blood sugar, and supporting a healthy gut, all critical factors for unlocking nerve health.

This key is arguably one of the most foundational and empowering. While nutrient deficiencies, covered in Chapter 2, highlight what you might lack, this chapter focuses on how your daily food choices can either perpetuate nerve damage or actively promote nerve healing and protection. We will explore the principles of an anti-inflammatory, nerve-supportive diet and look closely at the often-overlooked role of food sensitivities.

The Anti-Inflammatory Diet Principle: Reducing Triggers and Promoting Healing

Chronic inflammation is a silent assassin of nerves. It damages myelin, impairs blood flow, and directly irritates nerve fibers. The cornerstone of a neuropathy-healing diet is to systematically reduce inflammatory triggers while flooding your body with anti-inflammatory compounds.

Among the foods to embrace, your strongest allies are abundant vegetables and fruits. That means a wide array of colorful, non-starchy vegetables, especially leafy greens like spinach, kale, and collards, and cruciferous vegetables like broccoli, cauliflower, and Brussels sprouts, along with low-glycemic fruits like berries, avocados, lemons, and limes, all packed with antioxidants, vitamins, minerals, and fiber. Healthy fats are essential for nerve cell membranes and myelin, so include sources like extra virgin olive oil, avocados, nuts such as almonds, walnuts, and pecans, seeds such as chia, flax, and hemp, and fatty fish such as wild-caught salmon, mackerel, and sardines, rich in anti-inflammatory omega-3 fatty acids (EPA and DHA). Lean, high-quality proteins are necessary for tissue repair and neurotransmitter synthesis, so prioritize grass-fed or pasture-raised meats, wild-caught fish, organic poultry, and plant-based options like legumes, if tolerated, and quinoa. Fiber-rich foods, or prebiotics, feed beneficial gut bacteria, which produce short-chain fatty acids that reduce inflammation and support gut barrier integrity, so include legumes, whole grains if tolerated, and a variety of vegetables and fruits. And anti-inflammatory spices like turmeric (paired with black pepper for absorption), ginger, cinnamon, garlic, and rosemary contain potent compounds that combat inflammation.

On the other side of the ledger are the foods to limit or avoid, the common culprits that can fuel inflammation and undermine nerve health. Processed foods are often high in unhealthy fats, sugar, and artificial ingredients. Refined sugars and sweeteners directly contribute to blood sugar spikes, insulin resistance, and systemic inflammation, which are primary drivers of diabetic neuropathy and general nerve damage, and this includes the hidden sugars in sauces, dressings, and drinks. Unhealthy fats, trans fats (partially hydrogenated oils) and highly refined vegetable oils such as corn, soy, canola, sunflower, and safflower oil,

are high in pro-inflammatory omega-6s. Artificial ingredients, dyes, preservatives, and artificial sweeteners such as aspartame and sucralose, can trigger inflammation and neurological symptoms in sensitive individuals. And excessive alcohol can be directly neurotoxic and deplete vital nutrients.

Elimination Diet for Food Sensitivities: Unmasking Hidden Triggers

Beyond obvious allergies, many people suffer from delayed food sensitivities, where an immune reaction to certain foods causes low-grade inflammation that can manifest in various symptoms, including neuropathic pain. Identifying these triggers is crucial.

It helps to understand the difference between the two. A food allergy is an immediate, often severe, IgE-mediated immune reaction, for example hives or difficulty breathing. A food sensitivity, or intolerance, is a delayed, often IgG-mediated immune reaction that can cause a wide range of less immediate symptoms, such as digestive issues, headaches, joint pain, skin problems, and chronic inflammation impacting nerves.

The most common food sensitivities include gluten, found in wheat, barley, and rye, where gluten sensitivity or celiac disease can cause direct nerve damage, sometimes called gluten neuropathy, and significant systemic inflammation;²⁵ dairy, where casein and lactose can be inflammatory for many; soy, a common allergen and inflammatory food for some; corn, often genetically modified and inflammatory for sensitive individuals; and nightshades such as tomatoes, potatoes, peppers, and eggplant, which, while healthy for many, contain compounds called alkaloids that can aggravate inflammation and pain in sensitive individuals.

To conduct an elimination and reintroduction phase, start with a strict elimination phase of three to four weeks: remove all suspected inflammatory foods and common sensitivities, including gluten, dairy, soy, corn, eggs, peanuts, sugar, processed foods, and alcohol, and focus exclusively on whole, unprocessed, anti-inflammatory foods during this window. This gives your body a chance to calm down and symptoms to subside. After that, move into a reintroduction phase: bring back one food group at a time every two to three days, and monitor your symptoms closely, for example with a symptom journal. If symptoms worsen, neuropathic pain, fatigue, digestive upset, or brain fog, that food is likely a trigger for you. Once you identify your triggers, you can build a personalized diet that supports your nerve health without guilt or deprivation.

Hydration and herbal support round out the foundation. Water is vital for every bodily process, including nerve signal transmission and detoxification, so aim for half your body weight in ounces of filtered water daily. Certain herbs have neuroprotective, anti-inflammatory, or pain-relieving properties worth knowing about: turmeric and curcumin are potent anti-inflammatories and antioxidants; ginger reduces inflammation and pain; boswellia is anti-inflammatory, often used for joint pain but relevant for nerve inflammation too; and passionflower, California poppy, and valerian can help with nerve pain and improve sleep, indirectly supporting nerve repair. Always consult with your healthcare provider before adding herbal supplements, especially if you are on medications.

Sample 90-Day Food Guide Framework

This framework provides a structured approach to implementing your nerve-healing diet. It's designed to transition you from an inflammatory eating pattern to a sustainable, nerve-supportive lifestyle.

As a weekly framework, a useful guideline for your macronutrient breakdown is what we call 60/20/20: meals that stabilize blood sugar and provide sustained energy tend to draw about 60 percent of their content from healthy carbohydrates, primarily non-starchy vegetables, some fruits, and carefully selected whole grains if tolerated and after the elimination phase, focusing on complex carbohydrates with high fiber content; about 20 percent from healthy fats, from sources like avocado, olive oil, nuts, seeds, and fatty fish; and the remaining 20 percent from quality proteins, from lean meats, poultry, fish, eggs, and plant-based proteins. This is a general guideline. Your individual needs may vary, and a nutritionist can help fine-tune these percentages based on your specific health goals and activity level.

In the first phase, weeks one through four, the focus is elimination and detoxification. You strictly eliminate common inflammatory foods and sensitivities and emphasize whole, unprocessed foods, with the goal of reducing inflammation, stabilizing blood sugar, allowing the gut to heal, and identifying immediate triggers. Meals in this phase are built around abundant non-starchy vegetables, clean protein, and healthy fats, with minimal fruit and no grains initially. A representative day might start with a green smoothie made of spinach, plant-based protein powder, almond milk, and half an avocado; lunch could be a large mixed greens salad with grilled chicken or wild salmon, an olive oil and lemon dressing, and a handful of pumpkin seeds; dinner might be baked cod with steamed broccoli and sautéed asparagus; and a small handful of almonds makes a good snack.

In the second phase, weeks five through eight, the focus shifts to reintroduction and personalization. You systematically reintroduce one eliminated food group at a time, observing for symptoms, with the goal of identifying your individual food sensitivities and expanding your personalized diet. The menu philosophy continues with the anti-inflammatory base from phase one, but strategically adds foods back: breakfast can stay the same as phase one, lunch might become turkey and avocado lettuce wraps with a side of mixed berries as you reintroduce berries, and dinner might be gluten-free pasta, if you are reintroducing rice pasta, with lean ground turkey and a tomato-based sauce once tomatoes and other nightshades are back in.

In the third phase, weeks nine through twelve, the focus is maintenance and optimization. You solidify your personalized anti-inflammatory diet based on your findings and continue to prioritize nutrient density, with the goal of sustaining nerve healing, managing blood sugar, maintaining low inflammation, and feeling consistently better. Here you enjoy a diverse range of your newly identified safe and supportive foods: breakfast might be chia seed pudding with berries and walnuts, lunch could be leftover dinner or a large vegetable and lentil soup, and dinner might be a stir-fry with lean beef, a variety of colorful vegetables, and brown rice if tolerated.

Throughout all three phases, a handful of snacks reliably stabilize glucose between meals: hard-boiled eggs, avocado slices with a sprinkle of sea salt, a small handful of nuts such as almonds or walnuts or seeds such

as pumpkin or sunflower, celery sticks with almond or seed butter, berries with a dollop of unsweetened coconut yogurt, olives, and a small piece of wild-caught salmon or sugar-free turkey jerky.

Getting the logistics right makes the plan sustainable. Dedicate time each week to plan meals and snacks. Shop smart by focusing on the perimeter of the grocery store, where the produce and lean proteins live. Batch cook larger quantities of staples like roasted vegetables, cooked grains, and lean protein (chicken breast, ground turkey) so you can assemble meals easily throughout the week. Pre-chop vegetables so quick meals or snacks are always within reach. And keep a well-stocked pantry of healthy staples such as olive oil, vinegars, anti-inflammatory spices, nuts, seeds, and canned wild salmon.

By consciously choosing foods that nourish your nerves and avoiding those that inflame them, you are providing your body with the optimal environment for healing. This dietary shift is a powerful, sustainable lifestyle change rather than a temporary fix, and it is essential for reclaiming your nerve health.

Chapter 7: The Movement Plan, Restoring Function and Reducing Pain

While diet and targeted therapies provide crucial internal support for nerve health, movement is the indispensable key that bridges internal healing with external function. When you live with neuropathy, fear of falling, increased pain, or perceived weakness can lead to a sedentary lifestyle. However, inactivity often worsens symptoms by reducing circulation, decreasing muscle strength, and hindering neuroplasticity, the brain's ability to adapt and reorganize.

Conversely, strategic movement is a powerful therapeutic tool. It can directly benefit nerve health by improving blood flow, promoting the release of nerve growth factors, strengthening supportive muscles, and enhancing the communication between your brain and body. This chapter will outline research-backed exercise approaches that can reduce pain, improve sensation, and restore function, regardless of your current mobility level.

Benefits of Exercise for Neuropathy

Engaging in regular, appropriate physical activity offers a multitude of advantages for individuals with neuropathy. Movement increases blood flow to the extremities, delivering vital oxygen and nutrients to damaged nerves and helping to remove waste products, which is critical for nerve repair and function. Physical activity also stimulates the release of neurotrophic factors, like brain-derived neurotrophic factor (BDNF), proteins that promote the growth, survival, and differentiation of neurons and synapses, which can aid in nerve repair and regeneration; while research linking exercise-driven BDNF increases specifically to neuropathy is still developing, the broader link between exercise and BDNF is well established.²⁶ Exercise can activate your body's natural pain-relieving mechanisms, including the release of endorphins, and consistent, appropriate movement can also desensitize overactive pain pathways, an effect supported by research on exercise as a strategy for neuropathic pain management.²⁷ Stronger muscles from regular activity provide better support for joints and improved stability, reducing the risk of falls and compensating for nerve-related weakness. Many exercises specifically target the large diameter afferent pathways responsible for your sense of body position, directly addressing common neuropathy symptoms and improving balance and proprioception. Maintaining a healthy weight through exercise is crucial, especially for diabetic neuropathy, since excess weight exacerbates insulin resistance and inflammation. And exercise is a potent antidote to stress, anxiety, and depression, which often accompany chronic pain conditions like neuropathy, so the mood enhancement and stress reduction it brings positively impacts overall nerve health as well.

Types of Exercise to Embrace

The key is to start slowly, listen to your body, and progress gradually. Consistency is more important than intensity.

Low-impact aerobics aim to improve cardiovascular health, enhance circulation, and maintain overall fitness without excessive stress on joints or feet. Walking is the simplest place to start: even short, frequent walks can make a significant difference, so begin with 5 to 10 minutes a few times a day and gradually increase duration, wearing supportive, well-cushioned shoes. Swimming and water aerobics use the buoyancy of water to reduce weight-bearing stress, making them ideal for those with foot pain or balance issues, while the resistance of the water also provides gentle strengthening. A stationary or recumbent bike offers a non-weight-bearing way to improve leg circulation and strength, and an elliptical machine provides a similarly low-impact cardio workout. Aerobic exercise of this kind increases insulin sensitivity, helping cells absorb glucose more efficiently from the bloodstream, which directly benefits those with diabetic neuropathy or insulin resistance.

Resistance, or strength, training aims to build and maintain muscle mass, which improves strength and stability and can help manage blood sugar by increasing insulin sensitivity. This can take the form of bodyweight exercises like squats using a chair for support, shallow lunges, wall push-ups, and chair stands; resistance bands, which provide adjustable resistance for arms, legs, and core; light weights, dumbbells or household items like cans or water bottles, for bicep curls, tricep extensions, and shoulder presses; or machine weights, which provide controlled movements and support and are often a good starting point. Focus on proper form to prevent injury, and start with lighter weights and higher repetitions, gradually increasing resistance as strength improves. Resistance training builds muscle, and muscle is the primary site for glucose uptake, so more muscle mass means better glucose disposal and improved insulin sensitivity, even independent of weight loss.

Walking after meals is a simple yet highly effective strategy to immediately lower post-meal blood sugar spikes. Muscle activity immediately after eating helps to utilize circulating glucose for energy, preventing the large spikes that are damaging to nerves, so aim for a brisk 10 to 15 minute walk within 30 minutes after major meals; even short bursts count. Studies show that a short walk after a meal can significantly reduce postprandial (after-meal) glucose levels, sometimes more effectively than a longer walk at another time of day.

High-intensity interval training, in moderation, involves short bursts of intense exercise followed by brief recovery periods, and it can significantly improve cardiovascular fitness, insulin sensitivity, and mitochondrial function. For individuals with neuropathy, HIIT should be approached with extreme caution and only under professional guidance, especially if balance or pain is a major concern; a modified version can involve bursts of brisk walking followed by slower walking, or short periods of higher resistance on a stationary bike. It's not suitable for everyone, especially those with severe neuropathy, balance issues, or heart conditions, so always consult your doctor before attempting it. HIIT has been shown to improve insulin sensitivity and glucose metabolism even more effectively than moderate-intensity continuous exercise in some populations, due to its potent effects on muscle glucose uptake and mitochondrial biogenesis.

Functional movement routines for balance and flexibility directly enhance proprioception, balance, flexibility, and coordination, areas often severely impacted by neuropathy, and these exercises directly activate the large diameter afferent pathways discussed in Chapter 4. Tai chi and yoga are excellent for improving

balance, flexibility, strength, and body awareness, and many studios offer chair yoga or adaptive classes for those who need them. Balance drills, standing on one foot while holding onto support initially, tandem stance (heel-to-toe), or walking safely on uneven surfaces, build the same skills directly. Gentle stretching for ankles, calves, hamstrings, and feet helps maintain range of motion and reduce stiffness, and Pilates rounds this out by focusing on core strength, stability, and controlled movements.

Sample Weekly Movement Strategy for All Levels

This is a template meant to be adapted to your comfort level, ideally alongside a physical therapist or exercise physiologist experienced with neuropathy. A few principles apply at every level: start slow, beginning with very short durations of five to ten minutes if you are new to exercise and gradually increasing; listen to your body, since pain is a signal, and you should differentiate between muscle fatigue and neuropathic pain and never push through sharp or burning neuropathic pain; prioritize consistency, since daily, even short, bursts of activity are more effective than sporadic long sessions; build in variety by combining different types of exercise to challenge your body in multiple ways; and put safety first, making sure your environment is safe, wearing appropriate footwear, and considering supervision, especially for balance exercises.

Beginner level. For those new to exercise or with significant symptoms, a good starting point is 10 to 15 minutes of brisk walking daily, split into two or three sessions if needed, plus 5 minutes of gentle stretching; three times a week, add between 10 and 15 minutes of chair-based strength exercises such as chair squats or bicep curls with light weights; twice a week, add 5 to 10 minutes of supported balance exercises such as standing on one foot while holding a counter; and each day, close with a 10 to 15 minute walk after dinner.

Intermediate level. For those regularly active with moderate symptoms, aim for 30 minutes of brisk walking or cycling daily; three to four times a week, do 20 to 30 minutes of bodyweight strength training or light resistance bands; three times a week, add 15 to 20 minutes of balance training such as tai chi, yoga, or wobble board work; and again close each day with a 10 to 15 minute walk after dinner.

Advanced level. For those with good mobility and well-managed symptoms who are seeking optimization, aim for 30 to 45 minutes of moderate-intensity cardio daily, whether brisk walking, jogging, cycling, or swimming; three to four times a week, do between 30 and 45 minutes of resistance training with free weights, machines, or challenging bodyweight work; two to three times a week, add 15 to 20 minutes of dynamic balance and flexibility work such as advanced yoga, Pilates, or complex balance drills; once or twice a week, and only with caution, consider modified HIIT sessions, for example sprint intervals on a bike for 20 to 30 seconds followed by one to two minutes of recovery; and, as at every level, close each day with a 10 to 15 minute walk after dinner.

Always consult with your doctor or a physical therapist before starting any new exercise program, especially if you have existing health conditions or severe neuropathy. They can help tailor a plan that is safe and effective for your individual needs. Movement is not just about physical strength; it's about reconnecting with your body, fostering neuroplasticity, and empowering yourself on the path to nerve health.

Chapter 8: Mindset and Monitoring, the Unseen Influencers

Neuropathy is a physical condition, but its impact extends far beyond the nerves themselves. The persistent pain, numbness, and functional limitations can take a significant toll on your mental and emotional well-being. Chronic stress, anxiety, frustration, and even feelings of hopelessness can become entangled with the physical symptoms, creating a vicious cycle that can hinder healing.

This key acknowledges that true holistic recovery encompasses the psychological and emotional landscape as well as the biological. It's about cultivating a resilient mindset, reframing setbacks, celebrating small victories, and strategically monitoring your progress to stay motivated and informed. Your brain is intimately connected to your nervous system, and nurturing this connection is as vital as any diet or supplement.

The Brain-Nerve Connection: How Mindset Influences Neuropathy

Your brain and nervous system are not separate entities. The brain constantly processes sensory input from your nerves and influences how you perceive pain, and chronic stress, anxiety, and negative thought patterns can work against you in several ways. Stress hormones like cortisol can sensitize nerve endings, making pain feel more intense, and the brain, under chronic stress, can become “stuck” in a pain loop even with minimal physical stimulus. Persistent stress diverts energy away from repair processes, including nerve regeneration, and it can also suppress the immune system, making you more vulnerable to inflammation and infections. Chronic stress can lead to vasoconstriction, the narrowing of blood vessels, reducing vital blood flow to nerves. And imbalances in mood-regulating neurotransmitters like serotonin and dopamine can exacerbate pain perception and reduce your capacity to cope.

Cultivating a positive and proactive mindset isn't about ignoring your symptoms; it's about empowering yourself to influence your body's healing capacity and improve your quality of life.

Journaling Blood Sugar Trends

For those with diabetic neuropathy or insulin resistance, consistent glucose monitoring is not just about managing numbers; it's a powerful tool for understanding your body's unique responses and making informed dietary choices. While your fasting glucose and HbA1c give you a snapshot, daily logging reveals patterns. Each time, record the time of measurement (before meals, one to two hours after meals, before bed, upon waking), the glucose reading itself, meal details (what you ate, approximate portion sizes, and main ingredients, noting carbohydrate, protein, and fat content), any activity before or after the meal (a fifteen-minute walk after dinner, say), a simple stress level rating from one to five or a brief note about emotional state, sleep quality from the night before, and any changes in neuropathy symptoms, pain, numbness, or tingling, after specific foods or activities. Journaling helps you identify the triggers, foods,

stress, or lack of sleep, that cause blood sugar spikes and subsequent neuropathy flare-ups, and it empowers you to see the direct connection between your lifestyle choices and your symptoms.

Glucose Monitoring Tips

Consistency is key: test at consistent times so you can identify patterns. Testing before and after meals, so-called paired testing, is crucial to see how specific foods affect your blood sugar. If it's available and appropriate for your condition, a continuous glucose monitor can provide real-time data, revealing spikes and trends you might miss with finger-prick tests; this passive monitoring is particularly insightful for understanding the impact of stress, sleep, and subtle food choices. And share your journal and glucose trends with your doctor or nutritionist so you can fine-tune your diet and treatment plan together.

Reframing Setbacks and Plateaus

The journey to healing neuropathy is rarely a straight line. There will be days when symptoms flare, progress seems stalled, or motivation wanes, and these are not failures, they are opportunities for learning and recalibration. It's okay to feel frustrated or discouraged, and you can acknowledge and validate those feelings without letting them consume you. Instead of asking "why me," try asking what you can learn from this setback, such as whether you deviated from your diet, whether stress was particularly high, or whether you missed sleep. These insights are invaluable. Try reframing the narrative, too: instead of "I'm stuck," think "this is a temporary challenge, and I have the tools to navigate it." Focus on consistency, not perfection, since one bad day or meal won't derail your entire progress, you simply get back on track with the next choice. And avoid the comparison trap. Everyone's healing journey is unique, so focus on your own personal improvements instead of comparing your progress to others.

Tracking Wins: Non-Scale Victories

While objective data, blood markers and symptom scores, is important, celebrating "non-scale victories" is crucial for maintaining motivation and recognizing subtle but significant improvements. These are the daily signs that your efforts are paying off, even if the primary symptoms haven't fully resolved. They can be as simple as noticing, "I walked an extra five minutes today without increased pain," "my sleep was more restful last night," "I noticed a slight return of sensation in my toes," "I navigated a stressful situation without my symptoms flaring," "I cooked a healthy meal even when I felt tired," "I've consistently taken my supplements for a week," "my energy levels feel more stable," or "I feel less anxious about my condition." Dedicate a section of your journal, or a separate notebook, to record these wins as a victory journal, no matter how small they seem. Reviewing them regularly can provide a powerful boost to your morale and reinforce positive habits.

Sample Blood Marker Tracking Chart

Beyond daily glucose, keeping track of your broader health markers over time allows you to see the impact of your holistic interventions. This data, combined with your symptom journal, provides a comprehensive

picture of your progress. The chart below is designed for you to fill in as you retest over the following months, alongside the optimal range your practitioner is working toward with you.

Marker	Initial Value (Date)	Optimal Range	Value (Month 1)	Value (Month 3)	Value (Month 6)	Notes / Changes
HbA1c		< 5.4%				
Fasting Insulin		< 5 µIU/mL				
hs-CRP		< 1.0 mg/L				
Homocysteine		5 to 7 µmol/L				Lower is not better; below this window suggests overmethylation
Vitamin D (25-OH D)		40 to 80 ng/mL				
Vitamin B12		> 500 pg/mL (functional)				Look at MMA and homocysteine for true deficiency
RBC Magnesium		Optimal range is lab dependent				
Omega-3 Index		> 8%				
Thyroid Stimulating Hormone (TSH)		1.0 to 2.5 µIU/mL				Also check Free T3, Free T4, and antibodies
Other specific markers (e.g., autoantibodies)		As per lab and diagnosis				

Always discuss these results and optimal ranges with your healthcare practitioner.

Cultivating a positive mindset and diligently monitoring your progress are not passive activities; they are active components of your healing journey. They empower you to become an engaged participant in your own recovery, helping you to adapt, persevere, and ultimately, unlock greater well-being.

Your Path to Reclaiming Nerve Health

You have covered a lot of ground on neuropathy, moving beyond symptom management to embrace a holistic, root-cause approach. The aim of this guide was to empower you with knowledge, practical strategies, and the understanding that your body possesses an incredible capacity for healing, given the right support.

We have unveiled five critical keys to unlocking your neuropathy, each representing a vital area for investigation and intervention. Key 1, unmasking nutrient deficiencies, recognizes that optimal nerve function relies on a precise symphony of vitamins and minerals, and that deficiencies can be silent saboteurs. Key 2, addressing chronic infections and their impact, is about understanding how hidden or persistent pathogens can ignite systemic inflammation and autoimmune responses that directly harm your nervous system. Then comes Key 3, activating large diameter afferent pathways, which is about learning to stimulate specific nerve fibers through various therapies, from vibration and tactile stimulation to cutting-edge photobiomodulation and specialized electrical stimulation devices, to modulate pain, improve sensation, and foster neuroplasticity. Key 4, comprehensive testing beyond the conventional, means advocating for a deeper diagnostic dive into inflammation, metabolic imbalances, circulation, and toxicity to pinpoint the precise underlying drivers of your condition. And Key 5, diet and food sensitivities as tools for fueling nerve repair, means harnessing the immense power of an anti-inflammatory, nutrient-dense diet and identifying personal food triggers to create an optimal internal environment for nerve healing.

Beyond these core pillars, we also highlighted the indispensable role of the movement plan in Chapter 7, promoting circulation, strengthening muscles, and directly impacting nerve health through exercise. And in mindset and monitoring, Chapter 8, we explored how emotional well-being, stress management, and diligent tracking are powerful, often overlooked, components of a successful recovery journey.

Embracing the Holistic Journey

Reclaiming your nerve health is not a sprint; it is a marathon. There will be days of progress, and there may be days of frustration. The most crucial takeaway is this: you are not passive in this process. Every choice you make, from the food you eat to the way you manage stress and the movement you incorporate, sends a message to your nervous system.

This ebook has provided you with a comprehensive framework. However, this is a guide, not a prescription. Your body is unique, and your healing journey will be too. We strongly advocate for working closely with a qualified healthcare professional, ideally a functional medicine practitioner, neurologist, or physical therapist experienced in holistic neuropathy approaches. They can help you interpret test results, fine-tune supplement protocols, and tailor exercise routines to your specific needs and condition.

As you move forward from here, a few next steps are worth taking deliberately. Review and reflect. Revisit the keys that resonated most with you, and consider which areas you suspect might be most relevant to your neuropathy. Consult a professional. Seek out a healthcare provider who understands and practices the root-cause approach, and share this guide with them as a starting point for discussion. Implement gradually. Don't try to change everything at once; pick one or two areas to focus on first, such as beginning with the dietary principles or integrating simple daily movements. And be patient and persistent. Healing takes time, so celebrate every small victory, learn from setbacks, and remain committed to nurturing your body.

Neuropathy doesn't have to be a life sentence of pain and limitation. By understanding its many contributing factors and proactively addressing the root causes, you have the power to unlock your neuropathy and reclaim a life of greater comfort, vitality, and well-being. This is your path forward.

Part 4: The 90-Day Battle Plan, Meal Plan and Recipes

Embarking on a new dietary path can feel overwhelming, but with a structured plan, it becomes an exciting journey of discovery and healing. This 90-day battle plan is designed to transition you into a sustainable, nerve-supportive eating lifestyle. It's about giving you choices that fuel repair, reduce inflammation, and stabilize blood sugar, rather than about strict deprivation.

The plan is divided into three phases, each building upon the last, allowing your body to adapt and for you to identify your unique needs and sensitivities. Remember, consistency, not perfection, is the goal.

Weekly Framework: The 60/20/20 Guideline

This general guideline provides a macronutrient breakdown for balanced meals that promote stable blood sugar and sustained energy throughout your day. About 60 percent of your plate should be healthy carbohydrates, coming primarily from a wide variety of non-starchy vegetables, some low-glycemic fruits like berries, and carefully selected whole grains if reintroduced and tolerated, focusing on complex carbohydrates rich in fiber. About 20 percent should be healthy fats, essential for nerve cell membranes and myelin, incorporating sources like extra virgin olive oil, avocados, nuts such as almonds and walnuts, seeds such as chia, flax, and hemp, and fatty fish such as wild-caught salmon, mackerel, and sardines, rich in anti-inflammatory omega-3 fatty acids. And the remaining 20 percent should come from quality proteins, crucial for tissue repair, muscle maintenance, and neurotransmitter synthesis, prioritizing grass-fed or pasture-raised meats, wild-caught fish, organic poultry, and plant-based options like legumes, if tolerated, and quinoa. This 60/20/20 guideline is a general starting point. Your individual needs may vary based on activity level, specific health conditions, and personal tolerance, and consulting with a nutritionist or functional medicine practitioner can help you fine-tune these percentages.

Phase 1: Weeks 1 Through 4, Elimination and Detoxification

This initial phase focuses on removing common inflammatory foods and potential sensitivities to calm your system, reduce inflammation, and create a clean slate for healing. The goal is to reduce inflammation, stabilize initial blood sugar responses, begin gut healing, and identify any immediate dietary triggers. Meals are built around abundant non-starchy vegetables, clean protein, and healthy fats. During this phase, minimize fruits and eliminate all grains, dairy, legumes, corn, soy, and refined sugars. Keep hydration front of mind: drink at least half your body weight in ounces of filtered water daily, and consider starting your day with a glass of warm lemon water.

Phase 2: Weeks 5 Through 8, Reintroduction and Personalization

After the elimination phase, your body has had a chance to reset. This phase is about carefully reintroducing foods, one at a time, to identify what specifically triggers your symptoms, with the goal of pinpointing

individual food sensitivities and expanding your personalized diet based on clear symptom responses. The reintroduction protocol works like this: choose one eliminated food group, for example gluten-free grains like quinoa, and eat a small portion of it daily for two to three days while continuing your otherwise clean diet. Monitor your symptoms closely, including pain, numbness, tingling, digestive issues, fatigue, and brain fog. If symptoms worsen, that food is likely a trigger for you, and you eliminate it again; if no symptoms appear after two to three days, you can likely include that food in your diet in moderation. Either way, wait at least three days before reintroducing the next food group. The menu philosophy continues with the anti-inflammatory base from phase one, but strategically adds foods back in based on your findings.

Phase 3: Weeks 9 Through 12, Maintenance and Optimization

By this phase, you will have a clearer understanding of your body's specific needs and sensitivities. This is where you solidify your personalized anti-inflammatory diet, with the goal of sustaining nerve healing, maintaining stable blood sugar, keeping inflammation low, and optimizing your overall well-being with a sustainable eating pattern. Enjoy a diverse range of your newly identified safe and supportive foods, continue to prioritize nutrient density and mindful eating, and keep hydration regular. This phase becomes your new normal.

A short list of snacks reliably stabilizes glucose between meals throughout all three phases, designed to be nutrient-dense and to prevent blood sugar spikes while keeping your energy stable and supporting nerve health: hard-boiled eggs for portable protein, avocado slices with a sprinkle of sea salt, a small handful of nuts such as almonds or walnuts or seeds such as pumpkin or sunflower, celery sticks with almond or seed butter, berries such as strawberries, blueberries, or raspberries with a dollop of unsweetened coconut yogurt or a few pecans, a small portion of olives, and a small piece of wild-caught salmon or grass-fed, sugar-free beef or turkey jerky.

Grocery List and Meal Prep Hacks

Streamlining your meal preparation is key to sticking with a new dietary plan. Keep these staples on hand.

Produce

- Leafy greens (spinach, kale)
- Cruciferous vegetables (broccoli, cauliflower)
- Asparagus
- Bell peppers
- Zucchini
- Cucumber
- Carrots
- Onions

- Garlic
- Ginger
- Turmeric root
- Berries
- Lemons
- Limes
- Avocados

Proteins

- Wild-caught salmon, cod, and mackerel
- Organic chicken breast and thighs
- Grass-fed beef (ground or steak)
- Organic eggs

Healthy Fats

- Extra virgin olive oil, coconut oil, and avocado oil
- Raw nuts (almonds, walnuts, pecans)
- Raw seeds (chia, flax, hemp, pumpkin, sunflower)

Pantry

- Apple cider vinegar
- Various anti-inflammatory herbs and spices
- Unsweetened almond or coconut milk
- Organic chicken or beef broth
- Gluten-free rolled oats and quinoa (for phase two or three, if tolerated)

A handful of habits make meal prep painless. Plan ahead by dedicating 30 to 60 minutes each week to plan your meals and snacks, which reduces decision fatigue during busy days. Shop smart by focusing on the perimeter of the grocery store, where fresh, whole foods are located, and sticking to your list. Batch cook proteins by cooking larger quantities of lean protein, baked chicken breasts, roasted salmon, or ground turkey, that can be easily added to salads, wraps, or stir-fries throughout the week. Roast a large tray of mixed non-starchy vegetables, broccoli, cauliflower, bell peppers, and zucchini, at the beginning of the week for quick additions to any meal. Wash and dry your leafy greens and pre-chop other vegetables so they're ready for quick assembly. Prepare homemade dressings in advance using olive oil, vinegar, lemon juice, and herbs. And portion out nuts, seeds, or berries into small containers for grab-and-go snacks.

Recipes for Nerve Healing

These fifteen to twenty easy, balanced recipes align with the principles of the anti-inflammatory, blood sugar-stabilizing diet. They are designed to be easy to prepare, delicious, and rich in nutrients vital for nerve health. For the first week, in phase one, focus on the recipes below that are naturally free of common sensitivities like grains, dairy, corn, soy, and legumes; they are marked Phase 1 Approved throughout.

Breakfast Recipes

1. Berry-Spinach Power Smoothie

Phase 1 approved.

This smoothie is packed with antioxidants from the berries and greens, healthy fats from the avocado and seeds, and clean protein, all while staying low glycemic and anti-inflammatory. It yields one serving and takes about five minutes to prepare.

Ingredients

- 1 cup unsweetened almond or coconut milk
- 1 large handful fresh spinach (or de-stemmed kale)
- 1/2 cup mixed berries, fresh or frozen
- 1/4 medium avocado
- 1 scoop unflavored or vanilla plant-based protein powder (no added sugars or artificial sweeteners)
- 1 tbsp chia seeds or flax seeds
- Optional: 1/4 tsp ground turmeric plus a pinch of black pepper

Instructions

1. Combine all ingredients in a high-speed blender.
2. Blend until smooth and creamy, adding more liquid if needed for your desired consistency.
3. Pour into a glass and enjoy immediately.

2. Turmeric Scramble with Avocado

Phase 1 approved.

Eggs are a complete protein source, and turmeric is a potent anti-inflammatory, while the healthy fats from avocado support nerve cell membranes. It yields one serving, takes about five minutes to prep, and seven to ten minutes to cook.

Ingredients

- 2 organic, pasture-raised eggs
- 1 tbsp unsweetened almond or coconut milk (optional, for creaminess)
- 1/2 cup chopped non-starchy vegetables (spinach, bell peppers, or zucchini)
- 1/2 tbsp olive oil or coconut oil
- 1/2 tsp ground turmeric
- Pinch of black pepper
- Sea salt to taste
- 1/2 medium avocado, sliced

Instructions

1. In a bowl, whisk the eggs with the almond milk (if using), turmeric, black pepper, and sea salt.
2. Heat the oil in a non-stick pan over medium heat. Add the chopped vegetables and sauté for 3 to 5 minutes until tender-crisp.
3. Pour the egg mixture into the pan and scramble gently until cooked through.
4. Serve immediately with the sliced avocado on the side.

3. Chia Seed Pudding with Berries and Nuts

Phase 1 approved.

Chia seeds are rich in omega-3s and fiber, promoting gut health and reducing inflammation, and this pudding is low glycemic and highly customizable. It yields one serving and takes about five minutes to prepare, plus at least four hours, or overnight, to chill.

Ingredients

- 3 tbsp chia seeds
- 1 cup unsweetened almond or coconut milk
- 1/2 tsp vanilla extract (sugar-free)
- Optional: pinch of cinnamon
- 1/4 cup fresh mixed berries
- 1 tbsp chopped walnuts or almonds

Instructions

1. In a jar or bowl, whisk together the chia seeds, almond milk, vanilla extract, and cinnamon until well combined.

2. Cover and refrigerate for at least 4 hours, or preferably overnight, until thickened to a pudding-like consistency. Stir once or twice in the first hour to prevent clumping.
3. Before serving, top with the fresh berries and chopped nuts.

Lunch Recipes

4. Giant Anti-Inflammatory Salad with Grilled Chicken or Salmon

Phase 1 approved.

This salad maximizes nutrient intake from diverse greens and vegetables, provides clean protein and healthy fats, and is highly anti-inflammatory. It yields one serving, takes about fifteen minutes to prep, and ten to fifteen minutes to cook the protein.

Ingredients

- 2 to 3 cups mixed leafy greens (spinach, kale, romaine, or spring mix)
- 1 cup chopped non-starchy vegetables (cucumber, bell peppers, shredded carrots, or radish)
- 4 to 5 oz grilled organic chicken breast, wild-caught salmon fillet, or 1 can wild salmon or sardines
- 1/4 medium avocado, sliced or diced
- 1/4 cup toasted pumpkin seeds or sunflower seeds
- **For the lemon-herb vinaigrette:**
 - 2 tbsp extra virgin olive oil
 - 1 tbsp fresh lemon juice
 - 1/2 tsp dried Italian herbs or fresh chopped parsley or dill
 - Pinch of sea salt and black pepper

Instructions

1. Prepare your protein: grill the chicken or salmon, or drain the canned fish. Let it cool slightly and flake the salmon or sardines if using.
2. In a large bowl, combine the mixed greens and chopped vegetables.
3. Arrange the protein and avocado slices on top, then sprinkle with the toasted seeds.
4. In a small bowl, whisk together the olive oil, lemon juice, herbs, salt, and pepper for the vinaigrette.
5. Drizzle the dressing over the salad just before serving.

5. Turkey and Avocado Lettuce Wraps

Phase 1 approved.

This is a light, grain-free option packed with lean protein, healthy fats, and crisp vegetables, avoiding inflammatory grains entirely. It yields one serving, or two to three wraps, takes ten to fifteen minutes to prep, and five to seven minutes to cook the turkey.

Ingredients

- 4 to 5 large butter lettuce leaves or romaine lettuce hearts
- 4 oz lean ground turkey or organic ground chicken
- 1/2 cup finely diced bell peppers (any color)
- 1/4 cup celery, finely diced
- 1/4 cup cucumber, finely diced
- 1/4 medium avocado, mashed
- 1 tbsp olive oil
- 1 tsp apple cider vinegar
- 1/2 tsp garlic powder or fresh minced garlic
- Pinch of red pepper flakes (optional, for a kick)
- Sea salt and black pepper to taste

Instructions

1. Heat the olive oil in a skillet over medium heat. Add the ground turkey or chicken and cook, breaking it up with a spoon, until browned and cooked through. Drain any excess fat.
2. Remove the cooked turkey from the heat and let it cool slightly.
3. In a bowl, combine the cooled turkey with the diced bell peppers, celery, cucumber, mashed avocado, apple cider vinegar, garlic powder, red pepper flakes (if using), salt, and pepper. Mix well.
4. Spoon the mixture into the lettuce leaves and wrap them up. Serve immediately.

Dinner Recipes

6. Baked Salmon with Roasted Asparagus and Brussels Sprouts

Phase 1 approved.

Wild salmon is rich in omega-3s, EPA and DHA, crucial for nerve health and reducing inflammation, while the roasted vegetables provide fiber and antioxidants. It yields two servings, takes about ten minutes to prep, and twenty to twenty-five minutes to cook.

Ingredients

- 2 wild-caught salmon fillets (4 to 5 oz each)

- 1 bunch asparagus, woody ends trimmed
- 1 cup Brussels sprouts, halved
- 2 tbsp olive oil, divided
- 1 lemon, half sliced and half reserved for juicing
- 2 cloves garlic, minced (or 1 tsp garlic powder)
- 1/2 tsp dried dill (optional)
- Sea salt and black pepper to taste

Instructions

1. Preheat the oven to 400°F (200°C) and line a baking sheet with parchment paper.
2. On the baking sheet, toss the asparagus and Brussels sprouts with 1 tbsp of the olive oil, the minced garlic, salt, and pepper. Spread in a single layer.
3. Place the salmon fillets on the same baking sheet, in among the vegetables. Drizzle the salmon with the remaining 1 tbsp olive oil, sprinkle with dill (if using), salt, and pepper. Place lemon slices on top of each fillet.
4. Bake for 15 to 20 minutes, or until the salmon is cooked through and flakes easily and the vegetables are tender-crisp.
5. Serve immediately with a squeeze of fresh lemon juice.

7. Chicken and Vegetable Stir-Fry with Cauliflower Rice

Phase 1 approved.

This is a quick, versatile meal packed with protein and a rainbow of vegetables, and the cauliflower rice makes it a low-carb, grain-free alternative to a traditional stir-fry. It yields two servings and takes about fifteen minutes to prep and fifteen minutes to cook.

Ingredients

- 1 tbsp coconut oil or avocado oil
- 1 lb organic chicken breast or thighs, cut into 1-inch pieces
- 2 cups mixed stir-fry vegetables (broccoli florets, sliced bell peppers, shredded carrots, snap peas, sliced mushrooms, bok choy)
- 1 clove garlic, minced
- 1 tbsp fresh ginger, grated
- 2 cups cauliflower rice, fresh or frozen
- **For the sugar-free stir-fry sauce:**

- 2 tbsp coconut aminos
- 1 tbsp apple cider vinegar
- 1 tsp sesame oil (optional, for flavor)
- 1/4 cup filtered water or broth
- Pinch of red pepper flakes (optional)

Instructions

1. Heat the coconut oil in a large skillet or wok over medium-high heat. Add the chicken and cook until browned and cooked through, about 5 to 7 minutes. Remove from the pan and set aside.
2. Add the garlic and ginger to the pan and stir-fry for 30 seconds until fragrant.
3. Add the harder vegetables (broccoli and carrots) and stir-fry for 3 to 4 minutes. Then add the softer vegetables (bell peppers, mushrooms, snap peas, and bok choy) and stir-fry for another 3 minutes until everything is tender-crisp.
4. In a small bowl, whisk together all the stir-fry sauce ingredients.
5. Return the chicken to the pan with the vegetables. Add the cauliflower rice and pour the sauce over everything. Stir well to combine and heat through, about 2 to 3 minutes.
6. Serve hot.

8. Lean Ground Beef and Broccoli Bowls

Phase 1 approved.

This bowl provides quality protein and healthy fats, and broccoli brings its detoxifying, anti-inflammatory power to the plate. It yields two servings, takes about ten minutes to prep, and fifteen minutes to cook.

Ingredients

- 1 tbsp olive oil
- 1 lb grass-fed lean ground beef
- 1 large head broccoli, cut into small florets
- 1/2 onion, finely diced (optional)
- 2 cloves garlic, minced
- **For the sauce:**
- 3 tbsp coconut aminos
- 1 tbsp apple cider vinegar
- 1/2 tsp fresh grated ginger (optional)
- Pinch of red pepper flakes (optional)

- Sea salt and black pepper to taste

Instructions

1. Heat the olive oil in a large skillet or deep pan over medium-high heat. Add the ground beef and onion (if using) and cook, breaking up the meat with a spoon, until browned. Drain any excess fat.
2. Add the minced garlic and ginger and cook for 1 minute until fragrant.
3. Add the broccoli florets and cook, stirring occasionally, for 5 to 7 minutes until tender-crisp.
4. In a small bowl, whisk together the coconut aminos, apple cider vinegar, and red pepper flakes (if using).
5. Pour the sauce over the beef and broccoli. Stir well to coat and cook for another 2 to 3 minutes until the sauce slightly thickens.
6. Season with salt and pepper to taste and serve hot.

9. Quinoa Salad with Roasted Vegetables

Phase 2 or 3 approved, if quinoa is tolerated.

Quinoa is a complete protein and provides complex carbohydrates and fiber, while the roasted vegetables add antioxidants; make sure quinoa is tolerated after your reintroduction phase before relying on this one. It yields two servings, takes about fifteen minutes to prep, and twenty to twenty-five minutes to cook.

Ingredients

- 1 cup cooked quinoa, rinsed well before cooking
- 1 cup chopped mixed vegetables for roasting (zucchini, bell peppers, red onion, broccoli florets)
- 1 tbsp olive oil
- Pinch of sea salt and black pepper
- 1/2 cup chopped fresh parsley or cilantro
- Optional: 1/4 cup crumbled feta or goat cheese (Phase 3, if dairy is tolerated)
- **For the lemon-tahini dressing:**
 - 2 tbsp tahini
 - 2 tbsp fresh lemon juice
 - 1 tbsp water, or more to thin
 - 1 clove garlic, minced
 - Pinch of sea salt

Instructions

1. Preheat the oven to 400°F (200°C). Toss the chopped vegetables with the olive oil, salt, and pepper on a baking sheet. Roast for 15 to 20 minutes, or until tender and slightly caramelized.
2. In a large bowl, combine the cooked quinoa, roasted vegetables, and fresh herbs.
3. For the dressing, whisk together the tahini, lemon juice, water, minced garlic, and salt until smooth. Add more water if needed to reach your desired consistency.
4. Pour the dressing over the quinoa and vegetable mixture and toss to combine.
5. Serve warm or at room temperature, topped with the crumbled cheese if you're including it.

10. Lentil Soup with Root Vegetables

Phase 2 or 3 approved, if legumes are tolerated.

Lentils provide plant-based protein and fiber, promoting stable blood sugar and gut health, while the root vegetables add further nutrients; this one is for once legumes have been successfully reintroduced. It yields four servings, takes about fifteen minutes to prep, and forty-five to sixty minutes to cook.

Ingredients

- 1 tbsp olive oil
- 1 medium onion, diced
- 2 carrots, diced
- 2 celery stalks, diced
- 2 cloves garlic, minced
- 1 cup green or brown lentils, rinsed
- 6 cups vegetable broth, low sodium, no sugar added
- 1 can (14.5 oz) diced tomatoes, no sugar added, BPA-free can
- 1 tsp dried thyme
- 1/2 tsp dried rosemary
- Sea salt and black pepper to taste
- Fresh parsley for garnish (optional)

Instructions

1. Heat the olive oil in a large pot or Dutch oven over medium heat. Add the onion, carrots, and celery and sauté for 8 to 10 minutes until softened.
2. Add the minced garlic and cook for another minute until fragrant.
3. Stir in the rinsed lentils, vegetable broth, diced tomatoes, thyme, and rosemary. Bring to a boil, then reduce the heat to low, cover, and simmer for 30 to 45 minutes, or until the lentils are tender.

4. Season with sea salt and black pepper to taste.
5. Ladle into bowls and garnish with fresh parsley if desired.

Dinner Recipes, Continued

11. Baked Cod with Lemon-Garlic Herbs

Phase 1 approved.

Cod is a lean, easily digestible protein source, and the lemon and garlic add flavor without any inflammatory ingredients. It yields two servings, takes about ten minutes to prep, and twelve to fifteen minutes to cook.

Ingredients

- 2 cod fillets (6 oz each), or another white fish like halibut or tilapia
- 2 tbsp olive oil
- 2 cloves garlic, minced
- 1/2 lemon, thinly sliced
- 1 tbsp fresh chopped parsley (or 1 tsp dried)
- 1/2 tsp dried oregano
- Sea salt and black pepper to taste

Instructions

1. Preheat the oven to 400°F (200°C) and line a baking sheet with parchment paper. Place the cod fillets on the sheet.
2. In a small bowl, whisk together the olive oil, minced garlic, chopped parsley, oregano, salt, and pepper.
3. Pour the mixture over the cod fillets so they are well coated, and top each fillet with a few lemon slices.
4. Bake for 12 to 15 minutes, or until the fish is opaque and flakes easily with a fork.
5. Serve immediately with your favorite non-starchy steamed or roasted vegetables.

12. Turkey and Vegetable Chili, No Beans

Phase 1 approved.

This is a hearty, warming meal packed with lean protein and fiber from vegetables, highly customizable and anti-inflammatory without the typical bean load. It yields four to six servings, takes about fifteen minutes to prep, and thirty to forty minutes to cook.

Ingredients

- 1 tbsp olive oil

- 1 lb organic lean ground turkey
- 1 large onion, diced
- 2 celery stalks, diced
- 1 large bell pepper, any color, diced
- 2 cloves garlic, minced
- 1 can (28 oz) crushed tomatoes, no sugar added, BPA-free
- 1 can (14.5 oz) diced tomatoes, no sugar added, BPA-free
- 1 cup beef or vegetable broth, low sodium
- 2 tbsp chili powder, no sugar added
- 1 tsp ground cumin
- 1/2 tsp smoked paprika
- Pinch of cayenne pepper (optional, for heat)
- Sea salt and black pepper to taste
- Optional toppings (sliced avocado, fresh cilantro, green onions)

Instructions

1. Heat the olive oil in a large pot or Dutch oven over medium heat. Add the ground turkey and cook, breaking it up, until browned. Drain any excess fat and set the turkey aside.
2. Add the onion, celery, and bell pepper to the pot and sauté for 8 to 10 minutes until softened.
3. Add the minced garlic and cook for another minute until fragrant.
4. Return the turkey to the pot and stir in the crushed tomatoes, diced tomatoes, beef broth, chili powder, cumin, smoked paprika, and cayenne (if using).
5. Bring the chili to a simmer, then reduce the heat to low, cover, and cook for at least 20 to 30 minutes for the flavors to meld.
6. Season with salt and pepper to taste.
7. Serve hot, topped with sliced avocado, fresh cilantro, or green onions if desired.

13. Sheet Pan Chicken Fajita Bowls, No Tortilla

Phase 1 approved.

This is a simple, flavorful meal with lean protein and plenty of colorful, anti-inflammatory vegetables, and it's great for meal prepping. It yields two servings, takes about fifteen minutes to prep, and twenty-five to thirty minutes to cook.

Ingredients

- 1 lb organic chicken breast, sliced into thin strips
- 2 bell peppers, different colors, sliced
- 1 medium onion, sliced
- 2 tbsp olive oil
- **For the sugar-free fajita seasoning:**
- 1 tbsp chili powder
- 1 tsp ground cumin
- 1 tsp smoked paprika
- 1/2 tsp garlic powder
- 1/2 tsp onion powder
- Pinch of cayenne pepper (optional)
- Sea salt and black pepper to taste
- Optional for serving (sliced avocado, fresh cilantro, lime wedges)

Instructions

1. Preheat the oven to 400°F (200°C) and line a large baking sheet with parchment paper.
2. In a large bowl, combine the sliced chicken, bell peppers, and onion. Drizzle with the olive oil.
3. In a small bowl, mix all the fajita seasoning ingredients. Sprinkle the seasoning over the chicken and vegetables, tossing well to coat evenly.
4. Spread everything in a single layer on the prepared baking sheet.
5. Roast for 25 to 30 minutes, flipping halfway through, until the chicken is cooked through and the vegetables are tender-crisp and slightly charred.
6. Serve hot in bowls, topped with sliced avocado, fresh cilantro, and a squeeze of fresh lime juice.

Snack and Side Recipes

14. Roasted Brussels Sprouts with Balsamic Glaze

Phase 1 approved.

Brussels sprouts are a cruciferous vegetable, rich in compounds that support detoxification and reduce inflammation. It yields two servings, takes about ten minutes to prep, and twenty to twenty-five minutes to cook.

Ingredients

- 1 lb Brussels sprouts, trimmed and halved

- 1 tbsp olive oil
- Sea salt and black pepper to taste
- 2 tbsp balsamic vinegar, no added sugar

Instructions

1. Preheat the oven to 400°F (200°C) and line a baking sheet with parchment paper.
2. In a bowl, toss the Brussels sprouts with the olive oil, salt, and pepper until evenly coated.
3. Spread the Brussels sprouts in a single layer on the baking sheet and roast for 20 to 25 minutes, stirring halfway through, until tender inside and crisp and browned outside.
4. While the sprouts are roasting, pour the balsamic vinegar into a small saucepan. Bring it to a simmer over medium-low heat and cook for 5 to 7 minutes, stirring occasionally, until it reduces to a slightly thickened glaze. Be careful not to let it burn.
5. Once the sprouts are roasted, transfer them to a serving dish and drizzle with the balsamic glaze. Serve immediately.

15. Simple Guacamole with Veggie Sticks

Phase 1 approved.

Avocados are packed with healthy monounsaturated fats, crucial for nerve cell health, while the tomato, if tolerated, and onion add flavor and antioxidants; omit the tomato for a stricter phase one if nightshades are a concern. It yields two to three servings and takes about ten minutes to prep.

Ingredients

- 2 ripe avocados
- 1/4 cup red onion, finely diced (or green onion for a milder flavor)
- 1/4 cup tomato, finely diced (optional, omit for stricter Phase 1 if nightshades are a concern)
- 1 to 2 tbsp fresh lime juice
- 1 tbsp fresh chopped cilantro (optional)
- Pinch of red pepper flakes (optional)
- Sea salt to taste
- Cucumber slices, bell pepper strips, celery sticks and radish slices, for dipping

Instructions

1. In a medium bowl, mash the avocados with a fork to your desired consistency, whether chunky or smooth.
2. Add the diced red onion, tomato (if using), lime juice, chopped cilantro (if using), red pepper flakes (if using), and sea salt.

3. Mix gently to combine. Taste and adjust the seasoning as needed.
4. Serve immediately with the vegetable sticks for dipping.

16. Anti-Inflammatory Bone Broth

Phase 1 approved.

Rich in collagen, amino acids like glycine and proline, and minerals, bone broth supports gut health, reduces inflammation, and provides building blocks for tissue repair, including, potentially, nerve sheaths. Yields vary, since this makes a large batch; it takes about fifteen minutes to prep and eight to twenty-four hours to simmer.

Ingredients

- 2 to 3 lbs organic, grass-fed beef bones (marrow, knuckle, or joint bones), or organic chicken carcass and bones
- 1 gallon filtered water
- 2 tbsp apple cider vinegar (helps extract minerals)
- 1 large onion, quartered, skin on
- 2 carrots, roughly chopped
- 2 celery stalks, roughly chopped
- 2 bay leaves
- 1 tsp black peppercorns
- Optional, for extra flavor and nutrients (fresh ginger slices, turmeric root slices, parsley stems)

Instructions

1. If using beef bones, roast them first for richer flavor: preheat the oven to 400°F (200°C) and roast the bones on a baking sheet for 20 to 30 minutes until browned. Skip this step for chicken bones.
2. Place the bones in a large stock pot or slow cooker. Add the apple cider vinegar and cover with the filtered water. Let sit for 30 minutes before heating, since this helps draw out the minerals.
3. Add the onion, carrots, celery, bay leaves, peppercorns, and any optional additions you're using.
4. Stovetop method: bring the pot to a rolling boil, then immediately reduce the heat to the lowest simmer. Skim off any foam that rises to the surface during the first hour. Cover and simmer for at least 8 to 12 hours for chicken bones, or 12 to 24 for beef, the longer the better.
5. Slow cooker method: set it to low and cook for 12 to 24 hours instead.
6. Once done, remove the large bones and vegetables with tongs or a slotted spoon.
7. Strain the broth through a fine-mesh sieve or a cheesecloth-lined colander into clean jars or containers.

8. Let cool completely before refrigerating. The fat will solidify on top; you can skim it off or leave it.
9. Store in the fridge for up to 5 days, or freeze for up to 6 months. Sip warm plain or use as a base for soups.

Your Next Step: A Free Root Discovery Session

This ebook has shown you the what and the why behind your neuropathy, from the nutrient gaps and hidden infections to the nerve pathways, the testing, and the food choices that all play a part. That understanding alone puts you ahead of where you started, and it is real progress.

Finding your specific root cause, though, takes real testing and a plan built around your body, not a generic one. If you are ready to find out what is actually driving your neuropathy, the next step is a free, no-pressure conversation with me. We will talk through your history, your symptoms, and what a personalized path forward could look like for you. [Book your free Root Discovery Session here.](#) You can also reach the office at (919) 662-0520.

Consultations and ongoing care are both available wherever you are located. Being outside North Carolina is not a barrier to working with me. Tell me where you are when you reach out and I will lay out exactly how we would work together from there.

If you would rather start on your own, begin with the 90-day plan in this ebook and log your symptoms against your own baseline so the pattern is on paper before we talk.

Dr. Adam Seay, Root Health, Garner NC

For educational purposes only. This ebook does not constitute medical advice, diagnosis, or treatment, and reading it does not establish a patient-provider relationship.

References

1. Alvarez et al., "B Vitamin Deficiencies and Associated Neuropathies," *Current Nutrition Reports*, 2026. <https://pubmed.ncbi.nlm.nih.gov/41609902/>
2. Muhamad et al., "The Role of Vitamin B6 in Peripheral Neuropathy: A Systematic Review," *Nutrients*, 2023. <https://pubmed.ncbi.nlm.nih.gov/37447150/>
3. Koike et al., "Clinicopathologic features of folate-deficiency neuropathy," *Neurology*, 2015. <https://pubmed.ncbi.nlm.nih.gov/25663227/>
4. Sepassi et al., "Associations between long-term metformin use, vitamin B12 deficiency, and neuropathy," *Diabetes Research and Clinical Practice*, 2025. <https://pubmed.ncbi.nlm.nih.gov/40850508/>
5. Gilbody and Gilbody, "Vitamin D for Painful Diabetic Neuropathy: A Systematic Review and Meta-Analysis," *Endocrinology, Diabetes & Metabolism*, 2025. <https://pubmed.ncbi.nlm.nih.gov/41139799/>
6. Shin et al., "Magnesium and Pain," *Nutrients*, 2020. <https://pubmed.ncbi.nlm.nih.gov/32718032/>
7. Han et al., "A systematic review and meta-analysis of alpha-lipoic acid in the treatment of diabetic peripheral neuropathy," *European Journal of Endocrinology*, 2012. <https://pubmed.ncbi.nlm.nih.gov/22837391/>
8. Li et al., "Acetyl-L-carnitine in the treatment of peripheral neuropathic pain: a systematic review and meta-analysis of randomized controlled trials," *PLoS One*, 2015. <https://pubmed.ncbi.nlm.nih.gov/25751285/>
9. Zhang et al., "Omega-3 polyunsaturated fatty acid oral supplements for improving peripheral nerve health: a systematic review and meta-analysis," *Nutrition Reviews*, 2020. <https://pubmed.ncbi.nlm.nih.gov/31532492/>
10. Halperin, "Lyme disease and the peripheral nervous system," *Muscle & Nerve*, 2003. <https://pubmed.ncbi.nlm.nih.gov/12872317/>
11. "Peripheral neuropathy as a complication of neuroborreliosis," *International Journal of Infectious Diseases*, 2021. <https://pubmed.ncbi.nlm.nih.gov/33684563/>
12. "Bartonella- and Borrelia-related disease presenting as a neurological condition," *Microorganisms*, 2024. <https://pubmed.ncbi.nlm.nih.gov/38276194/>
13. Mallick-Searle et al., "Postherpetic neuralgia: epidemiology, pathophysiology, and pain management pharmacology," *Journal of Multidisciplinary Healthcare*, 2016. <https://pubmed.ncbi.nlm.nih.gov/27703368/>
14. Gavin et al., "Cranial and peripheral neuropathy due to Epstein-Barr virus infection," *Postgraduate Medical Journal*, 1997. <https://pubmed.ncbi.nlm.nih.gov/9338028/>
15. Nemni et al., "Peripheral neuropathy in hepatitis C virus infection with and without cryoglobulinaemia," *Journal of Neurology, Neurosurgery & Psychiatry*, 2003. <https://pubmed.ncbi.nlm.nih.gov/12933932/>
16. Pane et al., "Role of gut microbiota in neuropathy and neuropathic pain states," *Neurobiology of Disease*, 2022. <https://pubmed.ncbi.nlm.nih.gov/35623598/>
17. Daxboeck et al., "Mycoplasma pneumoniae central nervous system infections," *Current Opinion in Neurology*, 2006. <https://pubmed.ncbi.nlm.nih.gov/16914976/>
18. "Low level laser therapy for painful diabetic peripheral neuropathy," *Diabetes & Metabolic Syndrome*, 2019. <https://pubmed.ncbi.nlm.nih.gov/31405692/>
19. Holanda et al., "The mechanistic basis for photobiomodulation therapy of neuropathic pain," *Lasers in Surgery and Medicine*, 2017. <https://pubmed.ncbi.nlm.nih.gov/28075022/>

20. Sommer et al., "Inflammation in the pathophysiology of neuropathic pain," *Pain*, 2018. <https://pubmed.ncbi.nlm.nih.gov/29447138/>
21. Li et al., "The relationship between plasma homocysteine levels and diabetic peripheral neuropathy," *Zhonghua Nei Ke Za Zhi*, 2011. <https://pubmed.ncbi.nlm.nih.gov/21418881/>
22. Kirthi et al., "Prevalence of peripheral neuropathy in pre-diabetes: a systematic review," *BMJ Open Diabetes Research & Care*, 2021. <https://pubmed.ncbi.nlm.nih.gov/34006607/>
23. Bourque, "Autoimmune peripheral neuropathies," *Clinica Chimica Acta*, 2015. <https://pubmed.ncbi.nlm.nih.gov/25748038/>
24. Koszewicz et al., "The impact of chronic co-exposure to different heavy metals on small fibers of peripheral nerves. A study of metal industry workers," *Journal of Occupational Medicine and Toxicology*, 2021. <https://pubmed.ncbi.nlm.nih.gov/33858451/>
25. Hadjivassiliou et al., "Neuropathy associated with gluten sensitivity," *Journal of Neurology, Neurosurgery, and Psychiatry*, 2006. <https://pubmed.ncbi.nlm.nih.gov/16835287/>
26. Knaepen et al., "Neuroplasticity, exercise-induced response of peripheral brain-derived neurotrophic factor: a systematic review of experimental studies in human subjects," *Sports Medicine*, 2010. <https://pubmed.ncbi.nlm.nih.gov/20726622/>
27. Zhang et al., "Exercise for Neuropathic Pain: A Systematic Review and Expert Consensus," *Frontiers in Medicine*, 2021. <https://pubmed.ncbi.nlm.nih.gov/34901069/>