



ROOT CAUSE UNLOCKED · ULTIMATE GUIDE

Unlock Your *Neuropathy*, Ultimate Guide

A Holistic Guide to Reclaiming Nerve Health

A guide for anyone living with numbness, tingling, or burning in the hands and feet who has been offered medication for the symptom and never an explanation for the cause.

Dr. Adam Seay, DC

Root Cause Unlocked, Garner, NC · rootcauseunlocked.com · (919) 662-0520

Unlock Your Neuropathy, Ultimate Guide

A Holistic Guide to Reclaiming Nerve Health

BEFORE YOU START

Contents

INTRODUCTION

Stepping Beyond the Symptoms

Chapter 1: The Foundation, Understanding Neuropathy Beyond the Symptoms

Chapter 2: Key 1, Unmasking Nutrient Deficiencies

Chapter 3: Key 2, Addressing Chronic Infections and Their Impact

Chapter 4: Key 3, Activating Large Diameter Afferent Pathways

Chapter 5: Key 4, Comprehensive Testing Beyond the Conventional

Chapter 6: Key 5, Diet and Food Sensitivities for Nerve Repair

Chapter 7: The Movement Plan, Restoring Function and Reducing Pain

Chapter 8: Mindset and Monitoring, the Unseen Influencers

CONCLUSION

Your Path to Reclaiming Nerve Health

Part 4: The 90-Day Battle Plan, Meal Plan and Recipes

Your Next Step: A Free Root Discovery Session

References

From Understanding to Doing

What This Guide Is For

Panel Overview: Which Labs for Neuropathy Specifically, and Why

Before You Read: Four Rules for Interpreting These Results

Section 1: Neural Zoomer Plus, Peripheral Neuropathy Category

Section 2: Neural Zoomer Plus, Demyelination Category

Section 3: Systemic Context, Blood-Brain Barrier and Infections Categories

Section 4: Organic Acids Test, Markers Relevant to Nerve Symptoms

Section 5: Metabolic and Nutrient Panels Beyond the Zoomer Panels

Section 6: Additional Panels Worth Knowing About

Section 7: Testing Decision Framework

Section 8: Reading Your Results Together, Pattern Recognition

When to Seek Clinical Review

Track What Changes, From Anywhere

What This Guide Does Not Cover

Read this part first

What you are actually working toward

What the evidence actually shows, including the part usually left out

Part one: blood sugar, the biggest lever you have

Part two: alcohol, which is a cause and not a lifestyle nag

Part three: gluten, where the connection holds up but is narrower than the internet suggests

Part four: the foundation pattern

Part four and a half: absorption, which is where diet quietly fails

Part four and three quarters: eating for sleep, because sleep sets your pain threshold

Part five: the specific experiments worth running

Part six: a week you can actually cook

Running a fair diet experiment

Realistic expectations

When to seek professional care

Working with Root Health

References

Read this part first

Why nutrient evidence looks thinner than it is

What you are actually buying back

Supplements are supportive, not the main event

How to read this guide

Before you start: the medication conversation

The vitamin B6 problem, which most neuropathy formulas get wrong

Alpha-lipoic acid

Vitamin B12, but only if you are actually low

Vitamin D, if your level is low

Magnesium

Omega-3 fatty acids

What I deliberately left out, and why

Running a fair 12-week trial

Realistic expectations

When to seek professional care

My Favorite Support Products

Working with Root Health

References

Stepping Beyond the Symptoms

The persistent tingling, numbness, burning, and sharp pains of neuropathy can feel like an invisible cage, limiting your life and stealing your comfort. For millions worldwide, peripheral neuropathy isn't just a symptom, it's a chronic challenge that profoundly impacts daily activities, sleep, and overall quality of life. Whether you've been diagnosed with diabetic neuropathy, idiopathic neuropathy, or a specific form like post-herpetic neuralgia, the conventional approach often focuses on managing symptoms with medications that may come with their own set of side effects, without truly addressing the underlying cause.

But what if, instead of simply masking the discomfort, we could uncover the root imbalances contributing to your nerve damage and dysfunction? This ebook is your guide to that path, toward understanding, addressing, and potentially reversing the progression of your neuropathy.

Traditional medicine excels at acute care and crisis intervention, but chronic conditions like neuropathy often require a detective's eye and a comprehensive strategy. We believe that true healing comes from identifying and addressing the why behind your symptoms. Your body is an interconnected system, and nerve health is profoundly influenced by everything from your diet and nutrient status to your gut health, immune system, and even your stress levels.

In this guide, we will explore five fundamental “keys” to unlocking your neuropathy. These keys represent critical areas often overlooked in standard care, yet they hold immense power to influence nerve health and regeneration. The first key is unmasking nutrient deficiencies: discovering the vital vitamins and minerals essential for nerve function that may be missing from your diet or not adequately absorbed. The second is addressing chronic infections and their impact, understanding how hidden or persistent infections can trigger inflammation and autoimmune responses that attack your nerves. The third is activating large diameter afferent pathways, learning how to stimulate specific nerve fibers to improve sensory input, modulate pain, and re-educate your nervous system. The fourth is comprehensive testing that goes beyond the conventional, identifying the advanced diagnostics that can reveal deeper insights into inflammation, metabolic imbalances, circulation issues, and more. And the fifth is diet and food sensitivities, fueling nerve repair by harnessing the power of nutrition to reduce inflammation, stabilize blood sugar, and provide the building blocks for nerve regeneration.

This journey requires an open mind and a commitment to self-exploration. By the end of this ebook, you will be equipped with practical strategies, research-backed insights, and a detailed 90-day plan to embark on your path toward reclaiming nerve health and unlocking a life with less pain and greater vitality. It's time to stop just coping and start truly healing.

Chapter 1: The Foundation, Understanding Neuropathy Beyond the Symptoms

To truly unlock your neuropathy, we must first lay a solid foundation of understanding. Neuropathy is a term that describes damage to the peripheral nervous system, the vast network of nerves outside of your brain and spinal cord, rather than a single disease. These nerves are the communication highways that carry messages between your central nervous system and every other part of your body, including your muscles, skin, and internal organs.

When these nerves are damaged, their ability to transmit signals is compromised, leading to a wide array of symptoms, and the type of symptoms you experience depends on which nerves are affected. Sensory nerves transmit sensations like touch, pain, temperature, and vibration; damage to them often leads to numbness, tingling, burning, shooting pains, or an exaggerated sensitivity to touch, and this is the most common type of neuropathy. Motor nerves control muscle movement, and damage to them can result in muscle weakness, cramps, spasms, loss of balance, or difficulty with fine motor skills. Autonomic nerves control involuntary bodily functions like heart rate, blood pressure, digestion, bladder function, and sweating, so autonomic neuropathy can cause symptoms such as dizziness upon standing, digestive issues, bladder control problems, or abnormal sweating.

Defining Neuropathy: A Deeper Dive into Types

While symptoms can overlap, understanding the different classifications of neuropathy can help pinpoint potential causes. Peripheral neuropathy is the most common form, affecting nerves in the extremities, the feet, legs, hands, and arms; it often presents as a “stocking-glove” pattern, meaning symptoms begin in the toes and fingers and spread upwards. Diabetic neuropathy, the most prevalent cause of neuropathy worldwide, results from prolonged high blood sugar levels that damage nerve fibers; it often manifests as peripheral neuropathy but can also affect autonomic nerves. When no clear cause can be identified, it’s labeled idiopathic neuropathy. However, this simply means the underlying cause hasn’t been found yet, and our holistic approach is particularly vital here, as we aim to uncover those hidden factors. Compressive neuropathy is caused by pressure on a nerve, such as carpal tunnel syndrome or sciatica. Inflammatory neuropathy describes conditions like Guillain-Barré syndrome or chronic inflammatory demyelinating polyneuropathy (CIDP), where the immune system mistakenly attacks nerve tissue. And toxin-induced neuropathy is damage from exposure to heavy metals such as lead or mercury, certain industrial chemicals, or even some medications such as chemotherapy drugs.

The Root Cause Approach: Beyond Symptom Management

The conventional medical model often focuses on symptom management, with pain relievers for pain, anti-depressants for burning sensations, or medications to manage blood sugar in diabetics. While these

approaches can offer temporary relief, they rarely address why the nerves are being damaged in the first place. This is where the root cause approach comes in.

Imagine a leaky roof in your house. You could continually mop up the puddles, which is symptom management, but the leak would persist. To truly solve the problem, you need to find the hole in the roof and patch it: that is the root cause approach. Similarly, with neuropathy, understanding the underlying mechanisms of nerve damage is paramount. These mechanisms often involve chronic inflammation, a persistent low-grade inflammatory state that can damage nerve cells and their protective myelin sheaths; oxidative stress, an imbalance between free radicals and antioxidants that can lead to cellular damage, including nerve cells; poor circulation, since nerves require a steady supply of oxygen and nutrients and compromised blood flow can starve nerve tissue; metabolic imbalances, where high blood sugar, insulin resistance, and imbalanced lipid profiles can directly harm nerves; immune dysfunction, where autoimmune attacks on nerve tissue or chronic activation of the immune system by infections can lead to nerve damage; nutrient deficiencies, since nerves have high metabolic demands and require specific nutrients for their structure, function, and repair; and toxic burden, where exposure to environmental toxins or heavy metals can directly poison nerve cells.

By systematically investigating these areas, we can move beyond the “idiopathic” label and empower you with knowledge and actionable strategies. The upcoming chapters work through each of our five “keys,” providing you with the tools to identify potential root causes in your own body and implement a comprehensive plan for healing and restoring nerve health.

Chapter 2: Key 1, Unmasking Nutrient Deficiencies

Our nerves are not static wires; they are living, dynamic tissues with high metabolic demands. They constantly require a precise array of vitamins, minerals, and other micronutrients to build and maintain their structure, transmit signals efficiently, and repair themselves. When even one of these essential building blocks is in short supply, nerve function can falter, leading to the symptoms of neuropathy.

It's a common misconception that a "standard" diet provides all the necessary nutrients. Soil depletion, modern food processing, and individual genetic variations or medical conditions can all contribute to insidious nutrient deficiencies that silently undermine nerve health. This chapter covers what specific nutrients do for your nerves and how to identify and address common deficiencies linked to neuropathy.

The Vital Role of Nutrients for Nerve Health

Think of your nervous system as a complex electrical circuit, with nutrients acting as the insulation, the conductors, the energy source, and the repair crew. They are involved in forming the myelin sheath, the fatty insulation around nerve fibers that allows for rapid signal transmission; in synthesizing the neurotransmitters that allow nerve cells to communicate; in energy production, supplying the vast amount of ATP, or adenosine triphosphate, needed for nerve function; in antioxidant defense, protecting nerves from damaging oxidative stress; in modulating the body's inflammatory response; and in nerve growth and repair, providing the raw materials for nerve regeneration.

Common Deficiencies Linked to Neuropathy

Several key nutrients are repeatedly implicated in the development or worsening of neuropathy. Addressing these is often a critical first step in a holistic healing plan.

B Vitamins: The Nerve Powerhouses

The B-complex vitamins are a family of water-soluble nutrients that are indispensable for nervous system health, and their deficiency can directly lead to nerve damage.

Vitamin B1, or thiamine, is essential for glucose metabolism and energy production, particularly in nerve cells, which rely heavily on glucose, and it is crucial for the synthesis of neurotransmitters and the maintenance of the myelin sheath. Thiamine deficiency, a condition known as beriberi, is a classic cause of neuropathy, often seen in alcoholics or those with malabsorption, producing burning feet, muscle weakness, and impaired sensation.¹ Good dietary sources include whole grains, pork, fish, nuts, seeds, and legumes.

Vitamin B6, or pyridoxine, is involved in over one hundred enzymatic reactions, including neurotransmitter synthesis of serotonin and dopamine and the formation of myelin. Both deficiency and excess of B6 can cause neuropathy: deficiency can lead to sensory neuropathy with numbness and tingling, while high doses,

over 200 milligrams a day taken long term, can cause a toxic neuropathy, so careful monitoring is essential.² Good sources include chickpeas, salmon, tuna, chicken, potatoes, and bananas.

Vitamin B9, or folate and folic acid, is critical for DNA synthesis and repair and for the methylation cycle, which is vital for nerve health and detoxification, and it works synergistically with B12. Folate deficiency can lead to nerve damage, often presenting with symptoms similar to B12 deficiency.³ Good sources include leafy green vegetables, legumes, avocado, and fortified grains.

Vitamin B12, or cobalamin, is absolutely critical for the formation and maintenance of the myelin sheath, for DNA synthesis, and for the health of red blood cells. B12 deficiency is a very common and reversible cause of neuropathy, often presenting as numbness, tingling, weakness, balance issues, and cognitive difficulties, and it can be caused by inadequate dietary intake in vegans and vegetarians, by pernicious anemia (a lack of intrinsic factor), or by malabsorption issues, for example from certain medications like metformin or proton pump inhibitors, or from gut surgery.⁴ Good sources are animal products such as meat, fish, eggs, and dairy.

Vitamin D: Beyond Bone Health

Long known for its role in bone health, vitamin D, specifically D3, is now recognized as a critical neurosteroid with profound effects on nerve and immune function. It acts more like a hormone, influencing gene expression in nerves, reducing inflammation, supporting nerve growth factor production, and modulating immune responses. Deficiency is common in neuropathy patients, and studies suggest vitamin D supplementation can improve nerve function and reduce neuropathic pain, particularly in diabetic neuropathy, thanks to its anti-inflammatory and neuroprotective properties.⁵ Good sources include sunlight exposure, fatty fish such as salmon and mackerel, fortified foods, and supplements.

Magnesium: The Great Relaxer

Magnesium is an essential mineral involved in over three hundred enzymatic reactions, including those behind nerve transmission, muscle contraction, energy production, and pain modulation. It stabilizes nerve membranes, regulates neurotransmitter activity such as glutamate and GABA, and influences cellular energy in the form of ATP; it is a natural muscle relaxant and can help reduce nerve hyperexcitability. Deficiency can lead to muscle cramps, spasms, restless legs, and increased nerve pain sensitivity, and many people are deficient due to modern diets and stress.⁶ Good sources include leafy green vegetables, nuts, seeds, legumes, whole grains, and dark chocolate.

Alpha-Lipoic Acid (ALA): The Master Antioxidant

ALA is a powerful antioxidant naturally produced in the body and found in some foods. It is unique because it is both water soluble and fat soluble, which allows it to work throughout the body. It neutralizes a wide range of free radicals, regenerates other antioxidants such as vitamin C, vitamin E, and glutathione, improves glucose utilization, and supports mitochondrial function. Extensive research, particularly in diabetic neuropathy, shows ALA can significantly improve symptoms like burning, numbness, and pain by enhancing

nerve blood flow, reducing oxidative stress, and promoting nerve regeneration.⁷ It occurs naturally in red meat, organ meats, spinach, broccoli, and yeast, though supplementation is usually needed to reach therapeutic doses.

Acetyl-L-Carnitine (ALCAR): Nerve Repair Agent

An amino acid derivative, acetyl-L-carnitine is central to energy production within cells, especially in mitochondria, and has direct neuroprotective and neuroregenerative properties. It transports fatty acids into mitochondria for energy, promotes nerve growth factor production, supports nerve cell integrity, and has antioxidant and anti-inflammatory effects. Studies suggest ALCAR can help improve nerve conduction velocity, reduce neuropathic pain, and promote nerve fiber regeneration, particularly in chemotherapy-induced and diabetic neuropathies.⁸ It is found in red meat and dairy, and is often used therapeutically as a supplement.

Other Potential Deficiencies

Copper is essential for enzymes involved in nerve function and myelin, and both deficiency, often induced by excess zinc supplementation, and toxicity can cause neuropathy. Zinc is involved in nerve signaling and antioxidant defense, and balance is key. Omega-3 fatty acids, EPA and DHA, found in fatty fish, are powerful anti-inflammatory agents that are crucial for cell membrane integrity, including in nerve cells, and they can reduce nerve inflammation and support overall brain and nerve health.⁹

Testing for Deficiencies

Don't guess, test. Blood tests are crucial to identify specific nutrient deficiencies, and these are worth discussing with your healthcare provider. A complete vitamin B panel should look beyond just B12 to include B1, B6, and folate levels; for B12 specifically, measure both serum B12 and the more sensitive markers methylmalonic acid (MMA) and homocysteine, which rise when B12 is functionally deficient at the cellular level. The 25-hydroxy vitamin D test is the most accurate measure of your vitamin D status. Red blood cell, or RBC, magnesium is more reflective of cellular magnesium stores than serum magnesium. And a trace mineral analysis, whether blood or hair based, can assess levels of copper, zinc, and other important trace minerals.

Food First Approach and Targeted Supplementation

While comprehensive testing is vital, the “food first” principle should always be your foundation. A nutrient-dense, whole-food diet provides nutrients in their most bioavailable forms and with synergistic co-factors. Prioritize a wide variety of colorful vegetables, fruits, healthy fats, quality proteins, and whole grains (more on this in Chapter 6). Once deficiencies are identified through testing, targeted, high-quality supplementation can be transformative. Where possible, choose active forms, for example methylcobalamin or adenosylcobalamin for B12, L-methylfolate for folate, and magnesium glycinate or citrate for magnesium. Work with a knowledgeable practitioner to determine appropriate therapeutic dosages, especially for

nutrients like B6 where excess can be harmful, and opt for reputable brands that offer third-party testing for purity and potency.

Addressing nutrient deficiencies is not just about adding a supplement; it's about providing your nerves with the fundamental resources they need to heal, regenerate, and function optimally. This key lays the groundwork for all other healing strategies.

Chapter 3: Key 2, Addressing Chronic Infections and Their Impact

Imagine your immune system as a vigilant army, constantly patrolling your body to identify and neutralize threats. When it encounters an acute infection, like a cold or a cut, it mounts a swift, decisive attack and then returns to a state of calm. With chronic infections, however, this army remains perpetually on high alert, leading to persistent inflammation and immune dysfunction. This constant state of alert can wreak havoc on your entire system, and your delicate nerves are often caught in the crossfire.

The connection between chronic infections and neuropathy is often overlooked in conventional medicine, yet it represents a significant “key” to unlocking stubborn cases. These infections don’t necessarily directly invade and destroy nerve tissue, though some are capable of it; rather, they trigger systemic inflammation, provoke autoimmune responses, or deplete the body’s resources, all of which can damage nerves.

The Infection-Neuropathy Connection: How Persistent Invaders Damage Nerves

Chronic infections can contribute to neuropathy through several mechanisms. Chronic inflammation is one: the sustained immune response to a persistent pathogen releases inflammatory cytokines and chemical mediators that can directly irritate and damage nerve fibers and their surrounding supportive tissues. An autoimmune response, through what is called molecular mimicry, is another. In some cases, the immune system, trying to fight off an infection, mistakenly identifies healthy nerve tissue as being similar to the invading pathogen, and this molecular mimicry leads the immune system to attack its own nerves, resulting in autoimmune neuropathy. Nutrient depletion also plays a role, since fighting a chronic infection consumes vast amounts of the body’s energy and nutrient reserves, including B vitamins, zinc, and magnesium, exacerbating deficiencies that are already detrimental to nerve health. Some pathogens produce toxins that can be directly neurotoxic, impairing nerve function or causing damage. And chronic inflammation can damage blood vessels, compromising the essential blood supply to nerves.

Common Infections Implicated in Neuropathy

While any persistent infection could theoretically contribute to nerve issues, certain pathogens are more frequently linked to neuropathy.

Lyme disease, caused by *Borrelia burgdorferi* and its co-infections, is a notorious cause of neurological symptoms, including neuropathy.¹⁰ This tick-borne bacterial infection can cause a wide range of peripheral nerve problems, from numbness and tingling to burning pain and weakness, often fluctuating in severity and location, and Lyme can also trigger autoimmune responses.¹¹ Testing requires specialized blood tests, such as Western Blot, PCR, or specific culture tests, that often go beyond standard Lyme screens. Because Lyme so often travels with co-infections, the neurological picture can be genuinely difficult to sort out, which is a strong argument for better diagnostics than standard screens provide.¹²

The herpes viruses are another notable family. Varicella-zoster virus, the virus that causes chickenpox and shingles, can lie dormant in nerve roots for decades after the original chickenpox infection; its reactivation as shingles causes a painful rash and, in many cases, post-herpetic neuralgia, a severe, persistent neuropathic pain in the affected dermatome that continues even after the rash resolves.¹³ Epstein-Barr virus, a very common virus that can cause mononucleosis, is in some individuals linked, through chronic activation, to chronic fatigue, fibromyalgia, and various neurological symptoms, including peripheral neuropathy, possibly via immune dysregulation.¹⁴ Testing for both relies on antibody tests for active or past infection, looking at IgM and IgG.

Hepatitis C virus can cause a painful peripheral neuropathy, often linked to cryoglobulinemia, a condition where abnormal proteins in the blood clump together in cold temperatures, damaging small blood vessels and nerves.¹⁵ Testing involves blood tests for HCV antibodies and viral load.

Small intestinal bacterial overgrowth, or SIBO, and gut dysbiosis more broadly are not “classic” infections of the nerves. Even so, an imbalance in gut bacteria, or an overgrowth of bacteria in the small intestine, can lead to systemic inflammation, nutrient malabsorption (especially of B vitamins), and increased intestinal permeability, commonly known as leaky gut. This allows bacterial toxins and undigested food particles to enter the bloodstream, triggering systemic inflammation and potentially autoimmune responses that can impact nerve health.¹⁶ Testing includes lactulose or glucose breath tests for SIBO and comprehensive stool analysis for gut dysbiosis.

Mycoplasma pneumoniae, an atypical bacterium that commonly causes respiratory infections, can also lead to neurological complications, including neuropathy, often through autoimmune mechanisms where antibodies generated against the bacteria cross-react with nerve components.¹⁷ Testing looks for Mycoplasma antibodies in the blood.

Diagnostic Approaches: Uncovering Hidden Infections

Identifying chronic infections requires a thorough clinical history and often specialized laboratory testing, since standard blood tests might miss these subtle, persistent threats; this is a good reason to work with a functional medicine practitioner experienced in complex chronic infections. Expanded antibody panels look beyond acute infection markers to include IgG antibodies that indicate past exposure or chronic activation, such as EBV early antigen, VZV IgG, or specific Lyme co-infection panels. PCR testing, or polymerase chain reaction testing, can detect the genetic material of pathogens, indicating active infection, and in some cases specialized cultures may be used as well. A comprehensive stool analysis can identify bacterial, fungal, or parasitic overgrowths and imbalances in the gut microbiome. And environmental toxin screening, while not testing for an “infection” as such, matters too, since heavy metals and mold toxins can mimic and exacerbate infection symptoms and compromise the immune system, making the body more susceptible.

Holistic Strategies for Infection Management

Once a chronic infection is identified, the approach has several parts, focusing on supporting the immune system and reducing the pathogen's load without relying solely on long-term antibiotics, which can further disrupt the gut microbiome.

On the dietary side, an anti-inflammatory diet, as discussed in Chapter 6, reduces inflammatory foods such as processed foods, sugar, and refined oils, which is crucial to lower the systemic inflammatory burden and support immune function. Immune-boosting foods rich in vitamin C, zinc, vitamin D, and antioxidants, such as colorful fruits and vegetables, bone broth, garlic, onions, ginger, and turmeric, are worth incorporating regularly. Fermented foods like sauerkraut, kimchi, and kefir, along with prebiotic fibers like onions, garlic, and asparagus, support a healthy microbiome, which is foundational to a robust immune system.

Many natural compounds possess antimicrobial, antiviral, and antiparasitic properties that can help the body fight off chronic pathogens with fewer side effects than pharmaceuticals. Examples include olive leaf extract, oregano oil, berberine, garlic, artemisia, cat's claw, colloidal silver, and specific probiotics. These should always be used under the guidance of a qualified healthcare practitioner, given potential interactions and the importance of appropriate dosing.

Immune system modulation is another lever. Adaptogenic herbs like ashwagandha, rhodiola, and ginseng can help the body adapt to stress and support immune balance. Beta-glucans, found in medicinal mushrooms such as reishi and turkey tail, can prime the immune system to respond more effectively. Vitamin D and zinc, discussed in Chapter 2, remain critical for immune cell function. And glutathione, the body's master antioxidant, is vital for detoxification and immune health, with supplementation of precursors like N-acetyl cysteine (NAC) often beneficial.

Finally, stress reduction and sleep optimization matter more than they might seem to. Chronic stress suppresses the immune system, making the body more vulnerable to persistent infections, so techniques like meditation, yoga, and deep breathing are vital, and adequate, restorative sleep is non-negotiable for immune repair and regulation.

By identifying and strategically addressing chronic infections, you remove a significant burden on your immune system and reduce a potent source of inflammation that may be driving your neuropathy. This “key” is about supporting your body's innate ability to heal and restore balance.

Chapter 4: Key 3, Activating Large Diameter Afferent Pathways

Neuropathy often leads to a disruption in the delicate balance of your nervous system. While small nerve fibers transmit pain and temperature, it's the large diameter afferent pathways that carry crucial information about touch, pressure, vibration, and proprioception, your sense of body position in space. When these larger, myelinated fibers are damaged or their signals are overwhelmed by pain signals from smaller, unmyelinated fibers, the brain receives distorted or incomplete sensory information.

Activating and stimulating these large diameter afferent pathways is a powerful strategy in neuropathy management. This process, often referred to as sensory retraining or gating, helps in several ways at once. According to the Gate Control Theory of Pain, stimulating large afferent fibers can “close the gate” to pain signals from smaller fibers, reducing the perception of pain. Restoring accurate sensory input helps the brain better understand where your body is in space, improving coordination and reducing fall risk. By re-educating the nervous system and improving nerve signaling, numbness and tingling can gradually lessen as sensation returns or improves. And the process promotes neuroplasticity, the brain's ability to reorganize itself by forming new neural connections, which can help compensate for damaged areas.

This chapter explores various methods to activate these vital pathways, from simple at-home techniques to advanced therapeutic devices.

1. Vibration Therapy

Vibration is a powerful stimulus for large diameter afferent fibers, particularly Pacinian corpuscles, which are highly sensitive to vibration. Applying specific frequencies of vibration directly to affected areas, such as the feet or hands, stimulates these mechanoreceptors, sending strong signals up the large afferent pathways to the brain, which can help override pain signals and improve sensory perception. In practice, this can take the form of whole-body vibration platforms used for general stimulation and to improve balance, handheld vibrators applied to the soles of the feet, palms of the hands, or specific numb areas for direct stimulation, or medical-grade tuning forks, for example at 128 Hz, applied to bony prominences to assess and stimulate vibratory sensation. Vibration therapy is reported to reduce neuropathic pain, improve balance, and enhance sensory perception.

2. Tactile Stimulation

Simple touch and texture can re-engage sensory pathways. Light touch, brushing, or exposure to varied textures activates different types of mechanoreceptors, sending diverse sensory information to the brain, which helps to desensitize hypersensitive areas and re-awaken numb ones. This can look like walking barefoot on different textured surfaces such as grass, sand, or textured mats; using a soft sensory brush to gently stroke numb or painful areas; or receiving gentle massage, which can improve circulation and

stimulate nerve endings. Tactile stimulation helps to normalize sensory processing, reduce allodynia (pain from non-painful stimuli), and improve tactile discrimination.

3. Proprioceptive Exercises

These exercises focus on your body's awareness in space. Engaging in activities that challenge balance and coordination forces the brain to rely more heavily on proprioceptive input from joints and muscles, strengthening these pathways. Balance exercises such as standing on one leg, heel-to-toe walking, or using a wobble board or balance disc are one avenue; tai chi or yoga, which inherently improve balance, body awareness, and coordination, are another; and gait training, focused on proper walking mechanics, rounds this out. Together, these improve balance, reduce fall risk, enhance coordination, and strengthen the neural connections responsible for body awareness.

4. Photobiomodulation (Laser and Red Light Therapy)

Photobiomodulation, often referred to as low-level laser therapy or red light therapy, uses specific wavelengths of light to stimulate cellular function and promote healing. While it does not directly activate large diameter afferents in the same mechanical way as vibration, it creates an optimal environment for nerve health and repair, indirectly supporting the function of all nerve fibers, including large afferents.

Its mechanism works on several fronts. Red and near-infrared light penetrates tissues and is absorbed by chromophores, primarily cytochrome c oxidase, within the mitochondria of cells, and this absorption increases the production of adenosine triphosphate (ATP), the primary energy currency of cells, meaning nerve cells have more energy to function optimally and repair themselves. Photobiomodulation also helps modulate inflammatory pathways and reduce oxidative stress by neutralizing free radicals; since chronic inflammation and oxidative stress are major contributors to nerve damage, reducing them creates a healthier environment for nerve function and regeneration. Light therapy can promote vasodilation and angiogenesis, the widening of blood vessels and the formation of new ones, leading to increased blood flow to nerve tissues and ensuring nerves receive a vital supply of oxygen and nutrients. It has been shown to stimulate nerve growth factors, promote axonal sprouting, and support the regeneration of damaged nerve fibers and their myelin sheaths, which can lead to improvements in nerve conduction and overall function. And while it's not a direct activation of large afferents for gating, it can reduce pain by decreasing inflammation, improving nerve health, and potentially influencing nerve cell action potentials and neurotransmitter release.

For neuropathy specifically, the reported benefits include pain reduction (burning, shooting pain), improved sensation (reduced numbness and tingling), enhanced nerve conduction velocity, support for nerve regeneration, and reduced inflammation in nerve tissues. In application, clinic-based high-power therapeutic lasers (Class IV) are often used in clinical settings for deeper penetration and faster treatment times, while home-use red light devices, lower-power LED panels, wraps, and handheld units, are available for home use but require more consistent and longer application times. These devices typically use red light around 600 to 700 nanometers and near-infrared light around 780 to 1,100 nanometers for their ability to penetrate tissue. A systematic review found that low-level laser therapy has a positive effect in controlling diabetic neuropathic

pain,¹⁸ and research suggests that near-infrared laser light can rapidly block pain transmission and, with lower irradiances, offer prolonged pain relief by altering chronic inflammation.¹⁹

5. Electrical Stimulation Devices, Such As the ReBuilder Medical Device

Electrical stimulation has long been used in pain management and rehabilitation. Devices like the ReBuilder Medical Device are designed to deliver specific electrical signals to nerve pathways, aiming to re-educate nerves and improve their function.

The ReBuilder device is designed to generate a proprietary electrical waveform that reportedly mimics the natural electrical signals of healthy nerves, which is distinct from standard TENS (transcutaneous electrical nerve stimulation) units that often use simpler, less varied waveforms. By applying electrodes to specific areas, such as the soles of the feet or the hands, the device sends these signals along the nerve pathways, and the manufacturer states that it uses biofeedback to adjust its output to the individual's electrical parameters, aiming for more personalized and effective stimulation. By delivering precise electrical impulses, the ReBuilder aims to directly stimulate large diameter afferent nerve fibers, which can help “wake up” dormant nerves that have become less functional due to damage, improve nerve conduction by enhancing the speed and efficiency of signal transmission, and modulate pain through gating, since, similar to vibration, the strong, regular signals from the device can activate large afferents and help “close the gate” on pain signals from smaller, damaged nerves. The electrical stimulation can also induce muscle contractions and promote vasodilation, increasing local blood flow and oxygenation to nerve tissues, which is vital for nerve health and repair.

Reported benefits for neuropathy include a reduction in neuropathic pain (burning, aching, shooting pains), a decrease in numbness and tingling, improved sensation and proprioception, increased local blood circulation, relaxation of muscle spasms and reduction of cramps, and the potential for reduced reliance on pain medications. In practice, the ReBuilder device typically involves placing conductive pads on the soles of the feet, or the hands, lower back, or shoulders, or using a footbath with a conductive solution; treatments are usually 30 minutes, once or twice daily, with reported immediate relief that can last for several hours and cumulative improvements with continued use.

There are a few important considerations to state plainly. The ReBuilder Medical Device is FDA registered as a transcutaneous electrical nerve stimulator for pain relief. While anecdotal reports and manufacturer-conducted studies report high success rates, independent, large-scale, peer-reviewed clinical trials specifically on the ReBuilder device for neuropathy are less abundant in databases like PubMed compared to broader categories like TENS or photobiomodulation, so it's important to discuss this option with your healthcare provider to determine if it's appropriate for your specific condition. It is also contraindicated for individuals with certain pacemakers, especially those with defibrillator accessories, and for pregnant women.

Integrating Activation Strategies

The power of activating large diameter afferent pathways lies in consistency and combination. Incorporate a variety of these techniques into your daily routine: for example, combine targeted electrical stimulation with daily balance exercises, or use a handheld vibrator on numb areas while also receiving regular photobiomodulation treatments. By providing consistent, positive sensory input, you can help retrain your nervous system, reduce pain, and improve your overall nerve function. This key is about actively engaging your body's innate capacity for neuroplasticity and healing.

Chapter 5: Key 4, Comprehensive Testing Beyond the Conventional

When facing the complexities of neuropathy, relying solely on standard diagnostic tests can often leave crucial stones unturned. Conventional evaluations typically include nerve conduction studies (NCS) and electromyography (EMG) to assess nerve damage, along with basic blood work to rule out common culprits like overt diabetes or severe B12 deficiency. While these are important starting points, they often fail to uncover the nuanced, underlying imbalances that contribute to the majority of neuropathy cases.

The root cause approach to neuropathy demands a more sophisticated and comprehensive diagnostic strategy. This key involves looking beyond superficial markers to identify hidden inflammation, subtle metabolic dysregulation, circulatory compromise, and other systemic factors that may be silently driving your nerve damage. By truly understanding the physiological landscape of your body, you can create a truly targeted and effective treatment plan.

The Limitations of Standard Tests: Why a Deeper Dive Is Needed

Nerve conduction studies and electromyography assess the function of large, myelinated nerve fibers and muscle electrical activity. While invaluable for diagnosing the presence and type of nerve damage, they may not detect small fiber neuropathy, which involves the tiny nerves responsible for pain and temperature sensation, and more importantly, they tell you what nerve damage exists, but rarely why. Basic blood work, a fasting glucose test or HbA1c, may flag diabetes, and a basic B12 level might catch severe deficiency, but these often miss pre-diabetic states, functional B12 deficiencies, or other critical nutrient imbalances that aren't severe enough to show up on standard ranges but are still detrimental to nerve health. And when standard tests yield no clear cause, patients are often labeled with idiopathic neuropathy, which means the cause is unknown, not that there isn't a cause, and that is precisely where comprehensive testing becomes crucial.

Expanding the Diagnostic Lens: Key Areas for Deeper Investigation

Working with a functional medicine practitioner who specializes in neuropathy means accessing a broader array of diagnostic tools designed to uncover the physiological stressors impacting your nerves.

Inflammation markers are the silent nerve attackers, since chronic, low-grade inflammation is a major driver of nerve damage even without a diagnosed autoimmune condition, so identifying and quantifying this inflammation is paramount.²⁰ High-sensitivity C-reactive protein, or hs-CRP, is a sensitive but entirely nonspecific screening marker for systemic inflammation: an elevated result says an inflammatory load exists somewhere, not where it is coming from, and it confirms nothing on its own. Read it as a prompt to keep looking, not as an answer. Erythrocyte sedimentation rate is another general marker of inflammation, and less specific still. Homocysteine belongs in a different category, and it is one of the most consequential

numbers on this panel rather than a piece of trivia to wave off. It is a methylation-cycle marker. It rises when B12, folate, or B6 run short, and with thyroid or kidney trouble, and elevated levels are associated with increased inflammation, oxidative stress, and vascular changes that work against nerve health.²¹ Rather than chasing the number with blind supplementation, find which input is short (B12, folate, B6, thyroid, or kidney function), correct that one, and let homocysteine confirm the fix on retest. Fibrinogen, a protein involved in blood clotting and inflammation, can indicate chronic inflammation and increased risk of vascular issues affecting nerve blood supply when elevated. And more advanced cytokine testing, for markers like IL-6 and TNF-alpha, can measure specific inflammatory cytokines that directly contribute to nerve pain and damage, offering a deeper insight into immune system activation.

Blood sugar regulation matters well beyond a diabetes diagnosis, since many neuropathy cases are linked to blood sugar dysregulation even if a person isn't officially diagnosed with diabetes, and pre-diabetes or insulin resistance can cause nerve damage over time.²² Measuring fasting glucose and insulin together provides insight into how much insulin your body needs to keep blood sugar stable, and high fasting insulin indicates insulin resistance. Hemoglobin A1c gives an average blood sugar level over the past two to three months; while it's a standard test, looking at optimal ranges, below 5.4 percent, rather than just "non-diabetic" ranges, is important. The oral glucose tolerance test with insulin measures blood glucose and insulin levels at intervals after a glucose drink, and it can uncover subtle dysregulation that fasting tests miss. It is the most informative test that is realistically orderable for this question, which is not the same thing as a gold standard, and no single one of these numbers confirms insulin resistance on its own. And for some patients, wearing a continuous glucose monitor for a few days or weeks can provide invaluable real-time data on how different foods, stress, and activities impact blood sugar, revealing spikes and crashes that contribute to nerve stress.

Circulation assessment recognizes that nerves, like all tissues, depend on a robust blood supply for oxygen and nutrients, and compromised circulation is a direct pathway to nerve damage. A physical exam can reveal signs of poor circulation, and non-invasive tests like the ankle-brachial index can screen for peripheral artery disease, which often coexists with neuropathy. Blood tests for endothelial function markers like asymmetric dimethylarginine (ADMA) or endothelin-1 can provide insight into the health of your blood vessel lining, which is critical for healthy blood flow. And the inflammatory markers already discussed, high hs-CRP and homocysteine, also negatively impact vascular health.

Autoimmune markers matter because neuropathy can be an autoimmune condition where the immune system mistakenly attacks nerve components, and specific testing can help identify these cases.²³ Antinuclear antibodies (ANA) offer a general screen for autoimmune activity. Beyond that, specific autoantibody panels look at ganglioside antibodies, such as anti-GM1, anti-GD1a, and anti-GQ1b, which are associated with various autoimmune neuropathies like Guillain-Barré syndrome or multifocal motor neuropathy; MAG antibodies (myelin-associated glycoprotein antibodies), linked to demyelinating neuropathies; SSA and SSB antibodies associated with Sjögren's syndrome, which can cause significant small fiber neuropathy; and thyroid antibodies (TPOAb, TgAb), since autoimmune thyroid conditions like Hashimoto's can be associated with neuropathy.

Heavy metal toxicity testing addresses a hidden poison, since exposure to heavy metals can be profoundly neurotoxic, causing direct damage to nerve cells and their myelin sheaths. Hair mineral analysis can reflect long-term exposure to certain heavy metals. A urine heavy metal challenge test, which involves collecting urine before and after administering a chelating agent that mobilizes metals from tissues, provides a comprehensive assessment of body burden, while a blood heavy metal test is useful for recent or acute exposure. A study of metal industry workers with chronic occupational exposure to lead, cadmium, and arsenic found clear, measurable impairment of peripheral nerve function, most pronounced in the small sensory fibers that carry pain and temperature signals.²⁴

The Organic Acids Test, or OAT, offers metabolic insights by providing a snapshot of various pathways in the body, including mitochondrial function, neurotransmitter metabolism, and nutrient needs. It identifies imbalances in gut flora, such as yeast and bacterial overgrowth, nutrient deficiencies in B vitamins and antioxidants, and markers of oxidative stress that can impact nerve health; for example, high markers of certain organic acids can indicate issues with energy production or detoxification crucial for nerves. It helps reveal functional deficiencies that might not show up on simple blood tests and provides insight into the efficiency of various metabolic processes.

Comprehensive stool testing captures the gut-nerve connection, since your gut microbiome has a profound influence on your overall health, including your nervous system, and dysbiosis, an imbalance of gut bacteria, can lead to systemic inflammation, nutrient malabsorption, and the production of neurotoxic compounds. It analyzes the balance of beneficial and pathogenic bacteria, yeast, parasites, markers of inflammation, and digestive function, and it can identify issues like SIBO, candidiasis, or parasitic infections that contribute to systemic inflammation and nerve symptoms.¹⁶

The Value of Comprehensive Testing

Investing in these advanced diagnostic tests moves you from guessing to knowing. It allows your healthcare team to develop a highly personalized treatment plan that targets the precise mechanisms contributing to your neuropathy, rather than a generic approach. This key empowers you to peel back the layers of your condition and address the root causes, paving the way for more effective and lasting relief.

Chapter 6: Key 5, Diet and Food Sensitivities for Nerve Repair

Imagine your body as a high-performance vehicle. You wouldn't fill a sports car with low-grade fuel and expect it to run optimally. Similarly, the food you consume profoundly impacts every cell, tissue, and organ, especially your delicate nervous system. Diet is not merely about calories; it's about providing the specific nutrients for repair, reducing inflammation, stabilizing blood sugar, and supporting a healthy gut, all critical factors for unlocking nerve health.

This key is arguably one of the most foundational and empowering. While nutrient deficiencies, covered in Chapter 2, highlight what you might lack, this chapter focuses on how your daily food choices can either perpetuate nerve damage or actively promote nerve healing and protection. We will explore the principles of an anti-inflammatory, nerve-supportive diet and look closely at the often-overlooked role of food sensitivities.

The Anti-Inflammatory Diet Principle: Reducing Triggers and Promoting Healing

Chronic inflammation is a silent assassin of nerves. It damages myelin, impairs blood flow, and directly irritates nerve fibers. The cornerstone of a neuropathy-healing diet is to systematically reduce inflammatory triggers while flooding your body with anti-inflammatory compounds.

Among the foods to embrace, your strongest allies are abundant vegetables and fruits. That means a wide array of colorful, non-starchy vegetables, especially leafy greens like spinach, kale, and collards, and cruciferous vegetables like broccoli, cauliflower, and Brussels sprouts, along with low-glycemic fruits like berries, avocados, lemons, and limes, all packed with antioxidants, vitamins, minerals, and fiber. Healthy fats are essential for nerve cell membranes and myelin, so include sources like extra virgin olive oil, avocados, nuts such as almonds, walnuts, and pecans, seeds such as chia, flax, and hemp, and fatty fish such as wild-caught salmon, mackerel, and sardines, rich in anti-inflammatory omega-3 fatty acids (EPA and DHA). Lean, high-quality proteins are necessary for tissue repair and neurotransmitter synthesis, so prioritize grass-fed or pasture-raised meats, wild-caught fish, organic poultry, and plant-based options like legumes, if tolerated, and quinoa. Fiber-rich foods, or prebiotics, feed beneficial gut bacteria, which produce short-chain fatty acids that reduce inflammation and support gut barrier integrity, so include legumes, whole grains if tolerated, and a variety of vegetables and fruits. And anti-inflammatory spices like turmeric (paired with black pepper for absorption), ginger, cinnamon, garlic, and rosemary contain potent compounds that combat inflammation.

On the other side of the ledger are the foods to limit or avoid, the common culprits that can fuel inflammation and undermine nerve health. Processed foods are often high in unhealthy fats, sugar, and artificial ingredients. Refined sugars and sweeteners directly contribute to blood sugar spikes, insulin resistance, and systemic inflammation, which are primary drivers of diabetic neuropathy and general nerve damage, and this includes the hidden sugars in sauces, dressings, and drinks. Unhealthy fats, trans fats (partially hydrogenated oils) and highly refined vegetable oils such as corn, soy, canola, sunflower, and safflower oil,

are high in pro-inflammatory omega-6s. Artificial ingredients, dyes, preservatives, and artificial sweeteners such as aspartame and sucralose, can trigger inflammation and neurological symptoms in sensitive individuals. And excessive alcohol can be directly neurotoxic and deplete vital nutrients.

Elimination Diet for Food Sensitivities: Unmasking Hidden Triggers

Beyond obvious allergies, many people suffer from delayed food sensitivities, where an immune reaction to certain foods causes low-grade inflammation that can manifest in various symptoms, including neuropathic pain. Identifying these triggers is crucial.

It helps to understand the difference between the two. A food allergy is an immediate, often severe, IgE-mediated immune reaction, for example hives or difficulty breathing. A food sensitivity, or intolerance, is a delayed, often IgG-mediated immune reaction that can cause a wide range of less immediate symptoms, such as digestive issues, headaches, joint pain, skin problems, and chronic inflammation impacting nerves.

The most common food sensitivities include gluten, found in wheat, barley, and rye, where gluten sensitivity or celiac disease can cause direct nerve damage, sometimes called gluten neuropathy, and significant systemic inflammation;²⁵ dairy, where casein and lactose can be inflammatory for many; soy, a common allergen and inflammatory food for some; corn, often genetically modified and inflammatory for sensitive individuals; and nightshades such as tomatoes, potatoes, peppers, and eggplant, which, while healthy for many, contain compounds called alkaloids that can aggravate inflammation and pain in sensitive individuals.

To conduct an elimination and reintroduction phase, start with a strict elimination phase of three to four weeks: remove all suspected inflammatory foods and common sensitivities, including gluten, dairy, soy, corn, eggs, peanuts, sugar, processed foods, and alcohol, and focus exclusively on whole, unprocessed, anti-inflammatory foods during this window. This gives your body a chance to calm down and symptoms to subside. After that, move into a reintroduction phase: bring back one food group at a time every two to three days, and monitor your symptoms closely, for example with a symptom journal. If symptoms worsen, neuropathic pain, fatigue, digestive upset, or brain fog, that food is likely a trigger for you. Once you identify your triggers, you can build a personalized diet that supports your nerve health without guilt or deprivation.

Hydration and herbal support round out the foundation. Water is vital for every bodily process, including nerve signal transmission and detoxification, so aim for half your body weight in ounces of filtered water daily. Certain herbs have neuroprotective, anti-inflammatory, or pain-relieving properties worth knowing about: turmeric and curcumin are potent anti-inflammatories and antioxidants; ginger reduces inflammation and pain; boswellia is anti-inflammatory, often used for joint pain but relevant for nerve inflammation too; and passionflower, California poppy, and valerian can help with nerve pain and improve sleep, indirectly supporting nerve repair. Always consult with your healthcare provider before adding herbal supplements, especially if you are on medications.

Sample 90-Day Food Guide Framework

This framework provides a structured approach to implementing your nerve-healing diet. It's designed to transition you from an inflammatory eating pattern to a sustainable, nerve-supportive lifestyle.

As a weekly framework, a useful guideline for your macronutrient breakdown is what we call 60/20/20: meals that stabilize blood sugar and provide sustained energy tend to draw about 60 percent of their content from healthy carbohydrates, primarily non-starchy vegetables, some fruits, and carefully selected whole grains if tolerated and after the elimination phase, focusing on complex carbohydrates with high fiber content; about 20 percent from healthy fats, from sources like avocado, olive oil, nuts, seeds, and fatty fish; and the remaining 20 percent from quality proteins, from lean meats, poultry, fish, eggs, and plant-based proteins. This is a general guideline. Your individual needs may vary, and a nutritionist can help fine-tune these percentages based on your specific health goals and activity level.

In the first phase, weeks one through four, the focus is elimination and detoxification. You strictly eliminate common inflammatory foods and sensitivities and emphasize whole, unprocessed foods, with the goal of reducing inflammation, stabilizing blood sugar, allowing the gut to heal, and identifying immediate triggers. Meals in this phase are built around abundant non-starchy vegetables, clean protein, and healthy fats, with minimal fruit and no grains initially. A representative day might start with a green smoothie made of spinach, plant-based protein powder, almond milk, and half an avocado; lunch could be a large mixed greens salad with grilled chicken or wild salmon, an olive oil and lemon dressing, and a handful of pumpkin seeds; dinner might be baked cod with steamed broccoli and sautéed asparagus; and a small handful of almonds makes a good snack.

In the second phase, weeks five through eight, the focus shifts to reintroduction and personalization. You systematically reintroduce one eliminated food group at a time, observing for symptoms, with the goal of identifying your individual food sensitivities and expanding your personalized diet. The menu philosophy continues with the anti-inflammatory base from phase one, but strategically adds foods back: breakfast can stay the same as phase one, lunch might become turkey and avocado lettuce wraps with a side of mixed berries as you reintroduce berries, and dinner might be gluten-free pasta, if you are reintroducing rice pasta, with lean ground turkey and a tomato-based sauce once tomatoes and other nightshades are back in.

In the third phase, weeks nine through twelve, the focus is maintenance and optimization. You solidify your personalized anti-inflammatory diet based on your findings and continue to prioritize nutrient density, with the goal of sustaining nerve healing, managing blood sugar, maintaining low inflammation, and feeling consistently better. Here you enjoy a diverse range of your newly identified safe and supportive foods: breakfast might be chia seed pudding with berries and walnuts, lunch could be leftover dinner or a large vegetable and lentil soup, and dinner might be a stir-fry with lean beef, a variety of colorful vegetables, and brown rice if tolerated.

Throughout all three phases, a handful of snacks reliably stabilize glucose between meals: hard-boiled eggs, avocado slices with a sprinkle of sea salt, a small handful of nuts such as almonds or walnuts or seeds such

as pumpkin or sunflower, celery sticks with almond or seed butter, berries with a dollop of unsweetened coconut yogurt, olives, and a small piece of wild-caught salmon or sugar-free turkey jerky.

Getting the logistics right makes the plan sustainable. Dedicate time each week to plan meals and snacks. Shop smart by focusing on the perimeter of the grocery store, where the produce and lean proteins live. Batch cook larger quantities of staples like roasted vegetables, cooked grains, and lean protein (chicken breast, ground turkey) so you can assemble meals easily throughout the week. Pre-chop vegetables so quick meals or snacks are always within reach. And keep a well-stocked pantry of healthy staples such as olive oil, vinegars, anti-inflammatory spices, nuts, seeds, and canned wild salmon.

By consciously choosing foods that nourish your nerves and avoiding those that inflame them, you are providing your body with the optimal environment for healing. This dietary shift is a powerful, sustainable lifestyle change rather than a temporary fix, and it is essential for reclaiming your nerve health.

Chapter 7: The Movement Plan, Restoring Function and Reducing Pain

While diet and targeted therapies provide crucial internal support for nerve health, movement is the indispensable key that bridges internal healing with external function. When you live with neuropathy, fear of falling, increased pain, or perceived weakness can lead to a sedentary lifestyle. However, inactivity often worsens symptoms by reducing circulation, decreasing muscle strength, and hindering neuroplasticity, the brain's ability to adapt and reorganize.

Conversely, strategic movement is a powerful therapeutic tool. It can directly benefit nerve health by improving blood flow, promoting the release of nerve growth factors, strengthening supportive muscles, and enhancing the communication between your brain and body. This chapter will outline research-backed exercise approaches that can reduce pain, improve sensation, and restore function, regardless of your current mobility level.

Benefits of Exercise for Neuropathy

Engaging in regular, appropriate physical activity offers a multitude of advantages for individuals with neuropathy. Movement increases blood flow to the extremities, delivering vital oxygen and nutrients to damaged nerves and helping to remove waste products, which is critical for nerve repair and function. Physical activity also stimulates the release of neurotrophic factors, like brain-derived neurotrophic factor (BDNF), proteins that promote the growth, survival, and differentiation of neurons and synapses, which can aid in nerve repair and regeneration; while research linking exercise-driven BDNF increases specifically to neuropathy is still developing, the broader link between exercise and BDNF is well established.²⁶ Exercise can activate your body's natural pain-relieving mechanisms, including the release of endorphins, and consistent, appropriate movement can also desensitize overactive pain pathways, an effect supported by research on exercise as a strategy for neuropathic pain management.²⁷ Stronger muscles from regular activity provide better support for joints and improved stability, reducing the risk of falls and compensating for nerve-related weakness. Many exercises specifically target the large diameter afferent pathways responsible for your sense of body position, directly addressing common neuropathy symptoms and improving balance and proprioception. Maintaining a healthy weight through exercise is crucial, especially for diabetic neuropathy, since excess weight exacerbates insulin resistance and inflammation. And exercise is a potent antidote to stress, anxiety, and depression, which often accompany chronic pain conditions like neuropathy, so the mood enhancement and stress reduction it brings positively impacts overall nerve health as well.

Types of Exercise to Embrace

The key is to start slowly, listen to your body, and progress gradually. Consistency is more important than intensity.

Low-impact aerobics aim to improve cardiovascular health, enhance circulation, and maintain overall fitness without excessive stress on joints or feet. Walking is the simplest place to start: even short, frequent walks can make a significant difference, so begin with 5 to 10 minutes a few times a day and gradually increase duration, wearing supportive, well-cushioned shoes. Swimming and water aerobics use the buoyancy of water to reduce weight-bearing stress, making them ideal for those with foot pain or balance issues, while the resistance of the water also provides gentle strengthening. A stationary or recumbent bike offers a non-weight-bearing way to improve leg circulation and strength, and an elliptical machine provides a similarly low-impact cardio workout. Aerobic exercise of this kind increases insulin sensitivity, helping cells absorb glucose more efficiently from the bloodstream, which directly benefits those with diabetic neuropathy or insulin resistance.

Resistance, or strength, training aims to build and maintain muscle mass, which improves strength and stability and can help manage blood sugar by increasing insulin sensitivity. This can take the form of bodyweight exercises like squats using a chair for support, shallow lunges, wall push-ups, and chair stands; resistance bands, which provide adjustable resistance for arms, legs, and core; light weights, dumbbells or household items like cans or water bottles, for bicep curls, tricep extensions, and shoulder presses; or machine weights, which provide controlled movements and support and are often a good starting point. Focus on proper form to prevent injury, and start with lighter weights and higher repetitions, gradually increasing resistance as strength improves. Resistance training builds muscle, and muscle is the primary site for glucose uptake, so more muscle mass means better glucose disposal and improved insulin sensitivity, even independent of weight loss.

Walking after meals is a simple yet highly effective strategy to immediately lower post-meal blood sugar spikes. Muscle activity immediately after eating helps to utilize circulating glucose for energy, preventing the large spikes that are damaging to nerves, so aim for a brisk 10 to 15 minute walk within 30 minutes after major meals; even short bursts count. Studies show that a short walk after a meal can significantly reduce postprandial (after-meal) glucose levels, sometimes more effectively than a longer walk at another time of day.

High-intensity interval training, in moderation, involves short bursts of intense exercise followed by brief recovery periods, and it can significantly improve cardiovascular fitness, insulin sensitivity, and mitochondrial function. For individuals with neuropathy, HIIT should be approached with extreme caution and only under professional guidance, especially if balance or pain is a major concern; a modified version can involve bursts of brisk walking followed by slower walking, or short periods of higher resistance on a stationary bike. It's not suitable for everyone, especially those with severe neuropathy, balance issues, or heart conditions, so always consult your doctor before attempting it. HIIT has been shown to improve insulin sensitivity and glucose metabolism even more effectively than moderate-intensity continuous exercise in some populations, due to its potent effects on muscle glucose uptake and mitochondrial biogenesis.

Functional movement routines for balance and flexibility directly enhance proprioception, balance, flexibility, and coordination, areas often severely impacted by neuropathy, and these exercises directly activate the large diameter afferent pathways discussed in Chapter 4. Tai chi and yoga are excellent for improving

balance, flexibility, strength, and body awareness, and many studios offer chair yoga or adaptive classes for those who need them. Balance drills, standing on one foot while holding onto support initially, tandem stance (heel-to-toe), or walking safely on uneven surfaces, build the same skills directly. Gentle stretching for ankles, calves, hamstrings, and feet helps maintain range of motion and reduce stiffness, and Pilates rounds this out by focusing on core strength, stability, and controlled movements.

Sample Weekly Movement Strategy for All Levels

This is a template meant to be adapted to your comfort level, ideally alongside a physical therapist or exercise physiologist experienced with neuropathy. A few principles apply at every level: start slow, beginning with very short durations of five to ten minutes if you are new to exercise and gradually increasing; listen to your body, since pain is a signal, and you should differentiate between muscle fatigue and neuropathic pain and never push through sharp or burning neuropathic pain; prioritize consistency, since daily, even short, bursts of activity are more effective than sporadic long sessions; build in variety by combining different types of exercise to challenge your body in multiple ways; and put safety first, making sure your environment is safe, wearing appropriate footwear, and considering supervision, especially for balance exercises.

Beginner level. For those new to exercise or with significant symptoms, a good starting point is 10 to 15 minutes of brisk walking daily, split into two or three sessions if needed, plus 5 minutes of gentle stretching; three times a week, add between 10 and 15 minutes of chair-based strength exercises such as chair squats or bicep curls with light weights; twice a week, add 5 to 10 minutes of supported balance exercises such as standing on one foot while holding a counter; and each day, close with a 10 to 15 minute walk after dinner.

Intermediate level. For those regularly active with moderate symptoms, aim for 30 minutes of brisk walking or cycling daily; three to four times a week, do 20 to 30 minutes of bodyweight strength training or light resistance bands; three times a week, add 15 to 20 minutes of balance training such as tai chi, yoga, or wobble board work; and again close each day with a 10 to 15 minute walk after dinner.

Advanced level. For those with good mobility and well-managed symptoms who are seeking optimization, aim for 30 to 45 minutes of moderate-intensity cardio daily, whether brisk walking, jogging, cycling, or swimming; three to four times a week, do between 30 and 45 minutes of resistance training with free weights, machines, or challenging bodyweight work; two to three times a week, add 15 to 20 minutes of dynamic balance and flexibility work such as advanced yoga, Pilates, or complex balance drills; once or twice a week, and only with caution, consider modified HIIT sessions, for example sprint intervals on a bike for 20 to 30 seconds followed by one to two minutes of recovery; and, as at every level, close each day with a 10 to 15 minute walk after dinner.

Always consult with your doctor or a physical therapist before starting any new exercise program, especially if you have existing health conditions or severe neuropathy. They can help tailor a plan that is safe and effective for your individual needs. Movement is not just about physical strength; it's about reconnecting with your body, fostering neuroplasticity, and empowering yourself on the path to nerve health.

Chapter 8: Mindset and Monitoring, the Unseen Influencers

Neuropathy is a physical condition, but its impact extends far beyond the nerves themselves. The persistent pain, numbness, and functional limitations can take a significant toll on your mental and emotional well-being. Chronic stress, anxiety, frustration, and even feelings of hopelessness can become entangled with the physical symptoms, creating a vicious cycle that can hinder healing.

This key acknowledges that true holistic recovery encompasses the psychological and emotional landscape as well as the biological. It's about cultivating a resilient mindset, reframing setbacks, celebrating small victories, and strategically monitoring your progress to stay motivated and informed. Your brain is intimately connected to your nervous system, and nurturing this connection is as vital as any diet or supplement.

The Brain-Nerve Connection: How Mindset Influences Neuropathy

Your brain and nervous system are not separate entities. The brain constantly processes sensory input from your nerves and influences how you perceive pain, and chronic stress, anxiety, and negative thought patterns can work against you in several ways. Stress hormones like cortisol can sensitize nerve endings, making pain feel more intense, and the brain, under chronic stress, can become “stuck” in a pain loop even with minimal physical stimulus. Persistent stress diverts energy away from repair processes, including nerve regeneration, and it can also suppress the immune system, making you more vulnerable to inflammation and infections. Chronic stress can lead to vasoconstriction, the narrowing of blood vessels, reducing vital blood flow to nerves. And imbalances in mood-regulating neurotransmitters like serotonin and dopamine can exacerbate pain perception and reduce your capacity to cope.

Cultivating a positive and proactive mindset isn't about ignoring your symptoms; it's about empowering yourself to influence your body's healing capacity and improve your quality of life.

Journaling Blood Sugar Trends

For those with diabetic neuropathy or insulin resistance, consistent glucose monitoring is not just about managing numbers; it's a powerful tool for understanding your body's unique responses and making informed dietary choices. While your fasting glucose and HbA1c give you a snapshot, daily logging reveals patterns. Each time, record the time of measurement (before meals, one to two hours after meals, before bed, upon waking), the glucose reading itself, meal details (what you ate, approximate portion sizes, and main ingredients, noting carbohydrate, protein, and fat content), any activity before or after the meal (a fifteen-minute walk after dinner, say), a simple stress level rating from one to five or a brief note about emotional state, sleep quality from the night before, and any changes in neuropathy symptoms, pain, numbness, or tingling, after specific foods or activities. Journaling helps you identify the triggers, foods,

stress, or lack of sleep, that cause blood sugar spikes and subsequent neuropathy flare-ups, and it empowers you to see the direct connection between your lifestyle choices and your symptoms.

Glucose Monitoring Tips

Consistency is key: test at consistent times so you can identify patterns. Testing before and after meals, so-called paired testing, is crucial to see how specific foods affect your blood sugar. If it's available and appropriate for your condition, a continuous glucose monitor can provide real-time data, revealing spikes and trends you might miss with finger-prick tests; this passive monitoring is particularly insightful for understanding the impact of stress, sleep, and subtle food choices. And share your journal and glucose trends with your doctor or nutritionist so you can fine-tune your diet and treatment plan together.

Reframing Setbacks and Plateaus

The journey to healing neuropathy is rarely a straight line. There will be days when symptoms flare, progress seems stalled, or motivation wanes, and these are not failures, they are opportunities for learning and recalibration. It's okay to feel frustrated or discouraged, and you can acknowledge and validate those feelings without letting them consume you. Instead of asking "why me," try asking what you can learn from this setback, such as whether you deviated from your diet, whether stress was particularly high, or whether you missed sleep. These insights are invaluable. Try reframing the narrative, too: instead of "I'm stuck," think "this is a temporary challenge, and I have the tools to navigate it." Focus on consistency, not perfection, since one bad day or meal won't derail your entire progress, you simply get back on track with the next choice. And avoid the comparison trap. Everyone's healing journey is unique, so focus on your own personal improvements instead of comparing your progress to others.

Tracking Wins: Non-Scale Victories

While objective data, blood markers and symptom scores, is important, celebrating "non-scale victories" is crucial for maintaining motivation and recognizing subtle but significant improvements. These are the daily signs that your efforts are paying off, even if the primary symptoms haven't fully resolved. They can be as simple as noticing, "I walked an extra five minutes today without increased pain," "my sleep was more restful last night," "I noticed a slight return of sensation in my toes," "I navigated a stressful situation without my symptoms flaring," "I cooked a healthy meal even when I felt tired," "I've consistently taken my supplements for a week," "my energy levels feel more stable," or "I feel less anxious about my condition." Dedicate a section of your journal, or a separate notebook, to record these wins as a victory journal, no matter how small they seem. Reviewing them regularly can provide a powerful boost to your morale and reinforce positive habits.

Sample Blood Marker Tracking Chart

Beyond daily glucose, keeping track of your broader health markers over time allows you to see the impact of your holistic interventions. This data, combined with your symptom journal, provides a comprehensive

picture of your progress. The chart below is designed for you to fill in as you retest over the following months, alongside the optimal range your practitioner is working toward with you.

Marker	Initial Value (Date)	Optimal Range	Value (Month 1)	Value (Month 3)	Value (Month 6)	Notes / Changes
HbA1c		< 5.4%				
Fasting Insulin		< 5 µIU/mL				
hs-CRP		< 1.0 mg/L				
Homocysteine		5 to 7 µmol/L				Lower is not better; below this window suggests overmethylation
Vitamin D (25-OH D)		40 to 80 ng/mL				
Vitamin B12		> 500 pg/mL (functional)				Look at MMA and homocysteine for true deficiency
RBC Magnesium		Optimal range is lab dependent				
Omega-3 Index		> 8%				
Thyroid Stimulating Hormone (TSH)		1.0 to 2.5 µIU/mL				Also check Free T3, Free T4, and antibodies
Other specific markers (e.g., autoantibodies)		As per lab and diagnosis				

Always discuss these results and optimal ranges with your healthcare practitioner.

Cultivating a positive mindset and diligently monitoring your progress are not passive activities; they are active components of your healing journey. They empower you to become an engaged participant in your own recovery, helping you to adapt, persevere, and ultimately, unlock greater well-being.

Your Path to Reclaiming Nerve Health

You have covered a lot of ground on neuropathy, moving beyond symptom management to embrace a holistic, root-cause approach. The aim of this guide was to empower you with knowledge, practical strategies, and the understanding that your body possesses an incredible capacity for healing, given the right support.

We have unveiled five critical keys to unlocking your neuropathy, each representing a vital area for investigation and intervention. Key 1, unmasking nutrient deficiencies, recognizes that optimal nerve function relies on a precise symphony of vitamins and minerals, and that deficiencies can be silent saboteurs. Key 2, addressing chronic infections and their impact, is about understanding how hidden or persistent pathogens can ignite systemic inflammation and autoimmune responses that directly harm your nervous system. Then comes Key 3, activating large diameter afferent pathways, which is about learning to stimulate specific nerve fibers through various therapies, from vibration and tactile stimulation to cutting-edge photobiomodulation and specialized electrical stimulation devices, to modulate pain, improve sensation, and foster neuroplasticity. Key 4, comprehensive testing beyond the conventional, means advocating for a deeper diagnostic dive into inflammation, metabolic imbalances, circulation, and toxicity to pinpoint the precise underlying drivers of your condition. And Key 5, diet and food sensitivities as tools for fueling nerve repair, means harnessing the immense power of an anti-inflammatory, nutrient-dense diet and identifying personal food triggers to create an optimal internal environment for nerve healing.

Beyond these core pillars, we also highlighted the indispensable role of the movement plan in Chapter 7, promoting circulation, strengthening muscles, and directly impacting nerve health through exercise. And in mindset and monitoring, Chapter 8, we explored how emotional well-being, stress management, and diligent tracking are powerful, often overlooked, components of a successful recovery journey.

Embracing the Holistic Journey

Reclaiming your nerve health is not a sprint; it is a marathon. There will be days of progress, and there may be days of frustration. The most crucial takeaway is this: you are not passive in this process. Every choice you make, from the food you eat to the way you manage stress and the movement you incorporate, sends a message to your nervous system.

This ebook has provided you with a comprehensive framework. However, this is a guide, not a prescription. Your body is unique, and your healing journey will be too. We strongly advocate for working closely with a qualified healthcare professional, ideally a functional medicine practitioner, neurologist, or physical therapist experienced in holistic neuropathy approaches. They can help you interpret test results, fine-tune supplement protocols, and tailor exercise routines to your specific needs and condition.

As you move forward from here, a few next steps are worth taking deliberately. Review and reflect. Revisit the keys that resonated most with you, and consider which areas you suspect might be most relevant to your neuropathy. Consult a professional. Seek out a healthcare provider who understands and practices the root-cause approach, and share this guide with them as a starting point for discussion. Implement gradually. Don't try to change everything at once; pick one or two areas to focus on first, such as beginning with the dietary principles or integrating simple daily movements. And be patient and persistent. Healing takes time, so celebrate every small victory, learn from setbacks, and remain committed to nurturing your body.

Neuropathy doesn't have to be a life sentence of pain and limitation. By understanding its many contributing factors and proactively addressing the root causes, you have the power to unlock your neuropathy and reclaim a life of greater comfort, vitality, and well-being. This is your path forward.

Part 4: The 90-Day Battle Plan, Meal Plan and Recipes

Embarking on a new dietary path can feel overwhelming, but with a structured plan, it becomes an exciting journey of discovery and healing. This 90-day battle plan is designed to transition you into a sustainable, nerve-supportive eating lifestyle. It's about giving you choices that fuel repair, reduce inflammation, and stabilize blood sugar, rather than about strict deprivation.

The plan is divided into three phases, each building upon the last, allowing your body to adapt and for you to identify your unique needs and sensitivities. Remember, consistency, not perfection, is the goal.

Weekly Framework: The 60/20/20 Guideline

This general guideline provides a macronutrient breakdown for balanced meals that promote stable blood sugar and sustained energy throughout your day. About 60 percent of your plate should be healthy carbohydrates, coming primarily from a wide variety of non-starchy vegetables, some low-glycemic fruits like berries, and carefully selected whole grains if reintroduced and tolerated, focusing on complex carbohydrates rich in fiber. About 20 percent should be healthy fats, essential for nerve cell membranes and myelin, incorporating sources like extra virgin olive oil, avocados, nuts such as almonds and walnuts, seeds such as chia, flax, and hemp, and fatty fish such as wild-caught salmon, mackerel, and sardines, rich in anti-inflammatory omega-3 fatty acids. And the remaining 20 percent should come from quality proteins, crucial for tissue repair, muscle maintenance, and neurotransmitter synthesis, prioritizing grass-fed or pasture-raised meats, wild-caught fish, organic poultry, and plant-based options like legumes, if tolerated, and quinoa. This 60/20/20 guideline is a general starting point. Your individual needs may vary based on activity level, specific health conditions, and personal tolerance, and consulting with a nutritionist or functional medicine practitioner can help you fine-tune these percentages.

Phase 1: Weeks 1 Through 4, Elimination and Detoxification

This initial phase focuses on removing common inflammatory foods and potential sensitivities to calm your system, reduce inflammation, and create a clean slate for healing. The goal is to reduce inflammation, stabilize initial blood sugar responses, begin gut healing, and identify any immediate dietary triggers. Meals are built around abundant non-starchy vegetables, clean protein, and healthy fats. During this phase, minimize fruits and eliminate all grains, dairy, legumes, corn, soy, and refined sugars. Keep hydration front of mind: drink at least half your body weight in ounces of filtered water daily, and consider starting your day with a glass of warm lemon water.

Phase 2: Weeks 5 Through 8, Reintroduction and Personalization

After the elimination phase, your body has had a chance to reset. This phase is about carefully reintroducing foods, one at a time, to identify what specifically triggers your symptoms, with the goal of pinpointing

individual food sensitivities and expanding your personalized diet based on clear symptom responses. The reintroduction protocol works like this: choose one eliminated food group, for example gluten-free grains like quinoa, and eat a small portion of it daily for two to three days while continuing your otherwise clean diet. Monitor your symptoms closely, including pain, numbness, tingling, digestive issues, fatigue, and brain fog. If symptoms worsen, that food is likely a trigger for you, and you eliminate it again; if no symptoms appear after two to three days, you can likely include that food in your diet in moderation. Either way, wait at least three days before reintroducing the next food group. The menu philosophy continues with the anti-inflammatory base from phase one, but strategically adds foods back in based on your findings.

Phase 3: Weeks 9 Through 12, Maintenance and Optimization

By this phase, you will have a clearer understanding of your body's specific needs and sensitivities. This is where you solidify your personalized anti-inflammatory diet, with the goal of sustaining nerve healing, maintaining stable blood sugar, keeping inflammation low, and optimizing your overall well-being with a sustainable eating pattern. Enjoy a diverse range of your newly identified safe and supportive foods, continue to prioritize nutrient density and mindful eating, and keep hydration regular. This phase becomes your new normal.

A short list of snacks reliably stabilizes glucose between meals throughout all three phases, designed to be nutrient-dense and to prevent blood sugar spikes while keeping your energy stable and supporting nerve health: hard-boiled eggs for portable protein, avocado slices with a sprinkle of sea salt, a small handful of nuts such as almonds or walnuts or seeds such as pumpkin or sunflower, celery sticks with almond or seed butter, berries such as strawberries, blueberries, or raspberries with a dollop of unsweetened coconut yogurt or a few pecans, a small portion of olives, and a small piece of wild-caught salmon or grass-fed, sugar-free beef or turkey jerky.

Grocery List and Meal Prep Hacks

Streamlining your meal preparation is key to sticking with a new dietary plan. Keep these staples on hand.

Produce

- Leafy greens (spinach, kale)
- Cruciferous vegetables (broccoli, cauliflower)
- Asparagus
- Bell peppers
- Zucchini
- Cucumber
- Carrots
- Onions

- Garlic
- Ginger
- Turmeric root
- Berries
- Lemons
- Limes
- Avocados

Proteins

- Wild-caught salmon, cod, and mackerel
- Organic chicken breast and thighs
- Grass-fed beef (ground or steak)
- Organic eggs

Healthy Fats

- Extra virgin olive oil, coconut oil, and avocado oil
- Raw nuts (almonds, walnuts, pecans)
- Raw seeds (chia, flax, hemp, pumpkin, sunflower)

Pantry

- Apple cider vinegar
- Various anti-inflammatory herbs and spices
- Unsweetened almond or coconut milk
- Organic chicken or beef broth
- Gluten-free rolled oats and quinoa (for phase two or three, if tolerated)

A handful of habits make meal prep painless. Plan ahead by dedicating 30 to 60 minutes each week to plan your meals and snacks, which reduces decision fatigue during busy days. Shop smart by focusing on the perimeter of the grocery store, where fresh, whole foods are located, and sticking to your list. Batch cook proteins by cooking larger quantities of lean protein, baked chicken breasts, roasted salmon, or ground turkey, that can be easily added to salads, wraps, or stir-fries throughout the week. Roast a large tray of mixed non-starchy vegetables, broccoli, cauliflower, bell peppers, and zucchini, at the beginning of the week for quick additions to any meal. Wash and dry your leafy greens and pre-chop other vegetables so they're ready for quick assembly. Prepare homemade dressings in advance using olive oil, vinegar, lemon juice, and herbs. And portion out nuts, seeds, or berries into small containers for grab-and-go snacks.

Recipes for Nerve Healing

These fifteen to twenty easy, balanced recipes align with the principles of the anti-inflammatory, blood sugar-stabilizing diet. They are designed to be easy to prepare, delicious, and rich in nutrients vital for nerve health. For the first week, in phase one, focus on the recipes below that are naturally free of common sensitivities like grains, dairy, corn, soy, and legumes; they are marked Phase 1 Approved throughout.

Breakfast Recipes

1. Berry-Spinach Power Smoothie

Phase 1 approved.

This smoothie is packed with antioxidants from the berries and greens, healthy fats from the avocado and seeds, and clean protein, all while staying low glycemic and anti-inflammatory. It yields one serving and takes about five minutes to prepare.

Ingredients

- 1 cup unsweetened almond or coconut milk
- 1 large handful fresh spinach (or de-stemmed kale)
- 1/2 cup mixed berries, fresh or frozen
- 1/4 medium avocado
- 1 scoop unflavored or vanilla plant-based protein powder (no added sugars or artificial sweeteners)
- 1 tbsp chia seeds or flax seeds
- Optional: 1/4 tsp ground turmeric plus a pinch of black pepper

Instructions

1. Combine all ingredients in a high-speed blender.
2. Blend until smooth and creamy, adding more liquid if needed for your desired consistency.
3. Pour into a glass and enjoy immediately.

2. Turmeric Scramble with Avocado

Phase 1 approved.

Eggs are a complete protein source, and turmeric is a potent anti-inflammatory, while the healthy fats from avocado support nerve cell membranes. It yields one serving, takes about five minutes to prep, and seven to ten minutes to cook.

Ingredients

- 2 organic, pasture-raised eggs
- 1 tbsp unsweetened almond or coconut milk (optional, for creaminess)
- 1/2 cup chopped non-starchy vegetables (spinach, bell peppers, or zucchini)
- 1/2 tbsp olive oil or coconut oil
- 1/2 tsp ground turmeric
- Pinch of black pepper
- Sea salt to taste
- 1/2 medium avocado, sliced

Instructions

1. In a bowl, whisk the eggs with the almond milk (if using), turmeric, black pepper, and sea salt.
2. Heat the oil in a non-stick pan over medium heat. Add the chopped vegetables and sauté for 3 to 5 minutes until tender-crisp.
3. Pour the egg mixture into the pan and scramble gently until cooked through.
4. Serve immediately with the sliced avocado on the side.

3. Chia Seed Pudding with Berries and Nuts

Phase 1 approved.

Chia seeds are rich in omega-3s and fiber, promoting gut health and reducing inflammation, and this pudding is low glycemic and highly customizable. It yields one serving and takes about five minutes to prepare, plus at least four hours, or overnight, to chill.

Ingredients

- 3 tbsp chia seeds
- 1 cup unsweetened almond or coconut milk
- 1/2 tsp vanilla extract (sugar-free)
- Optional: pinch of cinnamon
- 1/4 cup fresh mixed berries
- 1 tbsp chopped walnuts or almonds

Instructions

1. In a jar or bowl, whisk together the chia seeds, almond milk, vanilla extract, and cinnamon until well combined.

2. Cover and refrigerate for at least 4 hours, or preferably overnight, until thickened to a pudding-like consistency. Stir once or twice in the first hour to prevent clumping.
3. Before serving, top with the fresh berries and chopped nuts.

Lunch Recipes

4. Giant Anti-Inflammatory Salad with Grilled Chicken or Salmon

Phase 1 approved.

This salad maximizes nutrient intake from diverse greens and vegetables, provides clean protein and healthy fats, and is highly anti-inflammatory. It yields one serving, takes about fifteen minutes to prep, and ten to fifteen minutes to cook the protein.

Ingredients

- 2 to 3 cups mixed leafy greens (spinach, kale, romaine, or spring mix)
- 1 cup chopped non-starchy vegetables (cucumber, bell peppers, shredded carrots, or radish)
- 4 to 5 oz grilled organic chicken breast, wild-caught salmon fillet, or 1 can wild salmon or sardines
- 1/4 medium avocado, sliced or diced
- 1/4 cup toasted pumpkin seeds or sunflower seeds
- **For the lemon-herb vinaigrette:**
 - 2 tbsp extra virgin olive oil
 - 1 tbsp fresh lemon juice
 - 1/2 tsp dried Italian herbs or fresh chopped parsley or dill
 - Pinch of sea salt and black pepper

Instructions

1. Prepare your protein: grill the chicken or salmon, or drain the canned fish. Let it cool slightly and flake the salmon or sardines if using.
2. In a large bowl, combine the mixed greens and chopped vegetables.
3. Arrange the protein and avocado slices on top, then sprinkle with the toasted seeds.
4. In a small bowl, whisk together the olive oil, lemon juice, herbs, salt, and pepper for the vinaigrette.
5. Drizzle the dressing over the salad just before serving.

5. Turkey and Avocado Lettuce Wraps

Phase 1 approved.

This is a light, grain-free option packed with lean protein, healthy fats, and crisp vegetables, avoiding inflammatory grains entirely. It yields one serving, or two to three wraps, takes ten to fifteen minutes to prep, and five to seven minutes to cook the turkey.

Ingredients

- 4 to 5 large butter lettuce leaves or romaine lettuce hearts
- 4 oz lean ground turkey or organic ground chicken
- 1/2 cup finely diced bell peppers (any color)
- 1/4 cup celery, finely diced
- 1/4 cup cucumber, finely diced
- 1/4 medium avocado, mashed
- 1 tbsp olive oil
- 1 tsp apple cider vinegar
- 1/2 tsp garlic powder or fresh minced garlic
- Pinch of red pepper flakes (optional, for a kick)
- Sea salt and black pepper to taste

Instructions

1. Heat the olive oil in a skillet over medium heat. Add the ground turkey or chicken and cook, breaking it up with a spoon, until browned and cooked through. Drain any excess fat.
2. Remove the cooked turkey from the heat and let it cool slightly.
3. In a bowl, combine the cooled turkey with the diced bell peppers, celery, cucumber, mashed avocado, apple cider vinegar, garlic powder, red pepper flakes (if using), salt, and pepper. Mix well.
4. Spoon the mixture into the lettuce leaves and wrap them up. Serve immediately.

Dinner Recipes

6. Baked Salmon with Roasted Asparagus and Brussels Sprouts

Phase 1 approved.

Wild salmon is rich in omega-3s, EPA and DHA, crucial for nerve health and reducing inflammation, while the roasted vegetables provide fiber and antioxidants. It yields two servings, takes about ten minutes to prep, and twenty to twenty-five minutes to cook.

Ingredients

- 2 wild-caught salmon fillets (4 to 5 oz each)

- 1 bunch asparagus, woody ends trimmed
- 1 cup Brussels sprouts, halved
- 2 tbsp olive oil, divided
- 1 lemon, half sliced and half reserved for juicing
- 2 cloves garlic, minced (or 1 tsp garlic powder)
- 1/2 tsp dried dill (optional)
- Sea salt and black pepper to taste

Instructions

1. Preheat the oven to 400°F (200°C) and line a baking sheet with parchment paper.
2. On the baking sheet, toss the asparagus and Brussels sprouts with 1 tbsp of the olive oil, the minced garlic, salt, and pepper. Spread in a single layer.
3. Place the salmon fillets on the same baking sheet, in among the vegetables. Drizzle the salmon with the remaining 1 tbsp olive oil, sprinkle with dill (if using), salt, and pepper. Place lemon slices on top of each fillet.
4. Bake for 15 to 20 minutes, or until the salmon is cooked through and flakes easily and the vegetables are tender-crisp.
5. Serve immediately with a squeeze of fresh lemon juice.

7. Chicken and Vegetable Stir-Fry with Cauliflower Rice

Phase 1 approved.

This is a quick, versatile meal packed with protein and a rainbow of vegetables, and the cauliflower rice makes it a low-carb, grain-free alternative to a traditional stir-fry. It yields two servings and takes about fifteen minutes to prep and fifteen minutes to cook.

Ingredients

- 1 tbsp coconut oil or avocado oil
- 1 lb organic chicken breast or thighs, cut into 1-inch pieces
- 2 cups mixed stir-fry vegetables (broccoli florets, sliced bell peppers, shredded carrots, snap peas, sliced mushrooms, bok choy)
- 1 clove garlic, minced
- 1 tbsp fresh ginger, grated
- 2 cups cauliflower rice, fresh or frozen
- **For the sugar-free stir-fry sauce:**

- 2 tbsp coconut aminos
- 1 tbsp apple cider vinegar
- 1 tsp sesame oil (optional, for flavor)
- 1/4 cup filtered water or broth
- Pinch of red pepper flakes (optional)

Instructions

1. Heat the coconut oil in a large skillet or wok over medium-high heat. Add the chicken and cook until browned and cooked through, about 5 to 7 minutes. Remove from the pan and set aside.
2. Add the garlic and ginger to the pan and stir-fry for 30 seconds until fragrant.
3. Add the harder vegetables (broccoli and carrots) and stir-fry for 3 to 4 minutes. Then add the softer vegetables (bell peppers, mushrooms, snap peas, and bok choy) and stir-fry for another 3 minutes until everything is tender-crisp.
4. In a small bowl, whisk together all the stir-fry sauce ingredients.
5. Return the chicken to the pan with the vegetables. Add the cauliflower rice and pour the sauce over everything. Stir well to combine and heat through, about 2 to 3 minutes.
6. Serve hot.

8. Lean Ground Beef and Broccoli Bowls

Phase 1 approved.

This bowl provides quality protein and healthy fats, and broccoli brings its detoxifying, anti-inflammatory power to the plate. It yields two servings, takes about ten minutes to prep, and fifteen minutes to cook.

Ingredients

- 1 tbsp olive oil
- 1 lb grass-fed lean ground beef
- 1 large head broccoli, cut into small florets
- 1/2 onion, finely diced (optional)
- 2 cloves garlic, minced
- **For the sauce:**
- 3 tbsp coconut aminos
- 1 tbsp apple cider vinegar
- 1/2 tsp fresh grated ginger (optional)
- Pinch of red pepper flakes (optional)

- Sea salt and black pepper to taste

Instructions

1. Heat the olive oil in a large skillet or deep pan over medium-high heat. Add the ground beef and onion (if using) and cook, breaking up the meat with a spoon, until browned. Drain any excess fat.
2. Add the minced garlic and ginger and cook for 1 minute until fragrant.
3. Add the broccoli florets and cook, stirring occasionally, for 5 to 7 minutes until tender-crisp.
4. In a small bowl, whisk together the coconut aminos, apple cider vinegar, and red pepper flakes (if using).
5. Pour the sauce over the beef and broccoli. Stir well to coat and cook for another 2 to 3 minutes until the sauce slightly thickens.
6. Season with salt and pepper to taste and serve hot.

9. Quinoa Salad with Roasted Vegetables

Phase 2 or 3 approved, if quinoa is tolerated.

Quinoa is a complete protein and provides complex carbohydrates and fiber, while the roasted vegetables add antioxidants; make sure quinoa is tolerated after your reintroduction phase before relying on this one. It yields two servings, takes about fifteen minutes to prep, and twenty to twenty-five minutes to cook.

Ingredients

- 1 cup cooked quinoa, rinsed well before cooking
- 1 cup chopped mixed vegetables for roasting (zucchini, bell peppers, red onion, broccoli florets)
- 1 tbsp olive oil
- Pinch of sea salt and black pepper
- 1/2 cup chopped fresh parsley or cilantro
- Optional: 1/4 cup crumbled feta or goat cheese (Phase 3, if dairy is tolerated)
- **For the lemon-tahini dressing:**
 - 2 tbsp tahini
 - 2 tbsp fresh lemon juice
 - 1 tbsp water, or more to thin
 - 1 clove garlic, minced
 - Pinch of sea salt

Instructions

1. Preheat the oven to 400°F (200°C). Toss the chopped vegetables with the olive oil, salt, and pepper on a baking sheet. Roast for 15 to 20 minutes, or until tender and slightly caramelized.
2. In a large bowl, combine the cooked quinoa, roasted vegetables, and fresh herbs.
3. For the dressing, whisk together the tahini, lemon juice, water, minced garlic, and salt until smooth. Add more water if needed to reach your desired consistency.
4. Pour the dressing over the quinoa and vegetable mixture and toss to combine.
5. Serve warm or at room temperature, topped with the crumbled cheese if you're including it.

10. Lentil Soup with Root Vegetables

Phase 2 or 3 approved, if legumes are tolerated.

Lentils provide plant-based protein and fiber, promoting stable blood sugar and gut health, while the root vegetables add further nutrients; this one is for once legumes have been successfully reintroduced. It yields four servings, takes about fifteen minutes to prep, and forty-five to sixty minutes to cook.

Ingredients

- 1 tbsp olive oil
- 1 medium onion, diced
- 2 carrots, diced
- 2 celery stalks, diced
- 2 cloves garlic, minced
- 1 cup green or brown lentils, rinsed
- 6 cups vegetable broth, low sodium, no sugar added
- 1 can (14.5 oz) diced tomatoes, no sugar added, BPA-free can
- 1 tsp dried thyme
- 1/2 tsp dried rosemary
- Sea salt and black pepper to taste
- Fresh parsley for garnish (optional)

Instructions

1. Heat the olive oil in a large pot or Dutch oven over medium heat. Add the onion, carrots, and celery and sauté for 8 to 10 minutes until softened.
2. Add the minced garlic and cook for another minute until fragrant.
3. Stir in the rinsed lentils, vegetable broth, diced tomatoes, thyme, and rosemary. Bring to a boil, then reduce the heat to low, cover, and simmer for 30 to 45 minutes, or until the lentils are tender.

4. Season with sea salt and black pepper to taste.
5. Ladle into bowls and garnish with fresh parsley if desired.

Dinner Recipes, Continued

11. Baked Cod with Lemon-Garlic Herbs

Phase 1 approved.

Cod is a lean, easily digestible protein source, and the lemon and garlic add flavor without any inflammatory ingredients. It yields two servings, takes about ten minutes to prep, and twelve to fifteen minutes to cook.

Ingredients

- 2 cod fillets (6 oz each), or another white fish like halibut or tilapia
- 2 tbsp olive oil
- 2 cloves garlic, minced
- 1/2 lemon, thinly sliced
- 1 tbsp fresh chopped parsley (or 1 tsp dried)
- 1/2 tsp dried oregano
- Sea salt and black pepper to taste

Instructions

1. Preheat the oven to 400°F (200°C) and line a baking sheet with parchment paper. Place the cod fillets on the sheet.
2. In a small bowl, whisk together the olive oil, minced garlic, chopped parsley, oregano, salt, and pepper.
3. Pour the mixture over the cod fillets so they are well coated, and top each fillet with a few lemon slices.
4. Bake for 12 to 15 minutes, or until the fish is opaque and flakes easily with a fork.
5. Serve immediately with your favorite non-starchy steamed or roasted vegetables.

12. Turkey and Vegetable Chili, No Beans

Phase 1 approved.

This is a hearty, warming meal packed with lean protein and fiber from vegetables, highly customizable and anti-inflammatory without the typical bean load. It yields four to six servings, takes about fifteen minutes to prep, and thirty to forty minutes to cook.

Ingredients

- 1 tbsp olive oil

- 1 lb organic lean ground turkey
- 1 large onion, diced
- 2 celery stalks, diced
- 1 large bell pepper, any color, diced
- 2 cloves garlic, minced
- 1 can (28 oz) crushed tomatoes, no sugar added, BPA-free
- 1 can (14.5 oz) diced tomatoes, no sugar added, BPA-free
- 1 cup beef or vegetable broth, low sodium
- 2 tbsp chili powder, no sugar added
- 1 tsp ground cumin
- 1/2 tsp smoked paprika
- Pinch of cayenne pepper (optional, for heat)
- Sea salt and black pepper to taste
- Optional toppings (sliced avocado, fresh cilantro, green onions)

Instructions

1. Heat the olive oil in a large pot or Dutch oven over medium heat. Add the ground turkey and cook, breaking it up, until browned. Drain any excess fat and set the turkey aside.
2. Add the onion, celery, and bell pepper to the pot and sauté for 8 to 10 minutes until softened.
3. Add the minced garlic and cook for another minute until fragrant.
4. Return the turkey to the pot and stir in the crushed tomatoes, diced tomatoes, beef broth, chili powder, cumin, smoked paprika, and cayenne (if using).
5. Bring the chili to a simmer, then reduce the heat to low, cover, and cook for at least 20 to 30 minutes for the flavors to meld.
6. Season with salt and pepper to taste.
7. Serve hot, topped with sliced avocado, fresh cilantro, or green onions if desired.

13. Sheet Pan Chicken Fajita Bowls, No Tortilla

Phase 1 approved.

This is a simple, flavorful meal with lean protein and plenty of colorful, anti-inflammatory vegetables, and it's great for meal prepping. It yields two servings, takes about fifteen minutes to prep, and twenty-five to thirty minutes to cook.

Ingredients

- 1 lb organic chicken breast, sliced into thin strips
- 2 bell peppers, different colors, sliced
- 1 medium onion, sliced
- 2 tbsp olive oil
- **For the sugar-free fajita seasoning:**
- 1 tbsp chili powder
- 1 tsp ground cumin
- 1 tsp smoked paprika
- 1/2 tsp garlic powder
- 1/2 tsp onion powder
- Pinch of cayenne pepper (optional)
- Sea salt and black pepper to taste
- Optional for serving (sliced avocado, fresh cilantro, lime wedges)

Instructions

1. Preheat the oven to 400°F (200°C) and line a large baking sheet with parchment paper.
2. In a large bowl, combine the sliced chicken, bell peppers, and onion. Drizzle with the olive oil.
3. In a small bowl, mix all the fajita seasoning ingredients. Sprinkle the seasoning over the chicken and vegetables, tossing well to coat evenly.
4. Spread everything in a single layer on the prepared baking sheet.
5. Roast for 25 to 30 minutes, flipping halfway through, until the chicken is cooked through and the vegetables are tender-crisp and slightly charred.
6. Serve hot in bowls, topped with sliced avocado, fresh cilantro, and a squeeze of fresh lime juice.

Snack and Side Recipes

14. Roasted Brussels Sprouts with Balsamic Glaze

Phase 1 approved.

Brussels sprouts are a cruciferous vegetable, rich in compounds that support detoxification and reduce inflammation. It yields two servings, takes about ten minutes to prep, and twenty to twenty-five minutes to cook.

Ingredients

- 1 lb Brussels sprouts, trimmed and halved

- 1 tbsp olive oil
- Sea salt and black pepper to taste
- 2 tbsp balsamic vinegar, no added sugar

Instructions

1. Preheat the oven to 400°F (200°C) and line a baking sheet with parchment paper.
2. In a bowl, toss the Brussels sprouts with the olive oil, salt, and pepper until evenly coated.
3. Spread the Brussels sprouts in a single layer on the baking sheet and roast for 20 to 25 minutes, stirring halfway through, until tender inside and crisp and browned outside.
4. While the sprouts are roasting, pour the balsamic vinegar into a small saucepan. Bring it to a simmer over medium-low heat and cook for 5 to 7 minutes, stirring occasionally, until it reduces to a slightly thickened glaze. Be careful not to let it burn.
5. Once the sprouts are roasted, transfer them to a serving dish and drizzle with the balsamic glaze. Serve immediately.

15. Simple Guacamole with Veggie Sticks

Phase 1 approved.

Avocados are packed with healthy monounsaturated fats, crucial for nerve cell health, while the tomato, if tolerated, and onion add flavor and antioxidants; omit the tomato for a stricter phase one if nightshades are a concern. It yields two to three servings and takes about ten minutes to prep.

Ingredients

- 2 ripe avocados
- 1/4 cup red onion, finely diced (or green onion for a milder flavor)
- 1/4 cup tomato, finely diced (optional, omit for stricter Phase 1 if nightshades are a concern)
- 1 to 2 tbsp fresh lime juice
- 1 tbsp fresh chopped cilantro (optional)
- Pinch of red pepper flakes (optional)
- Sea salt to taste
- Cucumber slices, bell pepper strips, celery sticks and radish slices, for dipping

Instructions

1. In a medium bowl, mash the avocados with a fork to your desired consistency, whether chunky or smooth.
2. Add the diced red onion, tomato (if using), lime juice, chopped cilantro (if using), red pepper flakes (if using), and sea salt.

3. Mix gently to combine. Taste and adjust the seasoning as needed.
4. Serve immediately with the vegetable sticks for dipping.

16. Anti-Inflammatory Bone Broth

Phase 1 approved.

Rich in collagen, amino acids like glycine and proline, and minerals, bone broth supports gut health, reduces inflammation, and provides building blocks for tissue repair, including, potentially, nerve sheaths. Yields vary, since this makes a large batch; it takes about fifteen minutes to prep and eight to twenty-four hours to simmer.

Ingredients

- 2 to 3 lbs organic, grass-fed beef bones (marrow, knuckle, or joint bones), or organic chicken carcass and bones
- 1 gallon filtered water
- 2 tbsp apple cider vinegar (helps extract minerals)
- 1 large onion, quartered, skin on
- 2 carrots, roughly chopped
- 2 celery stalks, roughly chopped
- 2 bay leaves
- 1 tsp black peppercorns
- Optional, for extra flavor and nutrients (fresh ginger slices, turmeric root slices, parsley stems)

Instructions

1. If using beef bones, roast them first for richer flavor: preheat the oven to 400°F (200°C) and roast the bones on a baking sheet for 20 to 30 minutes until browned. Skip this step for chicken bones.
2. Place the bones in a large stock pot or slow cooker. Add the apple cider vinegar and cover with the filtered water. Let sit for 30 minutes before heating, since this helps draw out the minerals.
3. Add the onion, carrots, celery, bay leaves, peppercorns, and any optional additions you're using.
4. Stovetop method: bring the pot to a rolling boil, then immediately reduce the heat to the lowest simmer. Skim off any foam that rises to the surface during the first hour. Cover and simmer for at least 8 to 12 hours for chicken bones, or 12 to 24 for beef, the longer the better.
5. Slow cooker method: set it to low and cook for 12 to 24 hours instead.
6. Once done, remove the large bones and vegetables with tongs or a slotted spoon.
7. Strain the broth through a fine-mesh sieve or a cheesecloth-lined colander into clean jars or containers.

8. Let cool completely before refrigerating. The fat will solidify on top; you can skim it off or leave it.
9. Store in the fridge for up to 5 days, or freeze for up to 6 months. Sip warm plain or use as a base for soups.

Your Next Step: A Free Root Discovery Session

This ebook has shown you the what and the why behind your neuropathy, from the nutrient gaps and hidden infections to the nerve pathways, the testing, and the food choices that all play a part. That understanding alone puts you ahead of where you started, and it is real progress.

Finding your specific root cause, though, takes real testing and a plan built around your body, not a generic one. If you are ready to find out what is actually driving your neuropathy, the next step is a free, no-pressure conversation with me. We will talk through your history, your symptoms, and what a personalized path forward could look like for you. [Book your free Root Discovery Session here.](#) You can also reach the office at (919) 662-0520.

Consultations and ongoing care are both available wherever you are located. Being outside North Carolina is not a barrier to working with me. Tell me where you are when you reach out and I will lay out exactly how we would work together from there.

If you would rather start on your own, begin with the 90-day plan in this ebook and log your symptoms against your own baseline so the pattern is on paper before we talk.

Dr. Adam Seay, Root Health, Garner NC

For educational purposes only. This ebook does not constitute medical advice, diagnosis, or treatment, and reading it does not establish a patient-provider relationship.

References

1. Alvarez et al., "B Vitamin Deficiencies and Associated Neuropathies," Current Nutrition Reports, 2026. <https://pubmed.ncbi.nlm.nih.gov/41609902/>
2. Muhamad et al., "The Role of Vitamin B6 in Peripheral Neuropathy: A Systematic Review," Nutrients, 2023. <https://pubmed.ncbi.nlm.nih.gov/37447150/>
3. Koike et al., "Clinicopathologic features of folate-deficiency neuropathy," Neurology, 2015. <https://pubmed.ncbi.nlm.nih.gov/25663227/>
4. Sepassi et al., "Associations between long-term metformin use, vitamin B12 deficiency, and neuropathy," Diabetes Research and Clinical Practice, 2025. <https://pubmed.ncbi.nlm.nih.gov/40850508/>
5. Gilbody and Gilbody, "Vitamin D for Painful Diabetic Neuropathy: A Systematic Review and Meta-Analysis," Endocrinology, Diabetes & Metabolism, 2025. <https://pubmed.ncbi.nlm.nih.gov/41139799/>
6. Shin et al., "Magnesium and Pain," Nutrients, 2020. <https://pubmed.ncbi.nlm.nih.gov/32718032/>
7. Han et al., "A systematic review and meta-analysis of alpha-lipoic acid in the treatment of diabetic peripheral neuropathy," European Journal of Endocrinology, 2012. <https://pubmed.ncbi.nlm.nih.gov/22837391/>
8. Li et al., "Acetyl-L-carnitine in the treatment of peripheral neuropathic pain: a systematic review and meta-analysis of randomized controlled trials," PLoS One, 2015. <https://pubmed.ncbi.nlm.nih.gov/25751285/>
9. Zhang et al., "Omega-3 polyunsaturated fatty acid oral supplements for improving peripheral nerve health: a systematic review and meta-analysis," Nutrition Reviews, 2020. <https://pubmed.ncbi.nlm.nih.gov/31532492/>
10. Halperin, "Lyme disease and the peripheral nervous system," Muscle & Nerve, 2003. <https://pubmed.ncbi.nlm.nih.gov/12872317/>
11. "Peripheral neuropathy as a complication of neuroborreliosis," International Journal of Infectious Diseases, 2021. <https://pubmed.ncbi.nlm.nih.gov/33684563/>
12. "Bartonella- and Borrelia-related disease presenting as a neurological condition," Microorganisms, 2024. <https://pubmed.ncbi.nlm.nih.gov/38276194/>
13. Mallick-Searle et al., "Postherpetic neuralgia: epidemiology, pathophysiology, and pain management pharmacology," Journal of Multidisciplinary Healthcare, 2016. <https://pubmed.ncbi.nlm.nih.gov/27703368/>
14. Gavin et al., "Cranial and peripheral neuropathy due to Epstein-Barr virus infection," Postgraduate Medical Journal, 1997. <https://pubmed.ncbi.nlm.nih.gov/9338028/>
15. Nemni et al., "Peripheral neuropathy in hepatitis C virus infection with and without cryoglobulinaemia," Journal of Neurology, Neurosurgery & Psychiatry, 2003. <https://pubmed.ncbi.nlm.nih.gov/12933932/>
16. Pane et al., "Role of gut microbiota in neuropathy and neuropathic pain states," Neurobiology of Disease, 2022. <https://pubmed.ncbi.nlm.nih.gov/35623598/>
17. Daxboeck et al., "Mycoplasma pneumoniae central nervous system infections," Current Opinion in Neurology, 2006. <https://pubmed.ncbi.nlm.nih.gov/16914976/>
18. "Low level laser therapy for painful diabetic peripheral neuropathy," Diabetes & Metabolic Syndrome, 2019. <https://pubmed.ncbi.nlm.nih.gov/31405692/>
19. Holanda et al., "The mechanistic basis for photobiomodulation therapy of neuropathic pain," Lasers in Surgery and Medicine, 2017. <https://pubmed.ncbi.nlm.nih.gov/28075022/>

20. Sommer et al., "Inflammation in the pathophysiology of neuropathic pain," *Pain*, 2018. <https://pubmed.ncbi.nlm.nih.gov/29447138/>
21. Li et al., "The relationship between plasma homocysteine levels and diabetic peripheral neuropathy," *Zhonghua Nei Ke Za Zhi*, 2011. <https://pubmed.ncbi.nlm.nih.gov/21418881/>
22. Kirthi et al., "Prevalence of peripheral neuropathy in pre-diabetes: a systematic review," *BMJ Open Diabetes Research & Care*, 2021. <https://pubmed.ncbi.nlm.nih.gov/34006607/>
23. Bourque, "Autoimmune peripheral neuropathies," *Clinica Chimica Acta*, 2015. <https://pubmed.ncbi.nlm.nih.gov/25748038/>
24. Koszewicz et al., "The impact of chronic co-exposure to different heavy metals on small fibers of peripheral nerves. A study of metal industry workers," *Journal of Occupational Medicine and Toxicology*, 2021. <https://pubmed.ncbi.nlm.nih.gov/33858451/>
25. Hadjivassiliou et al., "Neuropathy associated with gluten sensitivity," *Journal of Neurology, Neurosurgery, and Psychiatry*, 2006. <https://pubmed.ncbi.nlm.nih.gov/16835287/>
26. Knaepen et al., "Neuroplasticity, exercise-induced response of peripheral brain-derived neurotrophic factor: a systematic review of experimental studies in human subjects," *Sports Medicine*, 2010. <https://pubmed.ncbi.nlm.nih.gov/20726622/>
27. Zhang et al., "Exercise for Neuropathic Pain: A Systematic Review and Expert Consensus," *Frontiers in Medicine*, 2021. <https://pubmed.ncbi.nlm.nih.gov/34901069/>

From Understanding to Doing

Everything up to this point was about understanding what is happening in your nerves, why it may be happening, and which directions are worth investigating. That is the half a book can do well.

What follows is the other half, and it is a different kind of document on purpose. Three guides, written to be used rather than read straight through. The first tells you which lab panels answer which question and what the numbers mean, and it includes a request list you can hand to a clinician. The second covers what to eat and, more usefully, how to test a dietary change on yourself so you end up with an answer instead of a habit. The third walks through what has actually been studied for nerve symptoms, at what dose, and names the products whose trials came back negative.

Each one stands alone, so use them in whatever order matches what you are stuck on. If you are not sure where to begin, begin with the lab guide, because knowing what is driving your symptoms changes what you should do about the other two.

A note on repetition. A few points appear in both halves, the vitamin B6 ceiling in particular. That is deliberate. The ones repeated are the ones where getting it wrong causes harm, and they are worth meeting twice.

What This Guide Is For

You already know something is wrong. The numbness, the tingling, the burning in your feet or hands did not start for no reason, and it is not something you simply have to live with. This guide exists to help you understand what your own lab results actually mean: which panels look at which part of the problem, what a meaningful result on each one suggests, and how the pieces fit together once you can see them side by side.

This is not a treatment plan and it does not replace clinical judgment. What it does is supply the missing translation layer between a stack of lab results and a clear picture of what your nerves are actually dealing with. Understanding your labs is the first step in the Unlock Your Neuropathy framework: it is how you move from an unexplained symptom to a specific, investigable set of drivers.

What this guide does not do is just as important as what it does. It does not tell you what dose to take, what product to buy, or what your personal protocol should be. Those decisions depend on your full history and belong in a conversation with me or your treating provider, not in a self-serve document. If a result concerns you, or if you are experiencing any of the warning signs in the closing section, seek clinical review rather than waiting.

For educational purposes only. This guide does not constitute medical advice, diagnosis, or treatment, and reading it does not establish a patient-provider relationship.

Panel Overview: Which Labs for Neuropathy Specifically, and Why

Nerve symptoms can come from several different directions at once, which is why no single lab panel tells the whole story. Each of the panels in this guide looks at a different layer of the problem, and understanding what each one is built to see helps you understand why your provider might order more than one.

The Neural Zoomer Plus is the direct autoimmune and infection lens. It screens antibody reactivity across several neurological categories, including markers aimed squarely at peripheral neuropathy and at demyelination, the loss of the insulating sheath around nerve fibers, along with broader categories covering blood-brain barrier disruption and chronic infections. The core question this panel answers is whether your immune system is actively reacting against nerve or myelin tissue, and whether that reaction is isolated to the peripheral nerves in your hands and feet or part of a broader systemic pattern.

The Organic Acids Test, usually shortened to OAT, is the functional and metabolic lens. It measures urinary metabolites that reflect your B-vitamin status, your mitochondrial energy production, and your gut microbial activity in real time. Where the Neural Zoomer asks whether your immune system is attacking nerve tissue, the OAT asks something more basic: whether your cells currently have what they need, in terms of raw materials and usable energy, to build and maintain myelin and to keep nerve tissue running.

Standard blood chemistry, the kind drawn at any LabCorp or Quest location, fills in the metabolic-driver lens through HbA1c, fasting glucose, and fasting insulin. These three numbers, read together, are how you catch diabetic and prediabetic neuropathy, which remains the single most common cause of peripheral neuropathy worldwide. Vitamin B12 with methylmalonic acid, ordered either as its own blood draw or as part of the OAT, adds functional B12 status at the cellular level rather than just the circulating serum number, because B12 deficiency is a common and reversible cause of neuropathy and a normal serum B12 result can still coexist with a real functional deficiency.

Three additional panels round out the picture for specific circumstances. The Wheat Zoomer looks at gluten peptide reactivity and gut barrier markers, occludin and zonulin antibodies, and becomes relevant when Demyelination markers are elevated because of a documented molecular mimicry relationship between gliadin and myelin basic protein. The Heavy Metals Panel measures your body burden of neurotoxic metals, including arsenic and lead, both of which are documented causes of peripheral neuropathy entirely on their own, independent of any autoimmune or metabolic driver. And the Tickborne Diseases Panel screens for *Borrelia* species, *Bartonella*, and other tickborne and opportunistic co-infections, an infectious driver that standard two-tiered Lyme testing frequently misses; the panel's own intended use lists numbness, tingling, and Bell's palsy among the qualifying neurological symptoms.

You do not need every panel to start making sense of your symptoms. Section 7 walks through which panel to prioritize based on your own history.

Before You Read: Four Rules for Interpreting These Results

Read patterns, not single markers. One elevated antibody, especially a borderline one, carries less weight than several markers moving together across categories. A pattern across categories, for example a Peripheral Neuropathy marker plus a Blood-Brain Barrier marker, points toward a systemic process rather than an isolated local one, and that changes what is worth investigating next.

On the Neural Zoomer Plus, borderline results matter. The panel's reference ranges were established using a large healthy-adult population, and results are reported as In Control, Moderate Risk or borderline, or High Risk or positive. A borderline result is sub-threshold immune reactivity, not nothing, particularly when several categories are borderline at once or your symptom picture is significant.

The OAT reflects your current biochemical state, not a permanent deficit. Organic acids are urinary metabolites that shift as the underlying driver shifts. An elevated methylmalonic acid today does not mean your B12 status is permanently damaged. It means something is limiting B12 utilization right now, and the useful question is what.

Testing does not replace a clinical exam. Nerve conduction studies (NCS) and electromyography (EMG), the standard tests neurology uses to directly characterize nerve damage, answer a different question than the labs in this guide. NCS and EMG tell you what damage already exists. The panels here are aimed at why it is happening. Where red-flag symptoms are present, described in the closing section, NCS and EMG belong alongside this workup, not instead of it.

Section 1: Neural Zoomer Plus, Peripheral Neuropathy Category

This category targets ganglioside lipids and nerve-terminal proteins directly implicated in immune-mediated peripheral nerve disease, the category most directly relevant to numbness, tingling, and burning in the hands and feet.

Anti-GM1 targets a ganglioside lipid found in peripheral nerve and muscle-cell membranes. A positive result here is associated with multifocal motor neuropathy, present in most people who carry this diagnosis, along with Guillain-Barre syndrome and lower motor neuron syndromes. Its close relative, Anti-GM2, targets a related ganglioside lipid that also functions as a peripheral nerve antigen, and shows up in the same family of conditions: peripheral neuropathies presenting with tingling, numbness, sensation loss, burning in the feet or hands, and sharp stabbing pains, plus chronic motor neuropathy and Guillain-Barre syndrome. Two things are worth knowing if either of these came back elevated. *Campylobacter jejuni* infection is a documented trigger for Guillain-Barre syndrome through a mechanism called molecular mimicry, where the immune system's response to the bacteria cross-reacts with nerve tissue, so a recent significant gastrointestinal infection is relevant history worth mentioning to your provider. And these two antibodies carry a notably high co-occurrence with celiac disease, along with a recommendation to monitor blood sugar and HbA1c, given how common peripheral neuropathy is in that population. That is an explicit tie between this category and both the gluten and gut picture and the metabolic picture described later in this guide.

The remaining three markers in this category point toward rarer, more specific conditions that warrant direct physician discussion rather than a general nerve-support approach. Anti-Hu, also called ANNA-1, targets neuronal antigens involved in maintaining neuron identity, and is the antibody most frequently associated with paraneoplastic subacute sensory neuronopathy, a nerve condition that can arise in connection with an underlying cancer. Anti-Ri, or ANNA-2, targets neuronal nuclear antigens and is similarly linked to paraneoplastic neurological syndromes. Anti-Amphiphysin targets a membrane protein at nerve terminals involved in neurotransmitter release, and is associated with stiff-person syndrome as well as paraneoplastic neuropathy. None of these three should be read as routine nerve-support findings; a positive result on any of them belongs in front of a physician promptly.

If any marker in this category is elevated, the next question is whether it co-elevates with anything in the Demyelination, Blood-Brain Barrier, or Infections categories covered in the next two sections. Co-elevation across categories points toward a systemic neurological immune process rather than an isolated local nerve problem.

Section 2: Neural Zoomer Plus, Demyelination Category

Myelin is the fatty sheath that insulates your nerve fibers and lets electrical signals travel quickly from node to node, a process called saltatory conduction. When antibodies target the proteins that build or organize that sheath, signal transmission slows, which produces much of the numbness-and-tingling pattern common to neuropathy.

Anti-MAG targets myelin-associated glycoprotein, a protein involved in signaling between neurons and the glial cells that support them in both the central and peripheral nervous system. A positive result is associated with a specific, well-described condition, Anti-MAG peripheral neuropathy, marked by depressed reflexes, weakness, discriminative sensory loss, and slowed nerve conduction velocity. Anti-Neurofascin targets a protein that organizes the nodes of Ranvier, the small gaps along a nerve fiber that fast signal conduction depends on, and elevation here is associated with chronic inflammatory demyelinating polyneuropathy, Guillain-Barre syndrome, and multifocal motor neuropathy. Anti-MOG targets a minor protein sitting on the outermost surface of myelin, exposed and accessible to antibodies, and is linked to MOGAD, a demyelinating disorder distinct from multiple sclerosis whose symptoms include tingling in the arms and legs, muscle weakness, and impaired balance.

Anti-MBP targets myelin basic protein, the second most abundant protein in myelin and a marker of active demyelination when antibodies against it appear; because Anti-MBP is generally absent in healthy people, a positive result here carries real weight, and is associated with multiple sclerosis and other demyelinating disease. Anti-PLP targets a protein that makes up roughly half the protein content of adult central nervous system myelin and is essential for myelin compaction; elevation is linked to multiple sclerosis and optic neuritis, generally reflecting broader myelin autoimmunity rather than an isolated peripheral finding. Anti-Tubulin targets the structural protein of microtubules, the intracellular transport network inside nerve cells, and is associated with Guillain-Barre syndrome and chronic inflammatory demyelinating polyneuropathy; it also runs notably high in autoimmune thyroid disease, elevated in roughly half of Hashimoto's cases and close to that in Graves' disease, and mercury and other heavy metals are known to interfere with the way tubulin assembles into microtubules in the first place.

Two connections are worth carrying with you into later sections of this guide. The first is the gluten connection: myelin basic protein shares a documented molecular mimicry relationship with gliadin, the protein fraction of gluten, which is the basis for why gluten elimination shows up as a recurring theme in functional approaches to demyelinating neuropathy. If your Anti-MBP came back elevated, the Wheat Zoomer in Section 6 is the natural next test to evaluate gluten cross-reactivity. The second is the heavy-metal connection just mentioned under Anti-Tubulin: because mercury and other heavy metals can interfere with microtubule assembly, an elevated Anti-Tubulin result is a reasonable trigger to run the Heavy Metals Panel described later in this guide.

Section 3: Systemic Context, Blood-Brain Barrier and Infections Categories

On their own, these two categories are not the primary focus for most peripheral neuropathy cases. A co-elevation with either one, though, changes how a Peripheral Neuropathy or Demyelination finding should be read, from an isolated local nerve issue to part of a broader systemic process.

The Blood-Brain Barrier Disruption category includes Anti-S100B, Anti-Microglia, and Anti-GFAP, which reflect glial and astrocyte disruption more generally and are less specific to peripheral nerve symptoms on their own. One marker in this category deserves specific attention for neuropathy, however: Anti-GRP78 targets a stress protein found on brain microvascular endothelial cells, and is directly associated with numbness or loss of sensation throughout the body as a symptom, along with weakness in the limbs and painful spasms. That makes it one of the few markers in this category with a documented direct link to peripheral sensory symptoms rather than only central ones.

Within the Infections category, two viral markers have the clearest peripheral-nerve mechanisms. Anti-CMV is the more directly relevant of the two: cytomegalovirus infection is strongly associated with Guillain-Barre syndrome, and a majority of CMV-associated Guillain-Barre cases carry detectable Anti-GM2 antibodies, the same antibody covered in Section 1 that binds peripheral nerves and disrupts neuromuscular transmission. That is an explicit bridge between an Infections-category finding and the Peripheral Neuropathy category. Anti-EBV seropositivity, particularly to the EBNA antigen, correlates with molecular mimicry against myelin basic protein instead, tying it to the Demyelination category rather than to peripheral nerve tissue directly.

If your Peripheral Neuropathy or Demyelination markers are elevated alongside Blood-Brain Barrier or Infections markers, that combination is worth bringing to a Root Discovery Session as a pattern, rather than treating each marker as a separate, unrelated finding. One caveat belongs here as well: tick-borne infection is not covered on the Neural Zoomer Plus. If you have a history of tick exposure, that requires the separate Tickborne Diseases Panel discussed in Section 6.

Section 4: Organic Acids Test, Markers Relevant to Nerve Symptoms

Methylmalonic acid, usually abbreviated MMA, is the single most directly relevant OAT marker for nerve symptoms. Vitamin B12 is a required cofactor for the enzyme that converts methylmalonyl-CoA to succinyl-CoA. When B12 is insufficient or cannot be used at the cellular level, methylmalonyl-CoA gets redirected into methylmalonic acid instead, which then shows up as elevated MMA in urine. Because this reflects functional, cellular-level B12 utilization rather than the circulating serum number, MMA can be elevated even when a standard serum B12 test looks perfectly normal. Functional B12 deficiency specifically tends to travel with diabetes, pregnancy, cancer, renal insufficiency, and conditions marked by increased oxidative stress, and it is worth knowing that metformin, a common diabetes medication, can interfere with B12 absorption and elevate MMA on its own, an additional connection between the metabolic and nutrient-depletion pictures in neuropathy.

Folate, vitamin B9, works alongside B12 in the same pathway, the methylation cycle, and the two are worth understanding together rather than as separate line items on a lab report. The methylation cycle is also responsible for clearing homocysteine, an amino acid that becomes neurotoxic and inflammatory when it builds up in the blood rather than being cleared efficiently. When either B12 or folate runs short, methylation slows, homocysteine rises, and elevated homocysteine is associated with increased inflammation, oxidative stress, and vascular changes, all of which work against healthy nerve tissue. A standard serum folate test, drawn as part of routine blood work, tells you how much folate is circulating at that moment, and it is a genuinely useful starting point. But it does not always tell the full story, because circulating folate still has to be converted, through several enzymatic steps, into its active form, 5-methyltetrahydrofolate, sometimes written as methylfolate or 5-MTHF, before your cells can actually use it. A common variant in the MTHFR gene slows that conversion step. Someone carrying this variant can post an entirely normal serum folate number while still running functionally short on usable folate at the cellular level, the same normal-on-paper-but-insufficient-in-practice pattern already described above for B12 and methylmalonic acid. This is part of why methylation-pathway markers are worth reading together rather than one at a time: homocysteine gives a downstream readout of how well the whole methylation cycle is running, and on the OAT itself, urinary uracil serves a related purpose, because folate is required to convert uracil into thymine during DNA synthesis, so elevated uracil is a recognized signal of folate deficiency or impaired methylation even when serum folate looks adequate.

The practical takeaway, at the educational level this guide operates on, is that a normal folate result does not automatically rule folate out as a contributor to nerve symptoms, particularly if homocysteine is elevated, if uracil is elevated on the OAT, or if an MTHFR variant is known or suspected. Methylfolate, often paired with methyl-B12 and P5P, the active form of B6, is a form that bypasses the conversion step an MTHFR variant slows down, and it has been studied and used in supporting people with peripheral neuropathy. What that means for your own results, and whether it is worth pursuing, is exactly the kind of individualized decision this guide does not make; that conversation belongs in a Root Discovery Session.

Pyridoxic acid, a B6 marker, reflects a nutrient with an unusually broad job description: B6 is a cofactor for more than one hundred enzymatic reactions, including neurotransmitter synthesis and myelin formation, and it also supports B12 absorption. Both deficiency and long-term excess, generally meaning intake above 200 milligrams a day sustained over time, are documented causes of sensory neuropathy, which is why B6 status is worth tracking in either direction rather than assuming that more is automatically better.

Nerve cells, like brain cells, are highly energy-demanding, and the OAT measures several Krebs cycle intermediates, citric acid, cis-aconitic acid, alpha-ketoglutarate, succinic acid, and fumaric acid among others, that together describe how well your mitochondria are producing usable energy. A pattern of elevation across several of these markers points toward impaired mitochondrial energy production, a recognized contributor to both nerve dysfunction and the fatigue that so often

accompanies it. The OAT's gut microbial markers connect back to this same picture in an unexpected way: abnormal gut microbiota is itself a contributor to elevated methylmalonic acid, through increased propionic acid production that interferes with B12-related pathways in the liver, an internal link between the gut picture and the B12 picture that is easy to miss if each marker is read in isolation. The full interpretation of the gut-specific markers is covered in the companion guide, *Unlock Your Gut*.

Section 5: Metabolic and Nutrient Panels Beyond the Zoomer Panels

Diabetic and prediabetic neuropathy is the most common form of peripheral neuropathy worldwide, and the specialty panels described above are not built to measure blood sugar regulation directly. For this piece of the picture, standard chemistry panels, available through any LabCorp or Quest draw, fill the gap.

HbA1c reflects your average blood sugar over roughly the past two to three months. It is a standard test, but looking at an optimal range rather than only a non-diabetic cutoff is worth understanding: chronically elevated blood sugar drives the formation of advanced glycation end products, compounds that bind receptors and trigger inflammatory signaling capable of damaging nerve tissue over time, independent of a formal diabetes diagnosis. Fasting glucose and fasting insulin, measured together, show how much insulin your body needs to keep blood sugar stable. Elevated fasting insulin alongside normal fasting glucose is a sign of developing insulin resistance, often present years before glucose itself becomes abnormal, and this pairing can catch a metabolic driver of neuropathy that a glucose-only panel misses entirely. HOMA-IR is simply a calculated insulin resistance score derived from those same two numbers, a useful way to quantify insulin resistance from the same draw rather than ordering a separate test.

At Root Health, the ranges I consider optimal are deliberately tighter than the standard laboratory cutoff for normal, because the aim is to catch a developing metabolic driver of nerve damage long before it would ever meet the threshold for a diabetes diagnosis. A value drifting above the top of any of these ranges, even one still sitting inside the lab's wider normal range, is worth reading as an early metabolic signal rather than a clean result, and it is exactly the kind of finding that belongs in a conversation about your whole picture rather than a number to act on by itself.

These are clinical targets used to guide investigation. They are not validated diagnostic thresholds, and a result outside a functional window is not a diagnosis of anything.

Marker	Unit	Standard lab range	Root Health functional target
HbA1c	%	4.0 to 5.6	4.6 to 5.3
Glucose, fasting	mg/dL	70 to 99	75 to 86
Insulin, fasting	uIU/mL	2.0 to 10.0	2.6 to 5.0
HOMA-IR	index	0 to 2.5	0 to 1.4
Vitamin B12	pg/mL	200 to 1100	500 to 900
Methylmalonic acid (MMA)	nmol/L	0 to 243	0 to 162
Folate (RBC)	ng/mL	400 to 2000	600 to 1500
Homocysteine	umol/L	0 to 10.0	5 to 7
Vitamin D (25-OH)	ng/mL	20 to 100	40 to 80
Magnesium (RBC)	mg/dL	4.0 to 7.2	5.2 to 6.8
Ferritin	ng/mL	25 to 300	50 to 150
hs-CRP	mg/L	0 to 1.0	0 to 0.5
TSH	mIU/L	0.4 to 4.0	1.0 to 2.5

Four of these deserve a note. Homocysteine is the one where lower is not better: the 5 to 7 functional window is a window rather than a floor, and a homocysteine driven below it points toward overmethylation and B vitamin intolerance, which is its own problem rather than a better result. HOMA-IR has no meaningful lower bound; a low score reflects good insulin sensitivity, so only the upper edge of the window is the signal. Ferritin is the marker most likely to mislead here, because it rises with inflammation as well as with iron stores, so a ferritin sitting inside the functional window alongside an elevated hs-CRP does not reliably confirm adequate iron. And hs-CRP itself is a nonspecific screening marker: an elevated result tells you an inflammatory load exists somewhere, not where it is coming from, and it confirms nothing on its own. Treat it as a reason to keep reading the rest of the panel, never as an answer.

Vitamin B12 with methylmalonic acid, ordered as a standalone blood draw rather than through the OAT, uses the same functional-deficiency logic described in Section 4. Ordering serum B12 alongside MMA, rather than serum B12 alone, is what catches a functional deficiency that a normal serum number would otherwise hide. Serum folate and homocysteine belong in the same conversation. Ordered together, they extend that same functional lens from B12 to folate, and a homocysteine number that is elevated despite a normal serum folate is itself a reason to ask whether an MTHFR variant or another methylation bottleneck, rather than dietary intake, is the limiting factor, exactly as described in Section 4.

Section 6: Additional Panels Worth Knowing About

The Wheat Zoomer becomes relevant given the documented molecular mimicry between gliadin and myelin basic protein described in Section 2. Its gluten peptide reactivity testing, along with gut barrier markers occludin and zonulin antibodies, make it a natural companion test for anyone with elevated Demyelination markers. Zonulin specifically regulates the tight junctions between intestinal cells, and dysregulated zonulin signaling is a documented biomarker for increased gut barrier permeability, sometimes called leaky gut, which has its own downstream effects on systemic inflammation.

The Heavy Metals Panel is worth considering for two metals in particular that have a directly documented, independent peripheral neuropathy mechanism, separate from any autoimmune or metabolic driver. Chronic oral exposure to inorganic arsenic is documented to cause peripheral neuropathy along with anemia and skin changes, and lead is documented to commonly cause neuropathy in adults, in addition to its better-known effects on cognition and mood. This panel earns particular consideration when nerve symptoms have not responded to nutrient, metabolic, or immune-focused work, or where an exposure history, such as older housing, well water, or occupational exposure, is present.

The Tickborne Diseases Panel fills a gap that standard testing genuinely misses: standard two-tiered Lyme testing carries a documented sensitivity of only 30 to 40 percent in early disease, meaning a negative standard screen tells you far less than most patients assume. Neurological symptoms including numbness, tingling, dizziness, and Bell's palsy are listed as qualifying reasons to run the full tickborne panel rather than a standard Lyme screen alone, and *Borrelia garinii* in particular is associated with neurological presentations of Lyme disease. *Bartonella*, one of the tick-borne co-infections tested on this panel, produces its own symptom picture, fever, fatigue, muscle and joint pain, and swollen lymph nodes, that can overlap with and complicate a Lyme diagnosis rather than standing apart from it cleanly.

There is one further pattern worth naming, with a clear caveat about where it comes from. In my clinical experience, patients who turn out to have a *Bartonella* picture often describe their nerve sensations in a particular way: fizzing or electric rather than simply numb, along with buzzing, internal vibrations, or brief jolts. This is a clinical observation offered as something to mention to your provider, not a documented diagnostic sign. The published literature supports the general neurological symptom picture described in the paragraph above; it does not establish any specific sensation quality as *Bartonella*-specific. Treat it as a reason to raise the question, never as grounds to conclude the answer.

Section 7: Testing Decision Framework

You do not need to order everything at once, and where to start depends on your own history. If you have diabetes, prediabetes, or any history of blood sugar issues, start with HbA1c, fasting glucose, and fasting insulin from Section 5, run alongside a Neural Zoomer Plus, since the metabolic and autoimmune pictures are common enough together that checking both from the outset saves a second round of testing later. If your nerve symptoms are progressive or unexplained with no clear metabolic driver in your history, the Neural Zoomer Plus paired with the OAT is the core combination, giving you the immune and the functional-metabolic pictures side by side.

If you have a personal history of gluten sensitivity, a family history of autoimmune disease, or elevated Demyelination markers on the Neural Zoomer, add the Wheat Zoomer to evaluate the gluten connection described in Section 2. With a history of tick exposure, or nerve symptoms that present with prominent numbness, tingling, or a Bell's palsy-type pattern, go directly to the Tickborne Diseases Panel rather than relying on a standard Lyme screen, for the sensitivity reasons described in Section 6. If your history includes older housing, well water, or occupational exposure, the Heavy Metals Panel is worth prioritizing. And if you have a normal serum B12 result despite classic B12-deficiency symptoms, B12 with methylmalonic acid, ordered either as a standalone blood draw or as part of the OAT, is the test that can catch what the serum number alone missed.

Section 8: Reading Your Results Together, Pattern Recognition

Certain combinations of results are more informative read together than any one of them is alone. If your Peripheral Neuropathy category is elevated, particularly Anti-GM1 or Anti-GM2, alongside a positive Wheat Zoomer for gluten reactivity, that combination reflects a documented high co-occurrence between these two antibodies and celiac disease, and points toward the gut and immune pictures being connected rather than coincidental; Section 1 and Section 6 are the places to look next.

If Anti-MAG or Anti-MBP is elevated, the Wheat Zoomer is positive, and methylmalonic acid is elevated, the picture that emerges is molecular mimicry-driven demyelination compounded by a functional B12 deficit that is limiting your body's capacity to repair myelin, a combination worth reading across Sections 2, 4, and 6 together. If Peripheral Neuropathy or Demyelination markers are elevated alongside Anti-GRP78 or Anti-CMV, that points toward a systemic neurological immune process affecting both peripheral and central nervous tissue rather than an isolated local nerve problem, following the GRP78-sensation and CMV-GM2-Guillain-Barre mechanisms described in Section 3.

If Anti-GM1 or Anti-GM2 is elevated alongside elevated HbA1c or fasting insulin, that combination is a specific, documented reason to monitor blood sugar closely, given how common peripheral neuropathy is in people carrying these antibodies; Sections 1 and 5 cover this pairing. If your Neural Zoomer Plus comes back normal or negative but bacterial or fungal OAT markers and methylmalonic acid are elevated, your nerve symptoms are more likely metabolic and gut-driven than autoimmune, which points you toward Section 4 and the companion guide, *Unlock Your Gut*, rather than further immune workup. And if you have numbness or tingling with a history of tick exposure and a negative or inconclusive standard Lyme test, remember that standard two-tiered testing misses 60 to 70 percent of early Lyme disease; a dedicated Tickborne panel, covered in Section 6, is the appropriate next step, not a repeat of the same screen that already came back inconclusive.

When to Seek Clinical Review

This guide is self-serve education. It explains what your labs are showing you; it does not replace an individualized clinical evaluation, and it does not give dosing, sequencing, or treatment instructions. Bring your actual results to a Root Discovery Session or your appointment with me for interpretation specific to your case.

Seek clinical review without delay, rather than waiting on lab results, if you experience any of the following: a fall related to numbness or balance loss, rapidly progressing weakness in addition to numbness, symptoms spreading quickly over days rather than months, or any difficulty with breathing or swallowing alongside nerve symptoms. These are red-flag presentations where a conventional nerve conduction study or electromyography should be pursued alongside, not instead of, the functional workup in this guide.

Book a free Root Discovery Session to review your results and get clarity on which panel is the right next step for your specific case: rootcauseunlocked.com/discovery

Track What Changes, From Anywhere

Reading your labs once tells you where you are starting. What tells you whether anything is working is what happens to your symptoms over the following weeks, and that is worth recording rather than reconstructing from memory at your next appointment.

The Root Tracker is free, works on any device, and keeps your entries on your own device. Turn on recovery tracking if nerve symptoms are limiting what you can do day to day: rootcauseunlocked.com/tracker

Consultations and ongoing care are both available wherever you are located. Being outside North Carolina is not a barrier to working with me. Tell me where you are when you reach out and I will lay out exactly how we would work together from there.

Get in touch: rootcauseunlocked.com/contact

What This Guide Does Not Cover

This guide covers the lab panels most directly relevant to peripheral nerve symptoms. It is not a complete workup for every cause of nerve damage, nor a substitute for the exam and history that tell me which of these panels is worth your money in the first place.

Book a free Root Discovery Session to review your results: rootcauseunlocked.com/discovery

Root Health | Dr. Adam Seay, DC | Garner, NC | rootcauseunlocked.com/unlock-your-neuropathy

For educational purposes only. This guide does not constitute medical advice or establish a patient-provider relationship.

Read this part first

This guide is educational. It is not medical advice, it does not diagnose anything, and reading it does not make you my patient. Nothing here is a reason to change a medication on your own, and one section specifically warns against a change people make without telling their prescriber.

If you have rapidly progressing weakness, symptoms spreading over days rather than months, a fall related to numbness or balance, or any difficulty breathing or swallowing alongside nerve symptoms, seek care now rather than starting a diet.

What you are actually working toward

Most diet advice for neuropathy arrives as a list of anti-inflammatory foods, which is fine as far as it goes and tells you almost nothing about what to expect. You end up eating more salmon and blueberries, feeling vaguely virtuous, and six months later you still cannot tell whether any of it helped.

This guide is built the other way around. It starts with the one dietary change that has genuinely high-quality trial evidence behind it, tells you honestly where that evidence is strong and where it has been oversold, and then gives you a way to test the rest on yourself so you end up with answers instead of habits.

What the evidence actually shows, including the part usually left out

Blood sugar control is the intervention with real evidence, and the honest version has a catch worth knowing.

A Cochrane review examined enhanced glucose control for preventing and treating diabetic neuropathy.¹ In **type 1 diabetes**, pooling 1228 participants, tighter control significantly reduced the risk of developing clinical neuropathy, with an annualized risk difference of 1.84%. The review rates this as high-quality evidence.

In **type 2 diabetes**, pooling 6669 participants across four studies, the annualized risk difference was 0.58% and **it did not reach statistical significance**, at $P = 0.06$. Nerve conduction and vibration threshold measures did improve significantly in both groups.

This matters because most people with diabetic neuropathy have type 2, and the confident claim that tight glucose control prevents neuropathy is largely a type 1 finding being applied to a type 2 audience. The direction of effect in type 2 is favorable and the nerve measures do improve. It simply has not been demonstrated for clinical neuropathy at the level people assume.

And there is a real cost on the other side. The same review found that enhanced glucose control significantly increased severe adverse events including hypoglycemic episodes, in both types, and states plainly that this needs to be weighed when judging the risk and benefit.¹ Driving your numbers down aggressively is not a free action, and it is a decision to make with your prescriber rather than alone with a continuous glucose monitor and this guide.

What about the Mediterranean diet? A 2025 systematic review and meta-analysis of 14 studies found that adherence was associated with reduced risk of diabetic retinopathy and diabetic nephropathy. For **neuropathy specifically, the results were inconclusive due to study heterogeneity**, and the authors say so directly.² So a Mediterranean pattern is a defensible way to eat for several good reasons. Preventing neuropathy is not yet one of the proven ones, and anyone telling you otherwise is reading past the paper.

That is the honest state of the evidence. What follows is built on it rather than around it.

Part one: blood sugar, the biggest lever you have

Even with the caveat above, this is where the effort belongs, because it is the only dietary target with high-quality trial evidence behind it in any diabetic population, and because the mechanism is not in dispute.

The practical target is not a number, it is the size of the swing. Two people can have the same HbA1c, one running steadily and one spiking after every meal and crashing between. Post-meal spikes are the part a fasting number does not show you.

Four changes that flatten the curve, in order of how much they usually help:

Eat protein and fat before carbohydrate at the same meal. Same food, same portions, different order. This is the lowest-effort change in the guide and it requires you to buy nothing.

Anchor every meal with protein. Palm-sized portion, every time. This is the single most reliable way to reduce the size of a post-meal rise without counting anything.

Move for ten minutes after eating. A walk after your largest meal blunts the post-meal rise. It does not need to be exercise, it needs to be movement.

Cut liquid sugar entirely, first. Soda, juice, sweetened coffee drinks, and sports drinks hit faster and higher than any solid food. If you do one thing on this page, this is the one that changes the numbers fastest.

What not to do: do not start skipping meals or fasting aggressively to bring numbers down, particularly if you take insulin or a sulfonylurea. That is the fastest route to the severe hypoglycemia the Cochrane review flagged, and hypoglycemia has its own consequences.

Part two: alcohol, which is a cause and not a lifestyle nag

This section is short and it is not optional reading.

Alcohol causes peripheral neuropathy directly. A systematic review and meta-analysis of alcohol-related peripheral neuropathy examined prevalence, character, risk factors, and management across the literature.³ This is a recognized clinical entity with a well-described pattern: axonal degeneration with reduced density of both small and large nerve fibers, producing decreased sensation, heightened pain response, and weakness.⁴

Two mechanisms run in parallel. Alcohol is directly toxic to nerve tissue, and chronic use depletes thiamine, which nerves require.⁵

What this means for you. If you drink and you have unexplained neuropathy, alcohol is on the list of causes, not the list of aggravating factors. It is worth an honest accounting of actual intake with your clinician, because it changes what should be investigated and because it is one of the few causes where stopping changes the trajectory.

I am not going to give you a number of drinks that is safe with an existing neuropathy, because I do not have evidence for one.

Part three: gluten, where the connection holds up but is narrower than the internet suggests

Gluten neuropathy is a documented entity, not a wellness invention. It has its own literature, its own antibody profile, and its own HLA associations. A study of the electrophysiological progression of gluten neuropathy found that most patients presented with a length-dependent neuropathy, while a subset had sensory ganglionopathy, and that specific HLA types tracked with which pattern people got.⁶ Autonomic involvement is common too: in a separate study, two thirds of gluten neuropathy patients had abnormal sudomotor function in the hands or feet.⁷

But that is a specific diagnosis, not a general reason to avoid bread. The claim “gluten causes neuropathy” is different from “there is a form of neuropathy driven by gluten sensitivity in susceptible people.” The second is true. The first is not.

Get tested before you eliminate, not after. This is the single most common self-inflicted mistake in this area. Celiac testing depends on you currently eating gluten. Go gluten free for three months first and the testing becomes unreliable, and you have permanently lost the chance to get a clean answer without a deliberate gluten challenge, which is unpleasant.

So: if gluten is a serious question for you, ask for testing **while still eating gluten**, then eliminate. Not the other way around.

Part four: the foundation pattern

With the evidence-backed parts covered, this is the sensible background pattern. It is not a cure and I am not going to present it as one.

Build meals from:

Non-starchy vegetables, in volume and variety. Leafy greens, cruciferous vegetables, and anything with color.

Protein at every meal. Fish, poultry, eggs, meat, or legumes if they agree with you.

Fats from whole sources. Olive oil, avocado, nuts, and seeds. Fatty fish twice a week gives you EPA and DHA from food rather than a capsule.

Low-glycemic fruit. Berries especially, because they carry the least sugar for the most volume.

Herbs and spices generously. Turmeric, ginger, cinnamon, garlic, rosemary. Cheap, pleasant, and they cost you nothing to include.

Reduce:

Liquid sugar, covered above and worth repeating.

Ultra-processed food, which is the practical way to reduce refined carbohydrate, industrial oils, and additives all at once without auditing labels.

Alcohol, covered above.

A note on some common advice I am leaving out. You will see a rule that you should drink half your body weight in ounces of water daily. I cannot find evidence for that specific number, so it is not in this guide. Drink enough that you are not thirsty and your urine is pale. If you take a diuretic or have kidney or heart disease, fluid targets are a medical question, not a wellness one.

Part four and a half: absorption, which is where diet quietly fails

You can eat a nutrient and still not get it. For nerve symptoms this matters more than it does for most conditions, because the nutrients nerves depend on most are also the ones most easily blocked.

Vitamin B12 is the main one, and several very common medications interfere with it.

Proton pump inhibitors interfere with B12 absorption from food by slowing the release of gastric acid, and can lead to deficiency. H2 blockers work the same way, reducing the acid needed to separate B12 from food protein. Metformin might reduce B12 absorption and significantly lower serum concentrations.

None of that is a reason to stop any of those medications, and I want to be unambiguous about that. What it does justify is having B12 checked properly rather than assumed, and the lab guide in this kit explains why serum B12 alone can look fine while the cell is short.

Gut surgery changes the arithmetic entirely. After gastric bypass or sleeve gastrectomy the anatomy that absorbs B12 is altered, so dietary intake stops being a reliable route.⁸ If you have had bariatric surgery and you have nerve symptoms, this belongs near the top of the investigation, not the bottom.

One trap almost nobody knows about: zinc causes copper deficiency, and copper deficiency causes a neuropathy that looks exactly like B12 deficiency.

Copper deficiency polyneuropathy is described as virtually indistinguishable from the nerve damage caused by low B12.⁹ And excess zinc is a recognized driver of it, including after bariatric surgery and from sustained zinc supplementation.¹⁰ Zinc is in a great many immune-support products and people take it for months without thinking of it as a drug.

This is case-report level evidence rather than trial evidence, so treat it as a thing worth knowing rather than a common cause. But if you take zinc regularly and have unexplained neuropathy, it is a cheap question to ask and an easy one to miss.

What to do about all of this. None of it is solved by eating differently; testing is what solves it. That is the honest boundary of what a diet guide can do for you, not a dodge.

Part four and three quarters: eating for sleep, because sleep sets your pain threshold

This section is in a diet guide because the strongest lever food has on nerve pain in the short term may be indirect.

Sleep loss measurably lowers your pain threshold. In a study characterizing the brain mechanisms, acute sleep deprivation amplified pain reactivity in the primary somatosensory cortex while blunting activity in the regions that evaluate and contextualize pain. It expanded the temperature range at which a stimulus was classified as painful, specifically by lowering thresholds. And the researchers found that ordinary night-to-night changes in sleep within a single person predicted day-to-day changes in experienced pain.¹¹

That last part is the useful one. This is not just about severe insomnia. A normal bad night makes tomorrow hurt more.

Eating changes that protect sleep, in order of impact:

Stop drinking alcohol to fall asleep. It shortens the time to fall asleep and degrades the second half of the night. Given part two, this one is doing double duty.

Move caffeine earlier. Caffeine has a long half-life and afternoon coffee is still working at bedtime for many people. Try a hard stop at noon for two weeks before deciding it does not affect you.

Do not go to bed hungry or overfull. Both fragment sleep. A small protein-containing snack is better than either extreme if dinner was early.

Keep the last large meal away from bedtime. Reflux and blood sugar swings both worsen when you lie down soon after eating.

If nerve pain is what wakes you, fixing sleep hygiene will not fix that on its own, and the honest answer is that pain waking you is a treatment conversation, not a diet one.

Part five: the specific experiments worth running

The foundation pattern is background. These are the actual tests, and each one gives you a yes or a no.

Experiment one: liquid sugar, two weeks. Remove every sugar-sweetened drink including juice. Two weeks is enough because the effect on post-meal glucose is immediate. Watch afternoon energy and night-time symptom intensity.

Experiment two: meal sequencing, two weeks. Protein and vegetables first, starch last, same meals otherwise. If you use a glucose monitor, this one is directly measurable. If you do not, watch the two hours after your largest meal.

Experiment three: alcohol, four weeks off. Genuinely off, not reduced. Four weeks is a reasonable read on symptom intensity and sleep quality, both of which affect how much pain registers.

Experiment four: an elimination and reintroduction over six weeks, only if the first three did not settle it. Remove the common triggers for three to four weeks, then reintroduce one food group every two to three days while tracking symptoms. Gluten, dairy, soy, corn, eggs, and nightshades are the usual candidates.

The reason this guide uses elimination and reintroduction rather than a blood test is that it directly measures your own response, which is the thing you actually want to know. Blood panels marketed as food sensitivity tests are contested, and this guide does not rely on them. Do the celiac testing question first, per part three, because that one is a real diagnosis with real testing.

One at a time. Running three of these simultaneously produces a feeling, not a finding.

Part six: a week you can actually cook

Not a meal plan to follow forever. A worked example of what the pattern looks like when it stops being abstract.

Breakfasts, rotate any three

- Eggs with spinach and half an avocado
- Plain Greek yogurt, berries, walnuts, cinnamon
- Smoked salmon, sliced tomato, olive oil

Lunches

- Big salad with a tin of sardines or leftover chicken, olive oil and lemon
- Lentil soup with a side of greens
- Leftovers from the night before, which is what actually happens

Dinners

- Salmon, roasted broccoli, small sweet potato eaten last
- Chicken thighs, cauliflower, big green salad
- Beef and vegetable stir fry over cabbage instead of rice
- Bean and vegetable chili
- Whatever your household eats, with the protein and vegetables served first and the starch portion halved

Snacks if you need them

- A handful of nuts
- Apple with nut butter
- Hard boiled eggs

The pattern underneath. Protein anchors every meal, vegetables take the most space, starch appears last and in a smaller portion than habit suggests, and nothing arrives in liquid sugar form.

Running a fair diet experiment

Write down where you start. Before changing anything, record your symptoms now, noting the worst intensity, how often they happen, what they stop you doing, and how you slept. Diet changes are gradual and memory is unreliable about gradual change in both directions.

One change at a time. Yes, it is slower. It is also the only way to end up knowing something you can act on.

Give it the stated window. Two weeks for the blood sugar experiments, four for alcohol, six for a full elimination and reintroduction.

Decide in advance what counts as a result. “Maybe a bit better” is not a result. Sleeping through instead of waking twice is a result. Half as much burning by evening is a result.

Keep what worked, drop what did not. A restriction that produced nothing is a restriction you should stop, and this is the step almost everyone skips. People accumulate food rules the way they accumulate supplement bottles.

Track it: rootcauseunlocked.com/tracker

Turn on recovery tracking if nerve symptoms limit what you can do day to day. Free, works anywhere, and your entries stay on your device.

Realistic expectations

Here is what the evidence in this guide actually supports.

If you have type 1 diabetes, better glucose control has high-quality evidence for reducing your risk of developing clinical neuropathy, with an annualized risk difference of about 1.84%.¹ That benefit is genuine, and it is also a modest annual number that compounds over years rather than transforming a month.

If you have type 2 diabetes, better control improves nerve conduction and vibration threshold measures, while the reduction in clinical neuropathy did not reach statistical significance.¹ Worth doing. Not a guarantee, and not something to pursue so aggressively that you trade nerve symptoms for hypoglycemia.

If alcohol is part of your picture, stopping addresses a cause rather than a symptom, and that is a different category of change from anything else in this guide.³

If you turn out to have gluten neuropathy, dietary change is the treatment, which is why testing before eliminating matters so much.⁶

For everyone else, an anti-inflammatory pattern is a reasonable way to eat with modest and largely indirect support for nerve symptoms specifically. I would rather tell you that than sell you a food list as a cure.

Nothing in this guide reverses established nerve damage. What diet does is remove ongoing drivers so that whatever capacity for repair you have is not fighting uphill.

When to seek professional care

Immediately. Rapidly progressing weakness, symptoms spreading over days, a fall related to numbness or balance, or difficulty breathing or swallowing.

Before eliminating gluten. If celiac disease is a serious question, get tested while still eating gluten.

Before aggressive glucose lowering. If you take insulin or a sulfonylurea, talk to your prescriber about targets rather than setting your own.

At your next appointment. If you drink regularly and have unexplained neuropathy, or if you have been gluten free for months and never had celiac testing.

Working with Root Health

Book a free Root Discovery Session: rootcauseunlocked.com/discovery

Consultations and ongoing care are both available wherever you are located. Being outside North Carolina is not a barrier to working with me. Tell me where you are when you reach out and I will lay out exactly how we would work together from there.

The testing questions in part three and the glucose target conversation in part one also work just as well in your own clinician's office.

Get in touch: rootcauseunlocked.com/contact

Not ready for a conversation yet? Start tracking: rootcauseunlocked.com/tracker

References

1. Callaghan BC, Little AA, Feldman EL, Hughes RAC. Enhanced glucose control for preventing and treating diabetic neuropathy. *Cochrane Database Syst Rev.* 2012;(6):CD007543. PMID 22696371.
2. Mediterranean diet and diabetic microvascular complications: a systematic review and meta-analysis. *BMC Nutr.* 2025. PMID 40170125.
3. Julian T, Glasgow N, Syeed R, Zis P. Alcohol-related peripheral neuropathy: a systematic review and meta-analysis. *J Neurol.* 2019;266(12):2907-2919. PMID 30467601.
4. Peripheral systems: neuropathy. *Handb Clin Neurol.* PMID 25307593.
5. High-dose thiamine strategy in Wernicke-Korsakoff syndrome and related thiamine deficiency conditions associated with alcohol use disorder. *Indian J Psychiatry.* PMID 34194054.
6. Zis P, Sarrigiannis PG, Rao DG, Hadjivassiliou M. Gluten neuropathy: electrophysiological progression and HLA associations. *J Neurol.* 2021;268(1):199-205. PMID 32754830.
7. Sudomotor dysfunction in patients with gluten neuropathy. *Neural Sci.* 2022. PMID 34791566.
8. Vitamin B12 toxicity secondary to post-bariatric supplementation: a case report. 2026. PMID 42253685.
9. Copper Deficiency and Polyneuropathy: A Case Report. 2022. PMID 36158410.
10. A Perfect MDS Mimic: Zinc-Driven Copper Deficiency After Gastric Bypass. *Am J Case Rep.* 2026. PMID 42357854.
11. Krause AJ, Prather AA, Wager TD, Lindquist MA, Walker MP. The Pain of Sleep Loss: A Brain Characterization in Humans. *J Neurosci.* 2019;39(12):2291-2300. PMID 30692228.

Vitamin B12 medication interaction figures in part four and a half: National Institutes of Health, Office of Dietary Supplements, Vitamin B12 Fact Sheet for Health Professionals.

For educational purposes only. This guide does not constitute medical advice, diagnosis, or treatment, and reading it does not establish a patient-provider relationship.

Root Health | Dr. Adam Seay, DC | Garner, NC | rootcauseunlocked.com/unlock-your-neuropathy

Read this part first

This guide is educational. It is not medical advice, it does not diagnose anything, and reading it does not make you my patient. Nothing here is a reason to stop, start, or change a prescription on your own.

Two things make this guide different from most supplement lists written for nerve pain.

It tells you what did not work. Several products sold specifically for neuropathy have been tested properly and did not beat placebo. Those are in here by name, in their own section, with what the trial found. A list that only contains good news is not a list, it is an advertisement.

It puts the safety section before the shopping section. One of the most common ingredients in nerve-support formulas can cause the exact symptoms you are trying to treat. That is not a footnote, so it is near the front.

If you have rapidly progressing weakness, symptoms spreading over days rather than months, a fall related to numbness or balance, or any difficulty breathing or swallowing alongside nerve symptoms, stop reading and seek care now. Those presentations need evaluation, not a supplement plan.

Why nutrient evidence looks thinner than it is

A large outcome trial costs tens to hundreds of millions of dollars, and that money only exists when a patent can pay it back. You cannot patent magnesium, an herb, or a food. So the evidence hierarchy this guide grades against is not neutral: it systematically over-documents patentable drugs and under-documents nearly everything else. That is measurable, not rhetorical. A Cochrane methodology review found that manufacturer sponsorship of drug and device studies produces more favourable results and conclusions than independent funding, through a bias that standard quality checks do not catch (PMID 28207928).

Two honest consequences follow, and they pull in opposite directions.

First, “no evidence” has three different meanings, and this guide tries to always tell you which one applies. *Tested and failed:* a real trial ran and the answer was no. *Never properly tested:* nobody could afford the trial, so absence of evidence is not evidence of absence. *Tested small and won:* real but modest trials, usually measuring markers rather than outcomes, because markers are what a small budget can afford.

Second, the asymmetry earns nutrients humility, not automatic credit. When isolated nutrients have received large publicly funded trials, the results have been mixed, with some genuine wins, some nulls, and some harms. Small nutrient trials also carry their own publication bias toward positive results, just as drug trials tilt toward their sponsor. So this guide applies one rule in both directions. It names the funding and the evidence tier, and lets you see both.

What you are actually buying back

Most people arriving at a guide like this have already spent money. A bottle recommended in a forum, a nerve-support blend with a long ingredient list, a second bottle when the first one did nothing. The spending is rarely the real cost. The real cost is the months that went by while you waited to see whether it was working, with no way to tell.

What this guide is meant to buy back is that time. It tells you which handful of things have real trial evidence behind them, what size effect to expect, how long to give each one before you decide, and which ones to skip so you are not running six experiments at once and learning nothing from any of them.

Supplements are supportive, not the main event

The single most effective thing for the most common form of neuropathy is not in a bottle. Where nerve symptoms are driven by blood sugar, glycemic control does more than any supplement in this guide. With a B12 deficiency behind them, correcting the deficiency is the treatment and everything else is decoration. And when a medication, a toxin, or a chronic infection is the driver, the answer is upstream of the supplement aisle entirely.

Supplements earn their place as support around a cause that is being addressed. They are a poor substitute for finding the cause, which is what the lab guide in this kit is for.

How to read this guide

Everything is sorted into three tiers.

Foundation is the short list with the best evidence and the widest applicability. If you only do one thing, do something from here.

Targeted depends on what is actually driving your symptoms. These are not for everyone, and picking from here without knowing your driver is guessing.

Ask your practitioner first contains neither products nor links. It is a list of things worth raising with a clinician, with the specific questions to ask and what they will want to know before answering. Some things genuinely should not be bought off a page, and pretending otherwise would make the rest of this guide less trustworthy.

Every entry gives you what it is, what the evidence actually shows including where it is weak, a commonly studied amount, the form that was studied, and the cautions that matter.

Before you start: the medication conversation

This population is heavily medicated. Metformin, gabapentin, pregabalin, duloxetine, amitriptyline, SSRIs, tramadol, blood thinners, and proton pump inhibitors all show up routinely in people with nerve symptoms, and several of them interact with things in this guide.

Find your row before you buy anything.

If you	Avoid, cap, or watch	Do this instead
Take metformin or a proton pump inhibitor	Assuming your B12 is fine because it was checked once	Ask for B12 with methylmalonic acid , since serum B12 alone can look normal while the cell is short
Take an SSRI, SNRI, duloxetine, or tramadol	Serotonin-active products including 5-HTP, SAMe, St John's wort, and L-tryptophan	Ask before adding any of them, and learn the stop-and-call symptoms below
Take warfarin or another anticoagulant	High-dose vitamin E	Tell your prescriber about omega-3 rather than avoiding it. The bleeding warning is weaker than its reputation, but periodic INR checks are sensible
Are in or recently finished chemotherapy	Acetyl-L-carnitine	A randomized trial found it made chemotherapy-related neuropathy worse, with the effect still present two years later
Take insulin or another glucose-lowering drug	Unmonitored alpha-lipoic acid	ALA can lower blood glucose. Monitor more closely when starting, and know your hypoglycemia symptoms
Already take a B-complex or a multivitamin	Adding a second product containing B6	Add up B6 from all sources and keep the total under the ceiling described below
Have had gastric bypass, sleeve, or other gut surgery	Assuming an oral B12 will absorb	Ask about absorption testing and non-oral routes
Take levothyroxine	Taking minerals at the same time of day	Separate mineral supplements from thyroid medication by several hours
Are pregnant, nursing, or trying to conceive	Everything in this guide by default	Ask first. Almost nothing here was studied in pregnancy

Stop and call your prescriber if you develop agitation, confusion, a racing heart, shivering, muscle twitching or rigidity, heavy sweating, diarrhea, or fever after adding a serotonin-active supplement to an antidepressant. That combination is the top safety concern in this category.

The vitamin B6 problem, which most neuropathy formulas get wrong

This deserves its own section because it is the one place where a supplement marketed for nerve pain can cause nerve pain.

High vitamin B6 intake causes a sensory neuropathy. A 2023 systematic review in *Nutrients* reviewed twenty articles and concluded that higher B6 levels, which usually follow from taking supplements, can produce a predominantly and sometimes exclusively sensory neuropathy of the axonal type.¹ The symptoms are numbness, tingling, and burning in the hands and feet. If that sounds exactly like what you already have, that is the problem. It is difficult to tell B6 toxicity apart from the neuropathy you came in with.

The ceiling is 100 mg per day from all sources combined. That is the Tolerable Upper Intake Level for adults. Severe progressive sensory neuropathy with loss of coordination has been described with chronic intake of 1 to 6 grams daily over 12 to 40 months, and sensory neuropathy has been reported at doses below 500 mg per day, which is why the observed safe dose was halved to set the 100 mg limit.

Now the reassuring half, because the scary half alone is misleading. The same systematic review found that B6 supplementation, even as part of a nutritional multivitamin, has not been shown to be harmful at permitted daily doses in people who already have peripheral neuropathy.¹ A separate study of 261 patients with chronic idiopathic axonal polyneuropathy found that elevated plasma B6 levels were not related to neuropathy severity, nerve conduction results, or examination findings.² Nobody develops this from a standard multivitamin.

So the real risk is stacking, not any single product. A B-complex, plus a multivitamin, plus a nerve-support formula, each with a label that looks perfectly reasonable on its own, can quietly clear 100 mg between them. Nerve formulas in particular often carry high B6 precisely because it is marketed as a nerve vitamin.

What to do. Add up the B6 in everything you take, including anything labeled nerve support, energy, or stress formula. Keep the total under 100 mg per day. If you are already above it, that is worth a conversation, because B6 neuropathy generally improves after stopping when it is caught early.

And do not take B6 expecting it to treat your neuropathy. The same review found no firm evidence that low B6 causes peripheral neuropathy. It also found no data at all on B6 as a standalone treatment, only as part of combinations.¹

Tier 1: Foundation

Alpha-lipoic acid

What it is. An antioxidant made in the body that is soluble in both water and fat, which is unusual and lets it act in more tissue compartments than most antioxidants.

What the evidence actually shows. This is the best-studied supplement in the whole category, and the picture is genuinely good in the short term and genuinely unproven in the long term. Both halves matter.

In the SYDNEY 2 trial, 5 weeks of oral ALA at 600, 1200, or 1800 mg daily reduced total symptom score by 51%, 48%, and 52% respectively, compared with 32% on placebo, all statistically significant.³ The proportion of people achieving at least a 50% reduction in symptoms was 62%, 50%, and 56% across the three doses versus 26% on placebo. Stabbing pain, burning pain, and patients' own global assessment all improved.

The most useful finding in that trial is about dose. 600 mg performed as well as 1800 mg, and side effects rose with dose. Nausea, vomiting, and dizziness were dose-dependent.³ There is no reason to buy the megadose, and a good reason not to.

A systematic review pooling oral trials found a total symptom score reduction of 1.95 points versus placebo at 600 mg oral.⁴ A critical appraisal covering 1160 participants across the ALADIN, SYDNEY, ORPIL, SYDNEY 2, and ALADIN III trials concluded that 600 mg orally for up to 5 weeks offers benefit in symptoms and signs without significant side effects.⁵

Where it falls down, stated plainly. The NATHAN 1 trial ran 4 years and **missed its primary endpoint.**⁶ There was no significant difference between ALA and placebo on the primary composite measure. Some secondary measures did favor ALA, including neuropathy impairment scores, and fewer people on ALA progressed. But serious adverse event rates were higher on ALA. So the honest summary is that ALA has good evidence for reducing symptoms over weeks, and has not demonstrated that it changes the course of the disease over years.

Commonly studied amount: 600 mg daily, oral. **Form studied:** oral alpha-lipoic acid. Most trial evidence is on the racemic form, not R-lipoic acid, despite R-lipoic being marketed as superior. **Take it:** on an empty stomach, since food reduces absorption. **Cautions:** can lower blood glucose, which matters if you take insulin or another glucose-lowering medication. Nausea and dizziness rise with dose, which is another argument for 600 mg. Discuss with your prescriber if you have thyroid disease.

Vitamin B12, but only if you are actually low

What it is. A vitamin required to build and maintain myelin, the insulating sheath around nerve fibers.

What the evidence actually shows. B12 deficiency is one of the few genuinely reversible causes of peripheral neuropathy, which is exactly why it belongs at the foundation. But the value is in correcting a real deficiency, not in taking B12 because you have nerve symptoms.

For B vitamins as a treatment in people who are not deficient, a 2026 review and meta-analysis of 13 randomized trials covering 834 people with diabetic peripheral neuropathy found a mixed picture.⁷ Neuropathy screening questionnaire scores improved. Examination scores improved statistically but **did not reach the minimum clinically important difference**. Pain measured on standard numeric or visual analogue scales showed no significant effect at all. Sural nerve conduction velocity and amplitude improved. Peroneal and tibial measures were inconsistent, and tibial velocity actually favored the control group. The authors concluded that heterogeneity and risk of bias preclude firm recommendations for routine use.

So: correct a deficiency, yes. Expect B vitamins to treat neuropathy in someone who is replete, no.

Get tested properly. Serum B12 alone can sit inside the reference range while the cell is functionally short. Methylmalonic acid rises when B12 is inadequate at the tissue level, which is why it is ordered alongside. The lab guide in this kit covers what the numbers mean.

Who is at real risk: anyone on metformin or a proton pump inhibitor long term, anyone following a vegan or strict vegetarian diet, anyone with pernicious anemia, and anyone who has had gut surgery.

Commonly studied amount: varies enormously with the cause. Dietary shortfall, malabsorption, and pernicious anemia are treated very differently, and the last of those often needs a non-oral route. **Form:** methylcobalamin and hydroxocobalamin are the forms usually used in functional practice. **Cautions:** correcting B12 without also assessing folate can mask a folate problem. This is a test-first nutrient, not a guess-first one.

Vitamin D, if your level is low

What it is. A hormone more than a vitamin, with receptors throughout the nervous system.

What the evidence actually shows. A 2025 systematic review and meta-analysis of randomized trials found a significant short-term reduction in pain in painful diabetic neuropathy, with a standardized mean difference of 0.85, roughly equivalent to 11 points on common pain scales.⁸ That is a meaningful effect size.

The honest caveats are substantial and the authors state them. Only four trials totalling 320 people were pooled. Heterogeneity was moderate. **No study reported medium or long-term outcomes at all.** Study quality varied, with concerns about allocation concealment and selective outcome reporting. Their own conclusion is that larger and more rigorous trials with longer follow-up are needed before routine vitamin D testing or supplementation can be recommended in clinical practice for this indication.

So this is promising short-term evidence, not settled practice. Correcting a documented deficiency is defensible on general grounds regardless.

Commonly studied amount: varies by trial and by starting level, which is why this is a test-then-dose nutrient. **Form:** D3, cholecalciferol. **Cautions:** fat soluble and genuinely possible to overdo. Retest rather than escalating indefinitely.

Tier 2: Targeted, depending on what is driving it

Magnesium

What it is. A mineral involved in hundreds of enzymatic reactions, including nerve signal transmission and muscle relaxation.

What the evidence actually shows. Weaker than the tier 1 entries, and you should know that going in. The rationale is mechanistic and the clinical experience is favorable, particularly for the cramping, twitching, and restless legs that often accompany nerve symptoms, but there is not a body of randomized trial evidence in peripheral neuropathy comparable to what exists for ALA. It is included because it is inexpensive, low risk, and addresses a symptom cluster that genuinely bothers people, not because the trial evidence is strong.

Form: glycinate and malate are better tolerated than oxide, which is poorly absorbed and mostly acts as a laxative.

Cautions: reduce or avoid in significant kidney disease. Separate from thyroid medication and certain antibiotics by several hours. Loose stools are the usual sign of too much.

Omega-3 fatty acids

What it is. EPA and DHA, incorporated into cell membranes and involved in resolving inflammation.

What the evidence actually shows. Reasonable mechanistic grounding and general anti-inflammatory evidence, with limited neuropathy-specific randomized data. Treat this as supportive rather than targeted therapy, and be skeptical of anyone presenting it as a nerve treatment.

Form: triglyceride or re-esterified triglyceride form. **Cautions, and the usual warning here is overstated.** Fish oil is routinely described as a bleeding risk with warfarin. The actual evidence is more reassuring than the reputation: most research finds that 3 to 6 grams per day does not meaningfully affect anticoagulant status in people on warfarin, and a 2014 review concluded omega-3s do not increase clinically significant bleeding. FDA-approved omega-3 product labels say studies have not produced clinically significant bleeding episodes. Those same labels still recommend periodic INR monitoring, which is the sensible middle position. Tell your prescriber you are taking it and let them decide whether to check, rather than avoiding it out of a warning that the data does not support.

Tier 3: Ask your practitioner first

Nothing in this tier has a purchase link, on purpose.

These are things where the right answer depends on your history, your medication list, and often on a test result. Selling them from a page to someone I have never assessed would contradict everything else in this guide. What follows is what to ask and what your clinician will want to know.

Anything serotonin-active, if you take an antidepressant. 5-HTP, SAME, St John's wort, and L-tryptophan. Duloxetine and amitriptyline are both commonly prescribed for neuropathic pain, so this overlap is not hypothetical, it is the norm. *Ask:* "I take [medication]. Is there a serotonin syndrome risk if I add this, and what symptoms should make me stop and call you?" *They will want to know* every prescription, over-the-counter product, and supplement you take, including anything in a blend.

High-dose B vitamins of any kind. *Ask:* "Can we total up my B6 across everything I am taking, and is it under 100 mg?" *They will want to know* every label, not just product names. B6 hides in blends.

Alpha-lipoic acid, if you take glucose-lowering medication. *Ask:* "How should I monitor my blood sugar for the first few weeks after starting this?" *They will want to know* your current regimen and your recent readings or continuous monitor data.

Anything at all, if you take an anticoagulant. *Ask:* "Which of these affect bleeding risk, and do I need to stop anything before my procedure?" *They will want to know* your indication, your target range if applicable, and your procedure schedule.

Chelation or any product marketed for heavy metal detox. *Ask:* "Do my actual test results support this, and what are the risks of doing it without them?" *They will want to know* whether you have had genuine, unprovoked testing. Provoked or challenge urine metal testing is not a valid way to establish a body burden, and treatment decisions made on it are built on nothing.

If nobody is quarterbacking your case, that is the finding. The most common problem I see is six providers each managing one piece and nobody looking at the whole list, rather than the wrong supplement. If reading this tier made you realize you have no one to ask, that is worth a conversation.

Book a free Root Discovery Session: rootcauseunlocked.com/discovery

There is no obligation to continue, and if the answer is that your case belongs with someone else, I will tell you that.

What I deliberately left out, and why

This section exists because a supplement guide that only lists winners is not being honest with you.

Benfotiamine. This is the one that will surprise people, because benfotiamine is in a great many nerve-support formulas and is specifically marketed for neuropathy. The BOND study randomized people with type 2 diabetes and symptomatic polyneuropathy to benfotiamine 300 mg twice daily or placebo for **12 months**, and found **no significant effect** on corneal nerve fiber length, on any secondary morphometric measure, on skin biopsy parameters, on nerve conduction, on quantitative sensory testing, on autonomic function, on sudomotor function, or on quality of life.¹¹ Only the Neuropathy Symptom Score showed a trend that did not reach significance. The treatment did raise thiamine levels, so it was absorbed and doing something biochemically. It just did not translate into benefit. It was well tolerated, and that is the most positive thing that can honestly be said about it.

Acetyl-L-carnitine, and this one is the most important sentence in the guide if you have chemotherapy-related neuropathy. Acetyl-L-carnitine is widely sold for nerve support, including specifically to people going through chemotherapy. In SWOG S0715, a randomized, double-blind, multicenter trial of 409 women receiving taxane-based chemotherapy for breast cancer, acetyl-L-carnitine at 1000 mg three times daily for 24 weeks **made neuropathy worse, not better.**⁹ The trial's own words are that it "was unexpectedly found to increase CIPN." The difference was statistically significant, and it was still measurable at two years, long after the supplement had stopped. At 104 weeks, 39.5% of the carnitine group versus 34.4% of the placebo group reported a meaningful worsening from baseline.

There is no version of that result that supports selling acetyl-L-carnitine to this audience. If your neuropathy is chemotherapy-related, duloxetine remains the only pharmacological treatment recommended by ASCO,¹⁰ and that is a conversation with your oncology team.

High-dose vitamin B6 as a treatment. Covered above. There is no evidence for it as a standalone treatment, and a real risk of causing the symptoms you are treating.

Proprietary blends with undisclosed amounts. If a label says "nerve support complex 850 mg" without breaking out what is in it, you cannot total your B6, you cannot compare it to a studied dose, and you cannot tell your prescriber what you are taking. That is disqualifying on safety grounds alone.

Anything sold as reversing or curing neuropathy. No supplement has demonstrated this. ALA, the best-evidenced product in the category, missed its primary endpoint in the one trial designed to test disease modification over years.⁶

Products I could not find honest evidence for. Several ingredients common in nerve formulas have mechanistic rationale and no meaningful trial data in this condition. Their absence here is not an oversight.

Running a fair 12-week trial

The most common way a guide like this gets wasted is changing six things at once. Then something improves, or does not, and you have learned nothing you can act on.

One change at a time. Yes, it is slower. It is also the only way to end up knowing something.

Write down where you are starting. Before you take anything, record what your symptoms are like now, including how bad they are at their worst, how often they come, what they stop you doing, and how you slept. Memory is unreliable about gradual change in both directions, and this is what the free tracker is for.

Give it long enough, but not forever. Symptom-level effects in the ALA trials appeared over about five weeks.³ Nutrient correction takes longer, because you are rebuilding rather than modulating. Twelve weeks is a reasonable outer bound for deciding whether something earned its place.

Decide in advance what success looks like. “A bit better maybe” is not a result. Half as much burning at night, or sleeping through instead of waking twice, is a result.

If it did nothing, stop taking it. This is the step people skip. An unhelpful supplement that stays in the routine forever is how someone ends up on nine things, spending real money, with no idea which ones matter.

Track it: rootcauseunlocked.com/tracker

Turn on recovery tracking if nerve symptoms limit what you can do day to day. It is free, it works anywhere, and your entries stay on your own device.

Realistic expectations

Here is what the numbers in this guide actually mean for a person.

In the best short-term trial, 62% of people on 600 mg of ALA achieved at least a 50% reduction in symptom score, against 26% on placebo.³ That difference is genuine, and it is also not everyone. Roughly a quarter of people improved on placebo alone, which tells you something honest about how variable these symptoms are week to week.

Vitamin D reduced short-term pain by roughly 11 points on common pain scales in painful diabetic neuropathy, in four small trials, with no long-term data at all.⁸

B vitamins improved some measures and not others, and the improvement in examination scores did not reach the threshold considered clinically meaningful.⁷

None of these is a cure, none was tested for years with a positive result, and the one trial that did run for years missed its primary endpoint.⁶ What they add up to is a reasonable chance of meaningfully less burning and tingling, on top of addressing the cause, not instead of it.

If your neuropathy has a driver that can be found, finding it will do more than anything on this list.

When to seek professional care

Immediately, without waiting on anything here: rapidly progressing weakness, symptoms spreading over days rather than months, a fall related to numbness or balance loss, or any difficulty with breathing or swallowing alongside nerve symptoms.

Promptly: new or worsening symptoms after starting a supplement, any of the serotonin syndrome symptoms listed earlier, or nerve symptoms that began after starting a new medication.

At your next appointment: if you take metformin or a proton pump inhibitor and have never had B12 checked with methylmalonic acid, or if you are taking more than 100 mg of B6 daily across everything.

My Favorite Support Products

Everything above was written without reference to any product line, and the order of this guide reflects the strength of the evidence, not what is easy to buy.

Look for third-party testing first. NSF Certified for Sport and USP Verified both mean an independent lab confirmed the product contains what the label says and is free of specified contaminants. Supplements are regulated as food rather than as drugs, so label accuracy is not automatic. This matters more than brand.

Then check the four things this guide has given you. The amount and the form should both match what was studied, the full ingredient list should be disclosed rather than hidden in a proprietary blend, and the total B6 across everything you take should stay under 100 mg per day.

If you would like products already screened on those criteria, my dispensary is here:

us.fullscript.com/welcome/drseay

Disclosure: Root Health earns a portion of sales made through that link. Nothing in this guide was included, excluded, ranked, or dosed based on what it earns. Benfotiamine sells well and is in this guide's do-not-bother section because the twelve-month trial was negative. You can buy any of this anywhere, and if you already have a source you trust, use it.

Working with Root Health

Book a free Root Discovery Session: rootcauseunlocked.com/discovery

Consultations and ongoing care are both available wherever you are located. Being outside North Carolina is not a barrier to working with me. Tell me where you are when you reach out and I will lay out exactly how we would work together from there.

This guide's medication table and the tier 3 question list also work just as well in your own clinician's office as in mine.

Get in touch: rootcauseunlocked.com/contact

Not ready for a conversation yet? Start tracking: rootcauseunlocked.com/tracker

References

1. Kalyoncu M, et al. The Role of Vitamin B6 in Peripheral Neuropathy: A Systematic Review. *Nutrients*. 2023;15(13):2947. PMID 37447150.
2. Suh J, et al. Vitamin B6 levels do not correlate with severity of neuropathy in chronic idiopathic axonal polyneuropathy. *J Peripher Nerv Syst*. 2022;27(1):40-46. PMID 34931740.
3. Ziegler D, Ametov A, Barinov A, et al. Oral treatment with alpha-lipoic acid improves symptomatic diabetic polyneuropathy: the SYDNEY 2 trial. *Diabetes Care*. 2006;29(11):2365-2370. PMID 17065669.
4. Snedecor SJ, et al. Systematic review of treatments for diabetic peripheral neuropathy. *Diabet Med*. 2018;35(5):577-589. PMID 26822889.
5. Papanas N, Ziegler D. Critical appraisal of the use of alpha lipoic acid (thioctic acid) in the treatment of symptomatic diabetic polyneuropathy. *Ther Clin Risk Manag*. 2011;7:377-385. PMID 21941444.
6. Ziegler D, Low PA, Litchy WJ, et al. Efficacy and safety of antioxidant treatment with alpha-lipoic acid over 4 years in diabetic polyneuropathy: the NATHAN 1 trial. *Diabetes Care*. 2011;34(9):2054-2060. PMID 21775755.
7. Vitamin B Supplementation for Diabetic Peripheral Neuropathy: A Review and Meta-Analysis. *J Clin Med*. 2026. PMID 42452619.
8. Vitamin D for Painful Diabetic Neuropathy: A Systematic Review and Meta-Analysis of Randomised Controlled Trials. *Endocrinol Diabetes Metab*. 2025. PMID 41139799.
9. Hershman DL, Unger JM, Crew KD, et al. Two-Year Trends of Taxane-Induced Neuropathy in Women Enrolled in a Randomized Trial of Acetyl-L-Carnitine (SWOG S0715). *J Natl Cancer Inst*. 2018;110(6):669-676. PMID 29361042.
10. Pharmacological Treatment of Chemotherapy-Induced Neuropathy: A Systematic Review of Randomized Clinical Trials. *Pain Manag Nurs*. 2025. PMID 39952863.
11. Effects of benfotiamine treatment over 12 months on morphometric, neurophysiological and clinical measures in type 2 diabetes patients with symptomatic polyneuropathy: the BOND study. *BMJ Open Diabetes Res Care*. 2026. PMID 41571333.
12. Lundh A, Lexchin J, Mintzes B, et al. Industry sponsorship and research outcome. *Cochrane Database Syst Rev*. 2017;2(2):MR000033. PMID 28207928

Vitamin B6 Tolerable Upper Intake Level and toxicity figures: National Institutes of Health, Office of Dietary Supplements, Vitamin B6 Fact Sheet for Health Professionals.

For educational purposes only. This guide does not constitute medical advice, diagnosis, or treatment, and reading it does not establish a patient-provider relationship.

Root Health | Dr. Adam Seay, DC | Garner, NC | rootcauseunlocked.com/unlock-your-neuropathy