



ROOT CAUSE UNLOCKED

Unlock Your Mold *Toxicity*

*A Complete Guide to Understanding, Testing, and Healing from
Mycotoxin Exposure*

Dr. Adam Seay, DC

Seay Wellness / Root Health, Garner, NC · rootcauseunlocked.com · (919) 662-0520

Unlock Your Mold Toxicity

A Complete Guide to Understanding, Testing, and Healing from Mycotoxin Exposure

BEFORE YOU START

Contents

Table of Contents

INTRODUCTION

The Hidden Health Crisis

KEY 1

Understanding NRF2 Dysfunction in Mold Toxicity

KEY 2

Proper Lab Testing, Getting the Right Answers

KEY 3

Identifying Your Sources of Exposure

KEY 4

The Drainage-First Approach to Detoxification

KEY 5

Removing and Killing Mold in Your Environment

KEY 6

Strategic Supplementation for Recovery

KEY 7

Dietary Changes That Accelerate Healing

CONCLUSION

Your Next Steps with Root Health

References

Table of Contents

- Introduction: The Hidden Health Crisis
- Key 1: Understanding NRF2 Dysfunction in Mold Toxicity
- Key 2: Proper Lab Testing, Getting the Right Answers
- Key 3: Identifying Your Sources of Exposure
- Key 4: The Drainage-First Approach to Detoxification
- Key 5: Removing and Killing Mold in Your Environment
- Key 6: Strategic Supplementation for Recovery
- Key 7: Dietary Changes That Accelerate Healing
- Conclusion: Your Next Steps with Root Health

The Hidden Health Crisis

Mold toxicity is one of the most underdiagnosed and misunderstood health conditions seen in functional medicine today. Millions of people struggle with symptoms that stem directly from mycotoxin exposure (the toxic compounds produced by mold),¹ yet they are routinely told their labs are normal and their symptoms are caused by stress or anxiety. Meanwhile, their bodies are quietly fighting an invisible enemy that conventional medicine rarely looks for and almost never addresses.

If that story sounds familiar, you are in the right place.

At Root Health, we approach chronic illness differently. Our M4 Method (Movement, Manipulation, Medicinal Nutrition, and Metabolic Connection) is built on one foundational truth: symptoms are not the problem. They are signals pointing to a root cause. Mold toxicity demands this kind of systems-level thinking because mycotoxins do not damage just one body system. They disrupt detoxification, overwhelm the immune system, derange hormones, impair the nervous system, trigger histamine dysregulation, destroy the gut lining, and congest the very drainage pathways the body needs to heal.

That last point, drainage, is where most recovery attempts fail. You can take all the right supplements and eat a perfect diet, but if the body's extracellular matrix and lymphatic channels are congested, toxins have nowhere to go. They simply recirculate and redeposit, causing the same symptoms all over again. This is why drainage-first is the non-negotiable first step in everything we do at Root Health.

This guide walks you through seven keys to unlocking mold toxicity recovery. Each key builds on the one before it. Together they form a roadmap from confusion and suffering to clarity and restored health.

Understanding NRF2 Dysfunction in Mold Toxicity

To understand why mold toxicity is so debilitating, you first need to understand one critical molecule: NRF2.

NRF2 (Nuclear Factor Erythroid 2-Related Factor 2) is your body's master regulator of antioxidant defense and detoxification.² Think of it as the conductor of a large orchestra, coordinating hundreds of genes responsible for protecting your cells from oxidative damage and clearing harmful substances. When NRF2 is firing properly, your body produces powerful antioxidants like glutathione, catalase, and superoxide dismutase, and your liver processes toxins efficiently.

Here is where mold toxicity becomes particularly insidious: mycotoxins do not simply overwhelm your detoxification system. They actively suppress NRF2 activation. Several well-studied mycotoxins, including aflatoxins, ochratoxin A,³ and trichothecenes, directly inhibit NRF2 signaling through multiple mechanisms. The result is a vicious cycle: the toxins making you sick also dismantle the very system your body needs to remove them.

When NRF2 is suppressed, the downstream effects are far-reaching. Mitochondrial function deteriorates. Chronic inflammation accelerates. The immune system loses its regulatory balance. This explains the constellation of symptoms that mold toxicity patients typically present with: persistent fatigue, brain fog, joint pain, respiratory problems, and heightened sensitivity to chemicals and other environmental triggers.

Genetic variation also plays a role. Some individuals carry polymorphisms in NRF2 and related pathways that reduce their baseline capacity to activate this protective system. Those individuals may develop severe mold toxicity from exposures that others tolerate without difficulty, and they may need more intensive support to restore function.

The good news is that NRF2 suppression from mycotoxin exposure is reversible. Unlike a fixed genetic defect, this dysfunction can be corrected through targeted intervention. But NRF2 restoration is foundational, not optional. Without restoring the body's capacity to process toxins at the cellular level, detoxification attempts are like trying to drain a bathtub with a blocked pipe. Every strategy in this guide is oriented toward restoring that capacity.

Proper Lab Testing, Getting the Right Answers

The testing landscape for mold toxicity is full of misleading options and missed diagnoses. Understanding which tests actually provide useful clinical information can save you significant time, money, and frustration. Accurate testing does four things: it confirms mycotoxin exposure objectively, quantifies the burden, reveals the cascade of downstream damage mycotoxins have caused across multiple body systems, and provides the objective baseline needed to document treatment response.

Mold toxicity is not a single-organ problem. The testing approach at Root Health reflects this reality. We build a multi-panel picture that captures the mycotoxin load directly, maps the downstream effects on the gut, the histamine system, the nervous system, hormones, and micronutrient status, and identifies co-exposures and co-infections that alter the clinical picture. Below is the complete framework.

What Traditional Testing Misses

Standard allergy testing (skin prick tests and IgE antibody measurements for mold) provides virtually no useful information for mold toxicity. These tests measure an allergic immune response to mold proteins, not your body's burden of toxic mycotoxins or its capacity to clear them. It is entirely possible to have severe mold toxicity with completely normal allergy panels. This is one of the most common reasons patients spend years undiagnosed.

The Complete Mold Toxicity Lab Framework

Test & Platform	What It Measures	Why It Matters for Mold Toxicity
Vibrant Mycotoxins Profile (Vibrant Wellness)	29 mycotoxins by mass spectrometry: 5 aflatoxins, ochratoxin A, gliotoxin, zearalenone, fumonisins B1/B2/B3, trichothecenes (Roridin A/E/L2, Satratoxin G/H, T-2, Verrucarins A/J), citrinin, patulin, mycophenolic acid, chaetoglobosin A, sterigmatocystin, enniatin B1, dihydrocitrinone	The foundational test for mold toxicity. Uses parts-per-quadrillion sensitivity to detect both original mycotoxins and their metabolites, giving a complete picture of exposure load and detoxification capacity. Test before beginning any protocol to establish a true baseline.
Organic Acids Test (OAT) (Vibrant Wellness)	Fungal metabolites: tricarballic acid (Fusarium/fumonisin), tartaric acid (Candida/mold), arabinose, citric acid cycle markers, mitochondrial markers, neurotransmitter metabolites, B vitamin markers, oxalate burden	Tricarballic acid is a direct urinary biomarker of fumonisin-producing mold exposure. The OAT also captures Candida fermentation that frequently co-exists with mold illness, reveals mitochondrial dysfunction driven by mycotoxins, and flags oxalate accumulation, a common and painful consequence of both mold and Candida overgrowth that patients rarely connect to their exposure.
Gut Zoomer 3.0 (Vibrant Wellness)	300+ gut microbiome species by DNA sequencing; secretory IgA; calprotectin; lactoferrin; zonulin; intestinal permeability markers; Candida species; H. pylori; parasites	Mycotoxins damage the gut lining directly and devastate the microbiome. The Gut Zoomer surveys that picture: depleted Lactobacillus and Bifidobacterium, elevated pathobionts, barrier markers including zonulin, and inflammatory markers like calprotectin. Read zonulin as suggestive rather than confirmatory, since serum zonulin measured by commercial kit has been shown not to correlate with the physiologic lactulose-mannitol measure of permeability. ¹⁵ Low secretory IgA confirms that mucosal immune defense is compromised. This is also where Candida overgrowth, an extremely common mold co-condition, is directly identified.
Wheat Zoomer (Vibrant Wellness)	Wheat and gliadin antibodies; zonulin; intestinal permeability antibodies (anti-actin, anti-LPS); celiac genetics (HLA-DQ2/DQ8)	Mycotoxins and gluten both drive intestinal permeability through zonulin. ⁴ When both are present, the barrier damage is additive. The Wheat Zoomer quantifies how much permeability damage is present, identifies whether gluten is actively contributing to it, and screens for celiac genetics. For patients with both mold-driven and gluten-driven permeability, gluten elimination is structural: every

Test & Platform	What It Measures	Why It Matters for Mold Toxicity
		exposure reopens tight junctions the protocol is trying to close.
GI-MAP (Diagnostic Solutions)	Quantitative PCR for H. pylori, Candida albicans and other fungi, bacterial pathogens, parasites, zonulin, calprotectin, sIgA	Mold illness profoundly suppresses gut immune defenses, allowing co-infections to establish. H. pylori, parasites, and significant bacterial pathogen loads co-exist with mold toxicity far more commonly than in a healthy population and must be identified and addressed in protocol sequencing. The GI-MAP's quantitative PCR is highly sensitive for organisms that stool culture misses.
DAO / Histamine Panel (Precision Point Diagnostics)	Diamine oxidase (DAO) enzyme activity; histamine levels; DAO:histamine ratio; histamine metabolism markers	This is one of the most underappreciated labs in mold toxicity. Mycotoxins directly inhibit DAO, the primary enzyme responsible for breaking down histamine in the gut and bloodstream. ⁵ When DAO is impaired, patients develop histamine intolerance, reacting to foods, alcohol, and environmental triggers with symptoms virtually identical to mold toxicity itself: headaches, flushing, brain fog, heart pounding, nasal congestion, skin reactions, and anxiety. Many mold patients are simultaneously histamine intolerant without knowing it, and their symptoms persist or worsen on detoxification protocols because histamine-rich foods remain in their diet. DAO testing identifies this layer and changes the dietary prescription completely.
Intestinal Permeability / Leaky Gut Panel (Vibrant Wellness or Cyrex Array 2)	Actomyosin sIgA; occludin/zonulin antibodies; LPS (lipopolysaccharide) antibodies; bacterial endotoxin antibodies; anti-vinculin; anti-cytokeratin	This panel directly measures the immune response to gut barrier breakdown. Elevated LPS antibodies are consistent with bacterial endotoxin reaching systemic circulation, ⁴ which is a proposed driver of chronic inflammation. Worth knowing that these markers are not interchangeable and their evidence differs. LPS antibodies track bacterial translocation and separate from markers of intestinal damage, so they answer a different question rather than a confirming one. ¹⁶ IgA anti-actin is the best validated marker on this kind of panel, at 80% sensitivity and 85% specificity, though validated specifically for celiac disease with villous atrophy. ¹⁷ Anti-occludin and anti-zonulin antibodies are suggestive of tight junction disruption; I could not locate comparable human validation for the occludin antibody. When leaky gut testing is positive alongside mycotoxins, the protocol

Test & Platform	What It Measures	Why It Matters for Mold Toxicity
		must include gut barrier repair as a primary therapeutic target, not an afterthought.
Neural Zoomer Plus (Vibrant Wellness)	48 neurological antibodies: blood-brain barrier proteins (occludin, claudin-5, S100B); myelin markers; ganglioside antibodies; neurotransmitter receptor antibodies; GAD65; Purkinje cell antibodies	Mycotoxins have a particular affinity for nervous system tissue and are among the most potent neuroinflammatory compounds known. The Neural Zoomer Plus identifies whether mold exposure has triggered autoimmune neuroinflammation targeting the blood-brain barrier, myelin, or specific neuronal receptors. Brain fog, memory issues, anxiety, depression, and cognitive decline in mold patients are often being driven by active neuroimmune dysregulation that standard neurology workups never test for. This panel changes the urgency and direction of the neurological support in the protocol.
Hormone Zoomer (Vibrant Wellness)	Comprehensive urine hormone panel: estrogens (E1, E2, E3 and metabolites), progesterone metabolites, androgens (testosterone, DHEA, cortisol), estrogen metabolism pathways (2-OH, 4-OH, 16-OH), 5-alpha dominance ratio, adrenal rhythm	Mycotoxins are potent endocrine disruptors. Zearalenone, one of the most common mycotoxins found in food and water-damaged buildings, directly binds estrogen receptors and produces estrogen dominance patterns. ⁶ Ochratoxin A disrupts cortisol production and adrenal rhythm. The Hormone Zoomer quantifies how significantly mycotoxins have disrupted the endocrine system and identifies the specific patterns (estrogen dominance, androgen suppression, cortisol dysregulation) that require support. Patients with unexplained hormonal symptoms in the context of mold exposure should always run this panel.
Micronutrient Panel (Vibrant Wellness)	Intracellular levels of vitamins C, D, E, B1, B2, B3, B5, B6, B12, folate; magnesium; zinc; selenium; CoQ10; glutathione; antioxidant capacity; essential fatty acids	Mycotoxin detoxification is metabolically expensive. As the body attempts to clear mycotoxins, it consumes glutathione, B vitamins, vitamin C, zinc, selenium, and magnesium at an accelerated rate. Tricarballic acid from Fusarium mold binds magnesium tightly, creating magnesium deficiency that conventional serum magnesium will miss entirely. Supplementing without knowing what is depleted is guesswork. The intracellular panel shows what your cells actually contain, not just what is circulating in blood.

Test & Platform	What It Measures	Why It Matters for Mold Toxicity
Total Tox Burden (Environmental Toxins) (Vibrant Wellness)	38 environmental chemicals across 6 categories: organophosphates, organochlorines, heavy metals, plasticizers, parabens, BTEX compounds (benzene, toluene, xylene); plus mycotoxin overlap markers	Mold toxicity rarely exists in isolation. Patients with mycotoxin burden almost always carry a co-existing chemical toxin load from pesticide residues, plasticizers, solvents, and heavy metals that compete for the same detoxification pathways. The Total Tox Burden panel maps this complete chemical terrain and identifies whether heavy metal support, environmental chemical avoidance, or additional detoxification pathway support is required alongside the mold protocol.
Vibrant Tickborne Panel 2.0 (Vibrant Wellness)	Borrelia burgdorferi and co-infections (Bartonella, Babesia, Anaplasma, Ehrlichia, Rickettsia) by antibody and PCR; Western blot confirmation	Lyme disease and mold toxicity are the two most commonly co-occurring complex chronic illness presentations in functional medicine. Both suppress immune function, both drive neuroinflammation, and the symptoms overlap significantly. Mold exposure impairs the immune response to Lyme infections, making tickborne illness harder to clear in the presence of ongoing mycotoxin burden. Identifying this co-occurrence changes the sequencing of the protocol: drainage and mold clearance often must be prioritized before Lyme-specific treatment can be effective.
Shoemaker Biotoxin Panel (LabCorp or Quest)	MMP-9, C4a, C3a complement, MSH (Melanocyte Stimulating Hormone), VIP (Vasoactive Intestinal Peptide), TGF-beta-1, leptin, ADH/osmolality, VEGF	Dr. Ritchie Shoemaker's panel reveals the immune and neuroendocrine disruption that chronic biotoxin exposure produces. MMP-9 reflects tissue damage and vascular permeability. MSH reflects hypothalamic suppression. VIP reflects neuropeptide dysfunction driving sleep disruption and inflammation. C4a reflects chronic innate immune activation. Together these markers map the severity of systemic biotoxin injury and provide objective monitoring points for recovery. Many patients with longstanding illness and negative Mycotoxins Profile will still show abnormal Shoemaker markers, indicating ongoing biotoxin response even when urinary clearance appears normal.

Understanding What the Tests Show Together

The DAO Test: The Missing Piece Most Providers Never Order

Histamine intolerance is one of the most common and underrecognized layers of mold toxicity. Mycotoxins, particularly ochratoxin A and trichothecenes, directly suppress DAO enzyme production in the gut. DAO is

responsible for breaking down the histamine that comes from foods, from gut bacteria, and from your own immune cells. When DAO is impaired, histamine accumulates and produces a symptom picture that can look almost identical to mold toxicity itself: pounding headaches, facial flushing, racing heart, nasal congestion, anxiety, skin hives, digestive distress, and profound brain fog.

The clinical significance of this cannot be overstated. Many mold patients go through aggressive detoxification protocols and continue to react because nobody identified that they were simultaneously histamine intolerant. They eat fermented foods thinking they are supporting gut health. They drink bone broth for gut repair. They eat spinach and avocado because they are healthy anti-inflammatory foods. Every one of those is a high-histamine food that is driving symptoms while the mold protocol is trying to resolve them.

A low DAO activity result alongside elevated mycotoxins changes the dietary prescription completely and immediately. The low-histamine overlay must be added to the anti-mold diet, and DAO enzyme support is added to the supplement protocol, until both the mycotoxin burden is reduced and the gut lining has healed enough to restore DAO production. We cover this in detail in Key 7.

Leaky Gut Testing: Confirming the Systemic Cascade

Mycotoxins penetrate the gut epithelium and trigger both direct physical damage to tight junctions and immune-mediated attack on the gut barrier. The intestinal permeability panel and Wheat Zoomer together confirm whether this breach has occurred and how severe it is. When LPS antibodies are elevated, bacterial endotoxins are entering systemic circulation,⁴ and every meal that disrupts the gut barrier releases another wave of inflammatory endotoxin into the bloodstream. This is a major driver of the chronic inflammatory state that makes mold patients feel sick all the time, not just when they are in a moldy environment.

Identifying leaky gut objectively also changes the supplement protocol. L-glutamine, colostrum, and gut barrier repair agents move to higher priority when permeability markers confirm active breakdown. And it sharpens the dietary conversation. Positive barrier antibodies alongside gluten reactivity strengthen the case for a gluten trial. They do not by themselves establish a requirement, because the barrier markers are suggestive rather than confirmatory and their validation varies by marker. Where celiac disease is the question, test for it directly and do so while still eating gluten, because that is where dietary change stops being a trial and becomes the treatment.

The Neural Zoomer Plus: When Mold Has Reached the Brain

The brain fog, memory problems, word-finding difficulty, anxiety, and emotional dysregulation that mold patients experience are not just subjective complaints. The Neural Zoomer Plus provides the objective data showing whether mycotoxin exposure has triggered autoimmune antibody formation against the blood-brain barrier proteins, myelin sheaths, or specific neuronal structures. When these markers are elevated, the neurological symptoms are being driven by active neuroimmune dysregulation, not just toxin load. This changes the treatment prioritization and adds neural support products to the protocol.

Testing at Root Health

At Root Health, we do not run every panel on every patient. We use your clinical picture, FM Logic questionnaire findings, ZYTO bioenergetic scan, and QRA assessment to prioritize which labs will yield the most actionable information first. The Vibrant Mycotoxins Profile, DAO panel, and either Gut Zoomer or GI-MAP are almost always the starting point. Additional panels are added based on what the initial results show and which symptoms remain unexplained.

Important: Test before you treat.

Beginning a detoxification protocol before establishing baseline mycotoxin levels, DAO status, and gut barrier integrity removes your ability to document treatment response and may alter results. Work with your provider to time testing appropriately. Our Mycotoxin Lab Package at rootcauseunlocked.com is designed to give you the most clinically actionable starting point.

Identifying Your Sources of Exposure

This is where recovery efforts most often break down. Patients invest heavily in supplements and detoxification protocols while continuing to live or work in environments that perpetuate their toxic exposure. No protocol can provide lasting relief if ongoing exposure continues.

Where Mold Hides

The most dangerous mold exposures are rarely obvious. They occur inside wall cavities, beneath flooring, above ceiling tiles, inside HVAC ductwork, and in crawl spaces. The most toxic mold species often produce minimal visible growth or odor. Do not assume a clean-looking home or office is a safe one.

Water damage is the primary trigger. This does not require dramatic flooding. A slow roof leak, a weeping pipe fitting, or a window seal that failed years ago is enough to support extensive mold colonization behind walls. HVAC systems are particularly important to investigate because they can harbor mold and distribute mycotoxins throughout an entire building with every cycle.

Workplace and School Exposure

Sick building syndrome is often unrecognized mold toxicity affecting multiple occupants simultaneously. If several people in your workplace have developed similar symptoms, especially respiratory, neurological, or fatigue-related, mold contamination should be strongly suspected. Schools and healthcare facilities are especially prone to this problem and often slow to investigate or remediate.

Getting a Professional Assessment

Mold inspection quality varies dramatically. Look for inspectors who use multiple assessment methods, including moisture readings, visual inspection, air sampling, and dust sampling. Be cautious of inspectors who also offer remediation services, as this creates a conflict of interest.

Dust sampling reflects cumulative exposure over time⁷ and is more reliable than a single air sample. Keeping indoor relative humidity below 50% is one of the most effective long-term mold prevention strategies available.

The Drainage–First Approach to Detoxification

This is the key that separates effective mold toxicity recovery from failed attempts. Most people approach detoxification by jumping straight to binders, saunas, and antifungal supplements. The problem is that these interventions mobilize stored toxins, and if your drainage pathways are not open and functioning, those toxins have nowhere to go. They recirculate, redeposit in tissues, and often make symptoms significantly worse.

At Root Health, drainage-first is non-negotiable in every protocol we build.

Phase 1: Open the Drainage Pathways First

Before any detoxification strategy is introduced, the body's extracellular matrix (ECM) and lymphatic system must be adequately supported. The ECM is the fluid-filled connective tissue environment surrounding every cell. When it becomes congested, as it invariably does in chronic mold toxicity, cellular waste, toxins, and inflammatory byproducts become trapped.

Drainage Foundation, established before any detox or antifungal agents:

- A lymphatic drainage botanical, choose ONE strength based on chronicity:
- Acute strength: active inflammation, recent exposure
- Matrix strength: extracellular matrix congestion, xenobiotic accumulation (most common mold pattern)
- Chronic strength: longstanding illness, degeneration, deep cellular involvement
- Ionic trace minerals and electrolytes: supports extracellular fluid dynamics
- Methylated B12, liposomal or sublingual: methylation pathway support essential for toxin processing

These three are established and running for at least one week before Phase 2 begins.

Phase 2: Support the Body's Processing Capacity

Once drainage is established, the focus shifts to enhancing the body's innate ability to process and neutralize mycotoxins. This means supporting hepatic detoxification, boosting glutathione status, and addressing mitochondrial dysfunction.

- Liposomal glutathione: The master antioxidant that directly binds and neutralizes mycotoxins.⁸ Liposomal delivery ensures bioavailability that standard oral glutathione cannot match.
- N-acetyl cysteine: Primary glutathione precursor, supporting ongoing production as reserves are depleted during active detoxification.

- Liver support formula (milk thistle, dandelion root, artichoke, and the sulfur amino acids): Supports the Phase I and Phase II reactions responsible for transforming mycotoxins into excretable compounds.
- Active B complex: Methylated B vitamins as detoxification cofactors and for energy production.

Phase 3: Bind and Clear

Only after drainage is open and processing capacity is supported does it make sense to actively bind stored toxins. Introducing binders too early is one of the most common reasons patients feel dramatically worse when starting a mold protocol.

Binding agents are selected based on your specific mycotoxins identified on the Vibrant panel.⁹ Your provider will guide this choice.

Sauna therapy, when tolerated, accelerates mycotoxin elimination through sweat. Start short and low-temperature. Always replace electrolytes.

Lymphatic drainage (dry brushing, rebounding, manual massage, targeted movement) supports continued clearance through the ECM.

Removing and Killing Mold in Your Environment

No amount of clinical intervention can provide lasting recovery if you are returning each day to an environment that continues to expose you to mold and mycotoxins. Environmental remediation is a required component of healing.

Cosmetic Cleaning vs. True Remediation

Wiping visible mold from a surface may reduce what you can see without meaningfully reducing your exposure. The most significant mold growth is almost always in hidden areas: wall cavities, subfloor systems, inside ductwork. Surface cleaning that ignores these reservoirs provides false reassurance and no real protection. Moisture control is the foundation of all effective remediation. Without addressing the water source, mold returns regardless of how thoroughly visible growth was addressed.

Personal Protection During Removal

If you have mold toxicity, you are likely highly sensitized. Even brief contact during cleaning can trigger significant symptom flares. At minimum, wear an N95 or P100 respirator, eye protection, and disposable protective clothing before entering any area with known or suspected mold contamination.

Effective Approaches and Verification

For surface mold on non-porous materials, 3% hydrogen peroxide is generally more effective and safer than bleach. Bleach cannot penetrate porous materials where mold roots extend. For porous materials (drywall, carpet, insulation), the most effective approach is complete removal and replacement. HVAC systems require specialized professional cleaning. Post-remediation testing by an independent party confirms that spore levels have returned to acceptable ranges. Maintaining indoor humidity below 50% protects the work done.

Strategic Supplementation for Recovery

Supplement selection for mold toxicity recovery is where a lot of patients make expensive and sometimes counterproductive decisions. What is often missing is the clinical framework for sequencing them correctly. At Root Health, recommendations are built from single ingredients and standardized botanicals sequenced deliberately, so that drainage capacity is established before anything is mobilized. Nothing in the sequence is practitioner-exclusive, and every item can be obtained wherever you prefer to buy it.

The protocol below is organized by phase. Your specific protocol will be customized based on your Vibrant Mycotoxins results, DAO findings, Neural Zoomer results, ZYTO scan, and clinical presentation.

Foundation: Drainage (Phase 1, Begin Here)

Product	What It Does	Dose
Lymphatic drainage botanical (select one strength)	Extracellular matrix and lymphatic drainage foundation. The matrix strength is the most common starting point for mold patients (xenobiotic accumulation, deeper congestion). Chronic strength for longstanding degeneration. Acute strength 1 for acute. Never stacked, choose one by chronicity. Runs throughout entire protocol.	1 dropper 2x per day in warm water
Ionic trace minerals and electrolytes	Magnesium, potassium, and sulfate for extracellular fluid dynamics and cellular hydration. Alkalizes the terrain and is integral to every drainage protocol.	40 drops in 1 oz water daily; chronic: 2 full droppers 2x daily
Methylated B12	Sublingual or liposomal methylcobalamin for methylation pathways and drainage support. A non-negotiable part of the drainage foundation.	Per label, twice daily

Detoxification and Processing Support (Phase 2)

Product	What It Does	Dose
Liposomal glutathione	Master antioxidant and primary mycotoxin-binding molecule. ⁸ Liposomal delivery bypasses gut degradation. Depleted rapidly during mycotoxin exposure, must be actively replenished.	1 tsp sublingually daily (general); 1 tsp sublingually 3x daily (chronic)
N-acetyl cysteine	Primary glutathione precursor; replenishes stores consumed during active detoxification.	600 to 900 mg twice daily away from meals
Liver support formula	Milk thistle, dandelion root, artichoke, and sulfur amino acids for Phase I and II liver reactions. The liver bears primary responsibility for mycotoxin transformation.	Per label, once daily; up to three times daily in chronic cases
Active B complex	Methylated B vitamins as detoxification enzyme cofactors and for mitochondrial energy production. Relevant with MTHFR variants.	Per label, once daily; twice daily (chronic)
Magnesium BisGlycinate + L-Taurine	Tricarballic acid from Fusarium mold binds magnesium directly. Standard serum magnesium misses this depletion. Buffered BisGlycinate + L-Taurine form is preferred for absorption.	3 caps 2x daily (general); 6 caps 2x daily (chronic)

Immune, Adrenal, and Neural Support

Product	What It Does	Dose
Thymus glandular	Support for thymic immune regulation. Mycotoxins chronically suppress immune function, and glandular support is used to help restore immune responsiveness.	2 caps daily (general), 4 caps 2x daily (chronic); Thymus Code: 1 full dropper 2x daily in warm water
Vitamin D3 with K2	Cholecalciferol with MK-7 for immune modulation and anti-inflammatory support. Low vitamin D is nearly universal in chronic mold illness. Test, dose to the result, and retest.	3 to 5 sprays sublingually daily
Zinc	Zinc for immune support and antioxidant enzyme function. Depletion is common and worth confirming on a micronutrient panel.	25 to 30 mg elemental daily with food
Adaptogenic HPA support	Adaptogenic herbs such as ashwagandha and rhodiola, with adrenal glandular support where indicated. Chronic mold illness places real demand on the stress axis, and cortisol dysregulation affects detoxification capacity.	Per IF: 2 caps 2x daily; HPA Axis LF: 2 caps daily (general), 4 caps 2x daily (chronic)
ASuRA Brain	Neurological support for brain fog, cognitive function, and neuroinflammation identified on Neural Zoomer Plus. Used when neurological markers are elevated.	2 caps AM + 2 caps PM daily (general); 2 caps 3x daily (chronic)

Gut Restoration and Histamine Support

Product	What It Does	Dose
Saccharomyces boulardii Forte	Beneficial probiotic yeast that resists antibiotic disruption, supports gut mucosal immunity, and controls Candida, which commonly overgrows in the context of mold toxicity. DBVPG 6763 strain, 20 billion CFU/g.	2 caps 1 to 2x daily
a spore-based probiotic	Multi-strain probiotic for gut microbiome restoration. Mycotoxin exposure significantly disrupts Lactobacillus and Bifidobacterium populations.	1 to 2 caps 3x daily before meals (general); 3 caps 3x daily before meals (chronic)
Colostrum	Bovine colostrum supports gut-associated lymphoid tissue, rebuilding secretory IgA and mucosal immune defenses damaged by mycotoxin exposure.	1 to 3 caps daily between meals (general); 3 to 6 caps daily between meals (chronic)
L-glutamine with glycine and proline	L-glutamine is the primary enterocyte fuel for gut lining repair, and glycine and proline support structural collagen repair of the barrier. Essential when leaky gut testing is positive.	5 to 10 g in water 2x daily on empty stomach
DAO Enzyme Support (HistaGest-DAO)	When DAO testing confirms low diamine oxidase activity, supplemental DAO enzyme is added before meals to assist histamine breakdown while the gut lining heals. Used alongside low-histamine dietary protocol.	1 tablet before high-histamine meals

Key Reminders on Supplement Sequencing:

- Always establish Phase 1 drainage before Phase 2 or Phase 3. This is not optional.
- a multi-component binder, if included, must be taken 2 hours away from all other medications and supplements.
- Do not self-prescribe from this guide. Product selection and sequencing require clinical judgment based on your specific lab results, scan findings, and symptom presentation.
- Your protocol at Root Health will be built for your pattern, not a generic template.

Where to get these

Every item in this protocol is a single ingredient or a standardized botanical, available from any reputable retailer. Look for independent third-party testing (NSF, USP Verified, or equivalent) and match the form named here rather than the name on the front of the bottle.

If you would rather not evaluate that yourself, Root Health keeps a curated dispensary of professional-grade products, screened for those standards and organized by the phases above.

Root Health dispensary: us.fullscript.com/welcome/drseay

Disclosure: Root Health earns a portion of dispensary purchases. Nothing here was included, excluded, ranked, or dosed based on what it earns, and you can buy any of it anywhere.

Dietary Changes That Accelerate Healing

Diet is not a secondary consideration in mold toxicity recovery. What you eat directly influences the inflammatory burden your body carries, whether you are continuing to expose yourself to mycotoxins through food, whether histamine is accumulating faster than you can clear it, and whether the gut lining has any chance of healing. The mold toxicity recovery diet is not a single template. For most patients it is a layered approach that combines an anti-inflammatory foundation with mycotoxin-food avoidance and, when DAO testing confirms it, a low-histamine overlay.

Understanding which layer applies to you, and sequencing them correctly, is what separates dietary changes that feel vaguely helpful from changes that produce clear and measurable symptom improvement.

The Three Dietary Layers in Mold Toxicity

Layer 1: Anti-Inflammatory Foundation (all mold toxicity patients) - Remove processed foods, refined sugars, refined grains, industrial seed oils, and alcohol. - Build the plate around whole foods: clean protein, vegetables, healthy fats, limited low-glycemic carbohydrates.

Layer 2: Mycotoxin-Contaminated Food Avoidance (all mold toxicity patients) - Remove the highest-risk food sources of dietary mycotoxins that add to your total body burden. - See the complete list below.

Layer 3: Low-Histamine Overlay (patients with confirmed low DAO or histamine intolerance symptoms) - Additional restriction of high-histamine and histamine-releasing foods. - Add DAO enzyme support and histamine-clearing nutrients. - This layer is only fully implemented when DAO testing confirms deficiency, not every mold patient requires it.

Layer 1: Anti-Inflammatory Foundation

Mycotoxins drive chronic systemic inflammation, and dietary choices that amplify inflammation slow recovery. The foundation is straightforward: real, whole foods; high-quality protein; abundant non-starchy vegetables; healthy fats; minimal sugar; and no processed foods.

Remove These Foods Completely (Active Treatment Phase) - All added sugars: table sugar, honey, maple syrup, agave, high-fructose corn syrup, and all products containing them. Sugar feeds Candida (a nearly universal mold co-condition) and drives the inflammatory cascade mycotoxins are already sustaining. - Refined carbohydrates: white bread, white rice, pasta, crackers, pastries, and processed grain products. These are rapidly converted to glucose that fuels gut fungal overgrowth and spikes insulin, promoting the inflammatory state. - Alcohol in all forms. Alcohol is directly hepatotoxic and competes with mycotoxin detoxification for the same liver enzymes. It also liberates histamine and suppresses DAO activity, a double burden when histamine intolerance is already present. - Industrial seed oils (canola, soybean, sunflower, corn

oil). These are high in omega-6 fatty acids that amplify the arachidonic acid inflammatory pathway already activated by mycotoxins. - All artificial additives, preservatives, colorings, and flavor enhancers. Patients with mold toxicity are frequently chemically sensitive; synthetic additives trigger immune and histamine reactions that slow recovery.

Build the Plate Around These Foods - Non-starchy vegetables (the largest portion of every meal): leafy greens, broccoli, cauliflower, Brussels sprouts, asparagus, zucchini, cucumber, celery, artichokes, arugula, kale, chard. These are the anti-inflammatory backbone of the diet. Rich in fiber for beneficial bacteria, rich in sulforaphane and other NRF2-activating compounds, low in sugar, and supportive of Phase II liver detoxification. - Clean protein (adequate at every meal): pasture-raised poultry, grass-fed beef and lamb, wild-caught fish (especially cold-water fatty fish for omega-3s), pasture-raised eggs. Protein does not feed mold or yeast. Adequate protein is essential for glutathione synthesis, immune function, and gut barrier repair. Do not underestimate protein intake during recovery. - Healthy fats: avocado, olive oil, coconut oil, ghee from grass-fed butter. Healthy fats support cellular membrane repair damaged by mycotoxins, facilitate absorption of fat-soluble vitamins D and E, and provide anti-inflammatory fatty acids. Coconut oil's caprylic acid provides direct antifungal activity. - Low-glycemic carbohydrates in moderation: sweet potato, butternut squash, quinoa, millet, and buckwheat. These provide adequate carbohydrate without the rapid glucose spike that fuels Candida and drives insulin-mediated inflammation.

Layer 2: Mycotoxin-Contaminated Food Avoidance

This is the dietary layer that surprises most patients. Certain foods are common sources of mycotoxin exposure even in the absence of visible mold. If you are consuming these foods regularly during recovery, you are adding to the mycotoxin load your body is trying to clear. This is not permanent restriction for most patients; it is a therapeutic phase while the protocol reduces your total body burden.

High-Risk Mycotoxin Food Sources to Eliminate During Active Treatment - Corn and corn-derived products: corn is one of the most mycotoxin-contaminated crops globally, frequently carrying aflatoxins and fumonisins.¹⁰ This includes corn chips, popcorn, cornmeal, corn tortillas, cornstarch, and processed foods containing corn syrup or corn-derived ingredients. - Peanuts and peanut butter: peanuts are among the highest dietary sources of aflatoxins.¹⁰ Conventional peanut butter consistently tests positive for mycotoxins at concerning levels. This includes mixed nut butters containing peanuts. - Tree nuts with mycotoxin risk, pistachios, cashews, and almonds: pistachios in particular consistently test positive for aflatoxins and ochratoxin A. Walnuts and pecans carry lower but relevant risk. Macadamia nuts and coconut are generally lower risk. - Conventionally grown coffee: coffee beans are among the most commonly ochratoxin A-contaminated foods.¹¹ Many daily coffee drinkers carry ochratoxin A as their primary mycotoxin finding on lab testing, with coffee as the single source. During the active treatment phase, coffee elimination or switching to tested, certified mycotoxin-free coffee is clinically meaningful. - Dried fruits: the drying process concentrates sugars and creates conditions favorable to mycotoxin production. Raisins, dates, figs, apricots, and prunes should be eliminated during the treatment phase. - Conventionally grown grains, especially wheat, barley, rye, and oats: field molds contaminate grain crops and concentrate further during storage. Gluten-containing grains also drive zonulin-mediated gut permeability that compounds mycotoxin-driven barrier damage. -

Aged and fermented cheeses: the aging process involves mold cultures (*Penicillium* species in blue cheese, brie, and camembert) and generates mycotoxins in the final product. This applies particularly to blue cheese, aged cheddar, parmesan, brie, and camembert. - Mushrooms (during acute phase): wild and commercially grown mushrooms can carry mold contamination and mycotoxins depending on growing conditions. Medicinal mushroom supplements formulated in controlled conditions are generally acceptable; fresh or grocery mushrooms should be avoided during the acute phase. - Fruit juice and high-sugar fruits: grapes, mangoes, bananas, and dried fruit. High sugar content fuels *Candida* and drives glucose spikes even when the fruit itself is not mycotoxin-contaminated.

Lower-Risk Fruits Tolerated During Active Treatment - Lemons and limes: alkalizing, low sugar, support liver detox - Green apples in moderation - Berries in moderation: blueberries, raspberries, blackberries (organic when possible) - Pomegranate in moderation

Layer 3: Low-Histamine Overlay (When DAO Is Confirmed Low)

When DAO testing confirms low diamine oxidase activity, or when your symptom picture includes classic histamine intolerance features (headaches, flushing, racing heart, nasal congestion, anxiety, hives, or digestive distress after eating fermented or aged foods), the low-histamine layer must be added immediately.

The relationship between mold and histamine intolerance runs in both directions. Mycotoxins suppress DAO enzyme production, which reduces your capacity to clear histamine. At the same time, chronic gut inflammation from mycotoxin damage increases intestinal mast cell activation, which releases more histamine directly into the gut. The result is a histamine load that outpaces your clearing capacity, producing symptoms that are indistinguishable from mold reactions, and that continue until both the mycotoxin burden is reduced and the DAO system is restored.

High-Histamine Foods to Remove (With Confirmed Low DAO) - All fermented foods: kombucha, sauerkraut, kimchi, kefir, yogurt, fermented soy products (miso, tempeh, soy sauce, tamari), apple cider vinegar, wine vinegar, and fermented vegetables - Aged proteins: all aged cheeses, processed and cured meats (bacon, salami, pepperoni, hot dogs, smoked salmon), canned fish (tuna, sardines, anchovies), shellfish - Alcohol in all forms: alcohol is a potent histamine liberator and DAO inhibitor - Fermented beverages: wine (especially red wine), beer, kombucha, cider, fermented teas - Vinegar and vinegar-containing foods: pickles, relish, most mustards, ketchup, and commercial salad dressings - High-histamine vegetables: spinach, avocado, eggplant, tomatoes, and tomato products (sauces, paste, ketchup) - Strawberries, citrus fruits, kiwi, pineapple, bananas: histamine liberators even when not high-histamine themselves - Fermentation byproduct flavorings: yeast extract, nutritional yeast, bouillon cubes, and MSG - Chocolate and cocoa in all forms - Artificial additives: tartrazine (Yellow #5), benzoates, sulfites (also found in many wines and dried fruits)

Low-Histamine Foods That Are Safe and Encouraged - Fresh proteins: freshly cooked or frozen-from-fresh meats and fish are low in histamine. The histamine content of protein increases dramatically with time, cooking methods that involve browning or long cooking times (like slow cooker or stewing), and improper

storage. Buy fresh, cook same day, eat promptly, or freeze immediately. - Fresh vegetables (not from the high-histamine list): broccoli, cauliflower, Brussels sprouts, asparagus, zucchini, cucumber, sweet potato, butternut squash, beets, carrots, peas, leeks, garlic, and onion are all safe. - Most grains (if tolerated): rice, quinoa, millet, oats (certified gluten-free), and corn-free pasta. Avoid wheat if the Wheat Zoomer confirms gluten sensitivity. - Most fats and oils: coconut oil, olive oil, ghee, avocado oil, and most unprocessed nut/seed oils are low-histamine. - Herbs and spices (with exceptions): basil, oregano, thyme, rosemary, garlic, turmeric, and ginger are all low-histamine and anti-inflammatory. Avoid paprika, cayenne, chili powder, and cinnamon: these are histamine liberators in sensitive individuals. - Coconut milk and coconut-based products: good dairy alternatives for patients avoiding both dairy (histamine) and nut milks (mycotoxin risk). - Herbal teas: chamomile, peppermint, and rooibos are low-histamine and supportive. Nettle leaf tea actively inhibits histamine release and is a useful daily addition.

Supporting DAO Enzyme Recovery Through Diet and Nutrition

The gut lining produces DAO. When the gut lining heals, DAO production recovers. The dietary and supplement strategies that support gut barrier repair, glutamine, colostrum, zinc, vitamin B6, vitamin C, and copper, are also the strategies that restore DAO capacity over time. This is why the low-histamine layer is temporary for most patients, not permanent. As mycotoxin burden reduces, gut healing progresses, and DAO activity is restored, histamine-containing foods can be reintroduced carefully.

- Vitamin B6 (P5P): a required cofactor for DAO enzyme activity. B6 deficiency directly reduces DAO production. P5P (pyridoxal-5-phosphate), the active form, is preferred. The Micronutrient Panel will confirm whether B6 deficiency is present.
- Vitamin C: supports DAO activity and also acts as a direct histamine-degrading antioxidant. Camu camu and acerola cherry are food-form options with good bioavailability.
- Copper: a cofactor for DAO enzyme function. Copper deficiency is uncommon but impairs DAO activity when present.
- Quercetin: a natural mast cell stabilizer that reduces the amount of histamine released from gut mast cells. Found in onions, capers, and apples, and available as a supplement, often paired with zinc.

Foods That Actively Support Your Protocol

Beyond what to remove, these specific foods provide direct therapeutic benefit during mold toxicity recovery.

NRF2-Activating Foods

Because NRF2 suppression is the cellular foundation of mold toxicity (Key 1), foods that activate this pathway provide direct therapeutic benefit that no supplement fully replaces.

- Cruciferous vegetables (broccoli, cauliflower, Brussels sprouts, kale, arugula, bok choy): the richest dietary source of sulforaphane, the most potent food-derived NRF2 activator known. Sulforaphane induces Phase II liver detoxification enzymes and glutathione synthesis directly. Lightly steaming or

sauteing these vegetables improves sulforaphane bioavailability. Broccoli sprouts provide 10 to 100 times the sulforaphane concentration of mature broccoli¹² and are one of the most clinically meaningful foods in the mold recovery diet.

- Garlic (raw, freshly crushed): allicin, produced when garlic is crushed or chopped and allowed to rest for 10 minutes before cooking, activates NRF2 and provides direct antifungal activity against Candida. One to two cloves of raw garlic daily is a meaningful therapeutic food addition.
- Turmeric with black pepper: curcumin activates NRF2, reduces NF-kB inflammatory signaling, and has been shown to reduce ochratoxin A-mediated cell damage in research studies.¹³ Black pepper (piperine) increases curcumin absorption 20-fold.¹⁴ Use liberally in cooking or as a daily golden milk preparation.
- Green tea (if histamine is not a concern): EGCG in green tea activates NRF2 and has direct antifungal properties. If histamine intolerance is present, herbal teas are preferred initially.
- Beets: betalains in beets support Phase II liver detoxification and have NRF2-activating properties. Beet greens are particularly rich in these compounds.

Drainage and Lymphatic-Supporting Foods

- Celery and cucumber: high water content and natural electrolytes support lymphatic fluid movement and ECM hydration.
- Lemon water (first thing in the morning): stimulates bile production and liver detoxification enzymes. Add fresh lemon juice to warm water, not bottled lemon juice which may have sulfite preservatives.
- Ginger: supports lymphatic flow, reduces gut inflammation, and stimulates bile production. Fresh ginger tea or grated ginger in meals is a simple daily addition.
- Dandelion greens: a liver tonic food that supports bile flow and Phase I liver detoxification. Use raw in salads or brewed as dandelion tea.
- Burdock root: traditional lymphatic herb also used as a food in many cultures. Traditionally used for lymphatic support.

Gut-Healing Foods

- Bone broth (low-histamine preparation only): rich in glycine, proline, and collagen precursors that support gut lining repair. IMPORTANT for histamine-intolerant patients: bone broth must be prepared fresh, cooked for a short time (under 3 hours), and consumed immediately or frozen. Long-simmered or commercially prepared bone broth is extremely high in histamine and will worsen symptoms in DAO-deficient patients. If histamine intolerance is confirmed, use freshly prepared short-cook broth or substitute plain glycine powder instead.
- Cooked and cooled resistant starch (sweet potato, cooked-cooled rice): feeds Bifidobacterium species that produce butyrate, the primary fuel for colonocytes and a key gut lining repair nutrient. Cooling these starches after cooking increases their resistant starch content significantly.

- Flaxseeds (freshly ground): provide both soluble and insoluble fiber for bowel regularity essential for toxin elimination. Also provide lignans with mild estrogen-modulating activity relevant for mold-driven estrogen dominance.
- Artichokes: a prebiotic food that selectively feeds Bifidobacterium and Lactobacillus. Also supports gallbladder and liver function through cynarin content.

Hydration: More Important Than It Sounds

Many mycotoxins are eliminated through urine, and adequate hydration is a direct factor in clearance rate. The minimum target during active treatment is half your body weight in ounces of filtered water per day. Quality matters as much as quantity: filtered water free from chlorine, fluoride, and other contaminants provides the cleanest vehicle for toxin elimination. Chlorine in tap water impairs beneficial gut bacteria that are already depleted.

Adding electrolytes to water (through a mineral concentrate, a pinch of sea salt, or a squeeze of lemon) improves cellular uptake and lymphatic hydration compared to plain water alone. This is particularly important during sauna sessions or periods of active die-off when electrolyte losses are higher.

Meal Structure and Timing

Eat three structured meals per day: skipping meals causes blood sugar drops that drive carbohydrate cravings and elevate cortisol. Elevated cortisol suppresses the immune defenses needed for mycotoxin clearance and promotes gut permeability.

Front-load calories toward earlier in the day: circadian rhythm alignment improves liver detoxification efficiency. The liver's peak metabolic activity occurs between 1 and 3 am, a larger evening meal versus a substantial breakfast reverses this rhythm and reduces overnight detoxification capacity.

Consider time-restricted eating (12 to 16-hour fasting window) if well-tolerated: autophagy during the fasting period supports cellular cleanup and can accelerate toxin clearance. However, fasting is contraindicated when adrenal function is severely depleted, blood sugar dysregulation is present, or active die-off is producing significant symptoms. Confirm with your provider before implementing extended fasting windows.

Prepare food fresh when possible: leftovers accumulate histamine as proteins break down. For histamine-intolerant patients, same-day preparation is particularly important for meats and fish.

The Anti-Mold Diet Quick Start

Anti-Mold Diet, Remove - All added sugars and refined carbohydrates - Corn, peanuts, pistachios, cashews, and conventionally grown coffee - Alcohol and fermented beverages - Aged and moldy cheeses - Dried fruits, fruit juices, high-sugar fruits - Industrial seed oils (canola, soy, sunflower, corn) - Conventionally grown wheat, barley, rye - Processed foods with artificial additives

Anti-Mold Diet, Build the Plate Around - Non-starchy vegetables (half the plate), especially crucifers and NRF2-activating greens - Clean protein: pasture-raised poultry and eggs, grass-fed beef, wild-caught fish - Healthy fats: avocado, olive oil, coconut oil, ghee - Low-glycemic carbs in moderation: sweet potato, quinoa, millet, rice - Daily: fresh garlic, turmeric with black pepper, ginger, lemon water - Add when tolerated: broccoli sprouts (highest sulforaphane concentration)

If Low DAO Is Confirmed, Add These Restrictions - Avoid all fermented foods, aged proteins, alcohol, vinegar, spinach, avocado, tomatoes - Focus on freshly prepared proteins: buy fresh, cook same day, eat promptly or freeze - Add DAO enzyme support before meals - Add nettle leaf tea daily (natural histamine blocker) - Add vitamin B6 (P5P), vitamin C, and quercetin for DAO enzyme support - This layer is temporary: lifts as mycotoxin burden reduces and gut healing progresses

Diet Duration and Reintroduction

The strict anti-mold and anti-mycotoxin diet is maintained throughout the active treatment phase, typically 3 to 6 months for significant mold illness. Retesting with the Vibrant Mycotoxins Profile and OAT at the 3 to 4 month mark provides objective guidance for when reintroduction is appropriate, not just symptom improvement, which often precedes full mycotoxin clearance.

Reintroduction follows a deliberate sequence. Lower-risk foods are reintroduced first, one at a time over 3 to 4 days, monitoring for symptom return. Coffee is usually the last mycotoxin-risk food reintroduced, and only certified low-mycotoxin sources should be considered. Corn, peanuts, and conventionally stored grains remain a long-term consideration for patients with aflatoxin burden, as these represent ongoing dietary re-exposure risk.

Your Next Steps with Root Health

Recovery from mold toxicity is not a mystery when approached systematically. It is the predictable outcome of addressing the right things, in the right order, with appropriate support.

The seven keys in this guide give you the framework:

1. Understand that NRF2 dysfunction is the cellular foundation of why your body cannot clear these toxins on its own
2. Get the right labs: Vibrant Mycotoxins Profile, DAO panel, gut permeability markers, Neural Zoomer Plus, and supporting biomarkers, so every decision is based on data
3. Find and eliminate your sources of ongoing exposure, because no protocol overcomes active contamination
4. Open drainage pathways first, before any detoxification intervention is introduced
5. Remediate your environment properly, not cosmetically
6. Use a clinically guided, sequenced supplement protocol built around your actual pattern
7. Support your recovery through an anti-inflammatory, low-mycotoxin diet that addresses histamine intolerance when DAO testing confirms it and provides the raw materials your body needs to heal

What this guide cannot do is replace individualized clinical care. Mold toxicity is a complex, polysystemic condition that responds best when assessment and treatment are tailored to the individual. This is exactly what the M4 Method at Root Health is designed to do: Movement, Manipulation, Medicinal Nutrition, and Metabolic Connection, each playing a role in your recovery, all integrated based on what your body is actually showing us through clinical assessment, advanced lab testing, and bioenergetic scanning.

Ready to Take the Next Step?

If you are ready to move from this guide to a personalized recovery plan, here is how Root Health can help:

- **Mycotoxin Lab Package:** our curated Vibrant Wellness panel for mold toxicity, including the Mycotoxins Profile, DAO panel, and supporting biomarkers. Order at rootcauseunlocked.com.
- **Mold Toxicity Recovery Protocol:** a 3-month personalized protocol built on your lab results, clinical assessment, and clinical indicators. Includes product recommendations, dietary guidance (including low-histamine layer if indicated), drainage sequencing, and follow-up support. Available at rootcauseunlocked.com.
- **Root Health New Patient Evaluation:** for patients who want to start at the beginning with a comprehensive functional medicine workup. Schedule at rootcauseunlocked.com or call our Garner, NC office directly at 919-662-0520.

You have been searching for answers long enough. Let's find your root cause.

Dr. Adam Seay, DC | Root Health | rootcauseunlocked.com | Garner, NC

References

1. Hurraß J, Heinzow B, Walser-Reichenbach S, et al. AWMF mold guideline “Medical clinical diagnostics for indoor mold exposure,” update 2023. *Allergologie select*. 2024. <https://pubmed.ncbi.nlm.nih.gov/38756207/>
2. Kensler TW, Wakabayashi N, Biswal S. Cell survival responses to environmental stresses via the Keap1-Nrf2-ARE pathway. *Annual Review of Pharmacology and Toxicology*. 2007. <https://pubmed.ncbi.nlm.nih.gov/16968214/>
3. Limonciel A, Jennings P. A review of the evidence that ochratoxin A is an Nrf2 inhibitor: implications for nephrotoxicity and renal carcinogenicity. *Toxins*. 2014. <https://pubmed.ncbi.nlm.nih.gov/24448208/>
4. Fasano A. All disease begins in the (leaky) gut: role of zonulin-mediated gut permeability in the pathogenesis of some chronic inflammatory diseases. *F1000Research*. 2020. <https://pubmed.ncbi.nlm.nih.gov/32051759/>
5. Maintz L, Novak N. Histamine and histamine intolerance. *The American Journal of Clinical Nutrition*. 2007. <https://pubmed.ncbi.nlm.nih.gov/17490952/>
6. Rizvi S, Chhabra A, Sagurthi SR, et al. Zearalenone and its metabolites provoke contrasting modulatory responses on estrogen and androgen receptors: insights into the molecular basis of endocrine disruption in live cells. *Food and Chemical Toxicology*. 2025. <https://pubmed.ncbi.nlm.nih.gov/40484165/>
7. Vesper SJ, et al. The development and application of the Environmental Relative Moldiness Index (ERMI). *Critical Reviews in Microbiology*. 2025. <https://pubmed.ncbi.nlm.nih.gov/38651788/>
8. Deng J, Peng Z, Xia Z, et al. Five glutathione S-transferase isozymes played a crucial role in the detoxification of aflatoxin B1 in chicken liver. *Journal of Animal Science and Biotechnology*. 2025. <https://pubmed.ncbi.nlm.nih.gov/40197593/>
9. Kerkadi A, Barriault C, Tuchweber B, et al. Dietary cholestyramine reduces ochratoxin A-induced nephrotoxicity in the rat by decreasing plasma levels and enhancing fecal excretion of the toxin. *Journal of Toxicology and Environmental Health, Part A*. 1998. <https://pubmed.ncbi.nlm.nih.gov/9482354/>
10. Kuloba L, Wasik A, et al. Aflatoxins and fumonisins: assessment methods, biomarkers of exposure, modified forms, co-exposure, and impact on human health. *Molecules*. 2026. <https://pubmed.ncbi.nlm.nih.gov/42451647/>
11. Miller TJ, Adeuya A, Robin LP. U.S. FDA regulatory monitoring of ochratoxin A in human foods: 2008-2022. *Food Safety (Tokyo)*. 2026. <https://pubmed.ncbi.nlm.nih.gov/42371543/>
12. Fahey JW, Zhang Y, Talalay P. Broccoli sprouts: an exceptionally rich source of inducers of enzymes that protect against chemical carcinogens. *Proceedings of the National Academy of Sciences of the United States of America*. 1997. <https://pubmed.ncbi.nlm.nih.gov/9294217/>
13. Kövesi B, Worlanyo AP, Kulcsár S, et al. Curcumin mitigates ochratoxin A-induced oxidative stress and alters gene expression in broiler chicken liver and kidney. *Acta Veterinaria Hungarica*. 2024. <https://pubmed.ncbi.nlm.nih.gov/38536404/>
14. Shoba G, Joy D, Joseph T, Majeed M, Rajendran R, Srinivas PS. Influence of piperine on the pharmacokinetics of curcumin in animals and human volunteers. *Planta Medica*. 1998. <https://pubmed.ncbi.nlm.nih.gov/9619120/>
15. Serum Zonulin Measured by Commercial Kit Fails to Correlate With Physiologic Measures of Altered Gut Permeability in First Degree Relatives of Crohn’s Disease Patients. *Frontiers in Physiology*. 2021. PMID 33841181. *n=39 healthy first-degree relatives, one commercial kit. A caution, not a definitive refutation.*
16. Surrogate Markers of Intestinal Permeability, Bacterial Translocation and Gut-Vascular Barrier in Liver Disease. *Liver International*. 2025. PMID 40317887.

17. Carroccio A, et al. IgA anti-actin antibodies ELISA in coeliac disease: a multicentre study. *Digestive and Liver Disease*. 2007. PMID 17652043.