



ROOT CAUSE UNLOCKED

Unlock Your *Hashimoto's*

A Complete Functional Medicine Guide to Calming the Autoimmune Attack, Addressing the Root Triggers, and Reclaiming Your Thyroid Health

Dr. Adam Seay, DC

Seay Wellness / Root Health, Garner, NC · rootcauseunlocked.com · (919) 662-0520

Unlock Your Hashimoto's

*A Complete Functional Medicine Guide to Calming the Autoimmune Attack,
Addressing the Root Triggers, and Reclaiming Your Thyroid Health*

BEFORE YOU START

Contents

INTRODUCTION

The Medication Is Not Enough

KEY 1

What Hashimoto's Actually Is

KEY 2

The Complete Testing Framework

KEY 3

The Gut-Thyroid Connection

KEY 4

Lyme Disease and the Thyroid

KEY 5

Drainage First

KEY 6

The Anti-Inflammatory Diet for Hashimoto's

KEY 7

Strategic Supplementation for Hashimoto's

KEY 8

Your Next Steps with Root Health

References

The Medication Is Not Enough

There is a conversation that happens in functional medicine practices every week, repeated in thousands of variations but always carrying the same essential frustration. A patient sits down and explains that they have been on thyroid medication for two, five, sometimes ten years. Their TSH is controlled. Their endocrinologist is satisfied. But they are exhausted in a way that sleep does not touch. They cannot lose weight despite genuine effort. Their hair continues to thin. Their brain fog is worse than ever. Their mood is unpredictable. They feel nothing like themselves.

The medication is doing exactly what it was prescribed to do. It is replacing the thyroid hormone that the gland is no longer producing adequately. What it is not doing, and was never designed to do, is stop the immune system from attacking the thyroid. In Hashimoto's thyroiditis, the root problem is not the thyroid. The root problem is the immune system. Treating the thyroid in isolation while leaving the immune attack unaddressed addresses the hormone deficiency but not the process driving it.

Hashimoto's thyroiditis is the most common autoimmune disease in the United States and the primary cause of hypothyroidism in developed countries.¹ Despite its prevalence, the majority of patients who carry the diagnosis have never been told what drives the autoimmune attack, what maintains it, or what specifically can be done to reduce it. They have been given a medication and told their labs are managed. The progressive destruction of their thyroid tissue continues.

This guide takes a different position. The autoimmune attack in Hashimoto's is not random and it is not irreversible in all patients. It has specific, identifiable, and addressable triggers. Gut permeability allows antigen traffic that activates and sustains the immune response. Gluten cross-reactivity produces antibodies that attack both intestinal tissue and thyroid tissue through molecular mimicry. A common foodborne pathogen called *Yersinia enterocolitica* produces an outer membrane protein associated with cross-reactive immunity against the TSH receptor. *Helicobacter pylori* gastric colonization drives both molecular mimicry and secondary gut permeability that amplifies the autoimmune signal. Gut dysbiosis creates an LPS-driven inflammatory environment that sustains Th17 polarization. Lyme disease drives the same immune polarization that characterizes Hashimoto's lymphocytic thyroid infiltration. Nutrient deficiencies impair the enzyme systems that govern thyroid hormone production and conversion.

Every one of these drivers is testable. Every one is addressable. The patients who see meaningful reductions in their antibody levels, who reduce or eliminate their medication needs, who recover their energy and cognitive function, are the patients whose treatment addresses all of these upstream factors rather than only the downstream hormone deficiency.

This guide walks you through eight keys to unlocking Hashimoto's recovery. Read it as a clinical framework, not as a supplement shopping list. The goal is understanding. Understanding is what makes every subsequent clinical decision more effective.

What Hashimoto's Actually Is

The distinction between hypothyroidism and Hashimoto's thyroiditis is not merely semantic. It is the clinical difference between a treatment plan focused only on hormone replacement and one that also addresses the immune process driving the destruction. Understanding what Hashimoto's actually is, at the mechanistic level, is the foundation on which everything else in this guide rests.

Hypothyroidism vs. Hashimoto's

Hypothyroidism is a state: insufficient thyroid hormone production. Hashimoto's thyroiditis is a cause: an autoimmune disease in which the immune system generates antibodies against thyroid tissue, progressively destroying the gland's functional capacity. Most cases of hypothyroidism in the United States are caused by Hashimoto's, but the reverse is not true. Not all Hashimoto's patients are hypothyroid, at least not at diagnosis. Early in the disease process, the inflammatory destruction can actually cause transient hyperthyroid episodes as stored thyroid hormone leaks from damaged tissue into circulation. Over time, as destruction progresses, the gland loses functional mass and hypothyroidism becomes established.

This distinction matters because it determines what success looks like. For a physician focused on hypothyroidism, a controlled TSH is success. For a clinician addressing Hashimoto's, the target is reducing the autoimmune attack itself, measured by declining TPO and thyroglobulin antibodies over time. A patient whose TSH is controlled but whose antibodies remain at 800 is not a successfully treated Hashimoto's patient. The disease is progressing regardless of the hormone replacement.

The Autoimmune Mechanism

The immune system maintains a critical distinction between self and non-self. Tolerance to self-antigens is established during immune development through a process of eliminating or suppressing immune cells that react to the body's own proteins. When this tolerance breaks down, the immune system generates antibodies and cytotoxic cells that attack the body's own tissues. This is the definition of autoimmunity.

In Hashimoto's, the primary antibody targets are thyroid peroxidase (TPO), the enzyme essential for thyroid hormone synthesis, and thyroglobulin, the protein matrix in which thyroid hormones are stored. When the immune system produces TPO antibodies (TPOAb) and thyroglobulin antibodies (TgAb), it recruits lymphocytes into the thyroid gland in a process called lymphocytic infiltration. These lymphocytes release cytokines that drive chronic inflammation within the gland. Over years, the inflammatory destruction reduces the number of functioning follicular cells, and thyroid hormone output declines.

Crucially, this process can be active for years or decades before TSH rises above the conventional laboratory cutoff for hypothyroidism. Patients with significant antibody elevations and progressive thyroid destruction

may have entirely normal TSH results during this period. This is why antibody testing is not optional in a thyroid workup and why a normal TSH does not rule out active Hashimoto's disease.

The Thyroid-Immune Bidirectional Loop

One of the less-discussed aspects of Hashimoto's is that the destruction of thyroid tissue itself perpetuates the autoimmune response. As thyroid cells are damaged and lysed by lymphocytic attack, their contents, including TPO and thyroglobulin, are released into circulation. This ongoing antigen exposure continuously stimulates the immune system to produce more antibodies. The process is self-reinforcing. This is the mechanism that explains why, without addressing the upstream triggers that initiated the immune response, the autoimmune attack tends to progress over time rather than spontaneously resolving.

Addressing Hashimoto's clinically means two things simultaneously: reducing the upstream drivers that are activating and sustaining the immune response, and supporting the nutritional and metabolic conditions that allow the immune system to recalibrate toward tolerance. Neither intervention alone is sufficient. Both together are what produce measurable, sustained antibody reduction.

The Complete Testing Framework

The standard thyroid test, a TSH drawn at an annual physical, is the beginning of a workup for Hashimoto's, not the end. Understanding what each marker reveals, and which markers are routinely omitted that should not be, is essential for any patient who wants to understand the full picture of their autoimmune thyroid status.

The Core Thyroid Panel

TSH, or thyroid-stimulating hormone, is produced by the pituitary gland and reflects the brain's assessment of circulating thyroid hormone levels. It is a valuable screening marker, but its limitations are significant in the context of Hashimoto's. TSH can remain in the conventional normal range (0.4 to 4.5 mIU/L) for years while active autoimmune destruction is progressing. The functional medicine optimal range, 0.5 to 2.0 mIU/L, is considerably tighter and more clinically meaningful for patients with active Hashimoto's.

Free T4 is the primary hormone produced by the thyroid gland. It is inactive until converted to the active T3 form. Free T3 is the active thyroid hormone that binds to receptors on virtually every cell in the body. Low Free T3, even with normal TSH and normal Free T4, produces the full clinical picture of hypothyroidism at the cellular level and is among the most common explanations for persistent symptoms in medicated Hashimoto's patients. The conversion of T4 to T3 depends on selenium, zinc, adequate liver function, and low inflammatory burden, all of which are commonly compromised in Hashimoto's.

Reverse T3 is an inactive metabolite of T4 that competes with active T3 for receptor binding. When the body is under chronic inflammatory stress, as it is in active Hashimoto's, T4 is preferentially converted to Reverse T3 rather than active T3. Elevated Reverse T3 produces cellular hypothyroidism even when circulating hormone levels appear adequate. The Free T3 to Reverse T3 ratio is one of the most useful single numbers in a Hashimoto's workup.

TPO antibodies and thyroglobulin antibodies are the autoimmune markers that confirm the Hashimoto's diagnosis and quantify the degree of immune activity. Any elevation above the laboratory lower limit of detection is clinically meaningful. The goal of treatment is not to bring antibodies below the conventional cutoff but to drive them toward undetectable. An antibody level of 350 that drops to 90 over six months of treatment represents significant clinical progress even if both numbers are above the conventional normal threshold.

The Extended Hashimoto's Workup

Beyond the core thyroid panel, a complete Hashimoto's workup at Root Health includes several additional assessments that address the upstream drivers of the autoimmune process.

Test	What It Reveals	Clinical Significance
Vibrant Hormone Zoomer	Full thyroid panel (TSH, Free T3, Free T4, Reverse T3, TPOAb, TgAb), adrenal cortisol rhythm, comprehensive sex hormones	Single draw captures thyroid conversion, autoimmune activity, adrenal status. \$710 retail.
Vibrant Wheat Zoomer	Tight junction antibodies (occludin/zonulin), gluten and wheat peptide reactivity across 26 antigens	Identifies gut permeability and gluten cross-reactivity driving the autoimmune response. \$479 retail.
LabCorp H. pylori Stool Antigen or Urea Breath Test	Active H. pylori gastric colonization. Stool antigen or urea breath test are preferred over serology for detection of active infection.	Low-cost (\$50-80), high-yield first-line screen. Eradication studies show measurable TPO antibody reduction. Order before expensive panels.
LabCorp Yersinia Serology (IgG/IgA/IgM)	Antibodies to Yersinia enterocolitica, a foodborne pathogen associated with thyroid autoimmunity in cross-sectional research.	Consider in non-responders after initial gut and dietary intervention. Not on any standard thyroid or tick-borne panel. Order through the office.
Vibrant Tickborne 2.0 PCR	Borrelia burgdorferi (41 antigens), Babesia, Bartonella, Ehrlichia, Anaplasma	For patients in tick-endemic areas or with non-resolving antibodies after gut and dietary intervention. \$1,406 retail.
LabCorp EBV Panel (VCA IgG/IgM, EA IgG, EBNA IgG)	Distinguishes past resolved EBV infection, active reactivation (EA positive + VCA IgG positive + EBNA positive), and recent primary infection (VCA IgM positive)	EBV reactivation is a documented Hashimoto's trigger. Add when antibodies are not responding or when fatigue, lymph node history, or immune dysregulation are prominent. ~\$60-90 retail.
Vibrant Gut Zoomer Complete	Microbiome composition, pathogen detection, permeability markers, secretory IgA, short-chain fatty acids, Blastocystis hominis	Essential when gut symptoms are present, when antibodies are not responding, or to characterize dysbiosis before antimicrobial intervention.
Vibrant Micronutrient Panel	Selenium, zinc, iodine status, B vitamins, vitamin D, magnesium, iron	Nutrient deficiencies impair T4-to-T3 conversion and directly affect antibody production. \$850 retail.

The order of investigation should be guided by the clinical picture. For all confirmed Hashimoto's patients, the Hormone Zoomer and H. pylori stool antigen are the standard first-line tests, providing the highest yield at the lowest cost. Yersinia serology is added for patients who are not responding to initial gut and dietary

intervention. Wheat Zoomer is added when gluten reactivity is suspected. Tickborne 2.0 PCR is added for patients with tick exposure history or antibodies not responding after gut and infectious workup.

The Gut–Thyroid Connection

The relationship between gut health and thyroid autoimmunity is not metaphorical. It is mechanistic and specific. It is also addressable. Understanding the gut-thyroid pathways in Hashimoto's changes the entire treatment framework from one focused only on the thyroid to one that treats the upstream drivers where the autoimmune process actually begins.

Pathway One: Intestinal Permeability

The gut barrier, when functioning correctly, allows properly digested nutrients to enter circulation while blocking undigested proteins, bacterial endotoxins, and pathogenic organisms. When tight junction proteins are disrupted, this barrier becomes permeable, and substances that should remain in the gut lumen gain access to systemic circulation. The immune system encounters these foreign antigens and mounts a response.

In Hashimoto's, the specific consequence of this increased antigen exposure is activation of the autoimmune cascade through a process called bystander activation, in which immune cells activated against gut-derived antigens generate cross-reactive responses against thyroid tissue.² The persistent antigen traffic through a leaky gut maintains a chronic state of immune activation that keeps thyroid antibody production elevated even when other triggers are addressed.

This is why patients whose Hashimoto's does not respond to dietary changes alone often make progress when gut barrier repair is added to their protocol. The gluten-free diet removes the primary gliadin-zonulin trigger. The gut barrier repair restores the physical barrier that prevents the ongoing antigen traffic. Both are required for the autoimmune activation to be fully withdrawn.

Pathway Two: Gluten and Molecular Mimicry

Gliadin, the immunogenic fraction of gluten, shares structural epitopes with thyroid tissue. In patients with HLA-DQ2 or HLA-DQ8 genetic variants, the immune response to gliadin generates antibodies that cross-react with thyroid antigens through molecular mimicry. This is a well-documented immunological mechanism and stands among the most evidence-supported dietary interventions in Hashimoto's management.³

Multiple clinical studies have demonstrated measurable reductions in TPO antibody levels in Hashimoto's patients following strict gluten elimination. The key word is strict. The cross-reactive antibody production does not diminish with reduction. It requires complete elimination. Patients who believe they are gluten-free but are consuming hidden gluten through shared cooking surfaces, processed foods containing wheat starch, or soy sauce maintain the molecular mimicry response at a subclinical level.

The Wheat Zoomer panel is particularly valuable in this context because it tests reactivity to 26 different gluten and wheat peptides, many of which are missed by standard celiac panels. Non-celiac gluten sensitivity, in which the immune system reacts to specific gliadin peptides without producing the villous atrophy of celiac disease, is identified on the Wheat Zoomer in patients who test negative on standard celiac antibody screens. A negative celiac panel does not rule out clinically meaningful gluten reactivity.

Pathway Three: *Yersinia Enterocolitica*

Yersinia enterocolitica is a gram-negative foodborne bacterium transmitted through contaminated pork, unpasteurized dairy, and undercooked meats. It is not a tick-borne organism. It is not covered on any standard thyroid panel. Its association with Hashimoto's thyroiditis is one of the more studied infectious connections in thyroid autoimmunity research, with multiple cross-sectional studies documenting significantly elevated *Yersinia* antibodies in Hashimoto's patient populations compared to healthy controls.

The proposed mechanism involves a low molecular weight envelope protein (LP) produced by *Yersinia* that shares structural homology with the human TSH receptor.⁴ The hypothesis is that antibodies generated against this protein during *Yersinia* infection may cross-react with TSH receptor epitopes through molecular mimicry. This mechanism is biologically plausible and supported by the cross-sectional association data. A prospective cohort study (Efthymiou et al., 2011) did not find *Yersinia* seropositivity predictive of de novo thyroid autoimmunity development, which means the causal relationship has not been confirmed and the association may reflect shared genetic susceptibility.⁵ The clinical position at Root Health is that *Yersinia* is worth ruling out in patients who are not responding to dietary and gut barrier intervention, because identified *Yersinia* burden is addressable and the test is inexpensive.

Testing is a standalone LabCorp order: *Yersinia enterocolitica* IgG, IgA, and IgM serology. Elevated IgG indicates past or chronic exposure. Elevated IgA suggests active mucosal infection. Elevated IgM indicates recent or acute infection. The infection can be subclinical, producing no gastrointestinal symptoms while maintaining an antigenic burden that keeps thyroid antibodies elevated.

Clinical Note: *Elevated TPO antibodies that do not normalize despite dietary intervention and gut repair are a strong clinical signal that an upstream infectious trigger is maintaining the immune activation. H. pylori and Yersinia are the two most cost-effective investigations at that point. Do not stop investigating at the gut barrier.*

Pathway Four: *H. pylori* and Gastric Immune Activation

Helicobacter pylori is a gram-negative bacterium that colonizes the gastric mucosa in approximately 44 percent of the global population, making it one of the most prevalent chronic bacterial infections in the world. Its connection to Hashimoto's thyroiditis is supported by multiple cross-sectional studies showing significantly elevated *H. pylori* seroprevalence in Hashimoto's patients compared to healthy controls, and by

eradication studies demonstrating measurable reductions in TPO antibody levels following successful treatment.

The mechanisms are two-pronged. First, the *H. pylori* CagA protein (cytotoxin-associated gene A) shares structural similarities with thyroid antigens, generating cross-reactive antibodies through molecular mimicry.⁶ Second, *H. pylori* colonization produces chronic gastric inflammation that progressively impairs stomach acid production. Adequate gastric acid is required for protein denaturation in the stomach. When acid production is insufficient, intact food proteins reach the small intestine without proper denaturation, increasing the antigen load that reaches the gut-associated lymphoid tissue and promoting the permeability-driven immune activation described in Pathway One.

The clinical significance of *H. pylori* in Hashimoto's management is substantial because eradication data shows actual outcome benefit. Unlike the *Yersinia* association, which has cross-sectional support but limited prospective outcome data, *H. pylori* eradication studies have shown direct antibody reduction in Hashimoto's patients following successful treatment. This makes *H. pylori* one of the highest-yield investigations in Hashimoto's patients who are not responding to dietary and gut barrier intervention. Testing is inexpensive: LabCorp stool antigen or urea breath test runs approximately \$50 to 80 and detects active infection with high sensitivity. Serology is less preferred because it cannot distinguish active infection from past exposure.

Pathway Five: Gut Dysbiosis and the LPS-Autoimmunity Loop

Gut dysbiosis, an imbalance in the microbial ecosystem characterized by reduced diversity, depletion of beneficial short-chain fatty acid producers, and relative overgrowth of pro-inflammatory gram-negative species, creates a systemic inflammatory environment that sustains Hashimoto's autoimmunity through a mechanism distinct from gut permeability itself. The two pathways interact, but understanding them separately improves the precision of clinical intervention.

The primary mechanism is LPS endotoxemia. Lipopolysaccharide, the outer membrane component of gram-negative bacteria, ranks among the most potent activators of the innate immune system via Toll-like receptor 4 signaling. In a healthy microbiome with adequate gut barrier function, LPS is contained within the intestinal lumen. When dysbiosis increases the gram-negative bacterial load and gut permeability allows LPS translocation into systemic circulation, TLR-4 activation drives a pro-inflammatory cytokine cascade that sustains the Th17-dominant immune environment characteristic of Hashimoto's.⁷ The autoimmune thyroid attack is, at its base, a Th17-mediated process. Persistent LPS signaling from a dysbiotic gut is a consistent upstream driver of that polarization.

Short-chain fatty acids, particularly butyrate produced by Firmicutes species including *Faecalibacterium prausnitzii* and *Akkermansia muciniphila*, are the primary energy source for colonocytes and are essential for maintaining tight junction integrity and regulatory T-cell function. Treg cells, which suppress autoimmune reactivity, are butyrate-dependent. Dysbiosis reduces SCFA production, which simultaneously impairs gut barrier function and reduces the Treg activity that would normally suppress the autoimmune response. This

makes microbiome restoration a distinct intervention from gut barrier repair, addressing a different layer of the same problem.

One specific organism within the dysbiosis picture warrants individual attention: *Blastocystis hominis*, a parasitic protozoan found at significantly elevated rates in Hashimoto's patients compared to healthy controls in multiple studies. *Blastocystis* disrupts tight junction proteins, elevates zonulin, and generates antigen traffic through the gut barrier that sustains autoimmune activation. It does not produce prominent gastrointestinal symptoms in most carriers, making it easy to miss without targeted testing. *Blastocystis* is detected on the Gut Zoomer Complete, so no additional panel is required. Its presence should be treated as a contributing driver of gut permeability rather than incidental colonization in the context of active Hashimoto's.

The Gut Zoomer Complete is the appropriate assessment for dysbiosis characterization in Hashimoto's patients. It provides microbiome composition, short-chain fatty acid status, pathogen detection including *H. pylori*, *Blastocystis*, and other parasitic organisms, permeability markers, and secretory IgA in a single panel. The combination of dysbiosis profiling with permeability assessment in a single draw makes it the most clinically efficient gut workup for the Hashimoto's patient who is not responding to initial dietary intervention.

Pathway Six: Epstein-Barr Virus Reactivation

Epstein-Barr Virus (EBV) is the most significant viral trigger in Hashimoto's thyroiditis and the most consistently underinvestigated in clinical practice. Over 90 percent of adults carry EBV as a latent infection following primary exposure, typically in adolescence. The critical distinction for Hashimoto's is not past exposure but reactivation: the shift from latent viral dormancy to active viral replication, which generates the immune activation that drives thyroid autoimmunity through multiple overlapping mechanisms.⁸

The molecular mimicry evidence for EBV is more experimentally grounded than for most bacterial triggers. The EBV nuclear antigen EBNA-1 shares structural epitopes with thyroid autoantigens, and cross-reactive antibodies against EBNA-1 and TPO have been demonstrated in patient populations. EBV can also directly infect thyroid follicular cells, establishing a reservoir of viral replication within the thyroid gland itself that generates ongoing immune activation at the site of autoimmune attack. This direct thyroid tropism distinguishes EBV from most other infectious triggers, which operate through systemic immune mechanisms rather than local thyroid infection.

EBV reactivation drives the same Th17 polarization pathway shared by Lyme disease and active Hashimoto's. EBV-driven immune activation suppresses regulatory T-cell function, promotes IL-17 production, and creates the immunological environment in which thyroid autoimmunity escalates. In patients with both EBV reactivation and elevated thyroid antibodies, the viral reactivation is maintaining the immune polarization that makes antibody reduction difficult regardless of dietary and gut interventions.

Testing distinguishes four clinically distinct states. VCA IgM positive indicates acute or recent primary infection. VCA IgG positive with EBNA IgG negative indicates recent primary infection with incomplete seroconversion. VCA IgG positive with EBNA IgG positive and early antigen (EA) IgG positive indicates

active reactivation: the state most clinically relevant to Hashimoto's. VCA IgG positive with EBNA IgG positive and EA negative indicates past resolved infection with no current reactivation. The full four-marker panel is available through LabCorp for approximately \$60 to 90. EBV testing is indicated for any Hashimoto's patient with prominent fatigue, lymph node involvement history, a history of mononucleosis, or antibodies that are not responding after gut, dietary, and bacterial infectious workup has been completed.

Addressing EBV reactivation clinically involves antiviral immune support rather than the antimicrobial protocols used for bacterial triggers. Lysine, monolaurin, zinc, vitamin C, and adaptogenic immune modulators are the primary clinical tools. The complete EBV protocol is individualized based on reactivation markers and overall immune status, and is managed within the Root Restore program.

Other Infectious Considerations

Several additional infectious agents have documented or proposed associations with Hashimoto's that are clinically relevant in specific patient contexts.

Hepatitis C Virus: well-established bidirectional relationship. HCV can trigger thyroid autoimmunity through direct thyroid infection and molecular mimicry. Interferon-alpha treatment for HCV is a known thyroid autoimmunity trigger. Relevant for any patient with HCV history or prior interferon treatment. Standard HCV antibody testing applies.

Post-COVID Thyroiditis: SARS-CoV-2 infects thyroid follicular cells via ACE2 receptors, confirmed in autopsy studies.⁹ Post-COVID Hashimoto's-pattern thyroiditis is increasingly documented. For any patient whose antibody escalation followed COVID infection, this context affects treatment framing. No specific additional test; antibody trajectory tracking applies.

Klebsiella pneumoniae and Mycoplasma pneumoniae: cross-sectional association data exist for both. Klebsiella shares thyroid antigen homology through molecular mimicry. Mycoplasma produces immune activation via cold agglutinin mechanisms. Consider in patients with respiratory illness history preceding Hashimoto's onset. LabCorp Mycoplasma IgG/IgM available if clinically indicated.

Parvovirus B19: associated with multiple autoimmune conditions including thyroid autoimmunity. IgG/IgM available through LabCorp. Consider in patients with joint symptoms alongside thyroid antibody elevation.

Lyme Disease and the Thyroid

The biological overlap between Lyme disease and Hashimoto's thyroiditis is clinically important for a specific subset of patients: those in tick-endemic areas whose antibodies are not responding to gut and dietary intervention. The immunological mechanisms are plausible and the shared Th17 pathway is well-characterized. This section presents that evidence accurately, including where it is strong and where it remains at the level of working hypothesis.

The Borrelia-Thyroid Shared Immune Environment

A working hypothesis in functional immunology proposes that *Borrelia burgdorferi* outer surface proteins, particularly OspA and OspC, may share structural similarity with human thyroid tissue antigens. If confirmed, antibodies generated against these proteins during *Borrelia* infection could cross-react with thyroid proteins through molecular mimicry. This hypothesis is currently based on in silico sequence homology analysis and has not been confirmed through experimental demonstration of cross-reactive antibodies in patients or through animal models of Lyme-induced thyroiditis. It should be understood as a plausible but unvalidated mechanism, not as established molecular biology.

What is well-established is that chronic Lyme infection and Hashimoto's share a common immune polarization. *Borrelia* drives a sustained Th17-dominant immune environment, characterized by elevated IL-17, IL-21, and IL-6, with suppressed regulatory T-cell activity. This is the same immune environment that drives lymphocytic thyroid infiltration in Hashimoto's. The two conditions share an immune polarization pattern, which means that active Lyme disease creates and maintains conditions that are broadly permissive to autoimmune thyroid activity, regardless of whether direct molecular mimicry is operating.

The practical clinical implication is that in patients with concurrent Lyme and Hashimoto's, reducing the Lyme burden reduces the Th17 inflammatory environment that sustains the thyroid autoimmunity. This is a mechanistically coherent therapeutic target. The limitation is that no study has directly demonstrated TPO antibody reduction specifically attributable to Lyme treatment, so the claim should be framed as a rational clinical inference from shared immunology rather than a proven treatment outcome.

Symptom Overlap and Missed Diagnoses

Lyme disease and Hashimoto's share a significant symptom overlap. Fatigue, brain fog, cold intolerance, weight changes, hair thinning, anxiety, cognitive impairment, mood instability, and joint pain are features of both conditions. In patients presenting with this symptom cluster, each condition can mask the other. Providers who identify the Hashimoto's stop investigating, missing the Lyme. Providers who identify the Lyme stop investigating, missing the Hashimoto's. Patients end up being treated for half of their clinical picture.

The gut represents a third layer of interaction. *Borrelia* infection suppresses secretory IgA production, the gut's primary immune defense against pathogenic colonization. This suppression allows dysbiotic organisms to establish in the gut, promoting intestinal permeability that contributes to autoimmune thyroid activation through the mechanisms described in Key 3. Gut repair in Lyme-Hashimoto's patients is not optional because the Lyme infection is actively degrading the gut barrier that needs to be restored for the thyroid autoimmunity to quiet.

Who Should Be Tested

Tick-borne co-infection testing is indicated for Hashimoto's patients who meet any of the following criteria: have lived in or spent significant time in tick-endemic areas, have a history of a tick bite regardless of whether a bull's-eye rash was observed, have antibodies that are not responding to dietary intervention, gut repair, and *H. pylori* and *Yersinia* workup, have significant neurological or psychiatric symptoms alongside thyroid symptoms, or have been previously treated for Lyme disease but continue to have symptoms.

The Vibrant Tickborne 2.0 PCR panel tests *Borrelia burgdorferi* across 41 antigens, significantly more than the standard Western Blot, and covers the co-infections that most standard panels miss: *Babesia*, *Bartonella*, *Ehrlichia*, and *Anaplasma*. At \$1,406 retail, it is available through rootcauseunlocked.com/order-labs. Given its cost, it is most appropriately positioned as a secondary investigation after the less expensive gut and infectious workup has been completed.

Drainage First

Every treatment intervention for Hashimoto's, whether dietary, supplemental, or targeted at an infectious trigger, mobilizes waste products, inflammatory mediators, and cellular debris that must be processed and eliminated through the body's drainage systems. The lymphatic system, the liver, and the kidneys bear the primary burden of this clearance. When drainage capacity is insufficient for the load being generated, mobilized toxins recirculate, inflammatory burden increases, and the autoimmune process can actually flare rather than improve.

This is a leading explanation for why patients who begin aggressive Hashimoto's protocols, particularly antimicrobial protocols for *H. pylori*, *Yersinia*, or Lyme, experience a significant worsening of symptoms in the first weeks. The antimicrobial intervention is working. But the drainage capacity cannot handle the microbial debris load. The result is a Herxheimer reaction that the patient interprets as the treatment making them worse.

At Root Health, the drainage-first sequence is applied to every Hashimoto's case regardless of severity or protocol complexity. The drainage triad runs for a minimum of two weeks before any targeted intervention is introduced, and it is maintained throughout the full protocol.

The Drainage Triad for Hashimoto's

Product	Clinical Role	Dosing
Lymphatic botanical blend for matrix drainage	Extracellular matrix and lymphatic drainage for ECM congestion and xenobiotic accumulation. The appropriate selection for most Hashimoto's patients with dietary toxin burden and compensation history.	1 dropper 2x daily in warm water
Methylated B12, liposomal or sublingual	Methylation support for Phase 2 liver detoxification. Low methylation capacity impairs clearance of endotoxins, hormones, and the inflammatory byproducts generated by autoimmune activity.	5 sprays sublingual 2x daily
Electrolyte and trace mineral base	Electrolyte and mineral foundation for cellular hydration. Required for efficient intracellular waste transport into the lymphatic system.	Per label direction

For patients with significant chronicity, greater than three years of active symptoms, or confirmed Lyme co-infection, a longer drainage phase of three to four weeks before introducing antimicrobial support produces substantially better clinical outcomes. The investment of time in drainage preparation pays dividends in the form of protocols that are tolerated rather than abandoned.

Lymph Formula Selection

Acute strength: acute inflammation, recent exposure, active infection phase.

Matrix strength: extracellular matrix congestion, xenobiotic accumulation, most Hashimoto's cases.

Chronic strength: longstanding degeneration, heavy metal burden, symptoms present more than 3 years.

Never use more than one Lymph formula simultaneously.

The Anti-Inflammatory Diet for Hashimoto's

Diet in Hashimoto's is not a general health recommendation. It is a targeted clinical intervention that addresses specific mechanisms driving the autoimmune process. The dietary changes most consistently associated with measurable antibody reduction in clinical research are not general anti-inflammatory eating patterns, though those matter. They are specific to the molecular interactions between dietary components and the immune and gut pathways that sustain the autoimmune attack.

The Non-Negotiable Removals

Gluten elimination is the single highest-impact dietary intervention for Hashimoto's and the one with the most direct mechanistic evidence. As described in Key 3, gliadin directly triggers zonulin release, increasing gut permeability, and generates cross-reactive antibodies that attack thyroid tissue through molecular mimicry. Both mechanisms sustain the autoimmune attack independently. Strict elimination is required. Not reduction. Elimination. This means reading labels on every packaged product, asking about preparation methods when eating outside the home, and being aware that cross-contamination from shared cooking surfaces is sufficient to maintain the immune response.

The second critical removal is conventional dairy, specifically the casein A1 protein present in most commercially available cow's milk products. A1 casein is cleaved during digestion into beta-casomorphin-7, a peptide with opioid receptor activity that has been shown to impair gut barrier integrity and promote intestinal inflammation.¹⁰ In patients with Hashimoto's who are already managing a compromised gut barrier, continuing to consume A1 dairy maintains one of the upstream barrier-disrupting inputs that allows ongoing antigen traffic into systemic circulation. Some patients tolerate A2 dairy, from certain cow breeds, goats, and sheep, without the same permeability consequences.

Refined sugar and high-glycemic carbohydrates create an intestinal environment that favors the dysbiotic organisms most likely to maintain gut permeability and sustain the LPS-driven inflammatory signaling that activates the immune response. For Hashimoto's patients with concurrent *H. pylori* or *Yersinia* colonization, high sugar intake provides a substrate that supports bacterial survival in the gut ecosystem. Removal of refined carbohydrates is both a gut barrier intervention and an anti-pathogenic dietary strategy.

Seed oils, particularly canola, soybean, corn, sunflower, and safflower oils, promote the arachidonic acid cascade and generate pro-inflammatory prostaglandins that amplify the cytokine environment in which lymphocytic thyroid infiltration occurs. The omega-6 to omega-3 ratio in a standard Western diet is approximately 15 to 20:1. The ratio associated with low autoimmune reactivity is closer to 4:1. Replacing seed oils with olive oil, avocado oil, coconut oil, and fat from grass-fed animal sources shifts this ratio and reduces the inflammatory signaling that drives autoimmune activity at the thyroid.

What to Prioritize

Wild-caught fatty fish, particularly salmon, mackerel, and sardines, provide EPA and DHA that directly resolve inflammatory signaling through specialized pro-resolving lipid mediators. These are not merely anti-inflammatory in a general sense. They actively promote immune resolution. Consuming them three to four times per week during the active treatment phase provides a consistent substrate for inflammatory resolution that cannot be replicated with supplementation alone at the same clinical impact.

Selenium-rich foods deserve particular attention in Hashimoto's. Selenium is essential for the selenoprotein thioredoxin reductase, which regulates the redox environment of the thyroid gland and protects thyroid cells from the hydrogen peroxide generated during hormone synthesis. Brazil nuts are by far the most concentrated food source, with a single nut providing approximately 70 to 90 micrograms. Two to three Brazil nuts per day provides therapeutic selenium intake for most patients. The evidence for selenium in reducing TPO antibody levels is among the more consistent in Hashimoto's nutritional research, though the magnitude of antibody reduction varies across trials.¹¹

Cruciferous vegetables, including broccoli, cauliflower, Brussels sprouts, and cabbage, require individualized guidance in Hashimoto's. Raw cruciferous vegetables contain glucosinolates that can be converted to goitrogens, compounds that interfere with iodine uptake by the thyroid. For patients with adequate iodine status, this is not a significant concern in moderate amounts. Cooking deactivates the majority of the goitrogenic compounds and makes cruciferous vegetables safe for most Hashimoto's patients in standard dietary quantities.

Strategic Supplementation for Hashimoto's

The supplement protocol for Hashimoto's follows the same sequencing principle as every protocol at Root Health: drainage must be established before targeted interventions are introduced, and targeted interventions are selected based on lab findings rather than applied universally. The products named here are the practitioner-grade formulations used in clinical practice. The framework below explains the clinical reasoning so that you understand what each phase is accomplishing and why the sequence matters.

Phase One: Drainage (Weeks 1 through 2, continued throughout)

A lymphatic botanical, methylated B12, and an electrolyte and trace mineral base form the drainage foundation as described in Key 5. No targeted thyroid or immune interventions are introduced until the drainage triad has been running for a minimum of two weeks. For patients with significant Lyme co-infection or heavy metal burden, three to four weeks of drainage preparation is the clinical standard.

Phase Two: Gut Barrier Repair (Weeks 3 through 6)

Gut barrier repair is the second priority because restoring the physical barrier that prevents ongoing antigen traffic into systemic circulation is prerequisite to meaningful autoimmune modulation. L-glutamine, ideally in a powder that also supplies glycine and proline, provides the amino acid substrate for tight junction protein synthesis and enterocyte regeneration. Bovine colostrum supports the gut-associated lymphoid tissue function that regulates the immune response to gut microbial inhabitants. Bioavailable curcumin provides anti-inflammatory support for the gut environment that accompanies most Hashimoto's cases.

For patients with confirmed gluten reactivity on the Wheat Zoomer, strict gluten elimination runs concurrently with gut barrier repair from day one. The supplement support cannot compensate for ongoing gliadin-driven zonulin release. Both the dietary removal and the structural repair must run simultaneously.

Phase Three: Thyroid and Immune Support (Weeks 4 through 8)

A thyroid support formula supplying the nutritional cofactors of thyroid hormone production, chiefly selenium, zinc, iron, tyrosine, and vitamin A, is the primary matrix support here. It is preferred over iodine-focused formulas in Hashimoto's because excess iodine can paradoxically stimulate thyroid peroxidase activity and amplify the autoimmune response in susceptible patients. For patients with confirmed iodine deficiency, the clinical picture requires individualized assessment.

Methylated B12 continues from the drainage phase through the full protocol. Beyond its methylation and drainage role, B12 supports neurological function, mood stability, and energy metabolism, all of which are commonly compromised in Hashimoto's patients due to the metabolic burden of chronic autoimmunity.

Vitamin D3 with K2 as MK-7 addresses the vitamin D deficiency that is nearly universal in Hashimoto's patients. Vitamin D functions as an immune modulator, promoting Treg cell activity that suppresses autoimmune reactivity, and correlates inversely with TPO antibody levels in multiple clinical studies.¹² The target in Hashimoto's is 60 to 80 ng/mL, with ongoing monitoring for calcium metabolism in patients requiring extended high-dose supplementation to reach that level.

Phase Four: Targeted Antimicrobial Support (Weeks 6 through 12, based on lab findings)

When EBV reactivation is confirmed on the four-marker panel, antiviral immune support is introduced after drainage and gut barrier repair are established. The protocol centers on lysine (which competitively inhibits arginine-dependent EBV replication), monolaurin for direct antiviral activity, zinc, high-dose vitamin C, and adaptogenic immune modulators that support Th1/Treg balance and reduce the Th17 polarization driving both the viral reactivation and the thyroid autoimmunity. EBV is never eradicated but can be returned to stable latency as immune function is restored.

When *H. pylori* is identified, targeted eradication support is introduced after drainage and gut barrier repair are established. *H. pylori* eradication typically requires coordination with a primary care provider for antibiotic-based eradication protocols, or naturopathic botanical approaches under clinical supervision. The drainage phase is particularly important with *H. pylori* intervention because gastric die-off generates significant endotoxin load that must clear through an adequately functioning lymphatic and hepatic system.

When *Yersinia* serology is elevated, berberine has shown activity against *Yersinia enterocolitica* in vitro and provides broad-spectrum gut antimicrobial support. In vitro activity is a laboratory finding rather than a demonstrated clinical outcome, and it should be read that way. A broad-spectrum botanical antimicrobial provides general bacterial support for the broader dysbiotic environment that typically accompanies *Yersinia* colonization.

When Lyme co-infection is confirmed, the antimicrobial protocol is substantially more involved and requires individualized clinical management based on the specific organisms identified, their burden levels, and the patient's drainage and detoxification capacity. The complete Lyme antimicrobial protocol is addressed in the companion ebook *Unlock Your Lyme*. The sequencing principle is the same: drainage first, then barrier support, then targeted antimicrobial intervention with drainage maintained throughout.

Phase	Products	Duration	Goal
1. Drainage	Lymphatic botanical blend, methylated B12, electrolyte and trace mineral base	Weeks 1-2 minimum; continue throughout	Open lymphatic, liver Phase 2, cellular hydration
2. Gut Barrier	L-glutamine with glycine and proline, colostrum, bioavailable curcumin	Weeks 3-6	Restore barrier, reduce antigen traffic, repair GALT
3. Thyroid/Immune	Thyroid cofactor formula, methylated B12 (continuing), vitamin D3 with K2, selenium (food or supplemental)	Weeks 4-8	Support hormone conversion, modulate immune response, optimize D3
4. Antimicrobial / Antiviral	EBV: Lysine, Monolaurin, Zinc, Vitamin C, immune adaptogens. H. pylori: per eradication protocol. Yersinia: berberine plus broad-spectrum botanical antimicrobial support. Lyme: per Unlock Your Lyme.	Weeks 6-12, based on lab findings	Eliminate infectious triggers with drainage and barrier in place

Where to get these

Every item in this protocol is a single ingredient or a standardized botanical, available from any reputable retailer. Look for independent third-party testing (NSF, USP Verified, or equivalent) and match the form named here rather than the name on the front of the bottle.

If you would rather not evaluate that yourself, Root Health keeps a curated dispensary of professional-grade products, screened for those standards and organized by the phases above.

Root Health dispensary: us.fullscript.com/welcome/drseay

Disclosure: Root Health earns a portion of dispensary purchases. Nothing here was included, excluded, ranked, or dosed based on what it earns, and you can buy any of it anywhere.

Your Next Steps with Root Health

The framework in this guide is the clinical picture that most Hashimoto's patients never receive. It identifies the upstream drivers, provides a testing sequence that captures the full autoimmune picture, and lays out a treatment approach organized by mechanistic priority rather than symptom suppression. What it cannot do is replace individualized clinical care. Your specific antibody pattern, your gut findings, your H. pylori and Yersinia status, your nutrient deficiencies, and your history all determine how this framework applies to your case.

Start with the Right Testing

For every confirmed Hashimoto's patient, the minimum starting workup is the Vibrant Hormone Zoomer at \$710 and a LabCorp H. pylori stool antigen test at approximately \$50 to 80. These two tests together answer the two most immediately actionable questions: where is the thyroid conversion breaking down, and is H. pylori an active upstream trigger. Add the Wheat Zoomer at \$479 when gluten reactivity is suspected or when gut symptoms are present. Add Yersinia serology when antibodies are not responding to initial gut and dietary intervention. Add the Tickborne 2.0 PCR at \$1,406 when tick exposure history is present or when antibodies are not responding after the full infectious workup. All panels except H. pylori are available at rootcauseunlocked.com/order-labs.

Book a Root Discovery Session

A Root Discovery Session is a complimentary 30-minute clinical conversation with Dr. Seay. It is not a sales call. It is a focused clinical intake where your symptom history, prior testing, and current clinical picture are reviewed to identify the most targeted path forward. If you are uncertain which workup is appropriate for your case, or if you want clinical interpretation of prior results before investing in additional testing, this is the right starting point. Book at rootcauseunlocked.com/discovery.

The Root Restore Program

For patients who are ready for structured, supervised 1-on-1 clinical care, the Root Restore Program is a 6-month protocol that includes comprehensive testing, a personalized drainage-first supplement protocol, dietary guidance, and ongoing clinical oversight. Payment plans are available. Learn more at rootcauseunlocked.com/hashimotos.

A Note from Dr. Seay

If you have been on thyroid medication for years and you still do not feel well, that is information. It means the immune attack is continuing. It means the upstream drivers have not been addressed. It does not mean the

medication is wrong or that nothing can be done. It means the right investigation has not happened yet.

That investigation is exactly what Root Health is built for.

Dr. Adam Seay, DC · Root Health, Garner NC · rootcauseunlocked.com

Root Health · Garner, NC · rootcauseunlocked.com/hashimotos

References

1. Ralli et al., "Hashimoto's thyroiditis: An update on pathogenic mechanisms, diagnostic protocols, therapeutic strategies, and potential malignant transformation," *Autoimmunity Reviews*, 2020. <https://pubmed.ncbi.nlm.nih.gov/32805423/>
2. Cayres et al., "Detection of Alterations in the Gut Microbiota and Intestinal Permeability in Patients With Hashimoto Thyroiditis," *Frontiers in Immunology*, 2021. <https://pubmed.ncbi.nlm.nih.gov/33746942/>
3. Araújo et al., "Effects of Gluten-Free Diet in Non-Celiac Hashimoto's Thyroiditis: A Systematic Review and Meta-Analysis," *Nutrients*, 2025. <https://pubmed.ncbi.nlm.nih.gov/41228508/>
4. Wang et al., "Identification of outer membrane porin f protein of *Yersinia enterocolitica* recognized by antithyrotropin receptor antibodies in Graves' disease and determination of its epitope using mass spectrometry and bioinformatics tools," *The Journal of Clinical Endocrinology and Metabolism*, 2010. <https://pubmed.ncbi.nlm.nih.gov/20484489/>
5. Effraïmidis et al., "No causal relationship between *Yersinia enterocolitica* infection and autoimmune thyroid disease: evidence from a prospective study," *Clinical and Experimental Immunology*, 2011. <https://pubmed.ncbi.nlm.nih.gov/21488870/>
6. Shi et al., "Associations of *Helicobacter pylori* infection and cytotoxin-associated gene A status with autoimmune thyroid diseases: a meta-analysis," *Thyroid*, 2013. <https://pubmed.ncbi.nlm.nih.gov/23544831/>
7. Flower et al., "Investigating the Link between the Intestinal Microbiome and Th17/Treg Dysregulation in Hashimoto's Thyroiditis, and the Therapeutic Potential of Vitamin D," *Journal of the American Nutrition Association*, 2026. <https://pubmed.ncbi.nlm.nih.gov/41920977/>
8. Mertowska et al., "Does Epstein-Barr virus and intracellular Toll-like receptors affect the course of Hashimoto disease? Findings from studies on newly-diagnosed patients," *Polish Archives of Internal Medicine*, 2024. <https://pubmed.ncbi.nlm.nih.gov/39508448/>
9. Macedo et al., "Detection of SARS-CoV-2 infection in thyroid follicular cells from a COVID-19 autopsy series," *European Thyroid Journal*, 2022. <https://pubmed.ncbi.nlm.nih.gov/35900859/>
10. Robinson et al., "Effects of Different Cow-Milk Beta-Caseins on the Gut-Brain Axis: A Narrative Review of Preclinical, Animal, and Human Studies," *Nutrition Reviews*, 2025. <https://pubmed.ncbi.nlm.nih.gov/39024213/>
11. Huwiler et al., "Selenium Supplementation in Patients with Hashimoto Thyroiditis: A Systematic Review and Meta-Analysis of Randomized Clinical Trials," *Thyroid*, 2024. <https://pubmed.ncbi.nlm.nih.gov/38243784/>
12. Sun et al., "Vitamin D deficiency in Hashimoto's thyroiditis: mechanisms, immune modulation, and therapeutic implications," *Frontiers in Endocrinology*, 2025. <https://pubmed.ncbi.nlm.nih.gov/40822954/>