

# Morphofunctional changes at the active zone during synaptic vesicle exocytosis

This manuscript ([permalink](#)) was automatically generated from [aseedb/synaptic\\_tomo\\_ms@67b263b](#) on August 20, 2021.

## Authors

---

- **Julika Radecke\***

 [0000-0002-5815-5537](#) ·  [julikaradecke](#)

Institute of Anatomy, University of Bern, Bern, Switzerland; Department of Neuroscience, Faculty of Health and Medical Sciences, 2200 Copenhagen N, University of Copenhagen, Copenhagen, Denmark; Diamond Light Source Ltd, Didcot, Oxfordshire, United Kingdom · Funded by Grant XXXXXXXX

- **Raphaela Seeger\***

 [XXXX-XXXX-XXXX-XXXX](#) ·  [elatella](#)

Institute of Anatomy, University of Bern, Bern, Switzerland; Graduate School for Cellular and Biomedical Sciences, University of Bern

- **Ulrike Laugks**

 [0000-0003-4175-4354](#)

Max-Planck-Institute of Biochemistry, Am Klopferspitz 18, 82152 Martinsried, Germany

- **Kenneth N. Goldie**

 [0000-0002-7405-0049](#)

Center for Cellular Imaging and NanoAnalytics, Biozentrum, University of Basel, Basel, Switzerland

- **Henning Stahlberg**

 [0000-0002-1185-4592](#) ·  [sthennin](#)

Center for Cellular Imaging and NanoAnalytics, Biozentrum, University of Basel, Basel, Switzerland; EPFL; UNIL

- **Vladan Lučić**

 [0000-0003-3698-7436](#) ·  [vladanl](#)

Max-Planck-Institute of Biochemistry, Am Klopferspitz 18, 82152 Martinsried, Germany

- **Jakob B. Sørensen**✉

 [0000-0001-5465-3769](#) ·  [JBSorensen](#)

Department of Neuroscience, Faculty of Health and Medical Sciences, 2200 Copenhagen N, University of Copenhagen, Copenhagen, Denmark

- **Benoît Zuber**✉

 [0000-0001-7725-5579](#) ·  [aseedb](#)

Institute of Anatomy, University of Bern, Bern, Switzerland · Funded by Swiss National Science Foundation (Grant# xxxx)

✉ Address correspondence to [benoit.zuber@ana.unibe.ch](mailto:benoit.zuber@ana.unibe.ch) and [jakobbs@sund.ku.dk](mailto:jakobbs@sund.ku.dk).

\* These authors contributed equally.

# Abstract

---

The fusion of synaptic vesicles (SVs) with the plasma membrane (PM) proceeds through intermediate steps that remain poorly resolved. Additionally, the effect of persistent high or low exocytosis activity on intermediate steps remains unknown. Through time-resolved cryo-electron tomography, we ordered events into a sequence. Following stimulation, additional SVs are rapidly primed by forming tethers with the PM. Simultaneously, fusion initiation occurs by membrane curvature ('buckling') of the SV and PM. It is followed by the formation of a fusion pore, and the collapse of SV membrane. At this time, membrane-proximal, but not membrane-distal, vesicles lose their interconnections, allowing them to move towards the PM. A SNARE mutation that arrests spontaneous release caused vesicles to reside further from the membrane while forming more tethers, whereas a mutation stimulating spontaneous fusion caused a complete loss of membrane-proximal triple-tethered SVs, and a loss of intervesicle connectors. Overall, tether formation and connector dissolution is triggered by stimulation and adjusted to the spontaneous fusion rate. These morphological observations likely correspond to the transition of SVs from one functional pool to another.

- *Need to reformulate the part about 4E mutant (p value is not sufficient)*
- *For 4K mutant: check if the loss of tethered SVs restricted to those with more than only one, or to those with more than two tethers?*
- *Possibly mention here introduce the formation of distal connectors by stimulation*

## Introduction

---

In the central nervous system, neurons communicate through the release of neurotransmitters at synapses. This process relies on synaptic vesicle (SV) exocytosis, i.e. the fusion of SVs with the plasma membrane (PM). This in turn is eminently important for normal brain function such as movement coordination or memory formation. SV exocytosis involves a sequence of steps [1,2]. The vesicle is first docked to the active zone (AZ) PM. Subsequently the exocytosis machinery goes through a maturation process, termed priming, after which the SV is ready to fuse. These SVs form the readily releasable pool (RRP) of SVs. Finally, a calcium influx triggers fusion of the SV with the PM. Docked SVs are defined as the SVs in very close proximity or direct contact with the PM as observed by electron microscopy (EM), whereas priming refers to SV ability to undergo exocytosis immediately upon stimulation. Whether every docked SV is also primed has been debated [1,3]. A recent high-pressure freezing/freeze-substitution EM study of genetically modified synapses has indicated that vesicles that are in direct contact with the PM, i.e. docked, are primed and belong to the RRP and that this situation occurs downstream of vesicle tethering [4]. From a molecular perspective, priming involves several proteins, including the SNARE complex (SNAP-25, syntaxin-1, and synaptobrevin-2), Munc13, Munc18, synaptotagmin-1, and complexin [2,5]. All three SNAREs form a highly stable tight four-helix bundle, known as trans-SNARE complex. The surfaces of the SV and the PM, respectively, are negatively charged and therefore tend to repulse each other. The formation of the trans-SNARE complex counteracts this repulsion and brings the SV and the PM in high proximity [6]. Evidence has suggested that the SNARE complex is only partially zipped in primed SVs [7]. Furthermore, various studies have suggested that the formation of at least three SNARE complexes provides the necessary energy for a SV to become fusion-competent [8,9,10]. Yet in the absence of cytoplasmic  $\text{Ca}^{2+}$ , minimal spontaneous exocytosis takes place. When the presynaptic terminal gets depolarized by an action potential,  $\text{Ca}^{2+}$  flows in the cytoplasm and binds to synaptotagmin-1, which is localized at the SV surface. Upon  $\text{Ca}^{2+}$  binding, synaptotagmin-1 was proposed to insert between the head groups of the PM anionic phospholipids and trigger membrane curvature and destabilization, leading first to hemifusion and subsequently to fusion [11]. Interestingly, the trans-SNARE bundle surface is negatively charged, which contributes to the electrostatic barrier that minimizes spontaneous fusion and allows synaptotagmin-1 to act as an electrostatic switch that triggers exocytosis [12]. Introducing

negatively charged side chains by site-directed mutagenesis reduces the rate of spontaneous and evoked exocytosis whereas introducing more positive side chains enhances the rate of spontaneous exocytosis and depletes the RRP.

Cryo-electron tomography (cryo-ET), which preserves samples to atomic resolution, revealed that under resting conditions, no SV is in direct contact with the PM and the majority of AZ-proximal SVs are connected to the PM by a variable number of short tethers [13,14]. The observed gap between the SV and the PM is consistent with the model of an electrostatic barrier formed by the negative charges of the SV, the PM, and the trans-SNARE bundle [12]. In synaptosomes treated with hypertonic sucrose solution, which depletes the RRP, the majority of tethered vesicles had 1 or 2 tethers [13,15,16]. This observation suggested that the RRP consists of SV that are linked to the PM by 3 or more tethers. The RRP, as identified by morphological criteria, only represents a minority of AZ-proximal vesicles. This is in agreement with previous reports. In one of them the term pre-primed pool was used for the few vesicles (~1 vesicle at hippocampal synapses) that are rapidly released and another publication showed that the immediately releasable pool is made up of only 10-20% of the vesicles located on the AZ (equal to ~1 vesicle on hippocampal synapses) [17,18]. The ensemble of proximal vesicles that are not in the RRP have been termed non-RRP and presumably belong to the recycling pool that releases more slowly [13,19]. Farther away from the AZ, partially intermixed with the recycling pool, is the reserve pool containing vesicles that only release upon high frequency stimulation. Vesicles in the reserve pool are tightly clustered and well inter-connected by structures that were termed connectors [13,19]. It should be noted that the molecular nature of connectors is not known and is possibly heterogenous. Synapsin has been proposed as a molecular constituent but since the deletion of all forms of synapsin does not lead to the complete absence of connectors, it is clear that not all connectors contain synapsin [20,21]. The second row of SVs near the active zone, immediately after the proximal vesicles, is called the intermediate region. Resting state intermediate SVs are less densely packed and also less connected than proximal SVs [14]. This suggests that, after exocytosis of RRP SVs, intermediate SVs could be rapidly recruited in the RRP by diffusion [22]. Synaptic activity enhances the mobility of a fraction of SVs, whereas it induces synapsin dissociation from SVs in a synapsin phosphorylation-dependent manner [23,24]. The same mobility enhancement can be achieved through inhibition of synapsin dephosphorylation, which leads to synapsin dissociation from SVs, or by knocking out all three synapsin forms [25,26,27]. Interestingly, ribbon synapses do not express synapsin and show higher SV mobility than conventional synapses [28]. It is therefore conceivable that inter-SV connectors restrain SV diffusion and that synaptic activity influences the level of inter-SV connectivity and thereby their mobility.

To investigate this hypothesis and to better understand the impact of depolarization and synaptic activity on SV tethering, we designed two sets of cryo-ET experiments. On the one hand, we compared the morphology of wild-type rat synaptosome in resting state and a few milliseconds after depolarization. On the other hand, we imaged autapses in mice neuronal culture expressing either wild-type SNAP-25, a more positively charged SNAP-25 mutant, or a more negatively charged mutant of SNAP-25. The more positive SNAP-25 mutant, which is constitutively active and whose RRP is permanently depleted showed no triple-tethered SV, which confirmed the morphological definition of the RRP. Our experiments revealed that immediately after depolarization additional SVs are recruited to the RRP. Shortly after exocytosis the level of inter-SV connectivity was decreased among SVs situated in a 25 to 75-nm distance range from the AZ PM. Altogether, our results indicate that connectors regulate SV mobility and their recruitment at the AZ PM.

*Say something about the size of the RRP Do we want to talk about kiss-and-run? I have the feeling that we better not to keep the message nice and clear*

## Results

---

## Discussion

---

## Materials and methods

---

# References

---

1. **Vesicle Docking in Regulated Exocytosis**  
Matthijs Verhage, Jakob B Sørensen  
*Traffic* (2008-09) <https://doi.org/bjtx2n>  
DOI: [10.1111/j.1600-0854.2008.00759.x](https://doi.org/10.1111/j.1600-0854.2008.00759.x) · PMID: [18445120](https://pubmed.ncbi.nlm.nih.gov/18445120/)
2. **Neurotransmitter Release: The Last Millisecond in the Life of a Synaptic Vesicle**  
Thomas C Südhof  
*Neuron* (2013-10) <https://doi.org/f5gng4>  
DOI: [10.1016/j.neuron.2013.10.022](https://doi.org/10.1016/j.neuron.2013.10.022) · PMID: [24183019](https://pubmed.ncbi.nlm.nih.gov/24183019/) · PMCID: [PMC3866025](https://pubmed.ncbi.nlm.nih.gov/PMC3866025/)
3. **The readily releasable pool of synaptic vesicles**  
Pascal S Kaeser, Wade G Regehr  
*Current Opinion in Neurobiology* (2017-04) <https://doi.org/gbkfsd>  
DOI: [10.1016/j.conb.2016.12.012](https://doi.org/10.1016/j.conb.2016.12.012) · PMID: [28103533](https://pubmed.ncbi.nlm.nih.gov/28103533/) · PMCID: [PMC5447466](https://pubmed.ncbi.nlm.nih.gov/PMC5447466/)
4. **The Morphological and Molecular Nature of Synaptic Vesicle Priming at Presynaptic Active Zones**  
Cordelia Imig, Sang-Won Min, Stefanie Krinner, Marife Arancillo, Christian Rosenmund, Thomas C Südhof, JeongSeop Rhee, Nils Brose, Benjamin H Cooper  
*Neuron* (2014-10) <https://doi.org/gcvj2v>  
DOI: [10.1016/j.neuron.2014.10.009](https://doi.org/10.1016/j.neuron.2014.10.009) · PMID: [25374362](https://pubmed.ncbi.nlm.nih.gov/25374362/)
5. **The Synaptic Vesicle Release Machinery**  
Josep Rizo, Junjie Xu  
*Annual Review of Biophysics* (2015-06-22) <https://doi.org/gjnb9q>  
DOI: [10.1146/annurev-biophys-060414-034057](https://doi.org/10.1146/annurev-biophys-060414-034057) · PMID: [26098518](https://pubmed.ncbi.nlm.nih.gov/26098518/)
6. **Crystal structure of a SNARE complex involved in synaptic exocytosis at 2.4 Å resolution**  
RBryan Sutton, Dirk Fasshauer, Reinhard Jahn, Axel T Brunger  
*Nature* (1998-09) <https://doi.org/cwkm8k>  
DOI: [10.1038/26412](https://doi.org/10.1038/26412) · PMID: [9759724](https://pubmed.ncbi.nlm.nih.gov/9759724/)
7. **Sequential N- to C-terminal SNARE complex assembly drives priming and fusion of secretory vesicles**  
Jakob B Sørensen, Katrin Wiederhold, Emil M Müller, Ira Milosevic, Gábor Nagy, Bert L de Groot, Helmut Grubmüller, Dirk Fasshauer  
*The EMBO Journal* (2006-02-23) <https://doi.org/dhpwm3>  
DOI: [10.1038/sj.emboj.7601003](https://doi.org/10.1038/sj.emboj.7601003) · PMID: [16498411](https://pubmed.ncbi.nlm.nih.gov/16498411/) · PMCID: [PMC1409717](https://pubmed.ncbi.nlm.nih.gov/PMC1409717/)
8. **Single Vesicle Millisecond Fusion Kinetics Reveals Number of SNARE Complexes Optimal for Fast SNARE-mediated Membrane Fusion**  
Marta K Domanska, Volker Kiessling, Alexander Stein, Dirk Fasshauer, Lukas K Tamm  
*Journal of Biological Chemistry* (2009-11) <https://doi.org/chdxqp>  
DOI: [10.1074/jbc.m109.047381](https://doi.org/10.1074/jbc.m109.047381) · PMID: [19759010](https://pubmed.ncbi.nlm.nih.gov/19759010/) · PMCID: [PMC2797286](https://pubmed.ncbi.nlm.nih.gov/PMC2797286/)
9. **Fast Vesicle Fusion in Living Cells Requires at Least Three SNARE Complexes**  
R Mohrmann, H de Wit, M Verhage, E Neher, JB Sorensen  
*Science* (2010-09-16) <https://doi.org/c7q87p>  
DOI: [10.1126/science.1193134](https://doi.org/10.1126/science.1193134) · PMID: [20847232](https://pubmed.ncbi.nlm.nih.gov/20847232/)
10. **SNARE Proteins: One to Fuse and Three to Keep the Nascent Fusion Pore Open**  
L Shi, Q-T Shen, A Kiel, J Wang, H-W Wang, TJ Melia, JE Rothman, F Pincet

*Science* (2012-03-15) <https://doi.org/gk8gpf>  
DOI: [10.1126/science.1214984](https://doi.org/10.1126/science.1214984) · PMID: [22422984](https://pubmed.ncbi.nlm.nih.gov/22422984/) · PMCID: [PMC3736847](https://pubmed.ncbi.nlm.nih.gov/PMC3736847/)

11. **Membrane Curvature in Synaptic Vesicle Fusion and Beyond**  
Harvey T McMahon, Michael M Kozlov, Sascha Martens  
*Cell* (2010-03) <https://doi.org/d3gmvv>  
DOI: [10.1016/j.cell.2010.02.017](https://doi.org/10.1016/j.cell.2010.02.017) · PMID: [20211126](https://pubmed.ncbi.nlm.nih.gov/20211126/)
12. **An Electrostatic Energy Barrier for SNARE-Dependent Spontaneous and Evoked Synaptic Transmission**  
Marvin Ruiter, Anna Kádková, Andrea Scheutzw, Jörg Malsam, Thomas H Söllner, Jakob B Sørensen  
*Cell Reports* (2019-02) <https://doi.org/gfv5gd>  
DOI: [10.1016/j.celrep.2019.01.103](https://doi.org/10.1016/j.celrep.2019.01.103) · PMID: [30811985](https://pubmed.ncbi.nlm.nih.gov/30811985/)
13. **Quantitative analysis of the native presynaptic cytomatrix by cryoelectron tomography**  
Rubén Fernández-Busnadiego, Benoît Zuber, Ulrike Elisabeth Maurer, Marek Cyrklaff, Wolfgang Baumeister, Vladan Lučić  
*Journal of Cell Biology* (2010-01-11) <https://doi.org/b9c26b>  
DOI: [10.1083/jcb.200908082](https://doi.org/10.1083/jcb.200908082) · PMID: [20065095](https://pubmed.ncbi.nlm.nih.gov/20065095/) · PMCID: [PMC2812849](https://pubmed.ncbi.nlm.nih.gov/PMC2812849/)
14. **Molecular architecture of the presynaptic terminal**  
Benoît Zuber, Vladan Lučić  
*Current Opinion in Structural Biology* (2019-02) <https://doi.org/gk8gpd>  
DOI: [10.1016/j.sbi.2019.01.008](https://doi.org/10.1016/j.sbi.2019.01.008) · PMID: [30925443](https://pubmed.ncbi.nlm.nih.gov/30925443/)
15. **Definition of the readily releasable pool of vesicles at hippocampal synapses.**  
C Rosenmund, CF Stevens  
*Neuron* (1996-06) <https://www.ncbi.nlm.nih.gov/pubmed/8663996>  
DOI: [10.1016/s0896-6273\(00\)80146-4](https://doi.org/10.1016/s0896-6273(00)80146-4) · PMID: [8663996](https://pubmed.ncbi.nlm.nih.gov/8663996/)
16. **Properties of Synaptic Vesicle Pools in Mature Central Nerve Terminals**  
Anthony C Ashton, Yuri A Ushkaryov  
*Journal of Biological Chemistry* (2005-11) <https://doi.org/cb5883>  
DOI: [10.1074/jbc.m504137200](https://doi.org/10.1074/jbc.m504137200) · PMID: [16148008](https://pubmed.ncbi.nlm.nih.gov/16148008/)
17. **Vesicle release probability and pre-primed pool at glutamatergic synapses in area CA1 of the rat neonatal hippocampus**  
Eric Hanse, Bengt Gustafsson  
*The Journal of Physiology* (2001-03) <https://doi.org/fq7th3>  
DOI: [10.1111/j.1469-7793.2001.04811.x](https://doi.org/10.1111/j.1469-7793.2001.04811.x) · PMID: [11230520](https://pubmed.ncbi.nlm.nih.gov/11230520/) · PMCID: [PMC2278469](https://pubmed.ncbi.nlm.nih.gov/PMC2278469/)
18. **Reluctant Vesicles Contribute to the Total Readily Releasable Pool in Glutamatergic Hippocampal Neurons**  
KL Moulder  
*Journal of Neuroscience* (2005-04-13) <https://doi.org/dwqpxj>  
DOI: [10.1523/jneurosci.5231-04.2005](https://doi.org/10.1523/jneurosci.5231-04.2005) · PMID: [15829636](https://pubmed.ncbi.nlm.nih.gov/15829636/) · PMCID: [PMC6724923](https://pubmed.ncbi.nlm.nih.gov/PMC6724923/)
19. **Synaptic vesicle pools: an update**  
Denker  
*Frontiers in Synaptic Neuroscience* (2010) <https://doi.org/d74dd7>  
DOI: [10.3389/fnsyn.2010.00135](https://doi.org/10.3389/fnsyn.2010.00135) · PMID: [21423521](https://pubmed.ncbi.nlm.nih.gov/21423521/) · PMCID: [PMC3059705](https://pubmed.ncbi.nlm.nih.gov/PMC3059705/)
20. **The cytoskeletal architecture of the presynaptic terminal and molecular structure of synapsin 1.**  
N Hirokawa, K Sobue, K Kanda, A Harada, H Yorifuji

*Journal of Cell Biology* (1989-01-01) <https://doi.org/ddbqhb>  
DOI: [10.1083/jcb.108.1.111](https://doi.org/10.1083/jcb.108.1.111) · PMID: [2536030](https://pubmed.ncbi.nlm.nih.gov/2536030/) · PMCID: [PMC2115350](https://pubmed.ncbi.nlm.nih.gov/PMC2115350/)

21. **Three-Dimensional Architecture of Presynaptic Terminal Cytomatrix**  
L Siksou, P Rostaing, J-P Lechaire, T Boudier, T Ohtsuka, A Fejtova, H-T Kao, P Greengard, ED Gundelfinger, A Triller, S Marty  
*Journal of Neuroscience* (2007-06-27) <https://doi.org/bjw3mv>  
DOI: [10.1523/jneurosci.1773-07.2007](https://doi.org/10.1523/jneurosci.1773-07.2007) · PMID: [17596435](https://pubmed.ncbi.nlm.nih.gov/17596435/) · PMCID: [PMC6672225](https://pubmed.ncbi.nlm.nih.gov/PMC6672225/)
22. **Physical determinants of vesicle mobility and supply at a central synapse**  
Jason Seth Rothman, Laszlo Kocsis, Etienne Herzog, Zoltan Nusser, Robin Angus Silver  
*eLife* (2016-08-19) <https://doi.org/f9rmkt>  
DOI: [10.7554/elife.15133](https://doi.org/10.7554/elife.15133) · PMID: [27542193](https://pubmed.ncbi.nlm.nih.gov/27542193/) · PMCID: [PMC5025287](https://pubmed.ncbi.nlm.nih.gov/PMC5025287/)
23. **Activity-Dependence of Synaptic Vesicle Dynamics**  
Luca A Forte, Michael W Gramlich, Vitaly A Klyachko  
*The Journal of Neuroscience* (2017-11-01) <https://doi.org/gch7j3>  
DOI: [10.1523/jneurosci.0383-17.2017](https://doi.org/10.1523/jneurosci.0383-17.2017) · PMID: [28954868](https://pubmed.ncbi.nlm.nih.gov/28954868/) · PMCID: [PMC5666583](https://pubmed.ncbi.nlm.nih.gov/PMC5666583/)
24. **Synapsin dispersion and reclustering during synaptic activity**  
Ping Chi, Paul Greengard, Timothy A Ryan  
*Nature Neuroscience* (2001-10-29) <https://doi.org/cpdwc7>  
DOI: [10.1038/nn756](https://doi.org/10.1038/nn756) · PMID: [11685225](https://pubmed.ncbi.nlm.nih.gov/11685225/)
25. **Interactions of synapsin I with small synaptic vesicles: distinct sites in synapsin I bind to vesicle phospholipids and vesicle proteins.**  
F Benfenati, M Böhler, R Jahn, P Greengard  
*Journal of Cell Biology* (1989-05-01) <https://doi.org/d65t4v>  
DOI: [10.1083/jcb.108.5.1863](https://doi.org/10.1083/jcb.108.5.1863) · PMID: [2497106](https://pubmed.ncbi.nlm.nih.gov/2497106/) · PMCID: [PMC2115532](https://pubmed.ncbi.nlm.nih.gov/PMC2115532/)
26. **Visualization of Synaptic Vesicle Movement in Intact Synaptic Boutons Using Fluorescence Fluctuation Spectroscopy**  
Randolf Jordan, Edward A Lemke, Jurgen Klingauf  
*Biophysical Journal* (2005-09) <https://doi.org/cdpkky>  
DOI: [10.1529/biophysj.105.061663](https://doi.org/10.1529/biophysj.105.061663) · PMID: [15980175](https://pubmed.ncbi.nlm.nih.gov/15980175/) · PMCID: [PMC1366711](https://pubmed.ncbi.nlm.nih.gov/PMC1366711/)
27. **Synapsin Selectively Controls the Mobility of Resting Pool Vesicles at Hippocampal Terminals**  
A Orenbuch, L Shalev, V Marra, I Sinai, Y Lavy, J Kahn, JJ Burden, K Staras, D Gitler  
*Journal of Neuroscience* (2012-03-21) <https://doi.org/f3wq83>  
DOI: [10.1523/jneurosci.5058-11.2012](https://doi.org/10.1523/jneurosci.5058-11.2012) · PMID: [22442064](https://pubmed.ncbi.nlm.nih.gov/22442064/) · PMCID: [PMC3492757](https://pubmed.ncbi.nlm.nih.gov/PMC3492757/)
28. **High Mobility of Vesicles Supports Continuous Exocytosis at a Ribbon Synapse**  
Matthew Holt, Anne Cooke, Andreas Neef, Leon Lagnado  
*Current Biology* (2004-02) <https://doi.org/b9g9h2>  
DOI: [10.1016/j.cub.2003.12.053](https://doi.org/10.1016/j.cub.2003.12.053) · PMID: [14761649](https://pubmed.ncbi.nlm.nih.gov/14761649/)