Morphofunctional changes at the active zone during synaptic vesicle exocytosis

This manuscript (permalink) was automatically generated from aseedb/synaptic tomo ms@8f163ea on July 21, 2021.

Authors

• Julika Radecke*

© 0000-0002-5815-5537

Institute of Anatomy, University of Bern, Bern, Switzerland; Department of Neuroscience, Faculty of Health and Medical Sciences, 2200 Copenhagen N, University of Copenhagen, Copenhagen, Denmark; Diamond Light Source Ltd, Didcot, Oxfordshire, United Kingdom · Funded by Grant XXXXXXXX

Raphaela Seeger^{*}

ⓑ XXXX-XXXX-XXXX · **۞** elatella

Institute of Anatomy, University of Bern, Bern, Switzerland; Graduate School for Cellular and Biomedical Sciences, University of Bern

Ulrike Laugks

© 0000-0003-4175-4354

Max-Planck-Institute of Biochemistry, Am Klopferspitz 18, 82152 Martinsried, Germany

Kenneth N. Goldie

D 0000-0002-7405-0049

Center for Cellular Imaging and NanoAnalytics, Biozentrum, University of Basel, Basel, Switzerland

Henning Stahlberg

D 0000-0002-1185-4592 · **C** sthennin

Center for Cellular Imaging and NanoAnalytics, Biozentrum, University of Basel, Basel, Switzerland; EPFL; UNIL

Vladan Lučić

© 0000-0003-3698-7436 · ♥ vladanl

Max-Planck-Institute of Biochemistry, Am Klopferspitz 18, 82152 Martinsried, Germany

Jakob B. Sørensen[™]

© 0000-0001-5465-3769

Department of Neuroscience, Faculty of Health and Medical Sciences, 2200 Copenhagen N, University of Copenhagen, Copenhagen, Denmark

• Benoît Zuber[≌]

(D 0000-0001-7725-5579 ⋅ **(7** aseedb

Institute of Anatomy, University of Bern, Bern, Switzerland · Funded by Swiss National Science Foundation (Grant# xxxx)

✓ Address correspondence to benoit.zuber@ana.unibe.ch and jakobbs@sund.ku.dk. ::: {#equal_contribution} * These authors contributed equally.

Abstract

The fusion of synaptic vesicles (SVs) with the plasma membrane (PM) proceeds through intermediate steps that remain poorly resolved. Additionally, the effect of persistent high or low exocytosis activity on intermediate steps remains unknown. Through time-resolved cryo-electron tomography, we ordered events into a sequence. Following stimulation, additional SVs are rapidly primed by forming tethers with the PM. Simultaneously, fusion initiation occurs by membrane curvature ('buckling') of the SV and PM. It is followed by the formation of a fusion pore, and the collapse of SV membrane. At this time, membrane-proximal, but not membrane-distal, vesicles loose their interconnections, allowing them to move towards the PM. A SNARE mutation that arrests spontaneous release caused vesicles to reside further from the membrane while forming more tethers, whereas a mutation stimulating spontaneous fusion caused a loss of membrane-proximal SVs with more than two tethers, and loss of intervesicle connectors. Overall, tether formation and connector dissolution is triggered by stimulation and adjusted to the spontaneous fusion rate.

- Need to reformulate the part about 4E mutant
- For 4K mutant: is the loss of tethered SVs restricted to those with more than only one, or to those with more than two tethers?
- Possibly introduce the formation of distal connectors by stimulation

Introduction

In the central nervous system, neurons communicate through the release of neurotransmitters at synapses. This process relies on synaptic vesicle (SV) exocytosis, i.e. the fusion of SVs with the plasma membrane (PM). This in turn is eminently important for normal brain function such as movement coordination or memory formation. SV exocytosis involves a sequence of steps [1,2]. The vesicle is first docked to the active zone (AZ) PM. Subsequently the exocytosis machinery goes through a maturation process, termed priming, after which the SV is ready to fuse. These SVs form the readily releasable pool (RRP) of SVs. Finally, a calcium influx triggers fusion of the SV with the PM. Docked SVs are defined as the SVs in very close proximity or direct contact with the PM as observed by electron microscopy (EM), whereas priming refers to SV ability to undergo exocytosis immediately upon stimulation. Whether every docked SV is also primed has been debated [1,3]. A recent high-pressure freezing/freeze-substitution EM study of genetically modified synapses has indicated that vesicles that are in direct contact with the PM, i.e. docked, are primed and belong to the RRP and that this situation occurs downstream of vesicle tethering [4]. From a molecular perspective, priming involves several proteins, including the SNARE complex (SNAP-25, syntaxin-1, and synaptobrevin-2), Munc13, Munc18, synaptotagmin-1, and complexin [2,5]. All three SNAREs form a highly stable tight four-helix bundle, known as trans-SNARE complex. The surfaces of the SV and the PM, respectively, are negatively charged and therefore tend to repulse each other. The formation of the trans-SNARE complex conteracts this repulsion and brings the SV and the PM in high proximity [6]. Evidence has suggested that the SNARE complex is only partially zipped in primed SVs [doi:0.1038/sj.emboj.7601003?]. Furthermore, various studies have suggested that the formation of at least three SNARE complexes provides the necessary energy for a SV to become fusion-competent [7,8,9]. Yet in the absence of cytoplasmic Ca²⁺, minimal spontaneous exocytosis takes place. When the presynaptic terminal gets depolarized by an action potential, Ca²⁺ flows in the cytoplasm and binds to synaptotagmin-1, which is localized at the SV surface. Upon Ca²⁺ binding, synaptotagmin-1 was proposed to insert between the head groups of the PM anionic phospholipids and trigger membrane curvature and destabilization, leading first to hemifusion and subsequently to fusion[10]. Interestingly, the trans-SNARE bundle surface is negatively charged, which contributes to the electrostatic barrier that minimizes spontaneous fusion and allows synaptotagmin-1 to act as an electrostatic switch that triggers exocytosis [11]. Making the bundle more negative by site-directed mutagenesis reduces the

rate of spontaneous and evoked exocytosis whereas making it more positive enhances the rate of spontaneous exocytosis and depletes the RRP.

Cryo-electron tomography (cryo-ET), which preserves samples to atomic resolution, revealed that under resting conditions, no SV is in direct contact with the PM and the majority of AZ-proximal SVs are connected to the PM by a variable number of short tethers [12,13]. The observed gap between the SV and the PM is consistent with the model of an electrostatic barrier formed by the negative charges of the SV, the PM, and the trans-SNARE bundle [11]. In synaptosomes treated with hypertonic sucrose solution, which depletes the RRP, the majority of tethered vesicles had 1 or 2 tethers [12,14,doi:10.1016/s0896-6273?]. This observation suggested that the RRP consists of SV that are linked to the PM by 3 or more tethers. The RRP, as identified by morphological criteria, only represents a minority of AZ-proximal vesicles. This is in agreement with previous reports. In one of them the term pre-primed pool was used for the few vesicles (~1 vesicle at hippocampal synapses) that are rapidly released and another publication showed that the immediately releasable pool is made up of only 10-20% of the vesicles located on the AZ (equal to ~1 vesicle on hippocampal synapses) [15,16]. The ensemble of proximal vesicles that are not in the RRP have been termed non-RRP and presumably belong to the recycling pool that releases more slowly [12,17]. Farther away from the AZ, partially intermixed with the recycling pool, is the reserve pool containing vesicles that only release upon high frequency stimulation. Vesicles in the reserve pool are tightly clustered and well inter-connected by structures that were termed connectors [12,17]. It should be noted that the molecular nature of connectors is not known and is possibly heterogenous. Synapsin has been proposed as a molecular constituent but since the deletion of all forms of synapsin does not lead to the complete absence of connectors, it is clear that not all connectors contain synapsin [18,19]. The second row of SVs near the active zone, immediately after the proximal vesicles, is called the intermediate region. Resting state intermediate SVs are less densely packed and also less connected than proximal SVs [13]. This suggests that, after exocytosis of RRP SVs, intermediate SVs could be rapidly recruited in the RRP by diffusion [20]. Synaptic activity enhances the mobility of a fraction of SVs, whereas it induces synapsin dissociation from SVs in a synapsin phosphorylation-dependent manner [21,22]. The same mobility enhancement can be achieved through inhibition of synapsin dephosphorylation, which leads to synapsin dissociation from SVs, or by knocking out all three synapsin forms [23,24,25]. Interestingly, ribbon synapses do not express synapsin and show higher SV mobility than conventional synapses [26]. It is therefore conceivable that inter-SV connectors restrain SV diffusion and that synaptic activity influences the level of inter-SV connectivity and thereby their mobility.

To investigate this hypothesis and to better understand the impact of depolarization and synaptic activity on SV tethering, we designed two sets of cyro-ET experiments. On the one hand, we compared the morphology of wild-type rat synaptosome in resting state and a few milliseconds after depolarization. On the other hand, we imaged autapses in mice neuronal culture expressing either wild-type SNAP-25, a more positively charged SNAP-25 mutant, or a more negatively charged mutant of SNAP-25. The more positive SNAP-25 mutant, which is consitutively active and whose RRP is permantly depleted showed no triple-tethered SV, which confirmed the morphological definition of the RRP. Our experiments revealed that immediately after depolarization additional SVs are recruited to the RRP. Shortly after exocytosis the level of inter-SV connectivity was decreased among SVs situated in a 25 to 75-nm distance range from the AZ PM. Altogether, our results indicate that connectors regulate SV mobility and their recruitment at the AZ PM. Say something about the size of the RRP

Do we want to talk about kiss-and-run? I have the feeling that we better not to keep the message nice and clear

This manuscript is a template (aka "rootstock") for <u>Manubot</u>, a tool for writing scholarly manuscripts. Use this template as a starting point for your manuscript.

The rest of this document is a full list of formatting elements/features supported by Manubot. Compare the input (.md files in the /content directory) to the output you see below.

Basic formatting

Bold text

Semi-bold text

Centered text

Right-aligned text

Italic text

Combined italics and bold

Strikethrough

- 1. Ordered list item
- 2. Ordered list item
 - a. Sub-item
 - b. Sub-item
 - i. Sub-sub-item
- 3. Ordered list item
 - a. Sub-item
- List item
- List item
- List item

subscript: H₂O is a liquid

superscript: 2¹⁰ is 1024.

unicode superscripts⁰¹²³⁴⁵⁶⁷⁸⁹

unicode subscripts₀₁₂₃₄₅₆₇₈₉

A long paragraph of text. Lorem ipsum dolor sit amet, consectetur adipiscing elit, sed do eiusmod tempor incididunt ut labore et dolore magna aliqua. Ut enim ad minim veniam, quis nostrud exercitation ullamco laboris nisi ut aliquip ex ea commodo consequat. Duis aute irure dolor in reprehenderit in voluptate velit esse cillum dolore eu fugiat nulla pariatur. Excepteur sint occaecat cupidatat non proident, sunt in culpa qui officia deserunt mollit anim id est laborum.

Putting each sentence on its own line has numerous benefits with regard to <u>editing</u> and <u>version</u> <u>control</u>.

Line break without starting a new paragraph by putting two spaces at end of line.

Document organization

Document section headings:

Heading 1

Heading 2

Heading 3

Heading 4

Heading 5

Heading 6



Horizontal rule:

Heading 1's are recommended to be reserved for the title of the manuscript.

Heading 2's are recommended for broad sections such as Abstract, Methods, Conclusion, etc.

Heading 3's and Heading 4's are recommended for sub-sections.

Links

Bare URL link: https://manubot.org

<u>Long link with lots of words and stuff and junk and bleep and blah and stuff and other stuff and more stuff yeah</u>

Link with text

Link with hover text

Link by reference

Citations

Citation by DOI [27].

Citation by PubMed Central ID [28].

Citation by PubMed ID [29].

Citation by Wikidata ID [30].

Citation by ISBN [31].

Citation by URL [32].

Citation by alias [33].

Multiple citations can be put inside the same set of brackets [27,31,33]. Manubot plugins provide easier, more convenient visualization of and navigation between citations [28,29,33,34].

Citation tags (i.e. aliases) can be defined in their own paragraphs using Markdown's reference link syntax:

Referencing figures, tables, equations

Figure 1

Figure 2

```
Figure 3

Figure 4

Table 1

Equation 1

Equation 2
```

Quotes and code

Quoted text

Quoted block of text

Two roads diverged in a wood, and I—I took the one less traveled by, And that has made all the difference.

Code in the middle of normal text, aka inline code.

Code block with Python syntax highlighting:

```
from manubot.cite.doi import expand_short_doi

def test_expand_short_doi():
    doi = expand_short_doi("10/c3bp")
    # a string too long to fit within page:
    assert doi == "10.25313/2524-2695-2018-3-vliyanie-enhansera-copia-i-
        insulyatora-gypsy-na-sintez-ernk-modifikatsii-hromatina-i-
        svyazyvanie-insulyatornyh-belkov-vtransfetsirovannyh-geneticheskih-
        konstruktsiyah"
```

Code block with no syntax highlighting:

```
Exporting HTML manuscript
Exporting DOCX manuscript
Exporting PDF manuscript
```

Figures



Figure 1: A square image at actual size and with a bottom caption. Loaded from the latest version of image on GitHub.



Figure 2: An image too wide to fit within page at full size. Loaded from a specific (hashed) version of the image on GitHub.



Figure 3: A tall image with a specified height. Loaded from a specific (hashed) version of the image on GitHub.



Figure 4: A vector .svg image loaded from GitHub. The parameter sanitize=true is necessary to properly load SVGs hosted via GitHub URLs. White background specified to serve as a backdrop for transparent sections of the image.

Tables

Table 1: A table with a top caption and specified relative column widths.

Bowling Scores	Jane	John	Alice	Bob
Game 1	150	187	210	105
Game 2	98	202	197	102
Game 3	123	180	238	134

Table 2: A table too wide to fit within page.

	Digits 1-33	Digits 34-66	Digits 67-99	Ref.
pi	3.14159265358979323 846264338327950	28841971693993751 0582097494459230	78164062862089986 2803482534211706	piday.org
е	2.71828182845904523 536028747135266	24977572470936999 5957496696762772	40766303535475945 7138217852516642	nasa.gov

 Table 3: A table with merged cells using the attributes plugin.

	Colors		
Size	Text Color	Background Color	
big	blue	orange	
small	black	white	

Equations

A LaTeX equation:

$$\int_0^\infty e^{-x^2} dx = \frac{\sqrt{\pi}}{2} \tag{1}$$

An equation too long to fit within page:

$$x = a + b + c + d + e + f + g + h + i + j + k + l + m + n + o + p + q + r + s + t + u + v + w + x + y + z + 1 + 2 + 3 + 4 + 5 + 6 + 7 + 8 + 9$$
(2)

Special

▲ WARNING The following features are only supported and intended for .html and .pdf exports. Journals are not likely to support them, and they may not display correctly when converted to other formats such as .docx.

LINK STYLED AS A BUTTON

Adding arbitrary HTML attributes to an element using Pandoc's attribute syntax:

Manubot Manubot Manubot Manubot Manubot. Manubot Manubot Manubot Manubot. Manubot Manubot Manubot. Manubot Manubot. Manubot.

Adding arbitrary HTML attributes to an element with the Manubot attributes plugin (more flexible than Pandoc's method in terms of which elements you can add attributes to):

Manubot Manubo

Available background colors for text, images, code, banners, etc:

white lightgrey grey darkgrey black lightred lightyellow lightgreen lightblue lightpurple red orange yellow green blue purple

Using the **Font Awesome** icon set:



Light Grey Banner
useful for general information - manubot.org

1 Blue Banner

useful for important information - manubot.org

♦ Light Red Banner useful for *warnings* - <u>manubot.org</u>

References

1. Vesicle Docking in Regulated Exocytosis

Matthijs Verhage, Jakob B Sørensen

Traffic (2008-09) https://doi.org/bjtx2n

DOI: <u>10.1111/j.1600-0854.2008.00759.x</u> · PMID: <u>18445120</u>

2. Neurotransmitter Release: The Last Millisecond in the Life of a Synaptic Vesicle

Thomas C Südhof

Neuron (2013-10) https://doi.org/f5gng4

DOI: <u>10.1016/j.neuron.2013.10.022</u> · PMID: <u>24183019</u> · PMCID: <u>PMC3866025</u>

3. The readily releasable pool of synaptic vesicles

Pascal S Kaeser, Wade G Regehr

Current Opinion in Neurobiology (2017-04) https://doi.org/gbkfsd

DOI: 10.1016/j.conb.2016.12.012 · PMID: 28103533 · PMCID: PMC5447466

4. The Morphological and Molecular Nature of Synaptic Vesicle Priming at Presynaptic Active Zones

Cordelia Imig, Sang-Won Min, Stefanie Krinner, Marife Arancillo, Christian Rosenmund, Thomas C Südhof, JeongSeop Rhee, Nils Brose, Benjamin H Cooper

Neuron (2014-10) https://doi.org/gcvj2v

DOI: 10.1016/j.neuron.2014.10.009 · PMID: 25374362

5. The Synaptic Vesicle Release Machinery

Josep Rizo, Junjie Xu

Annual Review of Biophysics (2015-06-22) https://doi.org/ginb9q

DOI: 10.1146/annurev-biophys-060414-034057 · PMID: 26098518

6. Crystal structure of a SNARE complex involved in synaptic exocytosis at 2.4 Å resolution

RBryan Sutton, Dirk Fasshauer, Reinhard Jahn, Axel T Brunger

Nature (1998-09) https://doi.org/cwkm8k

DOI: 10.1038/26412 · PMID: 9759724

7. Single Vesicle Millisecond Fusion Kinetics Reveals Number of SNARE Complexes Optimal for Fast SNARE-mediated Membrane Fusion

Marta K Domanska, Volker Kiessling, Alexander Stein, Dirk Fasshauer, Lukas K Tamm

Journal of Biological Chemistry (2009-11) https://doi.org/chdxqp

DOI: 10.1074/jbc.m109.047381 · PMID: 19759010 · PMCID: PMC2797286

8. Fast Vesicle Fusion in Living Cells Requires at Least Three SNARE Complexes

R Mohrmann, H de Wit, M Verhage, E Neher, JB Sorensen

Science (2010-09-16) https://doi.org/c7q87p

DOI: <u>10.1126/science.1193134</u> · PMID: <u>20847232</u>

9. SNARE Proteins: One to Fuse and Three to Keep the Nascent Fusion Pore Open

L Shi, Q-T Shen, A Kiel, J Wang, H-W Wang, TJ Melia, JE Rothman, F Pincet

Science (2012-03-15) https://doi.org/gk8gpf

DOI: <u>10.1126/science.1214984</u> · PMID: <u>22422984</u> · PMCID: <u>PMC3736847</u>

10. Membrane Curvature in Synaptic Vesicle Fusion and Beyond

Harvey T McMahon, Michael M Kozlov, Sascha Martens

Cell (2010-03) https://doi.org/d3gmvv

DOI: 10.1016/j.cell.2010.02.017 · PMID: 20211126

11. An Electrostatic Energy Barrier for SNARE-Dependent Spontaneous and Evoked Synaptic Transmission

Marvin Ruiter, Anna Kádková, Andrea Scheutzow, Jörg Malsam, Thomas H Söllner, Jakob B Sørensen

Cell Reports (2019-02) https://doi.org/gfv5gd

DOI: <u>10.1016/j.celrep.2019.01.103</u> · PMID: <u>30811985</u>

12. Quantitative analysis of the native presynaptic cytomatrix by cryoelectron tomography

Rubén Fernández-Busnadiego, Benoît Zuber, Ulrike Elisabeth Maurer, Marek Cyrklaff, Wolfgang Baumeister, Vladan Lučić

Journal of Cell Biology (2010-01-11) https://doi.org/b9c26b

DOI: 10.1083/jcb.200908082 · PMID: 20065095 · PMCID: PMC2812849

13. Molecular architecture of the presynaptic terminal

Benoît Zuber, Vladan Lučić

Current Opinion in Structural Biology (2019-02) https://doi.org/gk8gpd

DOI: 10.1016/j.sbi.2019.01.008 · PMID: 30925443

14. Properties of Synaptic Vesicle Pools in Mature Central Nerve Terminals

Anthony C Ashton, Yuri A Ushkaryov

Journal of Biological Chemistry (2005-11) https://doi.org/cb5883

DOI: 10.1074/jbc.m504137200 · PMID: 16148008

15. Vesicle release probability and pre-primed pool at glutamatergic synapses in area CA1 of the rat neonatal hippocampus

Eric Hanse, Bengt Gustafsson

The Journal of Physiology (2001-03) https://doi.org/fq7th3

DOI: 10.1111/j.1469-7793.2001.0481i.x · PMID: 11230520 · PMCID: PMC2278469

16. Reluctant Vesicles Contribute to the Total Readily Releasable Pool in Glutamatergic Hippocampal Neurons

KL Moulder

Journal of Neuroscience (2005-04-13) https://doi.org/dwqpxi

DOI: 10.1523/jneurosci.5231-04.2005 · PMID: 15829636 · PMCID: PMC6724923

17. Synaptic vesicle pools: an update

Denker

Frontiers in Synaptic Neuroscience (2010) https://doi.org/d74dd7

DOI: 10.3389/fnsyn.2010.00135 · PMID: 21423521 · PMCID: PMC3059705

18. The cytoskeletal architecture of the presynaptic terminal and molecular structure of synapsin 1.

N Hirokawa, K Sobue, K Kanda, A Harada, H Yorifuji

Journal of Cell Biology (1989-01-01) https://doi.org/ddbqhb

DOI: <u>10.1083/jcb.108.1.111</u> · PMID: <u>2536030</u> · PMCID: <u>PMC2115350</u>

19. Three-Dimensional Architecture of Presynaptic Terminal Cytomatrix

L Siksou, P Rostaing, J-P Lechaire, T Boudier, T Ohtsuka, A Fejtova, H-T Kao, P Greengard, ED Gundelfinger, A Triller, S Marty

Journal of Neuroscience (2007-06-27) https://doi.org/bjw3mv

DOI: 10.1523/jneurosci.1773-07.2007 · PMID: 17596435 · PMCID: PMC6672225

20. Physical determinants of vesicle mobility and supply at a central synapse

Jason Seth Rothman, Laszlo Kocsis, Etienne Herzog, Zoltan Nusser, Robin Angus Silver *eLife* (2016-08-19) https://doi.org/f9rmkt

DOI: 10.7554/elife.15133 · PMID: 27542193 · PMCID: PMC5025287

21. Activity-Dependence of Synaptic Vesicle Dynamics

Luca A Forte, Michael W Gramlich, Vitaly A Klyachko

The Journal of Neuroscience (2017-11-01) https://doi.org/gch7j3

DOI: 10.1523/jneurosci.0383-17.2017 · PMID: 28954868 · PMCID: PMC5666583

22. Synapsin dispersion and reclustering during synaptic activity

Ping Chi, Paul Greengard, Timothy A Ryan

Nature Neuroscience (2001-10-29) https://doi.org/cpdwc7

DOI: <u>10.1038/nn756</u> · PMID: <u>11685225</u>

23. Interactions of synapsin I with small synaptic vesicles: distinct sites in synapsin I bind to vesicle phospholipids and vesicle proteins.

F Benfenati, M Bähler, R Jahn, P Greengard

Journal of Cell Biology (1989-05-01) https://doi.org/d65t4v

DOI: <u>10.1083/jcb.108.5.1863</u> · PMID: <u>2497106</u> · PMCID: <u>PMC2115532</u>

24. Visualization of Synaptic Vesicle Movement in Intact Synaptic Boutons Using Fluorescence Fluctuation Spectroscopy

Randolf Jordan, Edward A Lemke, Jurgen Klingauf

Biophysical Journal (2005-09) https://doi.org/cdpkkv

DOI: <u>10.1529/biophysj.105.061663</u> · PMID: <u>15980175</u> · PMCID: <u>PMC1366711</u>

25. Synapsin Selectively Controls the Mobility of Resting Pool Vesicles at Hippocampal Terminals

A Orenbuch, L Shalev, V Marra, I Sinai, Y Lavy, J Kahn, JJ Burden, K Staras, D Gitler

Journal of Neuroscience (2012-03-21) https://doi.org/f3wq83

DOI: 10.1523/jneurosci.5058-11.2012 · PMID: 22442064 · PMCID: PMC3492757

26. High Mobility of Vesicles Supports Continuous Exocytosis at a Ribbon Synapse

Matthew Holt, Anne Cooke, Andreas Neef, Leon Lagnado

Current Biology (2004-02) https://doi.org/b9g9h2

DOI: 10.1016/j.cub.2003.12.053 · PMID: 14761649

27. Sci-Hub provides access to nearly all scholarly literature

Daniel S Himmelstein, Ariel Rodriguez Romero, Jacob G Levernier, Thomas Anthony Munro, Stephen Reid McLaughlin, Bastian Greshake Tzovaras, Casey S Greene

eLife (2018-03-01) https://doi.org/ckcj

DOI: 10.7554/elife.32822 · PMID: 29424689 · PMCID: PMC5832410

28. Reproducibility of computational workflows is automated using continuous analysis

Brett K Beaulieu-Jones, Casey S Greene

Nature biotechnology (2017-04) https://www.ncbi.nlm.nih.gov/pmc/articles/PMC6103790/

DOI: <u>10.1038/nbt.3780</u> · PMID: <u>28288103</u> · PMCID: <u>PMC6103790</u>

29. **Bitcoin for the biological literature.**

Douglas Heaven

Nature (2019-02) https://www.ncbi.nlm.nih.gov/pubmed/30718888

DOI: <u>10.1038/d41586-019-00447-9</u> · PMID: <u>30718888</u>

30. Plan S: Accelerating the transition to full and immediate Open Access to scientific publications

cOAlition S

(2018-09-04) https://www.wikidata.org/wiki/Q56458321

31. **Open access**

Peter Suber MIT Press (2012)

ISBN: 9780262517638

32. Open collaborative writing with Manubot

Daniel S Himmelstein, Vincent Rubinetti, David R Slochower, Dongbo Hu, Venkat S Malladi, Casey S Greene, Anthony Gitter

Manubot (2020-05-25) https://greenelab.github.io/meta-review/

33. Opportunities and obstacles for deep learning in biology and medicine

Travers Ching, Daniel S Himmelstein, Brett K Beaulieu-Jones, Alexandr A Kalinin, Brian T Do, Gregory P Way, Enrico Ferrero, Paul-Michael Agapow, Michael Zietz, Michael M Hoffman, ... Casey S Greene

Journal of The Royal Society Interface (2018-04-04) https://doi.org/gddkhn
DOI: 10.1098/rsif.2017.0387 · PMID: 29618526 · PMCID: PMC5938574

34. Open collaborative writing with Manubot

Daniel S Himmelstein, Vincent Rubinetti, David R Slochower, Dongbo Hu, Venkat S Malladi, Casey S Greene, Anthony Gitter

PLOS Computational Biology (2019-06-24) https://doi.org/c7np

DOI: <u>10.1371/journal.pcbi.1007128</u> · PMID: <u>31233491</u> · PMCID: <u>PMC6611653</u>