

Spatial Dynamics of Soil Salinity across Khulna District, Bangladesh
(a) Period 1: Post-Monsoon Flushing (November)
(b) Period 2: Mid-Dry Season / Boro Stage (February)
(c) Period 3: Peak Dry Season / Marine Surge (April)
N
Phultala 0.5
Terokhada 0.4
Dighalia 0.6
Khulna Sadar 1.0
Dumuria 1.4
Batiaghata 2.0
Paikgachha 3.2
Dacope 4.1
Koyra 5.2
Phultala 0.9
Terokhada 0.8
Dighalia 1.0
Khulna Sadar 1.8
Dumuria 2.6
Batiaghata 3.6
Paikgachha 5.8
Dacope 7.4
Koyra 9.8
Phultala 1.6
Terokhada 1.4
Dighalia 1.8
Khulna Sadar 3.2
Dumuria 4.5
Batiaghata 6.4
Paikgachha 9.6
Dacope 12.8
Koyra 15.6
0 10 20 km
Northing [km]
Northing [metre]
2540 km
2520 km
2500 km
2480 km
2460 km
730 km 740 km 750 km 760 km 770 km 780 km
Easting [km]
0 2 4 6 8 10 12 14 16
Measured Soil Electrical Conductivity, EC (dS/m) — Saturated Paste Extract (ECe)
4.0 dS/m: Critical Boro Yield Reduction Threshold