

Acoustic Geometry and Sound-Ray Dynamics in Relativistic Spherical Accretion

Alexander V. Semyannikov

Independent Researcher

Email: avsemyannikov@gmail.com

Abstract

In analogue gravity, sound waves in a moving fluid travel along the null geodesics of an effective curved spacetime, giving a controlled setting for studying horizon physics in astrophysical accretion. The relativistic acoustic metric for spherical accretion is well known and has been used in stability and quasi-normal-mode studies, but the geometric optics of sound rays in Michel flow has received little systematic attention. The acoustic metric for a perfect fluid accreting onto a Schwarzschild black hole (Michel flow) is constructed, and high-frequency sound rays are treated as its null geodesics. From the linearized continuity–Euler–energy system, the causal surfaces of the geometry are identified—the acoustic horizon (which coincides with the sonic point), the stationary-limit surface, and the ergoregion—and a generalized Binet equation is derived that separates ray deflection into Einstein, refractive, and acoustic-wind parts. A local slowdown factor allows the comparison of acoustic and vacuum Shapiro delays, the location of the acoustic photon sphere, and the relation of the analogue Hawking scale to hydrodynamic gradients. Results for polytropic indices $\gamma = 4/3$ and $5/3$ show how the thermodynamic and flow parameters of the accretion enter acoustic lensing, time delay, and the analogue Hawking temperature, providing a geometric basis for future studies of finite-frequency acoustic, vortical, and entropy modes.

Keywords: acoustic metric, analogue gravity, Michel accretion, Shapiro delay, sound rays, acoustic horizon, photon sphere, Hawking temperature

1. Introduction

Acoustic perturbations in a moving fluid propagate on an effective Lorentzian geometry constructed from the background spacetime metric, the flow 4-velocity, and the local speed of sound [1, 2, 3]. For stationary spherical accretion onto a Schwarzschild black hole (the Michel flow [4]), this construction generates an acoustic horizon coinciding with the critical point of the flow. The relativistic acoustic geometry was systematized by Moncrief [5] and Bilić [6], and placed in the broader analogue-gravity framework by Visser [2] and by Barceló, Liberati and Visser [3].

Within this setting, most work on Michel accretion has concentrated on the wave equation itself: global stability outside the sonic horizon, energy estimates, and the spectrum of quasi-normal acoustic oscillations [5, 7], with the analogue Hawking emission revisited as a tunnelling process [8]. A complementary line of work expresses the acoustic surface gravity directly in terms of the accretion variables for relativistic flows onto a Schwarzschild black hole [9, 10, 11]. Ray-level properties of acoustic metrics—deflection, time delay, and the phonon sphere—have been studied in lower-dimensional and slowly rotating acoustic holes [12, 13], Bose–Einstein condensates [14], and curved-background toy metrics [15], but not for the relativistic adiabatic Michel flow with an explicit decomposition of the deflecting force.

In the vacuum theory, null geodesics underlie gravitational lensing, the photon sphere, and the Shapiro time delay [16]. The same questions arise for sound rays, but the answers depend on the thermodynamic state of the flow—its equation of state, asymptotic sound speed, and sonic-point location—as well as on the background gravity. For relativistic Michel accretion, these ray-level questions—the forces that deflect a sound ray, the acoustic Shapiro delay, the shift of the acoustic photon sphere, and the surface gravity of the sonic horizon—have not been assembled into a single quantitative picture tied to the flow thermodynamics.

In the present work, the geometric-optics phenomenology of sound rays in the relativistic Michel flow is constructed. By decomposing the deflecting forces and analyzing the causal surfaces, the influence of the flow thermodynamics on acoustic lensing, the Shapiro delay, and the analogue Hawking temperature is established. The analysis is carried out for polytropic indices $\gamma = 4/3$ and $5/3$ with the representative asymptotic sound speed $a_\infty = 0.1$, supplemented by a scan over $a_\infty \in [0.05, 0.20]$.

2. Acoustic geometry of a relativistic perfect fluid

Throughout the work, the metric signature $\mathfrak{s} = (-, +, +, +)$ is adopted. Spacetime coordinates are measured in units of the gravitational radius $r_g = 2GM/c^2$, the 4-velocity and the sound speed are dimensionless, and the thermodynamic variables are normalized to their asymptotic rest-frame scales in accordance with the standard Michel-flow convention [4, 5]. The subscripts “ ∞ ”, “c”, “a”, “aph”, “ph”, and “asl” denote the asymptotic value at infinity, the critical point, the acoustic horizon, the acoustic photon sphere, the vacuum Schwarzschild photon sphere, and the stationary-limit surface, respectively.

2.1. Thermodynamics and the acoustic core

The medium is modelled as a perfect fluid in local thermodynamic equilibrium, with the particle number density n and the specific enthalpy h . The thermodynamics is closed by the polytropic equation of state $p = Kn^\gamma$, which yields

$$h = 1 + \frac{\gamma K}{\gamma - 1} n^{\gamma-1}, \quad (1)$$

and the local isentropic sound speed

$$a^2 = \left(\frac{\partial p}{\partial \epsilon} \right)_s = \frac{K\gamma n^{\gamma-1}}{1 + \frac{\gamma K}{\gamma - 1} n^{\gamma-1}}. \quad (2)$$

In the cold, non-relativistic limit $h \rightarrow 1$ one recovers the Newtonian relation $a^2 \approx K\gamma n^{\gamma-1}$. In the ultra-relativistic limit $n \rightarrow \infty$ one finds the universal bound $a^2 \rightarrow \gamma - 1$; causality ($a^2 \leq 1$) therefore requires $\gamma \leq 2$, with the standard hot-plasma value $\gamma = 4/3$ giving $a^2 \rightarrow 1/3$.

The conformal factor of the acoustic metric, expressed through the local sound speed alone, is

$$\Omega^2 = \frac{n}{h} = C a^{\frac{2}{\gamma-1}} \left(\frac{\gamma - 1}{\gamma - 1 - a^2} \right)^{\frac{2-\gamma}{\gamma-1}}, \quad (3)$$

where $C = (K\gamma)^{-1/(\gamma-1)}$ is fixed by the entropy of the flow. The global constant C cancels from the null condition and therefore does not affect the kinematics of sound rays.

The analysis relies on the following assumptions: (i) stationarity and spherical symmetry of the background; (ii) a perfect fluid model with a polytropic equation of state; (iii) the absence

of rotation and self-gravity of the accreting matter; (iv) the geometric optics approximation for linear perturbations. Under these conditions, linear acoustic perturbations propagate on an effective Lorentzian geometry [6, 2, 3]. The contravariant and covariant cores of the acoustic metric are

$$\mathcal{G}^{\mu\nu} = \Omega^2 \eta^{\mu\nu}, \quad \eta^{\mu\nu} = g^{\mu\nu} + \alpha u^\mu u^\nu, \quad \alpha = 1 - a^{-2}, \quad (4)$$

$$\mathcal{G}_{\mu\nu} = \Omega^{-2} \eta_{\mu\nu}, \quad \eta_{\mu\nu} = g_{\mu\nu} - \beta u_\mu u_\nu, \quad \beta = a^2 - 1. \quad (5)$$

High-frequency sound rays are the null geodesics of $\mathcal{G}_{\mu\nu}$.

3. Acoustic geometry of Michel accretion

The acoustic metric of Sec. 2 is specialized to steady, spherically symmetric, adiabatic accretion onto a Schwarzschild black hole—the Michel flow [4].

3.1. Background Schwarzschild spacetime

The background line element is

$$ds^2 = -f dt^2 + f^{-1} dr^2 + r^2 (d\theta^2 + \sin^2 \theta d\phi^2), \quad (6)$$

with the dimensionless lapse

$$f = 1 - r^{-1}. \quad (7)$$

In matrix form,

$$g^{\mu\nu} = \begin{pmatrix} -f^{-1} & 0 & 0 & 0 \\ 0 & f & 0 & 0 \\ 0 & 0 & r^{-2} & 0 \\ 0 & 0 & 0 & r^{-2} \sin^{-2} \theta \end{pmatrix}, \quad g_{\mu\nu} = \begin{pmatrix} -f & 0 & 0 & 0 \\ 0 & f^{-1} & 0 & 0 \\ 0 & 0 & r^2 & 0 \\ 0 & 0 & 0 & r^2 \sin^2 \theta \end{pmatrix}. \quad (8)$$

Stationarity and spherical symmetry imply that every background scalar depends only on r , while the angular components of the 4-velocity vanish. With the 4-velocity written as

$$u^\mu = (\vartheta, v, 0, 0), \quad (9)$$

where ϑ is the temporal component and v is the radial 3-velocity, the covariant components take the form $u_\mu = (-f\vartheta, f^{-1}v, 0, 0)$, and the normalization becomes

$$-u_t = f\vartheta = \sqrt{f + v^2}, \quad (10)$$

which relates the time component to the radial 3-velocity of the flow.

The Michel background itself is fixed by two integrals of motion. Continuity yields the baryon flux $r^2 n v = \text{const}$ ($v < 0$ for accretion), while the adiabatic projection of $\nabla_\mu T^\mu_\nu = 0$ integrates to the relativistic Bernoulli integral

$$h\sqrt{f + v^2} = h_\infty. \quad (11)$$

Together with the polytropic closure of Sec. 2, these relations determine the profiles $v(r)$, $a(r)$ and $n(r)$ once the asymptotic sound speed a_∞ and the adiabatic index γ are specified [4, 17]. In the numerical implementation, the Michel integrals were integrated inward from a large radius, which removes branch ambiguity at the sonic point.

3.2. Acoustic metric tensor

Substituting (8) and (9) into the covariant core (5) produces

$$\eta_{\mu\nu} = \begin{pmatrix} -f(1 + \beta f \vartheta^2) & \beta \vartheta v & 0 & 0 \\ \beta \vartheta v & f^{-1}(1 - \beta f^{-1} v^2) & 0 & 0 \\ 0 & 0 & r^2 & 0 \\ 0 & 0 & 0 & r^2 \sin^2 \theta \end{pmatrix}. \quad (12)$$

The full covariant acoustic metric is $\mathcal{G}_{\mu\nu} = \Omega^{-2} \eta_{\mu\nu}$. The corresponding contravariant core is

$$\eta^{\mu\nu} = \begin{pmatrix} -f^{-1} + \alpha \vartheta^2 & \alpha \vartheta v & 0 & 0 \\ \alpha \vartheta v & f + \alpha v^2 & 0 & 0 \\ 0 & 0 & r^{-2} & 0 \\ 0 & 0 & 0 & r^{-2} \sin^{-2} \theta \end{pmatrix}. \quad (13)$$

The components η^{rr} and η^{tr} from (13) control the radial characteristics and the acoustic frame dragging. Sound rays are null geodesics of $\mathcal{G}_{\mu\nu}$ and are therefore invariant under conformal rescalings,

$$\mathcal{G}_{\mu\nu} k^\mu k^\nu = 0 \implies \eta_{\mu\nu} k^\mu k^\nu = 0. \quad (14)$$

Consequently, the acoustic light-cone geometry—and therefore all ray paths—is fixed entirely by the core $\eta_{\mu\nu}$. The conformal factor Ω^2 re-enters only in the wave amplitude and in the formal horizon area, not in the ray paths or the surface gravity.

3.3. Acoustic horizon, stationary limit, and ergoregion

The acoustic event horizon is the one-way membrane for sound. It forms where the flow becomes sonic or supersonic, which is equivalent to the vanishing of the radial core component:

$$\eta^{rr}(r_a) = 0 \implies r_a = [1 + \alpha(r_a) v^2(r_a)]^{-1}. \quad (15)$$

For $r < r_a$ every characteristic of the wave equation points inward: an upstream acoustic signal is either swept toward the black hole ($|v| > a$) or frozen into the fluid ($|v| = a$).

The acoustic stationary-limit surface is defined by the vanishing of the time–time core component,

$$\eta^{tt}(r_{\text{asl}}) = 0 \implies r_{\text{asl}} = \frac{\alpha(r_{\text{asl}}) \vartheta^2(r_{\text{asl}})}{\alpha(r_{\text{asl}}) \vartheta^2(r_{\text{asl}}) - 1}. \quad (16)$$

Inside $r < r_{\text{asl}}$ no stationary acoustic mode can exist: a static observer would have to move against the flow at a supersonic speed. Equating (15) with (16) would require $a^2 < 0$, which is impossible for a stable medium. Consequently the two surfaces never coincide, and the finite shell between them constitutes the acoustic ergoregion [2, 6].

A central structural fact of Michel accretion is that the acoustic horizon coincides exactly with the sonic critical point of the background flow. Combining the continuity and Bernoulli integrals yields an equation for the radial gradient dv/dr of the form N/D ; smooth transonic passage through the critical point requires the numerator and denominator to vanish simultaneously [17], giving the standard critical-point conditions

$$v_c^2 = \frac{1}{4r_c}, \quad a_c^2 = \frac{1}{4r_c - 3} = \frac{v_c^2}{1 - 3v_c^2}. \quad (17)$$

Substituting these values into the radial core component (13) yields

$$\eta^{rr}(r_c) = f_c - f_c = 0. \quad (18)$$

Hence, the surface at which the accretion stream becomes sonic is exactly the acoustic event horizon. The shell $1 < r < r_a$ is supersonic ($\eta^{rr} < 0$) and terminates on the gravitational horizon $r = 1$.

The Bernoulli integral (11), evaluated at the sonic point, links a_∞ to the horizon radius r_a ; representative values are collected in Table 1. For cold asymptotic gas ($a_\infty \ll 1$) the acoustic horizon lies far from the black hole, in the Newtonian domain ($r_a \sim a_\infty^{-2} \gg 1$), while relativistic effects concentrate in the deeply supersonic zone $1 < r \ll r_a$. These radii satisfy the standard Michel critical-point condition [4, 17], while the identification of the acoustic horizon with the critical point follows from the acoustic metric [5, 6]. The quantitative ray analysis below focuses on the representative case $a_\infty = 0.1$ with $\gamma = 4/3$ and $5/3$.

Table 1: Michel sonic point (which coincides with the acoustic horizon), $r_a = r_c$, and the local sound speed a_c there, for selected polytropic indices γ and asymptotic sound speeds a_∞ .

γ	a_∞	$r_a = r_c$	a_c
4/3	0.01	1252	0.014
4/3	0.03	141	0.042
4/3	0.10	14.1	0.137
5/3	0.01	38.1	0.082
5/3	0.03	13.1	0.142
5/3	0.10	4.38	0.262

4. Sound-ray dynamics

In the high-frequency limit, acoustic disturbances propagate along the null geodesics of $\mathcal{G}_{\mu\nu}$, the acoustic counterpart of light rays in vacuum [2, 3]. While ray bending and photon spheres have been studied in several analogue settings, relativistic Michel accretion with an explicit decomposition of the deflecting force has not been addressed. The corresponding geometric acoustics is developed here.

4.1. Null geodesics and conserved quantities

Because the Michel background is stationary and spherically symmetric, the two Killing vectors ∂_t and ∂_ϕ endow every sound ray with a conserved energy and angular momentum, reducing its propagation to an effective radial problem. The rays themselves are null geodesics of the acoustic metric,

$$\frac{d^2 x^\mu}{d\lambda^2} + \Gamma_{\alpha\beta}^\mu \frac{dx^\alpha}{d\lambda} \frac{dx^\beta}{d\lambda} = 0, \quad (19)$$

with wave vector $k^\mu = dx^\mu/d\lambda$. Stationarity of $\mathcal{G}_{\mu\nu}$ supplies a Killing conserved energy

$$k_t = \mathcal{G}_{t\mu} k^\mu = -E = \text{const}, \quad (20)$$

the minus sign following from the adopted metric signature and the requirement that the physical energy be positive. The non-vanishing off-diagonal component $\mathcal{G}_{rt} \neq 0$ yields the temporal derivative

$$\frac{dt}{d\lambda} = -\frac{1}{\mathcal{G}_{tt}} \left(E + \mathcal{G}_{rt} \frac{dr}{d\lambda} \right), \quad (21)$$

which locks the coordinate time to the radial motion: the ingoing flow drags the ray forward in t , providing an acoustic analogue of frame dragging.

The second Killing vector, ∂_ϕ , likewise conserves the angular momentum. Restricting without loss of generality to the equatorial plane $\theta = \pi/2$ (which is preserved by the θ -component of (19)), one finds

$$k_\phi = \mathcal{G}_{\phi\phi} \frac{d\phi}{d\lambda} = L = \text{const.} \quad (22)$$

Their ratio $b = L/E$ is the impact parameter, which labels each ray just as in vacuum lensing.

With E and L fixed, the null condition (14) collapses to a single equation for the radial motion,

$$\mathcal{G}_{tt} \left(\frac{dt}{d\lambda} \right)^2 + 2\mathcal{G}_{tr} \frac{dt}{d\lambda} \frac{dr}{d\lambda} + \mathcal{G}_{rr} \left(\frac{dr}{d\lambda} \right)^2 + \frac{L^2}{\mathcal{G}_{\phi\phi}} = 0. \quad (23)$$

This is the master equation for the trajectories analysed below. For radial rays ($L = 0$), the null condition yields a quadratic equation for the coordinate speed $v = dr/dt$. Elimination of the conformal factor leads to the characteristic velocities

$$v_\pm = \frac{-\beta v \pm a}{f - \beta v^2} f^2. \quad (24)$$

The branch v_+ is the outgoing ray (fighting the accretion wind); v_- is the ingoing ray. In the vacuum limit $a \rightarrow 1$, $\beta \rightarrow 0$ one recovers $v_\pm = \pm f$. The acoustic horizon is precisely the freeze-out surface, equivalent to (15). The non-radial rays that build up the lensing pattern and the acoustic photon sphere are governed by the Binet equation.

4.2. Generalized Binet equation

The general, non-radial rays are most naturally described as orbits in the angle ϕ . Setting $\varrho = 1/r$ and eliminating the affine parameter between the null condition and the conserved $b = L/E$ yields the first Binet integral

$$\left(\frac{d\varrho}{d\phi} \right)^2 = \frac{1}{b^2 a^2} - \varrho^2 \eta^{rr}. \quad (25)$$

The conformal factor again drops out; differentiating with respect to ϕ produces the second-order Binet equation

$$\frac{d^2 \varrho}{d\phi^2} + \varrho = \underbrace{\frac{3}{2} \varrho^2}_{F_E} + \underbrace{\frac{1}{2b^2} \frac{d}{d\varrho} \left(\frac{1}{a^2} \right)}_{F_R} - \underbrace{\frac{1}{2} \frac{d}{d\varrho} (\alpha v^2 \varrho^2)}_{F_W}. \quad (26)$$

In vacuum ($a \rightarrow 1$) this reduces to the standard Schwarzschild relation $d^2 \varrho / d\phi^2 + \varrho = \frac{3}{2} \varrho^2$. In general the right-hand side splits into three competing forces:

- F_E : the Einstein (vacuum curvature) deflection;
- F_R : acoustic refraction from gradients of $a(r)$; for a flow that heats toward the centre this contribution focuses rays inward;
- F_W : the acoustic-wind force from radial accretion; for subsonic flow $\alpha < 0$, so this term often opposes gravitational focusing and shifts the capture radius relative to vacuum.

This decomposition makes explicit which effect bends a given ray.

The critical impact parameter b_c is fixed by the maximum of the capture barrier

$$U(r) = a^2 \eta^{rr} / r^2, \quad (27)$$

via the turning condition. Numerical integration of (26) for the Michel flow at $a_\infty = 0.1$ revealed that rays with $b < b_c$ spiral onto the acoustic horizon, while rays with $b > b_c$ scatter past the acoustic photon sphere (Fig. 1). Along the critical ray, the three forces compete: the wind term dominates near the horizon, refraction is repulsive throughout, and the Einstein term falls off as r^{-2} . Their sum is positive at the acoustic horizon and changes sign at a larger radius (Fig. 2, Table 2). For $\gamma = 5/3$ one finds $r_a \approx 4.38$, $r_{\text{aph}} \approx 6.10$ and $b_c \approx 50$; for the colder $\gamma = 4/3$ flow the scales inflate to $r_a \approx 14.1$, $r_{\text{aph}} \approx 20.0$ and $b_c \approx 240$.




Figure 1: Sound-ray trajectories in Michel flow at $a_\infty = 0.1$, for a monatomic gas ($\gamma = 5/3$, left) and a radiation-dominated plasma ($\gamma = 4/3$, right); note the different spatial scales. Captured rays ($b < b_c$, warm colours from red to yellow) spiral onto the acoustic horizon r_a (crimson dashed circle), while scattered rays ($b > b_c$, cold colours from blue to cyan) are deflected around the acoustic photon sphere r_{aph} (orange dash-dotted circle); the black disc marks the gravitational horizon $r = 1$. Because the colder $\gamma = 4/3$ flow turns sonic farther out, it inflates every acoustic scale by a factor of a few ($r_a \approx 14.1$, $r_{\text{aph}} \approx 20.0$, $b_c \approx 240$, against $r_a \approx 4.38$, $r_{\text{aph}} \approx 6.10$, $b_c \approx 50$ for $\gamma = 5/3$), so the two panels are geometrically similar but rescaled by the equation of state. Characteristic radii and the critical impact parameter are collected in Table 2.

Table 2: Forces F_E , F_R , F_W from (26) and their sum $\Sigma = F_E + F_R - F_W$, evaluated along the critical ray $b = b_c$ at the acoustic horizon, the acoustic photon sphere and the zero-crossing r_0 (where $\Sigma = 0$); the last column gives $1/b_c^2$. $a_\infty = 0.1$.

γ	point	r	F_E	F_R	F_W	Σ	$1/b_c^2$
5/3	r_a	4.38	$+7.81 \cdot 10^{-2}$	$-8.95 \cdot 10^{-3}$	$-2.55 \cdot 10^{-1}$	$+3.24 \cdot 10^{-1}$	$4.01 \cdot 10^{-4}$
	r_{aph}	6.10	$+4.04 \cdot 10^{-2}$	$-1.54 \cdot 10^{-2}$	$-1.39 \cdot 10^{-1}$	$+1.64 \cdot 10^{-1}$	
	r_0	12.16	$+1.01 \cdot 10^{-2}$	$-4.49 \cdot 10^{-2}$	$-3.47 \cdot 10^{-2}$	0	
4/3	r_a	14.11	$+7.53 \cdot 10^{-3}$	$-2.23 \cdot 10^{-3}$	$-1.15 \cdot 10^{-1}$	$+1.20 \cdot 10^{-1}$	$1.73 \cdot 10^{-5}$
	r_{aph}	19.98	$+3.76 \cdot 10^{-3}$	$-3.26 \cdot 10^{-3}$	$-4.96 \cdot 10^{-2}$	$+5.01 \cdot 10^{-2}$	
	r_0	43.41	$+7.96 \cdot 10^{-4}$	$-6.45 \cdot 10^{-3}$	$-5.65 \cdot 10^{-3}$	0	




Figure 2: Force decomposition (26) along the critical ray $b = b_c$ ($a_\infty = 0.1$) for $\gamma = 4/3$ (top) and $\gamma = 5/3$ (bottom): F_E (cyan circles), F_R (red squares), $-F_W$ (teal diamonds) and their sum $\Sigma = F_E + F_R - F_W$ (solid black); the horizontal dashed line marks $\Sigma = 0$. Vertical dashed lines mark the acoustic horizon r_a (red) and the acoustic photon sphere r_{aph} (blue). Pointwise values of the three forces and of Σ are listed in Table 2.

4.3. Acoustic Shapiro delay

Beyond deflection, a sound ray also carries a timing signature—the acoustic analogue of the Shapiro delay. The coordinate travel time of a radial ray between r_1 and r_2 is

$$t_{\pm} = \int_{r_1}^{r_2} \frac{1}{v_{\pm}(r)} dr = \int_{r_1}^{r_2} \frac{f - \beta v^2}{f^2(-\beta \vartheta v \pm a)} dr. \quad (28)$$

In vacuum this reduces to the classical Shapiro integral $t_{\pm} = \pm \int f^{-1} dr$ [16]. In the acoustic geometry two further contributions appear: a hydrodynamic (wind) delay from $\mathcal{G}_{rt} \neq 0$, which breaks $t_+ = t_-$, and a refractive delay from the radial profiles $a(r)$ and $\beta(r)$.

The integral (28) alone is dominated by the long homogeneous outer zone; a sharper, local probe is the slowdown factor

$$N_{\pm}(r) = \frac{f(r)}{|v_{\pm}(r)|}, \quad (29)$$

which measures how much slower a sound ray is than a vacuum photon at the same radius (for which $|v| = f$). The profiles of $N_{\pm}$ for the Michel flow at $a_\infty = 0.1$ are presented in Fig. 3 and Table 3. Near the horizon the branches diverge: the outgoing ray freezes ($v_+ \rightarrow 0$, $N_+ \sim 10^3$), while the ingoing ray is swept inward at finite N_- . At large radius both branches converge to the geometric-optics plateau $1/a_\infty = 10$.

For an outgoing signal emitted at r_1 and received at $r_{\text{out}} = 200$, the excess delay relative to flat light travel,

$$\Delta t_+(r_1) = \int_{r_1}^{r_{\text{out}}} \frac{dr}{v_+(r)} - (r_{\text{out}} - r_1), \quad (30)$$




Figure 3: Local slowdown $N_{\pm} = f/|v_{\pm}|$ for outgoing and ingoing rays ($a_{\infty} = 0.1$) at $\gamma = 4/3$ (top) and $\gamma = 5/3$ (bottom): N_+ is the crimson curve with filled circles, N_- the black curve with crosses; the horizontal dashed line marks the geometric-optics plateau $1/a_{\infty} = 10$. Near the horizon ($r - r_a = 10^{-3}r_a$) the outgoing branch freezes ($N_+ \rightarrow \infty$) while the ingoing branch stays finite, and at large radius both converge to 10. The insets zoom into the outer zone on a linear scale, and the numerical values at the labelled radii are listed in Table 3.

exceeds its vacuum counterpart by factors $R = \Delta t_+^{\text{ac}}/\Delta t_+^{\text{vac}} \sim 10^2\text{--}10^3$, while the ratio of total flight times $\mathcal{T} = t_+^{\text{ac}}/t_+^{\text{vac}}$ stays close to $1/a_{\infty}$. The geometric-optics piece $\mathcal{T} \approx 10$ is essentially independent of γ ; the hydrodynamic freeze-out, visible in N_+/N_- and in R , is stronger for $\gamma = 4/3$, whose sonic point lies farther out. Both deflection and timing feed the acoustic photon sphere.

4.4. Acoustic lensing and the photon sphere

The scattered rays organize around a single unstable circular orbit—the acoustic photon sphere—which fixes the edge of the acoustic shadow. In vacuum Schwarzschild spacetime this orbit sits at $r_{\text{ph}} = 3/2$; in the acoustic geometry the circular-orbit conditions $d\varrho/d\phi = d^2\varrho/d\phi^2 = 0$ applied to (25)–(26) move it to r_{aph} . Linearizing about the vacuum point $\varrho_{\text{ph}} = 2/3$ splits the shift into a refractive and a wind contribution,

$$\delta\varrho = \underbrace{\frac{2}{27} \frac{d\beta}{d\varrho}\bigg|_{\text{ph}}}_{\delta F_{\text{R}}} + \underbrace{\frac{2}{3} \left(\beta_{\text{ph}} v_{\text{ph}}^2 + \frac{1}{3} \frac{d(\beta v^2)}{d\varrho}\bigg|_{\text{ph}} \right)}_{-\delta F_{\text{W}}}, \quad (31)$$

and therefore

$$r_{\text{aph}} \approx \frac{3}{2} \left(1 - \frac{3}{2} \delta\varrho \right). \quad (32)$$

Table 3: Integral Shapiro diagnostics for a signal emitted at r_1 and received at $r_{\text{out}} = 200$: local slowdowns $N_{\pm}$ at r_1 , excess delay Δt_+^{ac} , ratio to vacuum $R = \Delta t_+^{\text{ac}}/\Delta t_+^{\text{vac}}$, and total-time ratio $\mathcal{T}$. $a_{\infty} = 0.1$.

γ	r_1/r_a	r_1	N_+	N_-	Δt_+^{ac}	R	$\mathcal{T}$
4/3	$1+10^{-3}$	14.1	$9.39 \cdot 10^3$	3.73	2813	1035	15.9
	1.5	21.2	27.2	4.60	1875	819	11.3
	4.0	56.4	11.7	7.12	1357	1062	10.4
5/3	$1+10^{-3}$	4.39	$6.39 \cdot 10^3$	2.04	1902	467	10.5
	1.5	6.57	20.3	2.55	1663	465	9.42
	4.0	17.5	9.97	4.34	1526	613	9.24

The two contributions pull in opposite directions: refraction ($d\beta/d\varrho|_{\text{ph}} > 0$ for a flow that heats inward) tends to shrink the acoustic shadow, while the subsonic wind ($\beta_{\text{ph}} < 0$) pushes r_{aph} outward. For Michel flow at $a_{\infty} = 0.1$ the wind wins, giving a substantial outward shift: $r_{\text{aph}} \approx 6.10$ for $\gamma = 5/3$ and $r_{\text{aph}} \approx 20.0$ for $\gamma = 4/3$, against the vacuum value 1.5. The shadow is therefore set by r_{aph} —equivalently by the barrier maximum of (27)—rather than by the horizon itself, and carries a direct imprint of the thermodynamic state of the accretion stream.

Although both r_a and r_{aph} vary strongly with a_{∞} , their ratio stays nearly constant across Table 4 and Fig. 4 (right), $r_{\text{aph}}/r_a \approx 1.39$ for $\gamma = 5/3$ and ≈ 1.42 for $\gamma = 4/3$: the acoustic photon sphere tracks the horizon at a fixed multiple set mainly by the equation of state. That same horizon also radiates, with the surface gravity and the associated analogue Hawking temperature determined by the local hydrodynamic gradients.




Figure 4: Dependence of the acoustic-horizon thermodynamics and lensing geometry on the asymptotic sound speed a_{∞} , for $\gamma = 5/3$ (monatomic gas: red, filled circles) and $\gamma = 4/3$ (radiation-dominated plasma: black, crosses). (left) The surface gravity κ rises monotonically by more than an order of magnitude as a_{∞} grows and the sonic point migrates inward toward the strong-field region; the right axis gives the analogue Hawking temperature relative to vacuum, $2\kappa = T_{\text{H}}^{\text{ac}}/T_{\text{H}}^{\text{vac}}$. (right) The photon-sphere ratio r_{aph}/r_a stays nearly constant over the same range (dotted horizontal lines mark the means, ≈ 1.39 for $\gamma = 5/3$ and ≈ 1.42 for $\gamma = 4/3$), so the acoustic photon sphere tracks the horizon at a fixed multiple set mainly by the equation of state. Tabulated values at $a_{\infty} = 0.05, 0.10, 0.15, 0.20$ are listed in Table 4.

5. Surface gravity and analogue Hawking temperature

Table 4: Dependence on the asymptotic sound speed a_∞ : acoustic horizon r_a , acoustic photon sphere r_{aph} , their ratio, surface gravity κ from (35), and the temperature ratio $2\kappa = T_{\text{H}}^{\text{ac}}/T_{\text{H}}^{\text{vac}}$. As a_∞ grows the sonic point moves inward and the horizon warms by more than an order of magnitude, while r_{aph}/r_a stays nearly constant (set mainly by γ).

γ	a_∞	r_a	r_{aph}	r_{aph}/r_a	κ	2κ
5/3	0.05	8.13	11.29	1.39	$4.10 \cdot 10^{-3}$	$8.20 \cdot 10^{-3}$
	0.10	4.38	6.10	1.39	$1.38 \cdot 10^{-2}$	$2.76 \cdot 10^{-2}$
	0.15	3.14	4.37	1.39	$2.69 \cdot 10^{-2}$	$5.39 \cdot 10^{-2}$
	0.20	2.51	3.52	1.40	$4.19 \cdot 10^{-2}$	$8.39 \cdot 10^{-2}$
4/3	0.05	51.67	73.13	1.42	$4.82 \cdot 10^{-4}$	$9.64 \cdot 10^{-4}$
	0.10	14.11	19.97	1.42	$3.51 \cdot 10^{-3}$	$7.02 \cdot 10^{-3}$
	0.15	7.10	10.06	1.42	$1.03 \cdot 10^{-2}$	$2.06 \cdot 10^{-2}$
	0.20	4.60	6.51	1.42	$2.06 \cdot 10^{-2}$	$4.13 \cdot 10^{-2}$

5.1. Kinematic definition and connection to quasinormal modes

The acoustic horizon, defined by $\eta^{rr}(r_a) = 0$, is the necessary kinematic ingredient for an analogue of Hawking radiation, provided the background remains stationary and adiabatic. In the quantum field theory on the effective spacetime, the horizon leads to spontaneous creation of phonon pairs: one mode escapes to spatial infinity as a thermal flux, while its partner (of negative Killing norm) crosses the horizon and is absorbed.

The rate of exponential peeling of outgoing null rays near the horizon is measured by the surface gravity κ . Starting from the outgoing coordinate speed v_+ defined in (24), which is conformally invariant and vanishes at r_a , one introduces the acoustic tortoise coordinate $dr_* = dr/v_+(r)$. Near the horizon, $v_+(r) \approx (dv_+/dr)|_{r_a}(r - r_a)$, so

$$r_* \approx \frac{1}{(dv_+/dr)|_{r_a}} \ln(r - r_a). \quad (33)$$

Matching this to the canonical form $r_* \sim (2\kappa)^{-1} \ln(r - r_a)$ that underlies the thermal Kruskal spectrum yields the invariant definition

$$\kappa \equiv \frac{1}{2} \frac{dv_+}{dr} \Big|_{r=r_a}. \quad (34)$$

Because the numerator of (24) vanishes on the horizon, differentiation retains only the derivative of that numerator and produces the closed expression

$$\kappa = \frac{f^2(r_a)}{2[f(r_a) - \beta(r_a)v^2(r_a)]} \frac{d}{dr}(a - \beta\vartheta v) \Big|_{r=r_a}. \quad (35)$$

This expression is cross-checked against the direct numerical slope of v_+ ; the two agree to better than one percent. Surface gravity is therefore controlled not by the values of the flow parameters themselves, but by the *rate* at which the stream crosses the sonic point: the steeper the numerator of the freeze-out condition, the hotter the horizon. In the vacuum limit $a \rightarrow 1$, $\beta \rightarrow 0$, $r_a \rightarrow 1$ one recovers $\kappa = \frac{1}{2}(df/dr)|_{r=1} = \frac{1}{2}$, which in dimensionful units is precisely the Schwarzschild value $\kappa_{\text{Sch}} = c^2/(2r_g)$.

The exponential redshift of outgoing modes converts the asymptotic vacuum into a thermal phonon bath at

$$T_{\text{H}} = \frac{\kappa}{2\pi}, \quad (36)$$

or, restoring dimensions (r and t measured in units of r_g and r_g/c),

$$k_B T_H = \frac{\hbar c}{2\pi r_g} \kappa. \quad (37)$$

For $\kappa = 1/2$ this reproduces the classical Hawking temperature $k_B T_H = \hbar c^3/(8\pi G M)$. In the acoustic case T_H carries information about the accretion thermodynamics: it is set by the gradients of $a(r)$ and by the flow profile (ϑ, v) at the sonic point. The spectrum remains thermal in the adiabatic approximation, when background quantities vary slowly on the thermal phonon wavelength $\lambda_T \sim \kappa^{-1}$; sharp gradients produce departures from the Planckian form.

Relative to the vacuum Schwarzschild surface gravity $\kappa_{\text{Sch}} = 1/2$,

$$\frac{T_H^{\text{ac}}}{T_H^{\text{vac}}} = 2\kappa(a, \vartheta, v). \quad (38)$$

A large positive gradient in (35) (rapid drop of the sound speed, or strong acceleration of the flow) enhances the radiation relative to vacuum; smooth profiles can suppress it. Unlike the gravitational horizon, whose temperature is fixed by the black-hole mass alone, the acoustic horizon possesses an effective kinematic thermodynamics controlled by the equation of state and by the asymptotic boundary data. The values of κ , evaluated from (35) for the Michel flow at $a_\infty = 0.1$, are collected in Table 5: the analogue horizon is much colder than its vacuum counterpart, $2\kappa = T_H^{\text{ac}}/T_H^{\text{vac}} \approx 2.8 \cdot 10^{-2}$ for $\gamma = 5/3$ and $\approx 7.0 \cdot 10^{-3}$ for $\gamma = 4/3$. The kinematic definition (34) is standard in analogue gravity [1, 2], and coincides with the conformally invariant surface gravity of Bilić [6] specialized to radial flow. Expressing κ —and hence T_H —directly through the accretion variables at the sonic point follows the approach developed for relativistic accretion discs and for spherical and isothermal flows onto Schwarzschild black holes [9, 10, 11]; here it is carried out for the adiabatic polytropic Michel flow across the (γ, a_∞) grid.

Table 5: Surface gravity of the acoustic horizon from (35) and the analogue Hawking temperature relative to the vacuum Schwarzschild value $\kappa_{\text{Sch}} = 1/2$. $a_\infty = 0.1$; $T_H = \kappa/2\pi$. The last row is the vacuum benchmark $a \rightarrow 1$, $r_a \rightarrow 1$.

γ	$r_a = r_c$	κ	$T_H = \kappa/2\pi$	$2\kappa = T_H^{\text{ac}}/T_H^{\text{vac}}$
5/3	4.38	$1.38 \cdot 10^{-2}$	$2.20 \cdot 10^{-3}$	$2.76 \cdot 10^{-2}$
4/3	14.11	$3.51 \cdot 10^{-3}$	$5.58 \cdot 10^{-4}$	$7.02 \cdot 10^{-3}$
vacuum	1	1/2	1/4 π	1

The suppression reflects the smooth transonic crossing at the widely displaced sonic point ($r_a \approx 4.4$ and 14.1, respectively): the softer radiation-dominated flow ($\gamma = 4/3$) accelerates more slowly through a sonic point that lies farther out, giving the coldest horizon. This is not a feature of the single case $a_\infty = 0.1$: varying a_∞ from 0.05 to 0.20 (Table 4 and Fig. 4, left) revealed that κ rises monotonically by more than an order of magnitude as the asymptotic gas grows hotter and the sonic point migrates inward toward the strong-field region, so the analogue Hawking temperature is a continuous function of the asymptotic boundary data rather than an isolated value.

These tabulated values (Tables 5–4) quantify the scale that governs the acoustic ringdown of this background. Chaverra *et al.* [7] found that for a widely separated sonic horizon ($r_a \gg 1$) the quasi-normal frequencies of the Michel flow scale approximately with κ , so the entries of Table 4 set the natural frequency unit for that regime.

This link was made quantitative. A comparison of the tabulated fundamental frequencies $s = \sigma + i\omega$ with the surface gravity computed here (their normalization is twice the one adopted

here, $\kappa_{\text{Ch}} = 2\kappa$, and the present κ reproduces the tabulated values to better than 0.5% over $r_a = 3\text{--}30$) yielded the convention-independent ratios $\text{Re } s \approx -0.77 \kappa$ and $\text{Im } s \approx (0.42 + 1.21 \ell) \kappa$ for the multipole index $\ell \geq 1$, nearly constant for $r_a \gtrsim 10$. The tabulated κ therefore fixes both the decay rate and the oscillation frequency of the Michel ringdown; for example, at $a_\infty = 0.1$ ($\kappa = 3.5 \cdot 10^{-3}$, $r_a \approx 14$) the dipolar ($\ell = 1$) acoustic mode sits at $s r_g \approx (-2.7 + 5.7 i) \cdot 10^{-3}$.

5.2. Formal horizon area and thermodynamic limits

A formal area in the effective geometry may also be assigned to the horizon. The angular part of $\mathcal{G}_{\mu\nu}$ induced on the sphere $r = r_a$ gives

$$\mathcal{A}_a = \oint \sqrt{\mathcal{G}_{\theta\theta}\mathcal{G}_{\phi\phi}} d\theta d\phi = \frac{4\pi r_a^2}{\Omega^2(r_a)}, \quad (39)$$

which is the natural conformal generalization of the classical horizon-area formula [17]. A Bekenstein–Hawking “entropy” $S_a \propto \mathcal{A}_a$ can be written by analogy, but the analogy must be sharply limited. The temperature (36) is physically robust: it follows from the kinematics of the wave equation on the effective metric and requires no dynamical equations for $\mathcal{G}_{\mu\nu}$ itself. By contrast, $\mathcal{A}_a$ and any Clausius relation $\delta Q = T_H \delta S_a$ lack fundamental status here, because $\mathcal{G}_{\mu\nu}$ does not obey the Einstein equations and $\mathcal{A}_a$ need not be monotonic. The acoustic horizon therefore reproduces Hawking *radiation* as a kinematic effect, but not the full thermodynamics of black holes.

6. Conclusions

The relativistic acoustic geometry of perfect-fluid Michel accretion onto a Schwarzschild black hole was constructed, and high-frequency sound was analysed as null geodesics of the effective metric (4). On the Michel background the acoustic event horizon coincides exactly with the sonic critical point, while the stationary-limit surface remains distinct and bounds a genuine acoustic ergoregion. The geometric-optics phenomenology developed here—force decomposition, Shapiro diagnostics, photon-sphere shift and tabulated surface gravities—complements the existing wave-equation and quasinormal-mode analyses of the same background.

The geometric optics of sound rays was shown to be controlled by the generalized Binet equation (26) that decomposes deflection into Einstein, refractive and acoustic-wind forces. Numerical integration for $a_\infty = 0.1$ demonstrated that the wind dominates near the horizon, refraction is repulsive throughout, and the capture barrier maximum fixes an acoustic photon sphere that is strongly displaced outward relative to the vacuum value. Across $a_\infty = 0.05\text{--}0.20$ both radii vary strongly, yet their ratio remains nearly invariant, so the acoustic shadow tracks the horizon at a multiple set mainly by the equation of state. The acoustic Shapiro delay, diagnosed by the local slowdown (29), exhibits freeze-out of outgoing rays at the horizon and convergence of both branches to the plateau $1/a_\infty$ at large radii.

The surface gravity at the sonic horizon was expressed through hydrodynamic gradients, so that the analogue Hawking temperature probes the rate at which the flow crosses the sonic point. At $a_\infty = 0.1$ the horizon is much colder than vacuum; the surface gravity rises monotonically by more than an order of magnitude as a_∞ increases and the sonic point migrates inward. The temperature itself is a robust kinematic consequence of the wave equation; a formal horizon area can be written, but it does not endow the acoustic geometry with a full Bekenstein–Hawking thermodynamics.

Taken together, these results show that the thermodynamic and flow parameters of the accretion stream enter acoustic lensing, time delay and the analogue Hawking scale in a quantitatively controlled way. A natural extension, left for future work, would be a spatial-stability analysis of acoustic, vortical and entropy modes.

Acknowledgments

The work was performed by an independent researcher without institutional funding. The NASA Astrophysics Data System database was used during the preparation.

References

- [1] W. G. Unruh, Experimental black-hole evaporation, *Physical Review Letters* 46 (1981) 1351–1353. doi:10.1103/PhysRevLett.46.1351.
- [2] M. Visser, Acoustic black holes: horizons, ergospheres and Hawking radiation, *Classical and Quantum Gravity* 15 (6) (1998) 1767–1791, arXiv:gr-qc/9712010. doi:10.1088/0264-9381/15/6/024.
- [3] C. Barceló, S. Liberati, M. Visser, Analogue gravity, *Living Reviews in Relativity* 14 (1) (2011) 3. doi:10.12942/lrr-2011-3.
- [4] F. C. Michel, Accretion of matter by condensed objects, *Astrophysics and Space Science* 15 (1) (1972) 153–160. doi:10.1007/BF00649949.
- [5] V. Moncrief, Stability of stationary spherical accretion onto a Schwarzschild black hole, *The Astrophysical Journal* 235 (1980) 1038–1046. doi:10.1086/157707.
- [6] N. Bilic, Relativistic acoustic geometry, *Classical and Quantum Gravity* 16 (1999) 3953–3964. doi:10.1088/0264-9381/16/12/312.
- [7] E. Chaverra, M. D. Morales, O. Sarbach, Quasi-normal acoustic oscillations in the Michel flow, *Physical Review D* 92 (2015) 084023, arXiv:1501.01637. doi:10.1103/PhysRevD.92.084023.
- [8] N. R. Jangid, A. K. Ray, Acoustic hawking radiation as a tunnelling effect in Michel accretion, arXiv:2607.00037 (2026). arXiv:2607.00037, doi:10.48550/arXiv.2607.00037.
- [9] T. K. Das, N. Bilić, S. Dasgupta, Black-hole accretion disc as an analogue gravity model, *Journal of Cosmology and Astroparticle Physics* 2007 (06) (2007) 009, arXiv:astro-ph/0604477. doi:10.1088/1475-7516/2007/06/009.
- [10] P. Tarafdar, T. K. Das, Dependence of acoustic surface gravity on the geometric configuration of matter for axially symmetric background flows in the Schwarzschild metric, *International Journal of Modern Physics D* 24 (14) (2015) 1550096, arXiv:1305.7134. doi:10.1142/S0218271815500960.
- [11] M. A. Shaikh, I. Firdousi, T. K. Das, Relativistic sonic geometry for isothermal accretion in the Schwarzschild metric, *Classical and Quantum Gravity* 34 (15) (2017) 155008, arXiv:1612.07963. doi:10.1088/1361-6382/aa7b19.
- [12] B. Toshmatov, B. Ahmedov, Z. Stuchlík, Phonon motion around (2+1)-dimensional acoustic black hole, *European Physical Journal C* 83 (10) (2023) 981. doi:10.1140/epjc/s10052-023-12181-8.
- [13] A. E. Balali, A. Marrani, Lense–thirring acoustic black holes: Shadows and light, arXiv:2512.23113 (2025). arXiv:2512.23113, doi:10.48550/arXiv.2512.23113.

- [14] H. S. Vieira, K. Destounis, K. D. Kokkotas, Analog Schwarzschild black holes of Bose–Einstein condensates in a cavity: Quasinormal modes and quasibound states, *Physical Review D* 107 (2023) 104038, arXiv:2301.11480. doi:10.1103/PhysRevD.107.104038.
- [15] K. Pal, K. Pal, T. Sarkar, Analogue metric in a black bounce background, *Universe* 8 (4) (2022) 197, arXiv:2204.06395. doi:10.3390/universe8040197.
- [16] I. I. Shapiro, Fourth test of general relativity, *Physical Review Letters* 13 (1964) 789–791. doi:10.1103/PhysRevLett.13.789.
- [17] S. L. Shapiro, S. A. Teukolsky, *Black holes, white dwarfs, and neutron stars: The physics of compact objects*, Wiley-Interscience, 2004. doi:10.1002/9783527617661.