

Wild Boar, *Sus scrofa*, habitat use in the Danish
fenced nature area Tofte Skov

Mads Gammelgaard | 201905844^{1†}, Terese Bech Eriksen |
201804715^{1†}, Lærke Thrane Mikkelsen | 201607735^{1†},
Asger Svenning | 201907234^{1*†}

¹Department of Biology, Aarhus University, Ny Munkegade 114-116,
Aarhus C, DK-8000, Midtjylland, Denmark.

*Corresponding author(s). E-mail(s): asgersvenning@gmail.com;
Contributing authors: gammelgaardmads@gmail.com;
teresebecheriksen@gmail.com; laerkethrane@outlook.com;
†These authors contributed equally to this work.

Keywords: Wild Boar, Habitat Selection, Movement Behaviour, GPS, Ecology

Content

Preface

Paper

Wild Boar, *Sus scrofa*, habitat use in the Danish fenced nature area Tofte Skov

Appendix 1

Movement Ecology Guidelines for Authors

Appendix 2

Supplementary map of habitats and GPS-fix densities of wild boars in Tofte Skov

Appendix 3

1. Effect of Wild Boar (*Sus scrofa L.*) on Forests, Agricultural Lands and Population Management in Lithuania; Tarvydas et al. 2022[1].
2. Spatiotemporal behavioral plasticity of wild boar (*Sus scrofa*) under contrasting conditions of human pressure: primeval forest and metropolitan area; Podgorski et al. 2013[2].

Preface

The present paper constitutes a concise and original research study intended for publication in the journal Movement Ecology, published by Springer. It focuses on investigating Wild boars' movement and habitat use in Tofte Skov, Denmark. The data foundation for the project consisted of GPS and accelerometer data from GPS collars, however due to time constraints and for simplicity we opted not to use the accelerometer data and instead focused on the GPS collar data. The data spans from February 2019 to January 2021 and was collected from a total of 25 individual wild boar.

¹ Wild Boar, *Sus scrofa*, habitat use in the Danish
² fenced nature area Tofte Skov

³ Mads Gammelgaard | 201905844^{1†}, Terese Bech Eriksen |
⁴ 201804715^{1†}, Lærke Thrane Mikkelsen | 201607735^{1†},
⁵ Asger Svenning | 201907234^{1*†}

⁶ ¹Department of Biology, Aarhus University, Ny Munkegade 114-116,
⁷ Aarhus C, DK-8000, Midtjylland, Denmark.

⁸ *Corresponding author(s). E-mail(s): asgersvenning@gmail.com;
⁹ Contributing authors: gammelgaardmads@gmail.com;
¹⁰ teresebecheriksen@gmail.com; laerkethrane@outlook.com;
¹¹ [†]These authors contributed equally to this work.

¹² **Abstract**

¹³ **Background:** Wild boars (*Sus scrofa*) are potential contributors to increasing
¹⁴ biodiversity and are considered keystone ecosystem engineers. However, concerns
¹⁵ exist regarding their invasive nature and potential damage to native biodiversity.
¹⁶ Understanding the behaviour and habitat use of wild boars is crucial for effective
¹⁷ management and their potential role in nature restoration.

¹⁸ **Methods:** The study was conducted in the fenced nature area Tofte Skov in
¹⁹ North-Eastern Jutland, Denmark. GPS-collar data from **25** wild boars were
²⁰ collected over a two-year period. Land use maps were used to analyze habitat
²¹ selection, and a nested temporally blocked bootstrap analysis was employed to
²² estimate habitat selection coefficients. Behaviour traits, including turning angle
²³ and mean velocity between fixes, were also examined.

²⁴ **Results:** We found significant differences in habitat use by wild boars in Tofte
²⁵ Skov. Forests had the highest habitat use frequency, followed by other natural
²⁶ habitats such as wet nature and dry nature. Artificial areas, streams, non-paved
²⁷ roads, and lakes had the lowest use frequency. Habitat selection coefficients
²⁸ revealed that forests, wet forests, streams, non-paved roads, artificial areas, and
²⁹ extensive agriculture were preferred habitats. Wet nature (raised bog) had a
³⁰ significantly lower habitat selection coefficient. The movement behaviour of wild

31 boars varied during the day, with faster and more straight movements during the
32 day and slower and more turning movements during the night.

33 **Conclusion:** We conclude that wild boars in Tofte Skov prefer closed deciduous
34 forests as their main habitat and forage at night while traveling between habitats
35 during the day. They can adapt to open habitats if these areas provide high-
36 quality food. Wild boars could be included in fenced national parks, and improving
37 food availability in forests may reduce their reliance on agricultural fields. Future
38 research should explore wild boar behaviour when there is no supplementary
39 feeding and if there are predators, like wolves, present.

40 **Keywords:** Wild Boar, Habitat Selection, Movement Behaviour, GPS, Ecology

41 Background

42 In 2020, as part of the European Green Deal aimed at combatting the global loss of
43 biodiversity, the European Commission adopted a new strategy for the restoration and
44 conservation of biodiversity. One of the key goals of this strategy is to achieve 30%
45 protected nature areas on land and sea by 2030[3]. However, in Denmark, the Danish
46 Biodiversity Council estimates that the current extent of protected areas on land is
47 below 2.3%[4]. Therefore, there is an urgent need to establish more protected nature
48 areas in Denmark.

49 Large mammals, such as wild boars (*Sus scrofa*), can play a significant role in
50 increasing biodiversity within the future Nature-Nationalparks. They have been shown
51 to positively impact ecological processes[4]. However, the presence of wild boars in
52 European nature has been a subject of extensive discussion. Farmers are concerned
53 about substantial economic losses due to crop damage and the transmission of diseases,
54 especially African Swine Fever (ASF), to domesticated pigs. Furthermore, the invasive
55 capabilities of wild boars raise concerns among some conservation-/restoration-ists and
56 managers, as they can cause significant damage to native biodiversity and potentially
57 drive certain species to extinction[5, 6]. Although concerns about the invasive nature
58 of wild boar are valid by themselves, wild boars should be considered native in

59 Denmark under any conceivable definition of "native" considering that they have a
60 deep evolutionary history in Europe, which was only ended locally in Denmark in
61 1801 by intensive hunting[7, 8]. Today wild boar are common in most of their natural
62 range including Europe, Asia, and Northern Africa and have been introduced to novel
63 ecosystems in Australia, New Zealand, North and South America[9].

64 Simultaneously, wild boars is broadly considered as a keystone ecosystem engineers
65 by ecologists. They provide essential disturbances to habitats that have lost their
66 natural disturbance regimes, which were previously provided by extinct megafauna[4,
67 10]. The unique rooting behaviour of wild boars has been associated with increased
68 biodiversity[11–13], enhanced habitat heterogeneity[14, 15], and act as vectors for seed
69 and fungi dispersal[16, 17]. Therefore, the impact of wild boars on natural ecosystems
70 is complex and requires a deeper understanding. Gaining insights into the behaviour
71 and habitat use of wild boars is crucial for comprehending the circumstances in which
72 they can be utilized as a tool for nature restoration, as well as recognizing situations in
73 which they may cause habitat damage. Studies of the historical-natural habitat use of
74 wild boars in Europe have found that wild boar mostly inhabit old deciduous forests
75 containing some wet areas and open meadows[18]. However, in the modern European
76 landscape, wild boars have benefited immensely from the expansion of agriculture and
77 have adapted to exist in human dominated areas close to human settlement[1, 2, 19].

78 One reason for concern arises from wild boars' ability to adapt to a wide range of
79 habitats in which it is introduced. Consequently, they have established populations in
80 almost all habitats they have been introduced to[20]. Furthermore, wild boars exhibit
81 opportunistic omnivorous feeding habits and can adapt their feeding methods to grazing,
82 browsing, foraging, rooting, and even predation[20, 21]. Their rooting behaviour is
83 especially important for creating habitat heterogeneity by removing monocultures
84 of dominating plants and creating disturbed sites in the soil opening up areas for
85 disturbance adapted species[22–25]. Even though wild boars are able to access many




Fig. 1 Bootstrap distributions of *S. scrofa* habitat use frequencies in Tofte Skov.

86 different sources of food, wild boars will, to the farmers' dismay, feed almost exclusively
 87 on crops if possible. A review study on the wild boars diet[20] revealed that crops can
 88 make up 90% of their stomach contents, which contributes to the large conflict between
 89 farmers and wild boars. Additionally, authorities and farmers fear the transmission of
 90 African Swine Fever (ASF) from wild boars to domestic pigs[26], which would have
 91 detrimental consequences for the export of pork. Controlling the spread of ASF is
 92 strictly regulated for most countries in Europe with wild boar populations and is on the
 93 political agenda on an EU-level[27]. Some extreme measures implemented by countries
 94 like Denmark is to completely exterminate all wild boars in the country, to prevent
 95 transmission of ASF[28]. To conclude, the role of the wild boar in European nature
 96 is widely conflicted by, on one hand, being an effective tool for nature restoration
 97 in unnaturally undisturbed European ecosystems and on the other hand being an
 98 expensive pest for agricultural interests. To utilize these wild boars in nature restoration
 99 while reducing conflicts with human interests, some countries like Denmark, have
 100 opted to include wild boar populations in enclosed nature areas. This makes it possible
 101 to investigate the behaviour of wild boar in semi-natural conditions. Deepening our

102 understanding of wild boar behaviour and habitat use is detrimental to ensure optimal
103 management of wild boar, if they are to be used as a tool for nature restoration in the
104 future.

105 This study aims to further the understanding of how wild boar utilizes different
106 habitats, and discuss how this could be used to improve management practices. We
107 investigate wild boar habitat-use by utilizing GPS-collar data, along with land use
108 maps, to describe the differences in behaviour and usage of different habitats in the
109 fenced nature area Tofte Skov in North-Eastern Jutland. Because wild boars naturally
110 prefer forests and semi-open areas[18, 29], we hypothesize that the wild boars core
111 area will be closed deciduous forest. We predict that there will be higher habitat-use
112 frequency and higher selection coefficients for forest habitats compared to the other
113 habitat types. Furthermore, we hypothesize that in open areas, where the wild boars are
114 more exposed, wild boars will perform shorter and faster excursions compared to times
115 of lower exposure. We predict that the step length in open areas compared to closed
116 areas will be longer, and that the turning index will be closer to zero. Furthermore, we
117 predict, that wild boars will have longer step lengths and higher turn indexes during
118 the day, as they are more exposed.

119 Methods

120 Software

121 All data analysis was done in the R programming language[30]. General data wrangling
122 was done using the tidyverse packages[31], using the sf package[32] for simple spatial
123 features and the stars package[33] for spatial arrays. Figures were produced using the
124 ggplot2 package[34] and metR package[35] for contour plots.




Fig. 2 Visualization of the values of the turning behaviour trait α given a preceding line-segment following the direction of the green arrow, and a following line-segment in the direction of the circle periphery joined at the center of the circle (green dot). Grey lines are equally spaced (0.05) α -contours.

¹²⁵ Study area

¹²⁶ Tofte Skov is a 27km² fenced area in the southern part of Lille Vildmose in eastern
¹²⁷ Himmerland, Denmark. It was fenced in 1907 as part of the preservation of the area.
¹²⁸ Tofte Skov lies within the Natura 2000 area number 17, which includes the entire Lille
¹²⁹ Vildmose and Høstemark Skov. The area consists of numerous plains separated by
¹³⁰ different kinds of deciduous forests including beech, oak, birch and alder. Common
¹³¹ spruce and sitka spruce have previously been planted and widespread in the area, but
¹³² as part of the preservation of the area these species are now being removed in favor of
¹³³ the native deciduous species. Tofte Skov is also characterized by having large areas of

¹³⁴ raised bog and different populations of wildlife including 150 wild boars, wisents and
¹³⁵ red deer[36?].

¹³⁶ Collection of data

¹³⁷ During a two-year study conducted from February 2019 to January 2021, a total of 25
¹³⁸ wild boars were collared. The collaring process was divided into three rounds: the first
¹³⁹ round occurred in February and March 2019, the second round took place in November
¹⁴⁰ 2019, and the final round was conducted in November 2020. To facilitate the collaring
¹⁴¹ process, the wild boars, which were accustomed to supplementary feeding, were enticed
¹⁴² by driving vehicles and distributing feed.

¹⁴³ The collaring procedure involved anesthetizing the wild boars using a CO₂ rifle.
¹⁴⁴ The anesthesia induction was accomplished by darting, and the animals remained
¹⁴⁵ under anesthesia for a few minutes, during which they were collared and ear tagged to
¹⁴⁶ facilitate individual identification. Proper ventilation of the anesthetized animals was
¹⁴⁷ ensured during the process. During the initial collaring rounds, the population of wild
¹⁴⁸ boars exceeded expectations, resulting in significant weight losses due to insufficient
¹⁴⁹ food supply. Consequently, several collars did not fit properly and were not adequately
¹⁵⁰ secured, leading to their subsequent detachment. Additionally, interactions between
¹⁵¹ the boars also caused some collars to fall off. As a response to the suboptimal feeding
¹⁵² conditions, a significant portion of the boar population was euthanized and excluded
¹⁵³ from the study in early 2020. Following this intervention, the remaining population
¹⁵⁴ experienced weight gain, resulting in better collar retention. Towards the end of the
¹⁵⁵ study, the last five collars were manually removed ahead of schedule due to concerns that
¹⁵⁶ they might be too tight. It should be noted that the wild boars received supplementary
¹⁵⁷ feed in late 2019, but not in the subsequent year.

¹⁵⁸ Information regarding the data collection was obtained from Lars Haugaard
¹⁵⁹ (personal communication).




Fig. 3 Bootstrap distributions of *S. scrofa* habitat use anomalies in Tofte Skov. Anomalies are calculated as the difference between the observed habitat use frequency (see Figure 1) and the habitat frequencies in Tofte Skov.

160 Habitat types

161 To analyze habitat use, we used the land use classification Basemap04 produced by
 162 The institute for Environmental science in Aarhus University[37]. Basemap04 is a
 163 combination of remote sensing land cover classification maps and land cover census
 164 data. The map defines 34 land use / land cover categories. As many of the categories
 165 were redundant for our analysis, we joined all building categories, unused agriculture
 166 categories and other artificial land use as a single category called “Artificial”. We
 167 kept the non-nature categories: "Road, not paved", "Agriculture, extensive" and
 168 "Agriculture, intensive". We included all the nature categories to differentiate land use
 169 across different nature types.

170 Habitat Selection

The **Habitat Selection Coefficient** (abbreviated as HSC throughout this report), is an index which calculates the proportion between observed and expected habitat use

frequency:

$$HSC_H = \frac{O(F_H)}{E(F_H)}, \quad F_H = \frac{n_H}{\sum n_i}$$

Then we compare the HSC between habitats by division not subtraction:

$$\delta_{HSC}(i, j) = \frac{HSC_i}{HSC_j} = \frac{O(F_i)}{O(F_j)} \cdot \frac{E(F_i)}{E(F_j)}$$

171 In order to conduct a statistically sound analysis of habitat selection, we chose a nested
172 temporally blocked bootstrap analysis in which the habitat selection coefficient is
173 calculated in one week blocks stratified by individual. To estimate the habitat selection
174 coefficients and the pairwise differences in habitat selection coefficients, δ_{HSC} , we
175 estimated the empirical distribution of habitat selection by bootstrapping the stratified
176 blocks 100 000 times. This procedure is meant to provide an unbiased estimate for
177 habitat selection coefficients and pairwise factor differences, while taking into account
178 temporal autocorrelation and individual effects. By temporally blocking in one week
179 intervals, we eliminate most of the temporal autocorrelation as we expect GPS-fixes
180 to be almost independent after one week. The nested bootstrapping then provides
181 an assumption-free way of estimating the empirical distribution of habitat selection
182 coefficients, and allows for differences in variance and non-unimodal over time, between
183 habitats and individuals.

184 Behaviour traits

185 We wanted to investigate whether we could detect any differences in behaviour between
186 the different habitats. To do so we chose to investigate two parameters of behaviour
187 which are quantifiable, at least to some degree, from our GPS-data; (1) turning angle
188 and (2) mean velocity between fixes, both under the simplifying assumption that
189 individuals are walking in straight lines between the fixes. The latter assumption is
190 almost with complete certainty wrong, however it is not obvious that it would be

Procedure 1 Statistical procedure for nested temporally blocked bootstrapping.

Data Dataset of GPS-fixes annotated with habitat, time of recording and individual ID**Result** Nested temporally blocked bootstrap results with mean, quantiles, and p-values.

```
1: for each Individual-nested temporal block do
2:   Pre-calculate habitat frequencies
3: end for
4: for  $i \in \{1 \dots 100\,000\}$  do
5:   Bootstrap individuals ( $n = 25$ )
6:   for each individual do
7:     Bootstrap temporal blocks
8:   end for
9:   for each  $i \in \text{habitat}$  do
10:    HUFi = Average use frequency
11:    HSCi =  $\frac{\text{HUF}_i}{E(\text{HUF}_i)}$ 
12:   end for
13:   for each  $\{i, j\} \in \text{habitats}$  do
14:      $\Delta_{i,j} = \frac{\text{HSC}_i}{\text{HSC}_j}$                                 ▷ Pairwise differences in HSC
15:   end for
16: end for
17: Summarize mean, median, lower and upper quantiles and empirical p-values
18: Adjust p-values for multiple hypothesis testing
```

191 biased in either of the quantifiable behaviour traits across any kinds of stratifications
192 in our dataset. We have therefore limited our analysis to one of relative differences
193 between different types of habitats and times of day. The turning behaviour trait will
194 be quantified using $\alpha = 1 - \frac{\cos(\theta)+1}{2}$ such that 0 corresponds to straight movement
195 while 1 corresponds to turning around, while velocity is simply calculated by dividing
196 the distance between two consecutive points in space by their distance in time¹. The
197 cosine is used for quantifying turning, since it the cosine is equivalent to the lateral
198 component of the direction vector, i.e. if the cosine is 0.4 then if the animal moves
199 1km then it moves 400 meters either left or right of where it would have been, if it had
200 continued straight for the same amount of time with the same velocity.

Given that turning, α , is an attribute of an angle or the mid-point of three consecutive points, while velocity is an attribute of a line-segment or two consecutive

¹we use SI-units for velocity; m/s .




Fig. 4 Histograms of the behaviour traits (turning and velocity) on the raw data scale (first row, (A)) and transformed (second row, (B)). Velocity is simply log-transformed and scaled, while turning degree is transformed using a quantile-transformation between a fitted beta-distribution and a scaled normal distribution.

points, for this analysis velocity refers to the velocity leading up to a GPS-fix and turning is based on the angle created by the prior and next point. In order to incorporate both of these behavioural characteristics into a single index of behaviour suitable for our analysis, we then conducted a PCA on the data matrix consisting of all points for which turning and velocity could be quantified. However for this to statistically sound, the behaviour traits must first be examined and if necessary transformed to

approximately follow a scaled normal distribution as PCA assumes that the the data follows a multivariate normal distribution²[38]. As can be clearly seen in the first row (**A**) of [Figure 4](#) the behaviour traits are *not* normally distributed or scaled. However the velocity trait is easily transformed to an approximate scaled normal distribution by a log-transform followed by scaling by mean and standard deviation (see second row (**B**) of [Figure 4](#)). The turning trait is somewhat more complicated as it follows a bounded and bi-modal distribution ($\alpha \in [0, 1]$), however we have chosen to solve this problem by fitting a beta distribution on the calculated turning trait, α , values with the following parameters $a \approx 0.403$ and $b \approx 0.322$ using the `fitdistrplus` package[39]. By fitting a suitable distribution to the data, it is possible to do an invertible "quantile-normalization"[40, 41]:

$$f^*(\alpha) = \phi_{\mathcal{N}}(\phi_{\text{Beta}}^{-1}(\alpha))$$

201 This procedure produces an extremely appropriate transformation of α as can be seen
202 in the second row (**B**) of [Figure 4](#). After these preprocessing steps it was possible to
203 conduct an appropriate PCA analysis ([Figure 5](#)) with the first principal component,
204 **PC1**, explaining $\sim 61.1\%$ of the variance going from (positive) fast and straight
205 to (negative) slow and turning behaviour. We have chosen to refer to this principal
206 component as *Spurtiness*.

207 Results

208 Habitat Selection

209 We found significant differences in habitat use by wild boars in Tofte Skov. The largest
210 habitat use frequency of around 35% were, as expected, for "Forest". This is followed

²For simplicity I only focus on the marginal distributions.




Fig. 5 Biplot of PCA analysis of behaviour traits overlayed on top of a heatmap of the observed trait values on the appropriate transformed scales (explaining the awkward axis breaks on the turning-axis in particular). The percentages next to the principal component labels are the percentage of variance explained by each principal component.

211 by other natural habitats such as wet nature, dry nature and wet forests. The areas of
 212 lowest use frequency are: artificial areas, streams, non-paved roads and lakes (Figure 1).
 213 We found a significantly higher than expected habitat selection coefficient (HSC) for
 214 the habitats "Stream", "Road, not paved", "Nature, dry", "Missing", "Forest" and "Forest,
 215 wet", "Artificial" and "Agriculture, extensive" (former landing-strip, with feeding station
 216 and animal-release point), while "Nature, wet" (mostly raised bog) was the only habitat
 217 type with a significantly lower than expected HSC (Figure 3). Furthermore, we found
 218 that after Bonferroni correction for multiple testing "Agriculture, extensive" had a

219 significantly higher HSC than all other habitat types, while "Forest" had a significantly
 220 higher HSC than all other natural habitats, both open and closed, and unsurprisingly
 221 "Nature, wet" also had a significantly lower HSC than all other habitats. The only other
 222 habitat with a significantly different HSC from any other than the before mentioned is
 223 "Artifical" which had a very marginal higher HSC than "Nature, dry" (Table 1).

Table 1 Bootstrap estimates ($n = 100\,000$) of pairwise factor differences in habitat selection coefficients (HSC), only significant comparisons ($P \leq 0.05$) are included in the table. *P-values are Bonferroni adjusted ($n = 55$) and only one digit is displayed in the table. The full unrounded table can be found in the supplementary materials.

Habitat	Compared habitat	μ_Δ	$Q_{50\%}(\Delta)$	$Q_{5\%}(\Delta)$	$Q_{95\%}(\Delta)$	*P-value
Agriculture, extensive						
	Agriculture, intensive	∞	∞	5.2	∞	5.5×10^{-4} **
	Artificial	11.3	10.7	4.7	19.8	5.5×10^{-4} **
	Forest	12.2	11.8	6.1	19.7	5.5×10^{-4} **
	Forest, wet	22.0	21.1	11.0	36.0	5.5×10^{-4} **
	Lake	49.0	32.0	13.8	126.3	5.5×10^{-4} **
	Missing	16.9	15.9	8.0	29.3	5.5×10^{-4} **
	Nature, dry	22.5	21.7	11.4	36.6	5.5×10^{-4} **
	Nature, wet	159.5	151.6	85.2	260.7	5.5×10^{-4} **
	Road, not paved	16.4	15.1	7.2	30.2	5.5×10^{-4} **
	Stream	16.1	15.1	7.3	28.5	5.5×10^{-4} **
Artificial						
	Nature, dry	2.2	1.9	1.3	3.9	0.04 *
	Nature, wet	15.6	14.0	8.8	27.6	5.5×10^{-4} **
Forest						
	Forest, wet	1.8	1.8	1.4	2.3	8.2×10^{-3} *
	Nature, dry	1.9	1.8	1.5	2.4	2.7×10^{-3} **
	Nature, wet	13.4	13.1	9.3	18.5	5.5×10^{-4} **
Forest, wet						
	Nature, wet	7.5	7.3	5.0	10.8	5.5×10^{-4} **
Lake						
	Nature, wet	4.6	4.9	1.7	8.3	3.3×10^{-3} **
Missing						
	Nature, wet	10.0	9.6	6.1	15.2	5.5×10^{-4} **
Nature, dry						
	Nature, wet	7.3	7.2	4.9	9.9	5.5×10^{-4} **
Road, not paved						
	Nature, wet	9.6	10.2	5.8	16.7	5.5×10^{-4} **
Stream						
	Nature, wet	9.7	10.1	6.2	16.7	5.5×10^{-4} **

Time of Day	μ_{PC1}	μ_{Δ}	$Q_{50\%}(\Delta)$	$Q_{5\%}(\Delta)$	$Q_{95\%}(\Delta)$	P-value
[00-06)	-0.42	-0.02	-0.11	-0.85	1.04	0.322
[06-12)	0.25	-0.08	-0.09	-1.43	1.20	0.429
[12-18)	0.59	-0.10	-0.08	-1.20	1.00	0.383
[18-24)	0.07	0.05	0.05	-0.95	1.11	0.407

Table 2 Bootstrap results ($n = 100\,000$) of differences in spurtiness behaviour between open and closed habitats across different times of day. All values except μ_{PC1} are differences calculated $\Delta = \text{Spurtiness}_{\text{Open}} - \text{Spurtiness}_{\text{Closed}}$, such that positive values indicate more spury (fast and straight) behaviour in open habitats. P-values are not corrected for multiple testing. μ_{PC1} is simply the mean of PC1 across both "Open" and "Closed" habitats in the given time of day interval.

224 Habitat Behaviour

225 We then proceeded with an equivalent analysis of movement behaviour as we did
 226 with habitat selection (nested temporally blocked bootstrapping) stratified across
 227 four intervals of time of day ([00–06), [06–12), [12–18) and [18–24) and two habitat
 228 types "Open" and "Closed", where we chose to classify "Forest" and "Forest, wet" as
 229 "Closed" while the remaining habitats were classified as "Open". We found no significant
 230 differences between open and closed habitats at any time of day (Table 2). However,
 231 the movement behaviour of the wild boars differed during the day with faster and
 232 more straight movements at daytime, especially between 12–18, and slower and more
 233 turning movements at night between 00–06.

234 Conclusion

235 Habitat selection

236 As hypothesized, our findings revealed that the wild boars showed a clear preference
 237 for the areas with closed deciduous forests. Both when comparing frequencies and
 238 when taking the size of the different habitats into account, by investigating habitat
 239 selection coefficients. The forest habitats in Tofte Skov create preferable conditions
 240 for the wild boars of the area, as they provide some of the key conditions and food
 241 sources for ideal wild boar habitats as outlined in a scientific review of the European

wild boar population[18]. Amongst these conditions, forests provide: heavy brush for predator protection, roots, fruits, nuts, bark, shoots, leaves and other food items. The aversion of wet nature likely arises, as this habitat exclusively consists of nutrient poor marshlands with low quality food availability, high exposure and possibly difficult movement conditions caused by the deep sphagnum layer. Furthermore, investigating [Figure B1 in Appendix B](#) it is evident that the wild boars primarily use the wet nature habitat between the forest and the lake, and the wet nature closest to the forest edge. This indicates that wild boars only use the wet nature habitat, when they are traveling to and from the lake, likely to drink water, and will otherwise avoid walking in this habitat.

Furthermore, the extensive agriculture habitat had a significantly larger selection coefficient than the other habitats regardless of being an open area. This can be explained by this area being the place in which they are fed. Furthermore, the holding pen in which the wild boars are released from and caught for check-ups is close to the extensive agriculture (personal observations and communication with Thomas Holst Christensen). These interactions are generally positive for the wild boar, as they include feeding. An explanation for the preference for this area could therefore be that the wild boars associate this area with an abundant source of high quality food, and are thus not encouraged to search for food in other habitats. This is comparable to how European wild boar populations have adapted to prefer forest habitats close to agricultural fields containing high value food, that makes the benefit of the area higher than the potential cost[18, 20]. This is not surprising, as wild boars have a high degree of behavioural plasticity[2]. Furthermore, the predation pressure in Tofte Skov is very low. At the time of the data collection there were no top predators in the area. So the only predatory risk the population is exposed to is hunting from humans. The supplementary feeding changes the natural behaviour of how the wild boars use the different habitats, making it harder to broaden the results of this study to wild boars

²⁶⁹ in non-managed unfenced nature areas. Future studies should include areas with no
²⁷⁰ supplementary feeding to evaluate how the wild boars root in natural settings compared
²⁷¹ to areas with supplementary feeding, such as Tofte Skov.

²⁷² Behavioural differences between habitats

²⁷³ Our results showed that there were no significant differences in movement behaviour
²⁷⁴ between open and closed habitats ([Table 2](#)). Hence the wild boars do not move short
²⁷⁵ and fast in open areas, as hypothesized. They do not seem to be “scared” of the open
²⁷⁶ areas as predicted. As mentioned above this is likely because they have learned to
²⁷⁷ associate some of the open areas with feeding. Furthermore, the hunting pressure in
²⁷⁸ Tofte Skov at the time were almost non-existing. Today there is a wolf inside the fenced
²⁷⁹ area, that could potentially predate on wild boar, though since first registration of
²⁸⁰ the wolf in 2021 the employees have not registered any killings of wild boar (personal
²⁸¹ communication with Thomas Holst Christensen). Future studies of GPS collar data
²⁸² from Tofte Skov, could include comparison of the wild boars habitat use before and
²⁸³ after the wolf entered the area. Even though it does not seem like the wolf preys on the
²⁸⁴ wild boars, it could still have an effect on how the wild boars move around and use the
²⁸⁵ different habitats. Across all habitats the movement behaviour follows a clear tendency
²⁸⁶ with the boars having a higher amount of spurtiness during the day and less during
²⁸⁷ the night ([Table 2](#)). Spurtiness is characterised by being fast and straight movements
²⁸⁸ with a low amount of turning, which would indicate that the boars use the daytime to
²⁸⁹ move around between different areas or habitats. Conversely, the movements during the
²⁹⁰ night are characterised by being slower and with more turning, which could indicate
²⁹¹ rooting or other kind of foraging behaviour. This is consistent with the boars being
²⁹² nocturnal animals that forage during the night[42].

293 Management implications

294 As discussed, the preference of wild boars to use feeding stations in the extensive
295 agriculture area (the former landing strip) can be compared to the well documented
296 preference of wild boars, to forage on agricultural fields[20]. Wild boars are likely to have
297 the same preference in Denmark, considering the large availability of crops. Introducing
298 wild boars into Danish nature would therefore likely increase human-wildlife conflict
299 with farmers, as they could damage crops, and increase fear of transmission of African
300 Swine Fever. It is important to highlight that infection of AFS from wild boars to
301 domestic pigs is most likely in areas with many free range pigs such as Caucasus
302 and Russia. Furthermore, whilst wild boars can infect domestic pigs, AFS does not
303 persist in populations of wild boars, without continuous infection from other sources,
304 such as areas with infected domesticated pigs[43]. It is possible that proper fencing to
305 avoid contact between wild boars and domesticated pigs, would diminish infection risk.
306 However, whilst the risk of AFS spread from wild boars to domestic pigs, is contested,
307 it is still, in the current political landscape, unlikely that wild boars will be a tool for
308 nature management in unfenced nature areas. However, the issues relating to wild boar
309 contact with domestic pig production would be avoided, if wild boars were kept in
310 large fenced nature areas such as the proposed nature national parks in Denmark. This
311 would allow the wild boars to act as ecosystem engineers[10], creating much needed
312 disturbances[4] whilst minimizing human-wildlife conflict.

313 Concluding remarks & perspective

314 Our results provide valuable insights for the management of wild boar populations.
315 This study provides empirical evidence supporting the hypothesis that wild boars
316 prefer closed deciduous forests as their core habitat. We did not find any significant
317 differences in movement behaviour across different habitat types. We did however find
318 that wild boars move further and more straight during the day compared to during

the night. This suggests that they forage during the night and walk between habitats during the day. Furthermore, our results show that the wild boars in Tofte Skov can adapt a significant preference for open habitats, if they are associated with positive effects such as the provision of high quality food. This also indicates that if wild boars were to be set free in Danish nature, they would forage on fields. To avoid escalating the human-wildlife conflict, wild boars could be included in fenced nature national parks where they could forage naturally and create needed disturbances in the landscape. Furthermore, one could try to improve the natural food availability in areas with closed deciduous forest of high quality in terms of food for the wild boars. This could potentially decrease the wild boar's tendency to forage in agricultural fields. Further studies should include investigations on whether the wolf, which at the moment is the only potential predator in Denmark, can have a positive effect on avoiding wild boars going into agricultural fields to forage.

Supplementary information. All code and data is available on <https://github.com/asgersvenning/Vildsvinene>.

References

- [1] Tarvydas, A. & Belova, O. Effect of Wild Boar (*Sus scrofa* L.) on Forests, Agricultural Lands and Population Management in Lithuania. *Diversity* **14**, 801 (2022). URL <https://www.mdpi.com/1424-2818/14/10/801>.
- [2] Podgórski, T. *et al.* Spatiotemporal behavioral plasticity of wild boar (*Sus scrofa*) under contrasting conditions of human pressure: Primeval forest and metropolitan area. *Journal of Mammalogy* **94**, 109–119 (2013). URL <https://academic.oup.com/jmammal/article/94/1/109/844802>.

- 343 [3] European Union. Biodiversity strategy for 2030 (2023). URL https://environment.ec.europa.eu/strategy/biodiversity-strategy-2030{_}en.
- 344
345 [4] Normand, S. *et al.* Fra tab til fremgang - Beskyttet natur i Danmark i et
346 internationalt perspektiv (2022). URL <https://www.biodiversitetsraadet.dk/pdf/2022/12/Biodiversitetsraadet-2022-Fra-tab-til-fremgang-Final-hjemmeside.pdf>.
- 347
348 [5] Risch, D. R., Ringma, J. & Price, M. R. The global impact of wild pigs (*Sus*
349 *scrofa*) on terrestrial biodiversity. *Scientific Reports* **11** (2021). URL https://click.endnote.com/viewer?doi=10.1038%2Fs41598-021-92691-1&token=WzI5ODQzMjgsIJewLjEwMzgvczQxNTk4LTAyMS05MjY5MS0xIl0.xXS145O{_}iVfxkR7LTKkZeUqTkgA.
- 350
351
352
353 [6] Krull, C. R. & Egeter, B. Feral pig (*Sus scrofa*) predation of a green
354 and golden bell frog (*Litoria aurea*). *New Zealand Journal of Ecology*
355 **40**, 191–195 (2015). URL https://www.researchgate.net/publication/281448777{_}Feral{_}pig{_}Sus{_}scrofa{_}predation{_}of{_}a{_}green{_}and{_}golden{_}bell{_}frog
- 356
357 [7] Hald-Mortensen, P. Vildsvin (2012). URL <https://pattedyratlas.lex.dk/Vildsvin>.
- 358 [8] Ruvinsky, A., Rothschild, M. F., Larson, G. & Gongora, J. *Systematics and evolution of the pig* (CABI Publishing, 2011). URL https://www.researchgate.net/publication/259292293{_}Systematics{_}and{_}Evolution{_}of{_}the{_}Pig.
- 359
360
361
362 [9] Lewis, J. S. *et al.* Biotic and abiotic factors predicting the global distribution and
363 population density of an invasive large mammal. *Scientific Reports* **7**, 1–12 (2017).
364 URL <https://www.nature.com/articles/srep44152>.
- 365 [10] Fløjgaard, C. *et al.* *Biodiversitetseffekter af rewilding* 425 (2021). URL <http://dce2.au.dk/pub/SR425.pdf>.

- 367 [11] Dovrat, G., Perevolotsky, A. & Ne'eman, G. The response of Mediterranean
368 herbaceous community to soil disturbance by native wild boars. *Plant Ecology*
369 **215**, 531–541 (2014). URL <https://doi.org/10.1007/s11258-014-0321-3>.
- 370 [12] Sondej, I. & Kwiatkowska-Falińska, A. Effects of Wild Boar (*Sus scrofa* L.)
371 Rooting on Seedling Emergence in Białowieża Forest. *Polish Journal of Ecology*
372 **65** (2017).
- 373 [13] Brunet, J., Hedwall, P.-O., Holmström, E. & Wahlgren, E. Disturbance of the
374 herbaceous layer after invasion of an eutrophic temperate forest by wild boar.
375 *Nordic Journal of Botany* **34**, 120–128 (2016). URL <https://onlinelibrary.wiley.com/com/doi/abs/10.1111/njb.01010>.
- 377 [14] Horčíčková, E., Brůna, J. & Vojta, J. Wild boar (*Sus scrofa*) increases species
378 diversity of semidry grassland: Field experiment with simulated soil disturbances.
379 *Ecology and Evolution* **9**, 2765–2774 (2019). URL <https://onlinelibrary.wiley.com/doi/abs/10.1002/ece3.4950>.
- 381 [15] Welander, J. Spatial and temporal dynamics of wild boar (*Sus scrofa*)
382 rooting in a mosaic landscape. *Journal of Zoology* **252**, 263–271 (2000).
383 URL <https://zslpublications.onlinelibrary.wiley.com/doi/abs/10.1111/j.1469-7998.2000.tb00621.x>.
- 385 [16] Pokorny, B. Ecological importance and impacts of wild boar (*Sus scrofa* L.).
386 *Zlatorgov Zbornik* **2**, 2–30 (2013).
- 387 [17] Genov, P. V., Focardi, S., Morimando, F., Scillitani, L. & Ahmed, A. in *Ecological
388 Impact of Wild Boar in Natural Ecosystems* (eds Meijaard, E. & Melletti, M.)
389 *Ecology, Conservation and Management of Wild Pigs and Peccaries* 404–419 (Cam-
390 bridge University Press, Cambridge, 2017). URL <https://www.cambridge.org/>

- 391 core/books/ecology-conservation-and-management-of-wild-pigs-and-peccaries/
392 ecological-impact-of-wild-boar-in-natural-ecosystems/
393 B3993DF1AEF7CE5DEFFECCE289C82A57.
- 394 [18] Tack, J. *Wild boar (Sus scrofa) Populations in Europe: A scientific review of*
395 *population trends and implications for management* (2018).
- 396 [19] Bieber, C. & Ruf, T. Population dynamics in wild boar Sus scrofa: Ecol-
397 ogy, elasticity of growth rate and implications for the management of pulsed
398 resource consumers. *Journal of Applied Ecology* **42**, 1203–1213 (2005). URL
399 <https://onlinelibrary.wiley.com/doi/full/10.1111/j.1365-2664.2005.01094.x>
400 <https://onlinelibrary.wiley.com/doi/abs/10.1111/j.1365-2664.2005.01094.x>
401 [//besjournals.onlinelibrary.wiley.com/doi/10.1111/j.1365-2664.2005.01094.x](https://besjournals.onlinelibrary.wiley.com/doi/10.1111/j.1365-2664.2005.01094.x).
- 402 [20] Ballari, S. A. & Barrios-García, M. N. A review of wild boar Sus scrofa
403 diet and factors affecting food selection in native and introduced ranges
404 (2014). URL <https://onlinelibrary.wiley.com/doi/full/10.1111/mam.12015>
405 <https://onlinelibrary.wiley.com/doi/abs/10.1111/mam.12015>
406 <https://onlinelibrary.wiley.com/doi/10.1111/mam.12015>.
- 407 [21] Schley, L. & Roper, T. J. Diet of wild boar Sus scrofa in Western Europe, with
408 particular reference to consumption of agricultural crops. *Mammal Review* **33**,
409 43–56 (2003). URL <https://onlinelibrary.wiley.com/doi/full/10.1046/j.1365-2907.2003.00010.x>
410 <https://onlinelibrary.wiley.com/doi/abs/10.1046/j.1365-2907.2003.00010.x>
411 <https://onlinelibrary.wiley.com/doi/10.1046/j.1365-2907.2003.00010.x>.
- 412 [22] Sandom, C. & Hughes, J. Rewilding the Scottish Highlands: Do Wild Boar, Sus
413 scrofa, Use a Suitable Foraging Strategy to be Effective Ecosystem Engineers?
414 *Restoration Ecology* **21**, 336–343 (2013).

- 415 [23] Sims, N. K., John, E. A. & Stewart, A. J. A. Short-term response and recovery of
416 bluebells (*Hyacinthoides non-scripta*) after rooting by wild boar (*Sus scrofa*). *Plant
417 Ecology* **215**, 1409–1416 (2014). URL <https://doi.org/10.1007/s11258-014-0397-9>.
- 418 [24] Burrascano, S. *et al.* Quantifying Sus scrofa rooting effects on the understorey
419 of the deciduous broadleaf forests in Castelporziano Estate (Italy). *Rendiconti
420 Lincei* **26**, 317–324 (2015).
- 421 [25] Pankova, N. L., Markov, N. I. & Vasina, A. L. Effect of the Rooting Activity
422 of Wild Boar Sus scrofa on Plant Communities in the Middle Taiga of Western
423 Siberia. *Russian Journal of Biological Invasions* **11**, 363–371 (2020). URL <https://click.endnote.com/viewer?doi=10.1134%2Fs2075111720040116&token=WzI5ODQzMjgsIjEwLjExMzQvczIwNzUxMTE3MjAwNDAxMTYiXQ.46QwYxfkfUI6D5IyhjZl3sV0cio>.
- 427 [26] Cadenas-Fernández, E., Ito, S., Aguilar-Vega, C., Sánchez-Vizcaíno, J. M. &
428 Bosch, J. The Role of the Wild Boar Spreading African Swine Fever Virus in Asia:
429 Another Underestimated Problem. *Frontiers in Veterinary Science* **9**, 453 (2022).
- 430 [27] Europa-Komissionen. KOMMISSIONENS GENNEMFØRELSESFORORDNING
431 (EU) 2023/594 af 16. marts 2023 om særlige sygdomsbekæmpelsesforanstaltninger
432 vedrørende afrikansk svinepest og om opfævelse af gennemførelsesforordning (EU)
433 2021/605 (2023). URL <https://eur-lex.europa.eu/legal-content/DA/TXT/HTML/?uri=CELEX:32023R0594>.
- 435 [28] Naturstyrelsen. Regulering af vildsvin - Vildsvin kan bekæmpes i henhold til
436 BEK nr. 1006 af 14/6/2020 – Bekendtgørelse om vildtskader. (2020). URL
437 <https://naturstyrelsen.dk/naturoplevelser/jagt/vildsvin/>.

- 438 [29] Yokoyama, Y., Nakashima, Y., Yajima, G. & Miyashita, T. Simultaneous estimation
439 of seasonal population density, habitat preference and catchability of wild
440 boars based on camera data and harvest records. *Royal Society Open Science* **7**
441 (2020).
- 442 [30] R Core Team. R: A Language and Environment for Statistical Computing (2022).
443 URL <https://www.r-project.org/>.
- 444 [31] Wickham, H. *et al.* Welcome to the tidyverse. *Journal of Open Source Software*
445 **4**, 1686 (2019).
- 446 [32] Pebesma, E. Simple features for R: Standardized support for spatial vector data.
447 *R Journal* **10**, 439–446 (2018). URL <https://doi.org/10.32614/RJ-2018-009>.
- 448 [33] Pebesma, E. & Bivand, R. *Spatial Data Science: With applications in R* (Chapman
449 and Hall/CRC, 2023). URL <https://r-spatial.org/book/>.
- 450 [34] Wickham, H. *ggplot2: Elegant Graphics for Data Analysis* (Springer-Verlag New
451 York, 2016). URL <https://ggplot2.tidyverse.org>.
- 452 [35] Campitelli, E. *metR: Tools for Easier Analysis of Meteorological Fields* (2021).
453 URL <https://github.com/eliocamp/metR>.
- 454 [36] Nielsen, S. B. Lille Vildmose | Fredninger (2007). URL <https://www.fredninger.dk/fredning/lille-vildmose/>.
- 456 [37] Levin, G. Basemap. URL <https://envs.au.dk/faglige-omraader/samfund-miljoe-og-ressourcer/arealanvendelse-og-gis/basemap>.
- 458 [38] Jollife, I. T. & Cadima, J. Principal component analysis: A review and recent
459 developments (2016). URL <https://royalsocietypublishing.org/doi/10.1098/rsta.2015.0202>.

- 461 [39] Delignette-Muller, M. L. & Dutang, C. {fitdistrplus}: An {R} Package for Fitting
462 Distributions. *Journal of Statistical Software* **64**, 1–34 (2015).
- 463 [40] Müller, C. *et al.* Removing Batch Effects from Longitudinal Gene Expression -
464 Quantile Normalization Plus ComBat as Best Approach for Microarray Transcrip-
465 tome Data. *PLOS ONE* **11**, e0156594 (2016). URL <https://journals.plos.org/plosone/article?id=10.1371/journal.pone.0156594>.
- 467 [41] Peterson, R. A. & Cavanaugh, J. E. Ordered quantile normalization: a semi-
468 parametric transformation built for the cross-validation era. *Journal of Applied*
469 *Statistics* **47**, 2312–2327 (2020). URL <https://www.tandfonline.com/doi/abs/10.1080/02664763.2019.1630372>.
- 471 [42] Cahill, S., Llimona, F. & Gràcia, J. Spacing and nocturnal activity of
472 wild boar Sus scrofa in a Mediterranean metropolitan park. *Wildlife*
473 *Biology* **9**, 3–13 (2003). URL <https://www.researchgate.net/publication/239537275{ }Spacing{ }and{ }nocturnal{ }activity{ }of{ }wild{ }boar{ }Sus{ }scrofa{ }in{ }>
- 475 [43] Costard, S., Mur, L., Lubroth, J., Sanchez-Vizcaino, J. M. & Pfeiffer, D. U.
476 Epidemiology of African swine fever virus. *Virus research* **173**, 191–197 (2013).
477 URL <https://pubmed.ncbi.nlm.nih.gov/23123296/>.

478 **Appendix A Movement Ecology Guidelines for
479 Authors**



Submission Guidelines ▾

[Aims and scope](#)[Fees and funding](#)[Language editing services](#)[Copyright](#)

▼ [Preparing your manuscript](#)

[Comment](#)[Methodology](#)[Research](#)[Review](#)[Software](#)[Prepare supporting information](#)[Conditions of publication](#)[Editorial policies](#)[Peer-review policy](#)[Manuscript transfers](#)[Promoting your publication](#)

Research

Preparing your manuscript

The information below details the section headings that you should include in your manuscript and what information should be within each section.

Please note that your manuscript must include a 'Declarations' section including all of the subheadings (please see below for more information).

Title page

The title page should:

481

- present a title that includes, if appropriate, the study design e.g.:
 - "A versus B in the treatment of C: a randomized controlled trial", "X is a risk factor for Y: a case control study", "What is the impact of factor X on subject Y: A systematic review"
 - or for non-clinical or non-research studies a description of what the article reports
- list the full names and institutional addresses for all authors
 - if a collaboration group should be listed as an author, please list the Group name as an author. If you would like the names of the individual members of the Group to be searchable through their individual PubMed records, please include this information in the "Acknowledgements" section in accordance with the instructions below
 - Large Language Models (LLMs), such as [ChatGPT](#), do not currently satisfy our [authorship criteria](#). Notably an attribution of authorship carries with it accountability for the work, which cannot be effectively applied to LLMs. Use of an LLM should be properly documented in the Methods section (and if a Methods section is not available, in a suitable alternative part) of the manuscript.
- indicate the corresponding author

Abstract

The Abstract should not exceed 350 words. Please minimize the use of abbreviations and do not cite references in the abstract. Reports of randomized controlled trials should follow the [CONSORT](#) extension for abstracts. The abstract must include the following separate sections:

- **Background:** the context and purpose of the study
- **Methods:** how the study was performed and statistical tests used
- **Results:** the main findings
- **Conclusions:** brief summary and potential implications

- **Trial registration:** If your article reports the results of a health care intervention on human participants, it must be registered in an appropriate registry and the registration number and date of registration should be stated in this section. If it was not registered prospectively (before enrollment of the first participant), you should include the words 'retrospectively registered'. See our [editorial policies](#) for more information on trial registration

Keywords

Three to ten keywords representing the main content of the article.

Background

The Background section should explain the background to the study, its aims, a summary of the existing literature and why this study was necessary or its contribution to the field.

Methods

The methods section should include:

- the aim, design and setting of the study
- the characteristics of participants or description of materials
- a clear description of all processes, interventions and comparisons. Generic drug names should generally be used. When proprietary brands are used in research, include the brand names in parentheses
- the type of statistical analysis used, including a power calculation if appropriate

Results

This should include the findings of the study including, if appropriate, results of statistical analysis which must be included either in the text or as tables and figures.

Discussion

This section should discuss the implications of the findings in context of existing research and highlight limitations of the study.

483

Conclusions

This should state clearly the main conclusions and provide an explanation of the importance and relevance of the study reported.

List of abbreviations

If abbreviations are used in the text they should be defined in the text at first use, and a list of abbreviations should be provided.

Declarations

All manuscripts must contain the following sections under the heading 'Declarations':

- Ethics approval and consent to participate
- Consent for publication
- Availability of data and materials
- Competing interests
- Funding
- Authors' contributions
- Acknowledgements
- Authors' information (optional)

Please see below for details on the information to be included in these sections.

If any of the sections are not relevant to your manuscript, please include the heading and write 'Not applicable' for that section.

Ethics approval and consent to participate

Manuscripts reporting studies involving human participants, human data or human tissue must:

- include a statement on ethics approval and consent (even where the need for approval was waived)
- include the name of the ethics committee that approved the study and the committee's reference number if appropriate

Studies involving animals must include a statement on ethics approval and for experimental studies involving client-owned animals, authors must also include a statement on informed consent from the client or owner.

See our [editorial policies](#) for more information.

If your manuscript does not report on or involve the use of any animal or human data or tissue, please state "Not applicable" in this section.

Consent for publication

If your manuscript contains any individual person's data in any form (including any individual details, images or videos), consent for publication must be obtained from that person, or in the case of children, their parent or legal guardian. All presentations of case reports must have consent for publication.

You can use your institutional consent form or our [consent form](#) if you prefer. You should not send the form to us on submission, but we may request to see a copy at any stage (including after publication).

See our [editorial policies](#) for more information on consent for publication.

If your manuscript does not contain data from any individual person, please state "Not applicable" in this section.

Availability of data and materials

All manuscripts must include an 'Availability of data and materials' statement. Data availability statements should include information on where data supporting the results reported in the article can be found including, where applicable, hyperlinks to publicly archived datasets analysed or generated during the study. By data we mean the minimal dataset that would be necessary to interpret, replicate and build upon the findings reported in the article. We recognise it is not always possible to share research data publicly, for instance when individual privacy could be compromised, and in such instances data availability should still be stated in the manuscript along with any conditions for access.

Authors are also encouraged to preserve search strings on searchRxiv <https://searchrxiv.org/>, an archive to support researchers to report, store and share their searches consistently and to enable them to review and re-use existing searches. searchRxiv enables researchers to obtain a digital object identifier (DOI) for their search, allowing it to be cited.

Data availability statements can take one of the following forms (or a combination of more than one if required for multiple datasets):

- The datasets generated and/or analysed during the current study are available in the [NAME] repository, [PERSISTENT WEB LINK TO DATASETS]
- The datasets used and/or analysed during the current study are available from the corresponding author on reasonable request.
- All data generated or analysed during this study are included in this published article [and its supplementary information files].
- The datasets generated and/or analysed during the current study are not publicly available due [REASON WHY DATA ARE NOT PUBLIC] but are available from the corresponding author on reasonable request.
- Data sharing is not applicable to this article as no datasets were generated or analysed during the current study.
- The data that support the findings of this study are available from [third party name] but restrictions apply to the availability of these data, which were used under license for the current study, and so are not publicly available. Data are however available

from the authors upon reasonable request and with permission of [third party name].

- Not applicable. If your manuscript does not contain any data, please state 'Not applicable' in this section.

More examples of template data availability statements, which include examples of openly available and restricted access datasets, are available [here](#).

BioMed Central strongly encourages the citation of any publicly available data on which the conclusions of the paper rely in the manuscript. Data citations should include a persistent identifier (such as a DOI) and should ideally be included in the reference list. Citations of datasets, when they appear in the reference list, should include the minimum information recommended by DataCite and follow journal style. Dataset identifiers including DOIs should be expressed as full URLs. For example:

Hao Z, AghaKouchak A, Nakhjiri N, Farahmand A. Global integrated drought monitoring and prediction system (GIDMaPS) data sets. figshare. 2014.

<http://dx.doi.org/10.6084/m9.figshare.853801>

With the corresponding text in the Availability of data and materials statement:

The datasets generated during and/or analysed during the current study are available in the [NAME] repository, [PERSISTENT WEB LINK TO DATASETS].[Reference number]

If you wish to co-submit a data note describing your data to be published in [BMC Research Notes](#), you can do so by visiting our [submission portal](#). Data notes support [open data](#) and help authors to comply with funder policies on data sharing. Co-published data notes will be linked to the research article the data support ([example](#)).

Competing interests

All financial and non-financial competing interests must be declared in this section.

See our [editorial policies](#) for a full explanation of competing interests. If you are unsure whether you or any of your co-authors have a competing interest please contact the editorial office.

Please use the authors initials to refer to each authors' competing interests in this section.

487

If you do not have any competing interests, please state "The authors declare that they have no competing interests" in this section.

Funding

All sources of funding for the research reported should be declared. The role of the funding body in the design of the study and collection, analysis, and interpretation of data and in writing the manuscript should be declared.

Authors' contributions

The individual contributions of authors to the manuscript should be specified in this section. Guidance and criteria for authorship can be found in our [editorial policies](#).

Please use initials to refer to each author's contribution in this section, for example: "FC analyzed and interpreted the patient data regarding the hematological disease and the transplant. RH performed the histological examination of the kidney, and was a major contributor in writing the manuscript. All authors read and approved the final manuscript."

Acknowledgements

Please acknowledge anyone who contributed towards the article who does not meet the criteria for authorship including anyone who provided professional writing services or materials.

Authors should obtain permission to acknowledge from all those mentioned in the Acknowledgements section.

See our [editorial policies](#) for a full explanation of acknowledgements and authorship criteria.

If you do not have anyone to acknowledge, please write "Not applicable" in this section.

Group authorship (for manuscripts involving a collaboration group): if you would like the names of the individual members of a collaboration Group to be searchable through their individual PubMed records, please ensure that the title of the collaboration Group is included on the title page and in the submission system and also include collaborating author names as the last paragraph of the "Acknowledgements" section. Please add authors in the format First Name, Middle initial(s) (optional), Last Name. You can add institution or country information for each author if you wish, but this should be consistent across all authors.

Please note that individual names may not be present in the PubMed record at the time a published article is initially included in PubMed as it takes PubMed additional time to code this information.

Authors' information

This section is optional.

You may choose to use this section to include any relevant information about the author(s) that may aid the reader's interpretation of the article, and understand the standpoint of the author(s). This may include details about the authors' qualifications, current positions they hold at institutions or societies, or any other relevant background information. Please refer to authors using their initials. Note this section should not be used to describe any competing interests.

Footnotes

Footnotes can be used to give additional information, which may include the citation of a reference included in the reference list. They should not consist solely of a reference citation, and they should never include the bibliographic details of a reference. They should also not contain any figures or tables.

Footnotes to the text are numbered consecutively; those to tables should be indicated by superscript lower-case letters (or asterisks for significance values and other statistical data). Footnotes to the title or the authors of the article are not given reference symbols.

Always use footnotes instead of endnotes.

References

Examples of the Vancouver reference style are shown below.

See our [editorial policies](#) for author guidance on good citation practice

Web links and URLs: All web links and URLs, including links to the authors' own websites, should be given a reference number and included in the reference list rather than within the text of the manuscript. They should be provided in full, including both the title of the site and the URL, as well as the date the site was accessed, in the following format: The Mouse Tumor Biology Database.

<http://tumor.informatics.jax.org/mtbwi/index.do>. Accessed 20 May 2013. If an author or group of authors can clearly be associated with a web link, such as for weblogs, then they should be included in the reference.

Example reference style:

Article within a journal

Smith JJ. The world of science. Am J Sci. 1999;36:234-5.

Article within a journal (no page numbers)

Rohrmann S, Overvad K, Bueno-de-Mesquita HB, Jakobsen MU, Egeberg R, Tjønneland A, et al. Meat consumption and mortality - results from the European Prospective Investigation into Cancer and Nutrition. BMC Medicine. 2013;11:63.

Article within a journal by DOI

Slifka MK, Whitton JL. Clinical implications of dysregulated cytokine production. Dig J Mol Med. 2000; doi:10.1007/s801090000086.

Article within a journal supplement

Frumin AM, Nussbaum J, Esposito M. Functional asplenia: demonstration of splenic activity by bone marrow scan. *Blood* 1979;59 Suppl 1:26-32.

490

Book chapter, or an article within a book

Wyllie AH, Kerr JFR, Currie AR. Cell death: the significance of apoptosis. In: Bourne GH, Danielli JF, Jeon KW, editors. International review of cytology. London: Academic; 1980. p. 251-306.

OnlineFirst chapter in a series (without a volume designation but with a DOI)

Saito Y, Hyuga H. Rate equation approaches to amplification of enantiomeric excess and chiral symmetry breaking. *Top Curr Chem*. 2007. doi:10.1007/128_2006_108.

Complete book, authored

Blenkinsopp A, Paxton P. Symptoms in the pharmacy: a guide to the management of common illness. 3rd ed. Oxford: Blackwell Science; 1998.

Online document

Doe J. Title of subordinate document. In: The dictionary of substances and their effects. Royal Society of Chemistry. 1999. <http://www.rsc.org/dose/title of subordinate document>. Accessed 15 Jan 1999.

Online database

Healthwise Knowledgebase. US Pharmacopeia, Rockville. 1998.
<http://www.healthwise.org>. Accessed 21 Sept 1998.

Supplementary material/private homepage

Doe J. Title of supplementary material. 2000. <http://www.privatehomepage.com>. Accessed 22 Feb 2000.

University site

Doe, J: Title of preprint. <http://www.uni-heidelberg.de/mydata.html> (1999). Accessed 25 Dec 1999.

491

FTP site

Doe, J: Trivial HTTP, RFC2169. <ftp://ftp.isi.edu/in-notes/rfc2169.txt> (1999). Accessed 12 Nov 1999.

Organization site

ISSN International Centre: The ISSN register. <http://www.issn.org> (2006). Accessed 20 Feb 2007.

Dataset with persistent identifier

Zheng L-Y, Guo X-S, He B, Sun L-J, Peng Y, Dong S-S, et al. Genome data from sweet and grain sorghum (*Sorghum bicolor*). GigaScience Database. 2011.

<http://dx.doi.org/10.5524/100012>.

Figures, tables and additional files

See [General formatting guidelines](#) for information on how to format figures, tables and additional files.

[Submit manuscript](#)

Submit manuscript

[Editorial Board](#)

[Manuscript editing services](#)

[Instructions for Editors](#)

[Contact Support for Editors](#)

[Sign up for article alerts and news from this journal](#)

492

Affiliated with



Annual Journal Metrics

Citation Impact

5.253 - [2-year Impact factor \(2021\)](#)

5.313 - [5-year Impact factor \(2021\)](#)

1.610 - [Source Normalized Impact per Paper \(SNIP\)](#)

2.095 - [SCImago Journal Rank \(SJR\)](#)

Speed

42 days to first decision for all manuscripts (Median)

51 days to first decision for reviewed manuscripts only (Median)

Usage

344,604 Downloads (2022)

1,580 Altmetric mentions (2021)

More about our metrics



ISSN: 2051-3933

Contact us

Submission enquiries: cecille.calusa@springernature.com

General enquiries: info@biomedcentral.com

[Receive BMC newsletters](#)

[Accessibility](#) [Leave feedback](#)

[Manage article alerts](#)

[Press center](#) [Careers](#)

493

[Language editing for authors](#)

[Scientific editing for authors](#)

Follow BMC




By using this website, you agree to our [Terms and Conditions](#), [Your US state privacy rights](#), [Privacy statement](#) and [Cookies policy](#). [Your privacy choices/Manage cookies](#) we use in the preference centre.

SPRINGER NATURE

© 2023 BioMed Central Ltd unless otherwise stated. Part of [Springer Nature](#).

Fig. B1 Supplementary map of habitat types and density of wild boar GPS-fixes in Tofte Skov

494 Appendix B Supplementary map



495 Appendix C Included articles

Article

Effect of Wild Boar (*Sus scrofa* L.) on Forests, Agricultural Lands and Population Management in Lithuania

Arūnas Tarvydas and Olgirda Belova *

Institute of Forestry Lithuanian Research Centre for Agriculture and Forestry, Liepu Str. 1, Girionys, LT-53101 Kaunas, Lithuania

* Correspondence: olgirda.belova@lammc.lt

Abstract: The wild boar (*Sus scrofa* L., 1758) is one of the most common and widespread game species not only in the Baltic Sea region but also throughout its entire range. However, quantitative population management is required due to the danger of contagious diseases; the census is imprecise, and integrated management and monitoring are still missing in Lithuania. This study focused on the current condition of the wild boar population at the country level, species activity in forests and adjacent agricultural lands, the problem of damage to agriculture and forestry caused by wild boar, and methods for the management of the wild boar population. A methodology for the assessment of the effect of wild boar will help in reducing their impact. For the successful management of wild boar, an increase in the carrying capacity in conformity with animal density should be accomplished.

Keywords: wild boar; habitats; effect; damage; assessment; management



Citation: Tarvydas, A.; Belova, O. Effect of Wild Boar (*Sus scrofa* L.) on Forests, Agricultural Lands and Population Management in Lithuania. *Diversity* **2022**, *14*, 801. <https://doi.org/10.3390/d14100801>

Academic Editors: Friedrich Reimoser, Ursula Nopp-Mayr and Corrado Battisti

Received: 16 August 2022

Accepted: 21 September 2022

Published: 26 September 2022

Publisher's Note: MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Copyright: © 2022 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

1. Introduction

The wild boar (*Sus scrofa* L. 1758) is one of the most common and widespread game species not only in the Baltic Sea region but also throughout its entire range. It is native to forests ranging from Western and Northern Europe and North Africa to India, the Andaman Islands, and China and has been introduced to New Zealand and the United States [1,2]. Wild boars are an important and inherent component of the forest ecosystem in Lithuania.

The Lithuanian wild boar belongs to the central European subspecies *Sus scrofa*. In Lithuania, the spread of wild boars occurred no later than at the end of the early Holocene (i.e., 8–6 millennium B.C.), especially in the third to second millennium B.C [3]. The species was not rare: its bones make up 20–50% of ungulates' bones found at archaeological sites. In the period after World War II, the number of wild boars decreased because of an abundant population of predators (e.g., the wolf, *Canis lupus* L.), poaching, and other human disturbances (e.g., landscape conversion, forest operations, hunting, recreation, etc.). Historical sources indicate that [3] severe winters in the past negatively affected the annual increment of the population and caused the loss of at least a quarter of adults. Nevertheless, abundant supplementary feeding in winter, protection from poaching, a reduction in the predator numbers, and the improvement of habitat conditions (e.g., agricultural land mosaic; increase in the area under oak stands, and other measures to provide food and shelter) were suitable measures that allowed the species to recover and further increase in number. The behavioural plasticity of wild boars has allowed them to adapt to the feeding conditions of the anthropogenic landscape [4–6]. Consequently, the animals have gathered in the most suitable localities [4,6], causing a clumped distribution and inducing changes in their territorial and feeding behaviour while also becoming a prime risk factor for the spread of contagious diseases in a short time [7–16]. The distribution, movement, grouping, and abundance of wild boars depend on several key drivers, such as the available food supply, safety, and assurance of reproduction. When the animals' needs are unsatisfied [5,17–19], their adaptation to changes in their environment is disordered, and feeding requirements prevail [5,19,20]. Simultaneously, the damage caused to agricultural crops and forestry

increases [19–31]. Despite their omnivorous and highly opportunistic diet, wild boars mostly consume vegetative food, and a better food supply is found in deciduous and mixed forests [19,31,32]. Although wild boars are inseparable from forests, their effect on forests is not exceedingly significant, and their rooted area comprises only 0.9–2.9%, regardless of the high density of wild boar (e.g., density ranges from 15 to 82 individuals per 1000 ha) [33]. Wild boar rooting facilitates the sprouting and growth of tree seeds; moreover, the animals consume insects and rodents [32–35] and stimulate the decomposition of organic matter and the mobilization of soil nutrients [35]. Therefore, they have an important function in assisting nutrient transfer in native forest habitats [36–38]. Although wild boar rooting in forests reduces carbon (C) stocks in the forest soil, they have a negligible effect on large-scale emissions [37,38]. Wild boar mainly roots in older forests, and their rooting has little effect on the germination of seeds via the scarification of the topsoil layer [39]. One study found that rooting occurred in <1% of the area and caused negligible damage to the roots of trees with commercial value [39]. However, other authors have emphasized the considerable risk of damage to seedlings, especially during the first week after planting, with most damage occurring during the first four weeks [40,41]. Wild boars can be described as ecosystem engineers because they mechanically change their environment [42]. It has been found that the most damaged tree species are pine (*Pinus* spp.) or spruce (*Picea* spp.) [39–41] and hardwood species [41–46]. The damage caused by wild boar to seedlings in young forest plantations is site dependent and likely influenced by a combination of food availability, seedling accessibility, animal density, land cover, hunting pressure, and soil moisture [43]. Wild boar can also negatively affect forest litter (losses reaching 80–95%) [32]. Moreover, they contribute to the disappearance of rare plant species and damage tree roots, which cause slope erosion [44,45].

Wild boars cause hundreds of thousands of euros worth of damage [47–53] to almost all agricultural crops [20,21,28,45–47,54,55]. In Luxembourg, the damage caused by wild boar increased by 1600% from 1971 to 2004 [22]. The damage to crops differs in both space and time and is dependent on specific plant growth stages [22,47,48,54,56]. Wild boars spend less time in cultivated fields in March, April, and May than in the summer [48]. In mixed agricultural–forest landscapes, wild boars mostly use fields adjacent to forests [48,49,54]. Many authors have emphasized the severity of crop damage not only due to consumption but also due to trampling (e.g., [22,28,47]). Conflicts among landowners, hunters, and other stakeholders regarding wild boar have also increased. Damage assessment is one of the foremost objectives when addressing human/wildlife conflicts [25,46,50,53,57–60]. A methodology for the assessment of damage to agricultural crops, livestock, and forests caused by game animals, including wild boar, was approved in Lithuania in 2002, and a consolidated version has been valid since 20 October 2018 [59]. This methodology helps to mitigate the above-mentioned conflicts. It regulates the calculation of the monetary expression of the damage caused by wildlife to agricultural crops, livestock, forests, and hydraulic equipment for the owners, users, and managers of agricultural and forest lands and water bodies and lays out measures to protect crops against damage. Wild boar management techniques have been suggested in many countries, including the USA and Australia [61,62]. Problems caused by wild boar and the need for an assessment methodology and compensation have also been emphasized in different countries [40,59,63,64]. In eastern Germany, the needs of stakeholders were identified, and an integrated framework that leverages the individual strengths of previously isolated methodologies was proposed [65]. To assess the damage to agricultural crops caused by wild boar, unmanned aerial systems (drones) [66–68] are commonly used, or the assessments are based on surveys and questionnaires [40,69].

The overabundance of wild boar is a global issue of growing interest. To manage their numbers properly, knowledge of the density of local populations and any changes is needed. Density is an important indicator of animal living conditions and can be used as a tool for their monitoring, risk assessment, and management [19,28,54,70–78]. Various methods and models have been used to estimate density [70,71], including snow tracking, hunting-bag

data, drive hunts, pellet group counts, camera trapping, etc. A common practice with a long history is the counting of wild boar tracks in snow. Briefly, straight-line transects are located using a random or systematic sampling design and the location of each track or trail that crosses a transect is recorded [28,79–81]. However, snow tracking is available only for a short time, and snow conditions are often unsuitable for tracking [19,28,70,79]. Moreover, climate warming and drastic fluctuations in wild boar numbers due to contagious diseases [8–11] have encouraged the utilization of proper methods to estimate the density of wild boar populations. For example, in Lithuania, the wild boar population declined by 2–2.5 times from 1967 to 1990 due to outbreaks of classic swine fever (CSF) [82]; however, it recovered rapidly. The permissible quotas of wild boar densities were approved in 1995 [83] on the grounds of damage caused to agricultural crops and forests and the susceptibility of wild boar to contagious viral diseases. However, the quotas were cancelled in 2005 by the Ministry of Environment [84]. This shows that attention to wild boar management was abated in the absence of the direct emergency associated with zoonotic diseases. The abundance of wild boar has again exceeded permissible levels, and their environmental impact and the spread of contagious diseases [19,82,85] have become a challenge at the global level.

This situation causes great concern and questions about the further management of local populations of wild boar. Although the damage to landowners caused by wild boar is less visible today, the current situation encouraged us to present a long-term study on the effect of wild boar on agricultural crops and forests. We aimed to determine the occurrence of wild boar and assess the damage caused to forests and agricultural crops in different natural regions.

2. Materials and Methods

The study was conducted from 2010 to 2020 in different natural regions, including the model territory (MMMPV), with a total area of 5646 ha in the north-western part of Lithuania (Figure 1).

We employed an integrated method to assess the effect of wild boar on agricultural crops and forests as well as to determine animal distribution and habitat preferences. We estimate densities from indirect signs combining indirect snow-track surveys in the winter (in the presence of snow cover, considering the number of the ingoing and outgoing tracks both ways from the transect line within the grid of six parallel transects; the grid covered all forest stand) and counts of faeces taken within belt transects [31,70] in the early spring that was performed during the count of ungulates (belt transect unit 100 m × 2 m; each transect crosses all habitats and is established in parallel with the forest compartment line; habitat is characterized every 100 m) by pellet group counting (PGC + FGP) [6,28,31,85–90]. All counts were performed during the daytime. The density was calculated using the standard population density formula $D_p = N/A$, where N is the total number of individuals and A is the land area unit (1000 ha), considering the average defecation rate of the wild boar 4.95 for Lithuania [6]. The total number of counters was 7–12. The number of wild boars in the entire area of Lithuania was obtained from each hunting ground unit (HGU) ($n = 911$). Considering the delineation of territory for game animals [6,19,85,91], forests were divided into four categories as follows: (1) pure pine forests, including the maritime zone (western and southern Lithuania; other species comprise less than 10%; the study area 2736 ha and 5519 ha, respectively); (2) pine/spruce mixed forests (eastern and central Lithuania; deciduous species comprise 11–25%; the study area, 7505 ha); (3) mixed spruce/deciduous forests (deciduous species comprise 11–50%; the study area 3234 ha); and (4) deciduous forests with a mixture of spruce species (coniferous species comprise less than 10%; the study area 1043 ha), which were distributed in almost all natural regions. The data collected from the field studies and official statistics sets were stored in a database.



Figure 1. Location of Lithuania on a map of Europe (dark green colour) ($55^{\circ}10'24.96''$ N; $23^{\circ}53'41.29''$ E) and the model territory ($56^{\circ}01'54.9''$ N $21^{\circ}53'20.4''$ E (MMMPV) (red—borders of model territory and HGU names).

To estimate the number and age/sex structure of the local wild boar populations, we have used direct counting at baiting points reported during stand hunting, and camera traps, in the model territory. The camera traps were placed at the baiting points ($n = 10$). This method was combined with the tracking of wild boar herds, which stay outside of the attractive zone, using thermal imaging monoculars. To assess the age/sex structure of local wild boar populations, we distinguished three animal categories: (a) males older than 1 year; (b) females older than 1 year; and (c) juveniles aged up to 1 year. The total number of detected wild boar herds was 211. The herd index was defined on the grounds of the annual mean long-term recruitment, which reached 60%, considering species-specific susceptibility to diseases and mortality, differences in the age-related reproduction of females, and long-term observations [19,31]. To calculate herd indices for practical use, the summation of all observed animals in herds is divided by the number of herd observations in a certain territory.

The occurrence of wild boar and the damage caused to agricultural crops depending on animal density were studied during the summer–autumn period within a radius of 1 km from the forest edge. Belt transects were established every 100 metres from the forest edge for the whole agricultural crop area. The size of each belt transect unit was 0.1 ha ($100\text{ m} \times 10\text{ m}$). The number of transects depended on the length of the area. The total length of the transects was 920 km of the total area (921 ha). The intensity of the damage (i.e., “trampled”, “consumed”, etc.) to agricultural crops in the belt transects was estimated visually using a five-point scale as follows [85,92]:

- 0—no damage (intact);
- 1—sporadic damage, 5%;
- 2—less than half of all areas damaged, 30%;
- 3—more than half of all areas damaged, 75%;
- 4—all areas damaged, 100%.

500

The abundance of each agricultural crop (A) in their rotation was calculated using the formula below:

$$A = n \times 100/N\%$$

where n is the number of sample belt transects where each crop was found, and N is the total number of sample belt transects in all agricultural crops.

The average intensity (I) of the damage to each agricultural crop was calculated using the following formula:

$$I = i/N\%,$$

where i is the sum of the percentage of damage to each agricultural crop, and N is the total number of sample belt transects where each agricultural crop was found.

The index of the consumption (i.e., direct effects to plant tissues including mechanical damage) of each agricultural crop (trampled, eaten up, etc.) was calculated using the formula below:

$$C = A \times I\%,$$

The share of each agricultural crop consumed by animals was calculated using the following formula:

$$S = C \times 100/\Sigma c$$

where C is the consumption index of each agricultural crop in % and Σc is the sum of the percentage of the consumption indices of all agricultural crops.

The percentage (i.e., 5%, 30%, or 75%) of differently damaged agricultural crop area (D) was calculated using the following formula:

$$D = n \times 100/N\%,$$

where n is the number of belt transects, where the damage was 5%, 30%, or 75%, and N is the total number of sample belt transects.

The differently damaged (i.e., 5%, 30%, or 75%) agricultural crop area in hectares was calculated using the formula below:

$$D_{ha} = D_t \times D/100\%,$$

where D_t refers to the total area of each agricultural crop in hectares and D is the area of each agricultural crop differently damaged (5%, 30%, or 75%) by animals in hectares.

To study the wild boar diet, we analysed the stomach contents of 200 harvested wild boars. We collected 500 mL samples from the stomach and stored them in 5% formalin. The contents were washed with water in 1.7 mm and 5 mm mesh sieves. The food content of the samples was analysed using the standard volumetric method. We calculated and recorded the percentage by volume of each food item by displacing water in a volumetric beaker [91–93].

We assessed the effect of wild boar on the yield of different agricultural crops using a plot sampling method. The size of each sample plot unit was 1 m². The sample plots were established directly within the belt transects where crops were differently damaged (i.e., 5%, 30%, 75%, or 100%). Simultaneously, the control plots were established in areas with undamaged agricultural crops. The difference between the cereal mass of the control and sample plots allowed us to calculate the amount of trampled and consumed yield separately. This method enabled us to correctly ascertain the extent of the losses and predict certain preventive and precautionary means for the restoration and maintenance of the balance between animal numbers and the carrying capacity of their habitats.

Research outputs are the ground for population management. The quantitative method for the management of wild boar populations is based on the loss of agricultural

501

crops, population size, and annual recruitment. The ratio of population use is calculated using the following formula [47,60,85,94]:

$$N = IN + (RN - PN)/y,$$

where N refers to the ratio of use; IN is the wild boar annual recruitment in animal units; RN is the actual number of animals in the territory; PN is the permissible number of animals in the territory (the number of animals approved for the territory based on the category of the forests and their corresponding carrying capacity); and y is the number of years needed to recover the balance between the animal number and carrying capacity of the territory.

Data Analysis

All of the analyses were performed using MS Excel spreadsheets and the Statistica 8.0 software packages [95]. The relationships between variables (damage—density of wild boar, density—herd index, etc.) were determined using linear regression analysis. The significant differences between the means were determined using one-way ANOVA. The significance of the difference between the means of the obtained data was evaluated using Student's t -test, where $p < 0.05$ was considered significant.

3. Results

3.1. Wild Boar Distribution and Foraging in the Different Habitats

The distribution of wild boar depends on food availability; human disturbances, e.g., hunting intensity; and seasonal changes ([11], et al.). Wild boars are most abundant in areas with high soil fertility [6,19] in the northern, southwestern, and northern central parts of the territory of Lithuania.

Although wild boars are omnivorous animals, plant fodder predominates their diet, comprising almost 97%. In forest rooting sites, the ratio of plant food to animal food was 3:1, showing the dominance of plant food. The average shares of the rooting sites (% of the total study area) and foraging supply of different origins (plant or animal) for the different categories of forests are shown in Table 1. The total share of forest food was only 4% while agricultural crops were available. The analysis of the rooting sites showed that the rooting character depended on the forest stand parameters and forest sites ($r = 0.589$, $p \leq 0.05$).

Table 1. Average seasonal food supply of different origin (plant or animal) and the share of rooting sites in different forest categories in Lithuania.

Forest		Spring			Summer			Winter	
Category	Rooting	Food Supply		Rooting	Food Supply		Rooting	Food Supply	
	Site	g/1 m ²		Site	g/1 m ²		Site	g/1 m ²	
	%	Plant	Animal	%	Plant	Animal	%	Plant	Animal
Pure pine	0.1	0.1	27	0.3	0.2	2.3	0.003	2	2.8
Pine with spruce	2.1	144	7	0.8	22	15.2	0.01	237	0.4
Mixed spruce									
deciduous	0.5	144	14	0.3	155	15.6	0.09	303	0.3
Deciduous with									
spruce	1.0	320	31	1.3	204	28.5	0.09	299	-
Natural meadow	3.6	475	37	x	x	x	0.2	128	0.5

In pure pine forests, there was a low food supply, except for *Alnetum filipendulosum* black alder stands, where earthworm stores of up to 100 g per 1 square meter could be found. In spring, the rooted sites most often occurred in *Betuletum vaccinio-myrtillousum*, *Tremuletum vaccinio-myrtillousum*, and *Alnetum filipendulosum* birch and aspen stands in territories belonging to the pure pine forest category, while the area of rooted sites was the largest in the forest sites containing *Pinetum vacciniosum* and *Pinetum vaccinio-myrtillousum*.

In summer, when mushrooms start growing, the rooted sites were predominant in *Pinetum vacciniosum* and *Pinetum cladoniosum* pine stands. These were the largest rooted sites. In winter, the largest rooting area was found in *Alnetum filipendulosum* black alder stands, where food consumption fluctuated from 65% to 100% (Table 1).

The stands belonging to the pine with spruce forest category provided more plant food for the wild boar. The plant forage store amounted to 500–700 g per 1 m² of the rooted sites in *Pinetum myrtillo-oxalidosum* and *Piceetum myrtillo-oxalidosum* pine and spruce stands and in *Alnetum urticosum* and *Alnetum caricosum* black alder stands. Food consumption ranged from 57% to 100%. (Table 1). In spring, the greatest rooting was observed in *Alnetum caricosum* and *Alnetum urticosum* black alder stands and *Pinetum vaccinio-oxalidosum* and *Pinetum oxalidosum* pine stands, while the largest rooted sites occurred in *Pinetum oxalidosum*, *Pinetum vaccinio-myrtillousum*, and *Pinetum myrtillo-oxalidosum* pine stands and *Piceetum oxalidosum*, *Piceetum myrtillo-oxalidosum*, and *Piceetum vaccinio-myrtillousum* spruce stands. In summer, the most rooting occurred in *Alnetum urticosum*, *Piceetum oxalidosum*, and *Pinetum oxalidosum* black alder, spruce, and pine stands, whereas the largest rooted sites were found in *Pinetum oxalidosum* and *Piceetum oxalidosum* pine and spruce stands. In winter, more rooted sites occurred in the *Alnetum caricosum* black alder stands of pine with spruce forests, whilst the largest rooted sites were found in the *Alnetum caricosum*, *Pinetum myrtillousum*, *Pinetum myrtillo-oxalidosum*, *Piceetum myrtillousum*, and *Piceetum myrtillo-oxalidosum* black alder, pine, and spruce stands of the pine with spruce forests.

In the stands of mixed spruce/deciduous forests, the food supply was similar to that of the stands of the pine with spruce forest category, with the exception of *Myrtillo-oxalidosa*, *Urticosa*, and *Caricosa* forest sites with deciduous stands, where the foraging supply ranged from 230 to 930 g/m². Food consumption reached 32–100% (Table 1). In spring, the rooted sites were mostly found in *Alnetum caricosum* black alder stands and *Piceetum oxalidosum* spruce stands, and natural meadows. The largest rooting areas were found in *Piceetum oxalidosum* spruce stands and *Aegopodiosa* and *Oxalidosa* forest sites with deciduous stands. In summer, the wild boars mostly rooted in *Piceetum oxalidosum* spruce and *Alnetum urticosum* black alder stands, while the largest rooting area occurred in *Piceetum oxalidosum* spruce and deciduous stands. In winter, rooting was observed mostly in the natural meadows and in *Oxalidosa-myrtillosa* and *Oxalidosa nemorosa* forest sites with deciduous stands. The largest rooting areas were found in *Piceetum aegopodiosum* spruce stands, natural meadows, and *Myrtillo-oxalidosa* and *Caricosa* forest sites with deciduous stands.

In the deciduous spruce forests, the food supply was the greatest (Table 1), and food consumption comprised 50–90% [3,28]. In spring, the most rooting occurred in natural meadows, *Carico-nemorosa* forest sites with deciduous stands, and *Alnetum urticosum* and *Alnetum filipendulosum* black alder stands, while the largest rooted sites were found in *Carico-mixtoherbosa* forest sites with deciduous and mixed stands. In summer, rooting mostly occurred in *Alnetum urticosum* and *Alnetum filipendulosum* black alder stands and in *Carico-mixtoherbosa* forest sites with deciduous spruce stands, while the largest rooted sites were found in *Carico-mixtoherbosa* forest sites with deciduous stands, *Alnetum urticosum* and *Alnetum filipendulosum* black alder stands, and *Piceetum carico-mixtoherbosum* spruce stands. In winter, wild boar mostly rooted in *Alnetum urticosum* black alder stands, while the largest rooting areas were found in *Alnetum urticosum* black alder stands and *Piceetum nemorosa-oxalidosum* spruce stands.

Agricultural crops comprised a major part of the plant food consumed by wild boar (ca. 81%). The largest shares of agricultural crops were consumed in the summer (ca. 90%) and autumn (83%). The available residues of crops in fields were found to attract animals and still prevailed in their diet during the winter (ca. 79%) and winter–spring (ca. 66%) periods. Natural plant fodder comprised, on average, 17% of the animals' diet. This food was important in the spring–summer period, but its importance decreased in winter and autumn [2,6,96]. Above-ground food predominated in the animals' diet, comprising ca. 57% of their diet, and was particularly prevalent in the summer (more than two-thirds). The share of the underground food reached 40%. Cereal comprised 47% and potatoes 53%

of their annual diet. The third most preferable food after cereal and potatoes was beets (ca. 12%). Beets were rarely found in the stomach samples from the summer period, but their importance as supplemental fodder increased during autumn and winter. Green plants were found in 85% of wild boar stomachs; however, the share of this food was only found to be nearly 6% of the diet. A frequently observed food was different roots and bulbs, the share of which was 5%, while the frequency of occurrence was approximately 77%. Wild boar preferred the roots of plants such as *Pteridium aquilinum*, *Urtica dioica*, *Asarum europaeum*, *Oxalis acetosella*, *Anemone nemorosa*, *Ficaria verna*, *Caltha* sp., *Phragmites* sp., *Typha* sp., *Ranunculus* sp., and small shrubs, among others. Berries and fruits accounted for 1.5–2% of their annual diet. Another important food was acorns. This kind of food comprised 10–40% of their diet during the productive time of year and about 1.5% of their annual diet. Needles, leaves, and moss were observed less frequently. These types of food comprised 2.5% of the animals' diet in the winter and up to 0.2–0.4% during other seasons. Other plant foods, such as the roots and shoots of trees and shrubs and mushrooms, were rarer in the wild boar diet (0.4–0.7%). Foods of animal origin comprised only 3–4% of their diet; this share was negligibly more during the winter (5%), when small rodents and different carrion predominated than in the spring–summer period (3%). Earthworms were the main contributor to foods of animal origin. Different insects, including the larvae of the cockchafer (*Melolontha hippocastani* L.), flea-beetles (Elateridae), the chrysalides of sawflies (*Lyda nemoralis* Thoms., *Diprion pini* L., *Neodiprion sertifer* Geoffr., *Gilpinia polytoma*), and others, comprised a lower share of the wild boar diet. In spring and autumn, the share of food of animal origin was similar (ca. 2%).

As mentioned above, wild boars inhabit all forests; however, coniferous forests and spruce stands were found to be preferred in this study. Deciduous forests are more important during the summer, not only for foraging but also for bathing—there are a number of suitable bathing points in these forests. Coniferous forests protect wild boar against precipitation (i.e., snow and rain). Many wild boar dens were found under older spruces. Dens were rarely established in the undergrowth of spruce, under pine, or in the pine plantations, while several dens were found in the cuttings and under the roots of fallen trees. The dens were usually found in the southern, southeastern, or southwestern parts of trees (41%) and were exceedingly rare in the western part of trees (6.3%) because of the predominating western winds. In winter, most dens (60%) were covered by spruce twigs or sometimes by juniper, pine, and moss (14.5%) or herbs (1.4%). The dens without any bedding were considered temporary (ca. 23% of cases) [3]. During the warmer weather, the dens had no bedding except those of the females with sucking piglets.

In Lithuania, the largest herds of wild boar occurred in the spring. The average herd size was ca. six individuals; however, there could be up to twenty-two animals at the maximum. Generally, the females stayed with the juveniles, and separate herds sometimes joined. During summer and the beginning of autumn, the average herd size decreased to up to five animals. Simultaneously, more abundant herds of up to thirty-five individuals were also found. After the weather became cooler, the animals gathered into larger herds. The largest herds were observed in the winter, with up to twenty-seven animals, while their average size was 5.5 animals.

Although the wild boars were settled when the food supply was sufficient and the animals were not disturbed, they could move about 10–20 km per day because of human disturbances (e.g., four-wheel motor vehicles). Longer movements were characteristic of males. The average movement distance of wild boar is from 0.3 km to 10.5 km depending on the local conditions, such as the season, animal age, and sex [6,31,96,97]. The average herd of wild boar moves about 1–2 km in the wintertime and up to 10–12 km during the summer [96,97].

3.2. Effect of Wild Boar on Forests and Agricultural Crops and Management Issues

Wild boars visit the agricultural lands surrounded by forests depending on crop rotation and animal density. One individual can root 1 ha in the forest, ravage 0.3–0.4 ha

of crops from milky ripeness to harvest, and damage other crops yearly. The damage to agriculture caused by wild boar comprises 70–75% of total damage to agriculture; in comparison, damage by deer amounts to 25–30%. Wild boar trample, root, and consume winter and summer cereals, root crops, potatoes, damaged caraway, and other available species depending on their occurrence.

The total annual (excepting autumn) area of forest litter rooted by wild boar was, on average, 0.4% in pine forests, 2.9% in spruce/pine forests, 0.9% in mixed spruce/deciduous forests, and 2.4% in deciduous forests. The damage caused by wild boars was strongly and positively related to animal density ($r = 0.59$, $y = -0.140 + 0.120x$; $p < 0.001$), and stronger damage was observed in winter while less was found during summer ($r = 0.88$, $p < 0.05$).

The intensity of damage to rye and wheat was weakly but positively related to the distance from the forest. Animals are safer away from the forest in comparison to its edges because of possible disturbance factors (hunting, recreational activities, etc.). Wild boar can settle there for a long time. Similar but negative relationships were found between the damage caused by wild boar and potato and beet fields. The intensity of damage to oats was the greatest and was negatively related to the distance from the forest edge ($r = -0.80$, $p < 0.04$). The damage to potatoes and sugar beets was negatively related to the distance from the forest edge ($r = -0.48$ and $r = -0.41$, respectively; $p \leq 0.05$). The animals damaged more plants in smaller oat, potato, and beet fields. The principal factor of damage intensity was animal density (number of individuals per 1000 ha; $r = 0.59$; $p \leq 0.05$). The animals mostly trampled crops (ca. 80%), and less damage was due to the consumption of cereals and root crops (ca. 20%). Therefore, the damage caused by wild boar considerably changed the structure of the agricultural crops considered.

When the density of wild boar exceeds 100 animals per 1000 hectares (i.e., up to 115/1000 ha), the animals visit 23–52% of the agricultural crops within a radius of 1 km. At a density of 12 animals per 1000 ha, wild boars attend 6% of the agricultural crops, and when the density is only 6/1000 ha, this value is decreased to 3% [3,6,28].

4. Discussion

4.1. Effect of Wild Boar on Forests and Agricultural Crops

In Lithuania, foresters and game managers have long dealt with the problem of restraining and reducing the damage caused by wild boars, despite the considerable decline in the wild boar population since the first detection of African swine fever (ASF) in 2014. Today, ASF continues to expand its range in Lithuania, despite efforts to eradicate it and implement national disease control measures, including intensive hunting at the national level and in the model territory (MMMPV) (Figure 2).

Effective protection and a decrease in losses are possible if the reasons for wild boar damage and the factors necessary to improve methods for damage prediction and assessment are perfectly understood. Nevertheless, the assessment of the damage caused by wild boar and management strategies for practical application have not been sufficiently presented [40,50,51,59,63,64,98,99]. A methodology for the assessment of the damage caused by wild boar to agricultural crops and forests was included in the Wild Boar Management Programme, a legal act on the damage caused by wildlife to agricultural crops, facilities, and forests [19,100] in Lithuania, and is currently used to estimate crop losses.




Figure 2. Changes in wild boar harvesting and density in the model territory and entire country.
Note: MMMPV is the abbreviation of the model area.

4.2. Management of the Wild Boar Population

The management of local wild boar populations depends on wild boar numbers and their effect on agriculture and forestry. There are different management plans for wild boar populations [101–112]. Wild boar populations are managed territorially, quantitatively, and qualitatively [6,19,28,85]. The essence of territorial management is harvest planning and the regulation of numbers in the local wild boar populations despite any administrative borders, such as a region, forest enterprise, or single forest. The local wild boar population

should be the object of management. This approach provides scientific ground for the use of the population when considering the carrying capacity of a certain territory and maintaining an optimal population density. Territorial management is performed in forest stand complexes with an area from 5 to 10 thousand hectares when considering the size of the home range and seasonal migration of wild boars [27,30,84].

Direct observations are important measures for the qualitative management of wild boar. This method is suitable when considering the age/sex structure of the population. The scientifically grounded count method is necessary for the successful management of wild boar populations. The combination of several count methods can be used, for example, the count of animals during drive hunting; at baiting points; by faecal group (FPG), which can be combined with the usual deer count by pellet groups at the beginning of the growing season; or via the camera trapping method [68,71,72,113]. To manage local populations properly, an increase in the carrying capacity of the habitat and its conformity to animal density should be accomplished. In this respect, three categories of wild boar densities were distinguished in Lithuania [28,31,85]:

- Minimum density, when animal distribution is random, forage consumption is low, and the negative impact on the habitat is intangible. At this density, infection by parasitic diseases is not intensive, and the population size increases; immigration is common, while emigration is negligible;
- Permissible density, when the distribution of wild boar is spaced, forage consumption does not exceed the permissible limits, and the negative impact is insignificant;
- Ecological density, when animal distribution is clumped in favourable habitats. Here, the forage demand conforms to forage resources.

The permissible quotas for wild boar and other ungulates in the different forests were already legally approved (Table 2). The group of food supply is officially approved according to the forest category in Lithuania [114] (hunters pay taxes for the use of wildlife resources in each hunting ground unit HGU based on the group of food resources of their HGU).

Table 2. Permissible quotas for the local wild boar population density in the different forests [85,99,114].

Forest Category	Group of Food Supply	Density in All Forests		Wild Boar Density, n/1000 ha		
		Permissible	Ecological	Protective Forests	Managed, Recreational	Commercial Hunting
				Agroforests	Forests	Grounds
Pure pine (dominant forest sites: <i>Cladoniosa, Vacciniosa</i>)	IV	4–7	5–10	4	7	≥15
Pine with spruce (dominant forest sites: <i>Vacciniosa-myrtillosa, Myrtillosa</i>)-	III	6–11	10–15	6	11	≥20
Mixed spruce/deciduous (dominant forest sites: <i>Oxalidosa, Oxalido-myrtillosa</i>)	II	8–14	10–15	8	14	≥30
Deciduous with spruce (dominant forest sites: <i>Aegopodiosa, Carico-mixtoherbosa</i>)	I	10–15	15–20	10	15	≥30

Note: Variations in soil conditions were observed in *Piceetum myrtillo-sphagnosum*, *Piceetum oxalido-myrtillosum*, and *Piceetum asperulo-tiliosum* (+ = to).

The density quotas must correspond to the local conditions and consider the limiting ecological factors. In addition, the damage caused by wild boars to agricultural crops may not exceed 3–5%. The differences between the density of wild boar and harvesting in the model territory and the whole country are shown in Figure 2. The permissible density of wild boar should be kept in protected areas, forests used for recreational purposes, and agroforests. Ecological density would be suitable for commercial hunting ground units.

In these forests, wild boar density could exceed the ecological density depending on the foraging supply.

The qualitative management of the wild boar population is based on population structure, as shown in Table 3. The local wild boar population structure in Lithuania also depends on the feeding conditions during winter.

Table 3. Age and sex structure of the wild boar population in different forest stand categories.

Stand	Population Structure					
	Herd Index, av	Males Older than 1 Year, %	Females Older than 1 Year, %	Sex Ratio	Piglets below 1 Year of Age, %	Recruitment Coefficient
Pure pine	3.9	28.3 ± 3.8	27.6 ± 2.7	1:1.2	48.6 ± 4.1	0.9
Pine with spruce	4.9	18.7 ± 1.9	30.8 ± 1.2	1:1.6	50.5 ± 1.8	1.0
Mixed spruce/deciduous	5.5	16.4 ± 2.2	28.6 ± 1.9	1:1.7	55.0 ± 3.3	1.2
Deciduous with spruce	4.9	18.1 ± 1.3	33.7 ± 2.7	1:1.9	48.2 ± 3.1	0.9

If baiting during the winter was insufficient, the recruitment coefficient was only 0.9, while under the conditions of sufficient abundant feeding in the winter, it reached 1.7.

Until now, the herd index was not discussed in scientific publications, while it was emphasized several decades ago in Lithuania. However, its importance has emerged during the last decade in the context of the spread of ASF. As the herd index was not considered, the number of harvested animals exceeded 200–400%. A model of the relation between wild boar density and herd index can be expressed as follows: $y = -30.4 + 12.1x - 0.3x^2 - 0.03x^3$, where x is the herd index ($r = 0.89$, $t = 6.12$, $p \leq 0.05$). We recommend considering this useful parameter for wild boar management. The herd index for the different natural regions was determined as follows: 2–3 for pure pine forests in southern Lithuania and 3–4 for the same forests in western Lithuania; 3–4 for mixed coniferous forests in eastern Lithuania and 4–5 for the same forests in central Lithuania; 5–5.5 for mixed spruce/deciduous forests; and 5 for deciduous forests with spruce admixture, including a herd index of 4–5 for the same type of forests in northern Lithuania [19].

In pure pine forests, the herd index was 2–3, and the actual density of wild boar reached 27/1000 ha in 2013; however, because of ASF, it was reduced to 7/1000 ha in 2016, and long-term density stands at only 5/1000 ha.

In stands belonging to the pine with spruce forest category, the herd index was 3–4 in eastern Lithuania and 4–5 in the central part of the country.

In the stands of mixed spruce/deciduous forests, the herd index reached 5–5.5. Wild boar density was reduced to 14–22/1000 ha (2–4 times) because of the general measures against ASF. However, it should be less—around 8–12/1000 ha—considering the carrying capacity of these territories, the herd index, and the risk of ASF.

In the stands of deciduous forests with spruce, the average herd index was 5; however, it was less (4–5) in northern Lithuania.

To ascertain the carrying capacity of territories, game managers should consider the following criteria:

1. Forest species composition and forest habitats in the whole territory settled by local wild boar populations.
2. The distribution of the food supply.
3. The volume of losses to the agricultural crops and forests caused by wild boars.

The long-term annual recruitment of the local wild boar population in Lithuania is 50–60%, while the annual recruitment coefficient is 1.0–1.5. If animal foraging in winter is insufficient, a harvest volume of 80–100% is allowed. Following variable winters with severe and warm periods, harvesting should be reduced by up to 60–70%, which will allow the annual recruitment coefficient to decrease to 0.6–0.7 despite the high reproductive potential of wild boar [115]. To avoid sudden changes in animal numbers, the sex ratio of males to females should be 1.5:1.0. With respect to hunting, the share of juveniles below

1 year should be 70–80%, 15–20% for sub-adults, and 5–10% for adult animals. We must consider not only the high reproductive potential of wild boar but also their biological plasticity [116,117], sociality [4,118], and adaptability; damage to agriculture and forestry; habitat suitability; and other ecological features.

Traditional methods for wild boar hunting include drive hunting, excluding localities of ASF outbreaks, and stand hunting; stalking and chase are also popular, depending on the season. In Lithuania, before the detection of ASF in 2014, the hunting of males and second-year juveniles was allowed from 1 May to 1 March, and the hunting season for females was from 1 October to 1 March. The number of harvested animals was similar over 5 years (Figure 2). A total of 10.3 thousand animals were harvested in 1998/1999, while 10.84 thousand animals were harvested in the 2003 hunting season. However, this number reached 35.56 in the 2017/2018 season, representing an increase of 231.5%. It would be unrealistic to hunt more than 200 percent of the population. What is the reason for such an increase? All harvested animals and their number during the last hunting season must be reported until 5 April annually [114]. However, we recognized that hunters do not consider recruitment of the local population and the region-specific herd index. Since 2015, wild boar hunting has been allowed year-round. Maintaining the minimum permissible density depends on the category of the forest under consideration, i.e., pure pine, pine with spruce, mixed spruce/deciduous, or deciduous with spruce forests, and its functional purposes; its value can fluctuate from 4 to more than 30.

Wild boars are not welcome in agroforests of less than 300–500 ha. Near the forest edges, less preferable agricultural crops are sowed, and cultural pastures are established. In addition, in regions with the most damaged agricultural crops, hunter lookout towers have been established for hunting and frightening away wild boar. Moreover, electric fencing is also used to frighten away wild boar. These means are most notably used by landowners.

5. Conclusions

There is marked evidence of the importance of wild boar in Lithuanian game management, forestry, and agriculture. Considering the damage caused by wild boar, it is necessary to maintain the permissible density of the local wild boar population. This density depends on the local conditions of the forests and surrounding areas, in addition to the functional purposes of the land.

The wild boar population should be managed territorially, quantitatively, and qualitatively considering the density of the local population; the carrying capacity of the territory under consideration and the possibility of it increasing; the age-sex structure of the wild boar population; the annual recruitment of the population; and agricultural crop losses in the territory. In particular, we should seek to maintain the permissible density of the local population. Herd index could be a useful parameter to manage populations. The omission of this index causes incompatibility between the actual number of wild boars and the number of harvested animals (i.e., harvest can reach 120% or more, even reaching 400–550% in some localities). The herd index in the different natural regions was determined as follows: 2–3 for pure pine forests in southern Lithuania and 3–4 for the same forests in western Lithuania; 3–4 for mixed coniferous forests in eastern Lithuania and 4–5 for the same forests in central Lithuania; 5–5.5 for mixed spruce/deciduous forests; and 5 for deciduous forests with spruce admixture, including a herd index of 4–5 for the same type of forests in northern Lithuania.

Author Contributions: Conceptualization, O.B.; methodology, A.T. and O.B.; analysis, A.T. and O.B.; resources, O.B. and A.T.; data curation O.B.; writing—original draft preparation A.T. and O.B.; writing—review and editing, A.T. and O.B. All authors have read and agreed to the published version of the manuscript.

Funding: This research received no external funding.

Institutional Review Board Statement: Not applicable.

Informed Consent Statement: Not applicable.

Data Availability Statement: Not applicable.

Acknowledgments: This study was performed within the frames of the long-term research programme Sustainable Forestry and Global Changes, which was implemented by the Lithuanian Research Centre for Agriculture and Forestry. We are extremely grateful to our former Professor Vytautas Padaiga (passed away 24 March 2021) for his invaluable advice.

Conflicts of Interest: The authors declare no conflict of interest.

References

1. Anon. *Sus scrofa* [ISC] (feral pig). In *Invasive Species Compendium. Detailed Coverage of Invasive Species Threatening Livelihoods and the Environment Worldwide*; CAB International Wallingford: Oxon, UK, 2010. Available online: <https://www.cabi.org/isc/datasheet/119688#tosummaryOfInvasiveness> (accessed on 23 August 2022).
2. Spencer, P.B.S.; Hampton, J.O. Illegal translocation and genetic structure of feral pigs in Western Australia. *J. Wildl. Manag.* **2005**, *69*, 377–384.
3. Prūsaitė, J. *Fauna of Lithuania. Mammals*; Moksas: Vilnius, Lithuania, 1988; p. 293. ISBN 5420000547, 9785420000540. (In Lithuanian)
4. Belova, O. *Ethology of Game Animals*; Lututė: Kaunas, Lithuania, 2001; p. 280. ISBN 9955-452-48-X. (In Lithuanian with English Summary).
5. Bieber, C.; Ruf, T. Population dynamics in wild boar *Sus scrofa*: Ecology, elasticity of growth rate and implications for the management of pulsed resource consumers. *J. Appl. Ecol.* **2005**, *42*, 1203–1213.
6. Belova, O. Guidelines on Wild Boar (*Sus scrofa* L.) Population Qualitative, Quantitative and Territorial Management to Reduce Damage Caused to Agriculture and Forestry and to Prevent Contagious Diseases. In *Newest Recommendations for Agriculture and Forestry*; Semaškienė, R., Mikšys, V., Feiza, V., Kadžiulienė, Ž., Lazauskas, S., Ruzgas, V., Samuoliūnė, G., Eds.; LAMMC: Akademija, Lithuania, 2015; pp. 60–62.
7. Aliešiūnienė, A. Wild Boar (*Sus scrofa*) Feeding and Metastrongylus spp. Infection in the Feeding Points and the Main Epizootic Aspects. Master's Thesis, Lithuanian Veterinary Academy, Kaunas, Lithuania, 2010; p. 39.
8. Lange, M. *Alternative Control Strategies against ASF in Wild Boar Populations*; External Scientific Report; EFSA Supporting Publication: Parma, Italy, 2015; p. 29.
9. EFSA. Evaluation of possible mitigation measures to prevent introduction and spread of African swine fever virus through wild boar. Mitigation measures for African swine fever virus in wild boar. *EFSA J.* **2014**, *12*, 3616.
10. EFSA. African swine fever. EFSA Panel on Animal Health and Welfare (AHAW). *EFSA J.* **2015**, *13*, 4163. [[CrossRef](#)]
11. EFSA; SIGMA. *A Comprehensive Animal Disease Data Collection Approach. Harmonized Data Model for Domestic and Wild Animal Populations*; EFSA Supporting Publication: Parma, Italy, 2018; pp. 1–7. [[CrossRef](#)]
12. Risco, D.; Serrano, E.; Fernández-Llario, P.; Cuesta, J.M.; Gonçalves, P.; García Jiménez, W.L.; Martínez, R.; Cerrato, R.; Velarde, R.; Gómez, L.; et al. Severity of bovine tuberculosis is associated with co-infection with common pathogens in wild boar. *PLoS ONE* **2014**, *9*, e110123. [[CrossRef](#)]
13. Sorensen, A.; van Beest, F.M.; Brook, R.K. Impacts of wildlife baiting and supplemental feeding on infectious disease transmission risk—A synthesis of knowledge. *Prev. Vet. Med.* **2014**, *113*, 356–363.
14. Śmiertka, K.; Woźniakowski, G.; Kozak, E.; Niemczuk, K.; Frączyk, M.; Bocian, Ł.; Kowalczyk, A.; Pejsak, Z. African swine fever epidemic, Poland, 2014–2015. *Emerg. Infect. Dis.* **2016**, *22*, 1201–1207.
15. Oja, R.; Velström, K.; Moks, E.; Jokelainen, P.; Lassen, B. How does supplementary feeding affect endoparasite infection in wild boar? *Parasitol. Res.* **2017**, *116*, 2131–2137. [[CrossRef](#)]
16. Maslow, A.H. A Theory of Human Motivation. *Psychol. Rev.* **1943**, *50*, 370–396.
17. Maslow, A.H. *Motivation and Personality*; Harper and Row: New York, NY, USA, 1954.
18. Belova, O. The Main Behavioural Changes in the Herbivorous Game Animals in Managed Forests. *Balt. For.* **1999**, *5*, 64–68.
19. Belova, O.; Tarvydas, A.; Urbaitis, G. Wild boar distribution and habitat preference in Lithuania. In Proceedings of the 12th Intern. Symposium On Wild Boar and Other Suids, Lázně Bělohrad, Czech Republic, 4–7 September 2018; Kramer, J., Drimaj, J., Eds.; pp. 94–108.
20. Grodziński, W.; Maycock, L.; Weiner, J. (Eds.) *Forest Ecosystems in Industrial Regions*; Springer: Berlin/Heidelberg, Germany, 1984; pp. 241–268.
21. Schley, L.; Roper, T.J. Diet of wild boar *Sus scrofa* in Western Europe, with particular reference to consumption of agricultural crops. *Mammal. Rev.* **2003**, *33*, 43–56.
22. Schley, L.; Dufrêne, M.; Krier, A.; Frantz, A.C. Patterns of crop damage by wild boar (*Sus scrofa*) in Luxembourg over a 10-year period. *Eur. J. Wildl. Res.* **2008**, *54*, 589–599.
23. Baubet, E.; Bonenfant, C.; Brandt, S. Diet of the wild boar in the French Alps. *Galemys* **2004**, *16*, 99–111.
24. Cervo, I.B.; Guadagnin, D.L. Wild boar diet and its implications on agriculture and biodiversity in Brazilian forest–grassland ecoregions. *Anim. Biodivers. Conserv.* **2020**, *43*, 123–136. [[CrossRef](#)]

510

25. Amici, A.; Serrani, F.; Rossi, C.M.; Primi, R. Increase in crop damage caused by wild boar (*Sus scrofa* L.): The “refuge effect”. *Agron. Sustain. Dev.* **2012**, *32*, 683–692. [CrossRef]
26. Hegel, C.G.Z.; Marini, M.Á. Impact of the wild boar, *Sus scrofa*, on a fragment of Brazilian Atlantic Forest. *Neotrop. Biol. Conserv.* **2013**, *8*, 17–24. [CrossRef]
27. Honda, T.; Sugita, M. Environmental factors affecting damage by wild boar (*Sus scrofa*) to rice fields in Yamanashi Prefecture, central Japan. *Mammal. Study* **2007**, *32*, 173–176.
28. Padaiga, V. *Biological Essentials of the Game Management*; Žiburio leidykla: Vilnius, Lithuania, 1996; p. 212. (In Lithuanian)
29. Ficetola, G.F.; Bonardi, A.; Mairota, P.; Leronni, V.; Padoa-Schioppa, E. Predicting wild boar damages to croplands in a mosaic of agricultural and natural areas. *Curr. Zool.* **2014**, *60*, 170–179. [CrossRef]
30. Fonseca, C. Winter habitat selection by wild boar *Sus scrofa* in south-eastern Poland. *Eur. J. Wildl. Res.* **2008**, *54*, 361–366. [CrossRef]
31. Belova, O. Long-term changes in the wild boar and population control. In *Newest Recommendations for Agriculture and Forestry*; LAMMC: Akademija, Lithuania, 2018; pp. 52–54.
32. Janulaitis, Z.; Padaiga, V. Effect of wild boar on the forest biogeocoenosis. In Proceedings of the Scientific Conference Wildlife Management in Problem-Oriented Forestry, Kaunas-Girionys, Lithuania, November 1983; pp. 45–46. (In Russian).
33. Mori, E.; Ferretti, F.; Lagrotteria, A.; La Greca, L.; Solano, E.; Fattorini, N. Impact of wild boar rooting on small forest-dwelling rodents. *Ecol. Res.* **2020**, *35*, 675–681. [CrossRef]
34. Torre, I.; Cahill, S.; Grajera, J.; Raspall, A.; Vilella, M. Small mammal sampling incidents related to wild boar (*Sus scrofa*) in natural peri-urban areas. *Anim. Biodivers. Conserv.* **2022**, *45*, 33–42. [CrossRef]
35. Lacki, M.J.; Lancia, R.A. Changes in Soil Properties of Forests Rooted by Wild Boar. *Proc. Annu. Cont. Southeast. Assoc. Fish Wildl. Agencies* **1983**, *37*, 228–236.
36. Wirthner, S. The Role of Wild Boar (*Sus scrofa* L.) Rooting in Forest Ecosystems in Switzerland. Ph.D. Thesis, ETH Zurich, Zurich, Switzerland, 2011. [CrossRef]
37. Wirthner, S.; Schütz, M.; Page-Dumroese, D.S.; Busse, M.D.; Kirchner, J.W.; Risch, A.C. Do changes in soil properties after rooting by wild boars (*Sus scrofa*) affect understory vegetation in Swiss hardwood forests? *Can. J. For. Res.* **2021**, *42*, 585–592. [CrossRef]
38. Risch, A.C.; Wirthner, S.; Busse, M.D.; Page-Dumroese, D.S.; Schütz, M. Grubbing by wild boars (*Sus scrofa* L.) and its impact on hardwood forest soil carbon dioxide emissions in Switzerland. *Oecologia* **2010**, *164*, 773–784. [CrossRef]
39. Haaverstad, O.; Hjeljord, O.; Wam, H.K. Wild boar rooting in a northern coniferous forest—Minor silviculture impact. *Scand. J. For. Res.* **2014**, *29*, 90–95. [CrossRef]
40. Skoták, V.; Drimaj, J.; Kamler, J. Evaluation of damage to forest tree plantations by wild boar in the Czech Republic. *Hum.-Wildl. Interact.* **2021**, *15*, 66–73. [CrossRef]
41. Zeman, J.; Hrbek, J.; Drimaj, J.; Plhal, R.; Kamler, J.; Adamec, Z.; Heroldova, M. Wild Boar Impact to the Natural Regeneration of Oak and Acorn Importance in its Diet. *Acta Univ. Agric. Et Silvic. Mendel. Brun.* **2016**, *64*, 579–585.
42. Sims, N.K.E. The ecological impacts of wild boar rooting in East Sussex. Ph.D. Thesis, School of Biological Sciences University of Sussex, Falmer, UK, 2005; p. 276. Available online: <http://www.britishwildboar.org.uk/The%20ecological%20impacts%20of%20wild%20boar%20rooting%20in%20East%20Sussex.pdf> (accessed on 16 March 2022).
43. Fern, M.P.; Armstrong, J.B.; Barlow, R.J.; Kush, J.S. Ecological factors influencing wild pig damage to planted pine and hardwood seedlings. *Hum.-Wildl. Interact.* **2020**, *14*, 228–238.
44. Bratton, S. The effect of the European wild boar (*Sus scrofa*) on the high-elevation vernal flora in Great Smoky Mountains National Park. *Bull. Torrey Bot. Club* **1974**, *101*, 198–206. [CrossRef]
45. Howe, T.; Singer, F.J.; Ackerman, B.B. Forage relationships of European wild boar invading northern hardwood forest. *J. Wildl. Manag.* **1981**, *45*, 748–754. [CrossRef]
46. Massei, G.; Genov, P.V. The environmental impact of wild boar. *Galemys* **2004**, *16*, 135–145.
47. Janulaitis, Z.; Padaiga, V. Some fundamentals of the damage caused by wild boar to agricultural crops. In Proceedings of the Scientific Conference Wildlife Management in the Problem-Oriented Forestry, Kaunas-Girionys, Lithuania, November 1983; pp. 43–44.
48. Gerard, J.F.; Cargnelutti, B.; Spitz, F.; Valet, G.; Sardin, T. Habitat use of wild boar in a French agroecosystem from late winter to early summer. *Acta Theriol.* **1991**, *36*, 119–129.
49. Genov, P. Significance of natural biocenoses and agrocenoses as the source of food for the wild boar (*Sus scrofa* L.). *Ekol. Pol.* **1981**, *29*, 117–136.
50. Schön, T. The cost of having wild boar: Damage to agriculture in South-Southeast Sweden. Master’s Thesis, Swedish University of Agricultural Sciences, Umeå, Sweden, 2013; p. 40.
51. Engeman, R.M. Economic Considerations of Damage Assessment. In *Human Conflicts with Wildlife: Economic Considerations*; Volume 4, pp. 36–41. Available online: <https://digitalcommons.unl.edu/nwrchumanconflicts/4> (accessed on 14 June 2021).
52. Fern, M.; Barlow, R.; Slootmaker, C.; Kush, J.; Shwiff, S.; Teeter, L.; Armstrong, J. Economic Estimates of Wild Hog (*Sus scrofa*) Damage and Control Among Young Forest Plantations in Alabama. *Small-Scale For.* **2021**, *20*, 503–516. [CrossRef]
53. Kotulski, Y.; König, A. Conflicts, crises and challenges: Wild boar in the Berlin city—A social empirical and statistical survey. *Nat. Croat. Zagreb* **2008**, *17*, 233–246.

511

54. Fattorini, N.; Ferretti, F. Estimating wild boar density and rooting activity in a Mediterranean protected area. *Mamm. Biol.* **2020**, *100*, 241–251. [CrossRef]
55. Statistical Database SBS. 2021. Available online: <https://www.statistikdatabasen.scb.se/pxweb/en/ssd/> (accessed on 20 April 2021).
56. Boyce, C.M.; VerCauteren, K.C.; Beasley, J.C. Timing and extent of crop damage by wild pigs (*Sus scrofa* Linnaeus) to corn and peanut fields. *Crop Prot.* **2020**, *133*, 105–131. [CrossRef]
57. Lombardini, M.; Meriggi, A.; Fozzi, A. Factors influencing wild boar damage to agricultural crops in Sardinia (Italy). *Curr. Zool.* **2017**, *63*, 507–514. [CrossRef]
58. Schlageter, A. Preventing Wild Boar *Sus scrofa* Damage—Considerations for Wild Boar Management in Highly Fragmented Agroecosystems. Ph.D. Thesis, University of Basel, Basel, Switzerland, 2015; p. 97.
59. Bergman, T.E. *An Investigation of Human-Wild Boar Conflict—The Perceived Need for Economical Compensation among Farmers Due to Crop Damage Caused by Wild Boars—A Case Study in Arboga, Sweden*; Department of Social and Economic Geography, Arbetsrapporter, Uppsala University: Uppsala, Sweden, 2014; p. 30.
60. Anon. *On the Methodology for Assessment of Damage Caused by Game Animals to Agricultural Crops, Livestock and Forest*; Consolidated Version from 28-09-2021. Code Posted: Official Gazette, 2002, No. 97-4303, id 102301MISAK0486/359; New Edition: Since 2015-01-24: No. D1-69/3D-36, 2015-01-22, publ. TAR 2015-01-23, id 2015-01000; The Ministry of Environment and The Ministry of Agriculture of the Republic of Lithuania: Vilnius, Lithuania, 2018. Available online: <https://e-seimas.lrs.lt/portal/legalAct/lt/TAD/TAIS.187999/PqnIxigXak> (accessed on 23 August 2022).
61. West, B.C.; Cooper, A.L.; Armstrong, J.B. Managing wild pigs: A technical guide. *Hum. -Wildl. Interact. Monogr.* **2009**, *1*, 55.
62. Mitchell, B.; Balogh, S. *Monitoring Techniques for Vertebrate Pests. Feral Pigs*; NSW Department of Primary Industries, Bureau of Rural Sciences: Canberra, Australia; Natural Heritage Trust: Canberra, Australia, 2007; p. 32.
63. Wilson, C.J. *Feral Wild Boar in England. Status, Impact and Management*; A Report on behalf of Defra European Wildlife Division; Defra, RDS National Wildlife Management Team: Exeter, UK, 2005; p. 56.
64. Putman, R.; Apollonio, M. *Behaviour and Management of European Ungulates*; Whittles Publishing: Scotland, UK, 2014; p. 304.
65. König, H.J.; Ceausu, S.; Reed, M.; Kendall, H.; Hemminger, K.; Reinke, H.; Ostermann-Miyashita, E.-F.; Wenz, E.; Eufemia, L.; Hermanns, T.; et al. Integrated framework for stakeholder participation: Methods and tools for identifying and addressing human–wildlife conflicts. *Conserv. Sci. Pract.* **2021**, *3*, e399. [CrossRef]
66. Michez, A.; Morelle, K.; Lehaire, F.; Widar, J.; Authelet, M.; Vermeulen, C.; Lejeune, P. Use of unmanned aerial system to assess wildlife (*Sus scrofa*) damage to crops (*Zea mays*). *J. Unmanned Veh. Syst.* **2016**, *4*, 266–275. [CrossRef]
67. Rutten, A.; Casaer, J.; Vogels, M.; Addink, E.A.; Borre, J.V.; Leirs, H. Assessing agricultural damage by wild boar using drones: Wild Boar Damage Assessment. *Wildl. Soc. Bull.* **2018**, *42*, 568–576. [CrossRef]
68. Kim, M.; Chung, O.-S.; Lee, J.-K.A. Manual for Monitoring Wild Boars (*Sus scrofa*) Using Thermal Infrared Cameras Mounted on an Unmanned Aerial Vehicle (UAV). *Remote Sens.* **2021**, *13*, 4141. [CrossRef]
69. Rutten, A.; Casaer, J.; Onkelinx, T.; De Smet, L.; Witters, N.; Huysentruyt, H.L. Using an online survey to assess the spatial distribution of wild boar (*Sus scrofa* L.) crop damage and factors influencing this distribution and severity in Limburg province, Belgium. *Belg. J. Zool.* **2019**, *149*, 1–13. [CrossRef]
70. Belova, O. Wildlife Census Methods: Reliability and Application. In *Wildlife Census Methods: Reliability and Application*; Žemaitija National Park: Plateliai, Lithuania, 2010; pp. 21–25.
71. ENETWILD Consortium; Grignolio, S.; Apollonio, M.; Brivio, F.; Vicente, J.; Acevedo, P.; Palencia, P.; Petrovic, K.; Keuling, O. *Guidance on Estimation of Abundance and Density Data of Wild Ruminant Population: Methods, Challenges, possibilities*; EFSA Supporting Publication: Parma, Italy, 2020; p. 54. [CrossRef]
72. ENETWILD-Consortium; Acevedo, P.; Aleksovski, V.; Apollonio, M.; Berdión, O.; Blanco-Aguiar, J.A.; del Rio, L.; Ertürk, A.; Fajdiga, L.; Escribano, F.; et al. *Wild Boar Density Data Generated by Camera Trapping in Nineteen European Areas*; EFSA Supporting Publication: Parma, Italy, 2022; p. 21. [CrossRef]
73. Keeping, D.; Pelletier, R. Animal Density and Track Counts: Understanding the Nature of Observations Based on Animal Movements. *PLoS ONE* **2014**, *9*, e96598. [CrossRef]
74. Kie, J.G. *Performance in Wild Ungulates: Measuring Population Density and Condition of Individuals*; Gen. Tech. Rep. PSW-106; Pacific Southwest Forest and Range Experiment Station, Forest Service, U.S. Department of Agriculture: Berkeley, CA, USA, 1988; p. 17.
75. Gilbert, S.L.; Hundertmark, K.J.; Lindberg, M.; Person, D.; Boyce, M.S. The Importance of Environmental Variability and Transient Population Dynamics for a Northern Ungulate. *Front. Ecol. Evol.* **2020**, *8*, 531027. [CrossRef]
76. Chauvenet, A.L.M.; Gill, R.M.A.; Smith, G.C.; Ward, A.I.; Massei, G. Quantifying the bias in density estimated from distance sampling and camera trapping of unmarked individuals. *Ecol. Model.* **2017**, *350*, 79–86. [CrossRef]
77. Santini, L.; Benítez-López, A.; Dormann, C.F.; Huijbregts, M.A.J. Population density estimates for terrestrial mammal species. *Glob. Ecol. Biogeogr.* **2022**, *3*, 978–994. [CrossRef]
78. Focardi, S.; La Morgia, V.; Montanaro, P.; Riga, F.; Calabrese, A.; Ronchi, F.; Aragno, P.; Scacco, M.; Calmant, R.; Franzetti, B. Reliable estimates of wild boar populations by nocturnal distance sampling. *Wildlife Biol.* **2020**, *2020*, 1–15. [CrossRef]
79. D’Eon, R.G.; Wilson, S.F.; Hamilton, D. *Ground-Based Inventory Methods for Ungulate Snow-Track Surveys. Standards for Components of British Columbia’s Biodiversity*; No. 33a; Resources Information Standards Committee: Victoria, BC, Canada, 2006.

80. Bobek, B.; Merta, D.; Furtek, J. Use of a Line Intercept Snow Track Index and Plot Sampling for Estimating Densities of Wild Boar (*Sus scrofa*) in Southwestern Poland. *Wildl. Biol. Pract.* **2014**, *10*, 7–16. [CrossRef]
81. Dzięciołowski, R. Estimating ungulate numbers in a forest by track counts. *Acta Teriol.* **1976**, *21*, 217–222.
82. Malakauskas, A.; Karvelienė, B. Dangerous pig diseases: Classical and African swine fever. *Mano Ūkis* **2010**, *4*, 77–79.
83. Anon. *On the Approval of Permissible Quotas of Ungulate Density in Forests of the Republic of Lithuania*; The Order No. 86 15/05/1995; TAR; Parliament of the Republic of Lithuania: Vilnius, Lithuania. (In Lithuanian)
84. Anon. *On the Approval of the Preparation, Submission, and Approval of the Schedule of Procedures of Wildlife Management Plans*; The Order No. D1-162 21/03/2005. Available online: <https://e-seimas.lrs.lt/portal/legalAct/lt/TAD/TAIS.252979?jfwid=-11tbwd089g> (accessed on 23 August 2022). (In Lithuanian)
85. Padaiga, V. Project of the Programme for the Integration and Reasoned Management of the Forest and Game Management. Vilnius, Lithuania, 1994; p. 84. (In Lithuanian)
86. Bennett, L.J.; English, P.F.; McCain, R. A study of deer populations by use of pellet-group counts. *J. Wildl. Manag.* **1940**, *4*, 398–403.
87. Eberhardt, L.; van Etten, R.C. Evaluation of the pellet group count as a deer census method. *J. Wildl. Manag.* **1956**, *20*, 70–74.
88. Neff, D.J. The pellet-group count technique for big game trend, census, and distribution: A review. *J. Wildl. Manag.* **1968**, *32*, 597–614.
89. Acevedo, P.; Vicente, J.; Höfle, U.; Cassinello, J.; Ruiz-Fons, F.; Gortazar, C. Estimation of European wild boar relative abundance and aggregation: A novel method in epidemiological risk assessment. *Epidemiol. Infect.* **2007**, *135*, 519–527. [CrossRef]
90. Plhal, R.; Kamler, J.; Homolka, M. Faecal pellet group counting as a promising method of wild boar population density estimation. *Acta Theriol.* **2014**, *59*, 561–569. [CrossRef]
91. Belova, O. Delineation of the Territory of Lithuania for the Hares Leporidae: Estimation of the Habitat Suitability. *Balt For.* **1999**, *5*, 49–59.
92. Janulaitis, Z. Wild boar foraging in the agricultural landscape of Lithuanian SSR. In Proceedings of the Scientific Conference Wildlife Management in the Problem-Oriented Forestry, Kaunas-Girionys, Lithuania, November 1983; pp. 41–42.
93. Merta, D.; Mocała, P.; Pomykacz, M.; Frąckowiak, W. Autumn-winter diet and fat reserves of wild boars (*Sus scrofa*) inhabiting forest and forest-farmland environment in south-western Poland. *Folia Zool.* **2014**, *63*, 95–102. [CrossRef]
94. Janulaitis, Z.; Padaiga, V. Control of wild boar population in Lithuanian SSR. In *LMŪMTI Mokslo Darbai*; LMŪMTI: Vilnius, Lithuania, 1987; Volume 23, pp. 163–178. (In Lithuanian)
95. StatSoft. *Statistica 8.0, an Advanced Analytics Software Package*, StatSoft Inc.: Tulsa, OK, USA, 2008. Available online: www.statsoft.com (accessed on 27 March 2018).
96. Belova, O. ASF management strategy in wild boar population. In *Newest Recommendations for Agriculture and Forestry*; Semaškienė, R., Mikšys, V., Feiza, V., Kadžiulienė, Ž., Lazauskas, S., Ruzgas, V., Samuolienė, G., Eds.; LAMMC: Akademija, Lithuania, 2020; Volume 10, pp. 59–60.
97. Belova, O.; Gedminas, A.; Urbaitis, G. GPS technology to track wild boars in Lithuania: The tool for research and management. In Proceedings of the 34 IUGB Congress, Kaunas, Lithuania, 26–30 August 2019; pp. 37–38.
98. Piekarczyk, P.; Tajchman, K.; Belova, O.; Wójcik, M. Crop damage by wild boar (*Sus scrofa* L.) depending on the crop composition in Central-Eastern Poland. *Balt* **2021**, *27*, 86–96. [CrossRef]
99. Vercauteren, K.C.; Dolbeer, R.A.; Gese, E.M. Identification and Management of Wildlife Damage. In *The Wildlife Techniques Manual*; Silvy, N.J., Ed.; Paper 1204; Johns Hopkins University Press: Baltimore, MD, USA; USDA National Wildlife Research Center–Staff Publications: Fort Collins, CO, USA, 2010; pp. 232–269.
100. Anon. *On the Approval of the Methodology on Estimation of Damage Caused by Wildlife to Crops, Facilities, and Forests*; The Order No. 486/359, No. D1-69/3D-36, 2015-01-2; The Parliament of the Republic of Lithuania: Vilnius, Lithuania, 2002. Available online: <https://www.e-tar.lt/portal/legalAct.html?documentId=73ea58b0a2f611e4a82d9548fb36f682> (accessed on 23 August 2022)
101. Baleišis, R.; Bluzma, P.; Balčiauskas, L. *Ungulates of Lithuania*; Mokslo: Vilnius, Lithuania, 1987; p. 200. (In Lithuanian)
102. Kairiūkštis, L.; Padaiga, V. Peculiarities of wildlife management in the problem-oriented forestry. In Proceedings of the Scientific Conference Wildlife Management in the Problem-Oriented Forestry, Kaunas-Girionys, Lithuania, November 1983; pp. 45–46.
103. Reimoser, F.; Reimoser, S. Ungulates and their management in Austria. In *European Ungulates and their Management in the 21st Century*, 1st ed.; Apollonio, M., Andersen, R., Putman, R., Eds.; Cambridge University Press: Cambridge, UK, 2010; Volume 16, pp. 338–356. ISBN 10 0521760615.
104. Magnusson, M. Population and management models for the Swedish wild boar (*Sus scrofa*). Master's Thesis, Swedish University of Agricultural Sciences Grimsö Wildlife Research Station, Uppsala, Sweden, 2010; p. 26.
105. Keuling, O. Managing Wild Boar—Considerations for Wild Boar Management Based on Game Biology Data. Ph.D. Thesis, Dresden University of Technology, Dresden, Germany, 2009; p. 91.
106. Boitani, L.; Mattei, L.; Nonis, D.; Corsi, F. Spatial and activity patterns of wild boar in Tuscany, Italy. *J. Mammal.* **1994**, *75*, 600–612.
107. Lemel, J.; Truvé, J.; Söderberg, B. Variation in ranging and activity behaviour of European wild boar *Sus scrofa* in Sweden. *Wildl. Biol.* **2003**, *9*, 29–36.
108. Vild, O.; Hédl, R.; Kopecký, M.; Szabó, P.; Suchánková, S.; Zouha, V. The paradox of long-term ungulate impact: Increase of plant species richness in a temperate forest. *Appl. Veg. Sci.* **2017**, *20*, 282–292. [CrossRef]

513

109. Schmidt, R.H.; Timm, R.M. Vertebrate impacts on oak regeneration in California: A review of management options. In Proceedings of the 10th Great Plains Wildlife Damage Conference, Lincoln, Nebraska, 15–18 April 1991; Hygnstrom, S.E., Case, R.M., Johnson, R.J., Eds.; University of Nebraska: Lincoln, Nebraska, 1991; pp. 134–144.
110. Rutten, A.; Casaer, J.; Strubbe, D.; Leirs, H. Agricultural and landscape factors related to increasing wild boar agricultural damage in a highly anthropogenic landscape. *Wildlife Biol.* **2020**, *2020*, 1–11. [CrossRef]
111. Cappa, F.; Bani, L.; Meriggi, A. Factors affecting the crop damage by wild boar (*Sus scrofa*) and effects of population control in the Ticino and Lake Maggiore Park (North-western Italy). *Mamm. Biol.* **2021**, *101*, 451–463. [CrossRef]
112. Ballari, S.A.; Barrios-García, M.N. A review of wild boar *Sus scrofa* diet and factors affecting food selection in native and introduced ranges. *Mamm. Rev.* **2013**, *44*, 124–134. [CrossRef]
113. Rowcliffe, J.M.; Field, J.; Turvey, T.; Carbone, C. Estimating animal density using camera traps without the need for individual recognition. *J. Appl. Ecol.* **2008**, *45*, 1228–1236. [CrossRef]
114. Anon. *On the Confirmation of the Rules of Hunting in the Republic of Lithuania*, The Order No. 258; Consolidated version No. D1-610 Official Gazette Valstybės žinios; TAR: Vilnius, Lithuania, 2019; p. 16390. Available online: <https://e-seimas.lrs.lt/portal/legalAct/lt/TAD/TAIS.104124/asr> (accessed on 23 August 2022).
115. Servanty, S.; Gaillard, J.-M.; Ronchi, F.; Focardi, S.; Baubet, E.; Gimenez, O. Influence of harvesting pressure on demographic tactics: Implications for wildlife management. *J. Appl. Ecol.* **2011**, *48*, 835–843. [CrossRef]
116. Podgórski, T.; Baś, G.; Jedrzejewska, B.; Sönnichsen, L.; Śnieżko, S.; Jedrzejewski, W.; Okarma, H. Spatiotemporal behavioral plasticity of wild boar (*Sus scrofa*) under contrasting conditions of human pressure: Primeval forest and metropolitan area. *J. Mammal.* **2013**, *94*, 109–119. [CrossRef]
117. Gordigiani, L.; Viviano, A.; Brivio, F.; Grignolio, S.; Lazzeri, L.; Marcon, A.; Mori, E. Carried away by a moonlight shadow: Activity of wild boar in relation to nocturnal light intensity. *Mamm. Res.* **2022**, *67*, 39–49. [CrossRef]
118. Maselli, V.; Rippa, D.; Russo, G.; Ligrone, R.; Soppeisa, O.; D’Aniello, B.; Raia, P.; Fulgione, D. Wild boars’ social structure in the Mediterranean habitat. *Ital. J. Zool.* **2014**, *81*, 610–617. [CrossRef]

Spatiotemporal behavioral plasticity of wild boar (*Sus scrofa*) under contrasting conditions of human pressure: primeval forest and metropolitan area

TOMASZ PODGÓRSKI,* GRZEGORZ BAŚ, BOGUMIŁA JĘDRZEJEWSKA, LEIF SÖNNICHSEN, STANISŁAW ŚNIEŻKO, WŁODZIMIERZ JĘDRZEJEWSKI, AND HENRYK OKARMA

Mammal Research Institute, Polish Academy of Sciences, 17–230 Białowieża, Poland (TP, BJ, LS, WJ)

Institute for Nature Conservation, Polish Academy of Sciences, Mickiewicza 33, 31-120 Kraków, Poland (GB, SS, HO)

Leibniz Institute for Zoo and Wildlife Research, Alfred-Kowalek-Str. 17, 10315 Berlin, Germany (LS)

* Correspondent: t_podgorski@ibs.bialowieza.pl

The aim of this study was to examine the effects of sex and age of individuals, season, and human pressure (both human presence and habitat structure) on spatiotemporal behavior of wild boar (*Sus scrofa*). Specifically, we compared wild boar behavior under contrasting conditions of human pressure, within the primeval temperate forest of Białowieża (eastern Poland) and the metropolitan area of Cracow (Poland). It was predicted that, compared with the forest, wild boar within the urban area will have smaller home ranges due to restricted space, will show longer daily movements due to patchy resources, and will exhibit increased nocturnal activity and ranging behavior in an attempt to avoid human interference. We used radiotracking data from 35 wild boar. Animals inhabiting the urban area had smaller home ranges, yet covered almost twice as much distance on a daily basis than individuals inhabiting primeval forest. Daily duration of activity was similar in the 2 study areas. However, distribution of activity throughout the day differed considerably. In the urban area wild boar were almost exclusively nocturnal, whereas in the primeval forest wild boar activity was evenly distributed throughout the day. Additionally, in the urban area, activity was strongly associated with traveling speed, whereas in the primeval forest active wild boar moved about to little extent. Seasonal effects were stronger in the primeval forest and affected daily distance traveled, duration of activity, and level of diurnality. This study showed that wild boar can adjust their spatiotemporal behavior to local conditions and this may be one factor explaining recent rapid demographic expansion in Europe.

Key words: activity, daily distance, home range, Poland, space use, urban area

© 2013 American Society of Mammalogists

DOI: 10.1644/12-MAMM-A-038.1

The ranging behavior and spatiotemporal activity of mammals reflect interactions between individuals and their environment. Numerous factors have been shown to affect home range size: body size, sex, and age (McLoughlin and Ferguson 2000; McNab 1963; Mysterud et al. 2001), food abundance and distribution (Clutton-Brock and Harvey 1978; Taitt 1981), population density and social structure (Damuth 1981; Jetz et al. 2004), and risk of predation (Tufto et al. 1996). However, the importance of each factor depends on the scale of analysis, i.e., species, population, or individual level (McLoughlin and Ferguson 2000). Similarly, observed patterns of temporal activity are shaped by intrinsic and extrinsic factors. Internally controlled activity rhythms are influenced by a number of cyclic environmental factors, such as photoperiod, temperature fluctuations, regular feeding bouts (Daan and

Aschoff 1982), and temporal disturbances (e.g. predation, human interference—Keuling et al. 2008a; Kitchen et al. 2000).

In addition to these factors that, often interactively, shape spatiotemporal behavior of mammals, human activity modifies behavior and home ranges of animals. Human pressure (e.g., hunting, tourism) can directly affect behavior and may result in shifts toward nocturnal activity (Kitchen et al. 2000) or influence movements and space use patterns (Frank and Woodroffe 2001; Jeppesen 1987). In contrast, human activity may indirectly influence behavior because human-induced



habitat fragmentation has a profound effect on the spatial behavior of mammals due to the creation of patchy resource distribution and barriers that constrain animal movements (Campbell-Smith et al. 2011; Tolon et al. 2009; Vistnes et al. 2004).

Studies have highlighted considerable intraspecific variation in home range size of wild boar (*Sus scrofa*), but with no clear pattern with regard to sex and age-related differences, seasonal changes, and geographic location (Baber and Coblenz 1986; Boitani et al. 1994; Keuling et al. 2008b; Lemel et al. 2003; Massei et al. 1997; Russo et al. 1997; Singer et al. 1981). In mammals, it is predicted that changes in home range size are inversely related to population density ("density-dependent hypothesis"—Abramsky and Tracy 1980; Wood and Brenneman 1980) and food availability ("food-exploitation hypothesis"—Larter and Gates 1994). However, studies testing these hypotheses in wild boar populations reached conflicting conclusions (Bertolotto 2010; Massei et al. 1997).

Besides considerable variation in spatial behavior, temporal patterns can also be modified by several factors. Wild boars are active for approximately 12 h a day and their activity is mainly confined to dusk, dawn, and night (Boitani et al. 1994; Keuling et al. 2008a; Lemel et al. 2003; Massei et al. 1997; Russo et al. 1997; Singer et al. 1981). The duration and distribution of activity is affected by weather conditions (Lemel et al. 2003), habitat type (Boitani et al. 1994; Singer et al. 1981), and seasonality (daylight length, food availability; Keuling et al. 2008a). However, these studies were conducted in areas with high levels of human interference (e.g., hunting) and/or areas transformed by human activities, which may alter the natural behavior of the species. Information on wild boar ranging behavior and activity in undisturbed ecosystems is lacking.

In response to human activities, wild boar can adjust their spatiotemporal activity in an attempt to avoid human contact or to take advantage of the human-induced changes within the landscape. For example, wild boars have been shown to increase their nocturnal activity in response to increased hunting pressure and in proximity to human settlements (Keuling et al. 2008a). Furthermore, hunting pressure affects home range size (Maillard and Fournier 1995; Sodekat and Pohlmeyer 2001) and can cause temporary range shifts to ensure access to seasonal food resources offered by agricultural crops (Keuling et al. 2009).

In recent decades the wild boar numbers within Europe have increased significantly (Apollonio et al. 2010; Sáez-Royuela and Telleria 1986). This creates potential for serious conflicts between humans and wild boar in agricultural regions as well as densely populated rural and urban areas (Geisser and Reyer 2004). Several factors have been proposed that may have contributed to increase of wild boar densities, such as increasing global temperatures, more frequent mast fruiting events, as well as management practices that are ineffective in regulating population numbers (Bieber and Ruf 2005). However, the expansion of populations and their growth may be partially attributed to the behavioral plasticity of the species.

In this paper, we compared the activity and spatiotemporal behavior of wild boar populations in 2 contrasting environments differing in human pressure, namely one population inhabiting the primeval temperate forest of Białowieża (eastern Poland) and another inhabiting the metropolitan area of Cracow, the second largest city in Poland. These 2 areas differ in human pressure (both human presence and habitat structure), with high exposure in Cracow and low in primeval forest. This study is the first direct comparison between populations to investigate wild boar behavioral plasticity under contrasting conditions of human pressure and habitat structure. We hypothesized that, compared with primeval forest, wild boar within urban areas will have smaller home range sizes due to restricted space; show increased daily movement due to patchy resources; and exhibit increased nocturnal activity and ranging behavior to avoid human interference.

MATERIALS AND METHODS

Study area.—The study was conducted in 2 areas of Poland—Białowieża Primeval Forest in the eastern part of the country and a metropolitan area of Cracow in the south—characterized by profound differences in human pressure and habitat structure (Fig. 1). The Białowieża Primeval Forest (BPF)—located on the Polish–Belarusian border ($52^{\circ}47'N$, $23^{\circ}48'E$; $\sim 1,450 \text{ km}^2$)—is the last remnant of the European temperate lowland forest. Within Poland, BPF (600 km^2) is dominated by 3 main forest types, including a high proportion of old growth: pedunculate oak (*Quercus robur*), small-leaved lime (*Tilia cordata*), European hornbeam (*Carpinus betulus*) stands, mixed coniferous stands, and European ash (*Fraxinus excelsior*), black alder (*Alnus glutinosa*) stands.

The study area was located in the center of the Polish part of the BPF. The study area consisted of 92.6% forest, 7.2% open land (meadows, river valleys), and 0.2% anthropogenic areas (e.g., allotments)—Corine Land Cover database (European Environment Agency, <http://www.eea.europa.eu/>). Two-thirds of the study area was within the borders of the Białowieża National Park and its buffer zone, where hunting is prohibited, motorized traffic is allowed only under permission from the National Park, and forest managements are limited to sanitary cutting and fencing off regeneration areas. Tourists are restricted to day hikes on designated trails and overnight camping is prohibited. The remaining part of the study area is within the commercial part of the BPF, where limited logging and hunting occur. However, hunting from fixed locations is only permitted at a few designated sites. Motorized traffic in the commercial part of the BPF is allowed only for forestry service vehicles. Human density in the Polish part of the BPF is about 7 inhabitants/ km^2 and the density of roads accessible for 2-wheel-drive is about 1.2 km/km^2 in the commercial part of the forest (Theuerkauf et al. 2003). However, there was no human settlement in the close vicinity of the study area.

The density of the wild boar population is estimated to be 2.35 ind./ km^2 . In addition to wild boar, four other ungulate species occur in the forest, including red deer (*Cervus elaphus*),




Fig 1.—Maps detailing location of the study areas in Poland and radiolocations of wild boars (*Sus scrofa*; all individuals) in the study areas. In Cracow City, areas occupied by radiotracked wild boar consisted of 18% forest, 43% arable/open land, and 39% estates and industrial areas, whereas in Białowieża Primeval Forest 92.6%, 7.2%, and 0.2%, respectively.

roe deer (*Capreolus capreolus*), moose (*Alces alces*), and the European bison (*Bison bonasus*). Two large carnivores, Eurasian lynx (*Lynx lynx*) and wolf (*Canis lupus*), also occur in stable populations (Jędrzejewska and Jędrzejewski 1998). The climate is transitional between Atlantic and continental types, with stronger influence of the latter (Jędrzejewska and Jędrzejewski 1998). Mean January temperature is -4.1°C and the mean number of days with snow cover is 105 (Institute of Soil Science and Plant Cultivation, <http://www.zazi.iung.pulawy.pl/>). Annual precipitation ranges from 550 to 600 mm (Institute of Geography and Spatial Planning, Polish Academy of Sciences, <http://www.igipz.pan.pl/>).

The second study area, Cracow metropolitan area, is located in the south of Poland ($50^{\circ}1'N$, $19^{\circ}53'E$). Cracow is the second biggest city of the country, with an estimated human population of 750,000. Approximately one-third of the city area is covered with meadows, pastures, orchards, arable lands, parks, and woods. The wild boar population in this area is located in the peripheral part of the city and its habitat is a mosaic of woods, parks, open land (meadows, gardens, and agricultural fields), and anthropogenic areas (housing estates, warehouses, and industrial estates). Forests, covering 4% of the city, are mainly concentrated to the western part and largely consist of European beech (*Fagus sylvatica*) and pedunculate oak-dominated stands. Radiolocations were taken over a 40-km^2 area in the western part of the city and consisted of 18% forest, 43% open land, and 39% anthropogenic area (Corine

Land Cover database). Roe deer is the only other ungulate sympatric species within the city limits. Single-hunter hunts are permitted in the outskirts of the city and nearly one-half of the wild boar population is harvested annually. Wild boar population density is estimated to be 0.15 ind./km^2 within this area (unpublished data of the Institute for Nature Conservation, Polish Academy of Sciences, Cracow). Mean January temperature is -2.1°C and the mean number of days with snow cover is 62. Annual precipitation ranges from 650 to 700 mm (Institute of Climatology, Jagiellonian University, Cracow, <http://www.klimat.geo.uj.edu.pl/>).

Trapping and radiotracking.—Data were collected between February 2006 and December 2008. The same method was used to capture wild boar in both study areas: cage and wooden traps ($1.5 \times 1 \times 2 \text{ m}$) and large drop-net traps (Jędrzejewski and Kamler 2004) baited with maize. A combination of Zoletil (tilotamide and zolazepam) and Domitor (medetomidine) mixture (1:0.025 ratio) was administered intramuscularly to immobilize captured wild boar. Atipemazole hydrochloride (Antisedan) was used as an antidote (Kreeger 1997). In some instances a combination of ketamine and xylazine was used (1:0.05 ratio). Animals weighing less than 30 kg were only immobilized with ketamine (0.2 ml/kg) and were handled without being fully anesthetized. Finally, captured animals were fitted with ear-tag radiotransmitters (Advanced Telemetry Systems, Isanti, Minnesota, and Wagener Telemetriealanlagen, Cologne, Germany). Procedures used for capture and

immobilization followed guidelines of the American Society of Mammalogists (Sikes et al. 2011), and our research and handling protocol was approved by the Local Ethical Commission for Experiments on Animals in Białystok and Cracow, Poland.

Upon capture, wild boars were classified as juveniles (<12 months old; hereinafter referred to as juv.), subadults (12–24 months; referred to as subad.), or adults (>24 months; referred to as ad.) dependent on tooth eruption (Matschke 1967). However, because trapping mainly took place during winter months, when some animals were about to enter the next age class, individuals were assigned to their respective age classes during the tracking period for analytical purposes (e.g., juveniles captured at age of 10 months and tracked for 7 months would be considered as subadults). A total of 35 wild boar was radiotagged (BPF: 9 ad. and 8 subad. females, 3 ad. and 9 subad. males; Cracow: 2 ad. and 1 subad. females; 2 ad. and 1 subad. males), 26 of which belonged to 7 marked social groups (BPF: 24 ind. in 6 groups; Cracow: 2 ind. in 1 group). The remaining 9 individuals (BPF: 5; Cracow: 4) were either solitary or members of unmarked groups.

Radiotagged individuals were located 2–4 times per week with equal intensity during the day and night (G-test: $G_1 = 0.48$, $P = 0.49$). Consecutive locations were separated by at least 12 h to ensure independence of observations (Swihart and Slade 1985). The locations of individuals were determined on foot by recording at least 3 bearings for each triangulation using a 3-element Yagi antenna (Titley Scientific, Lawnton, Australia) and Yaesu FT-817 transceiver (Yaesu Musen Co., Tokyo, Japan). A vehicle was used to move about the study areas. The location of an individual was calculated from a given set of bearings and using the maximum likelihood estimator method described by Lenth (1981). Only location estimates with error ellipse ≤ 2 ha were included for further analysis, and the program LOAS (Ecological Software Solutions) was used to calculate positions from triangulation. Accuracy of triangulation was determined in the field by locating transmitters in known location (Harris et al. 1990). Mean estimated error between the known transmitter location and those obtained from telemetry was 153 ± 107 m ($\bar{X} \pm SD$, $n = 120$).

Throughout the study tagged individuals were monitored during 24-h continuous tracking sessions, when both location and the activity of individuals were recorded at 30-min intervals. Animals were tracked by a single person (6-h shifts) taking bearings approximately every 10 min from forest roads (BPF) or streets and park trails (Cracow). Continuous tracking followed methodology used by Kamler et al. (2007a,b). Individual activity was determined by assessing irregularities in signal strength when each bearing was taken (Boitani et al. 1994; Kenward 1987; Keuling et al. 2008a; Massei et al. 1997). Researchers listened to the signal for 1 min and classified wild boar as active if the signal was alternating in strength, or inactive when the signal was constant. Direct observations of radiotagged individuals confirmed that changes in signal strength were attributed to both traveling and on-site activity

(e.g., foraging), whereas constant signals indicated a lack of movement (e.g., resting).

Data analysis.—Wild boars were followed for 9.7 ± 3.16 ($\pm SD$, $n = 34$, min = 6, max = 18) months (BPF: 9.5 ± 2.8 , $n = 29$; Cracow: 11.0 ± 5.1 , $n = 5$) and 86 ± 39.4 (min = 35, max = 177) locations were collected per individual (BPF: 74 ± 27.7 , Cracow: 153 ± 28.8). To avoid underestimation of home range size, only those individuals tracked for >3 months and that yielded >30 locations were used in analyses. Using this threshold value we found no effect of increased tracking time or number of locations on home range size (Spearman's $r = 0.06$, $n = 34$, $P = 0.74$ and $r = 0.04$, $P = 0.81$, respectively).

We estimated home range size using two methods: local-convex-hull (LoCoH—Getz and Wilmers 2004) and minimum convex polygon (MCP). The LoCoH method has an advantage over classical methods, e.g., MCP, because hard boundaries and inaccessible areas within the home range (e.g., rivers, highways, buildings) are identified (Getz et al. 2007; Huck et al. 2008). Thus, home ranges are constructed in such a way that only the areas truly utilized by animals were included, which is of great importance for accurate estimation of home ranges in fragmented habitats, e.g., urban or agricultural areas. However, we included results obtained with MCP to allow for comparison with other studies (Table 1). Specifically, the adaptive LoCoH method (a -LoCoH) was used, because it is remarkably robust to deviations from the optimal value of the parameter a , and, compared with other LoCoH methods, is considered to be the superior method of estimating home range size (Getz et al. 2007). Value of a parameter was set at the greatest distance between any 2 points in a set of an individual's locations, as suggested by Getz et al. (2007). Huck et al. (2008) showed that a -LoCoH estimates home range size reliably with number of locations comparable with our sample sizes. Ninety percent isopleths were used for home range estimation as it was recently demonstrated that using isopleths greater than 90% can result in unreliable area estimates biased by sample size and sensitive to outliers (Börger et al. 2006).

During the study, 26 wild boars were monitored 71 times during continuous, 24-h tracking sessions. The number of sessions was distributed evenly across sex and age classes (ad. M: 20; subad. M: 15; ad. F: 20; subad. F: 16). Seasons were defined as winter (December–February), spring (March–May), summer (June–August), and autumn (September–November) to reflect changes in climate, vegetation, and life cycle of wild boar. Continuous tracking was conducted during every season (winter: 12 sessions; spring: 23; summer: 17; autumn: 17). Using data collected from 24-h tracking sessions, the following parameters of the daily behavior were calculated: daily range size: using a -LoCoH and MCP. All locations were used under the assumption that they would reflect the actual area utilized within the 24-h period; daily distance traveled: sum of distances between consecutive locations; duration of activity: sum of active readings recorded at 10-min intervals (assuming each active reading = 10 min) over the 24-h period; diurnality: proportion of active readings during daylight (time from

Table 1.—Mean values of home range size, daily range size, daily distance traveled, duration of activity, and diurnality of wild boar (*Sus scrofa*) in Cracow City and Białowieża Forest study areas, Poland, during 2006–2008.

	Białowieża Primeval Forest			Cracow City			<i>P</i> -value ^a
	<i>n</i> ^b	$\bar{X} \pm SD$	Min–max	<i>n</i>	$\bar{X} \pm SD$	Min–max	
Home range size (km ²)							
α -LoCoH 90%	29	2.2 ± 0.94	0.8–4.6	5	1.0 ± 0.60	0.6–2.1	0.044
MCP 90%	29	4.2 ± 2.51	0.9–9.9	5	3.9 ± 3.16	1.0–8.7	—
Daily range size (km ²)							
α -LoCoH 100%	20 (35)	1.0 ± 0.73	0.1–3.6	6 (36)	0.9 ± 0.64	0.1–2.5	—
MCP 100%	20 (35)	1.3 ± 1.36	0.1–7.3	6 (36)	2.4 ± 2.19	0.1–8.8	—
Daily distance traveled (km)	20 (35)	6.8 ± 2.56	2.5–16.4	6 (36)	12.9 ± 6.51	3.1–26.7	0.002
Activity duration (h/day)	20 (35)	11.2 ± 5.23	3.3–22.8	6 (36)	11.4 ± 1.70	7.2–14.9	0.299
Diurnality ^c	20 (35)	0.52 ± 0.24	0.0–1.0	6 (36)	0.1 ± 0.08	0.0–0.3	<0.001

^a If shown, *P*-value refers to parameter Area in the most parsimonious model for the given response variable.

^b Number of animals, in parentheses: number of 24-h continuous tracking sessions.

^c Percent of active fixes during daylight hours.

sunrise to sunset) compared with the total number of active readings during each 24-h period. Data were controlled for seasonal changes in day length, daylight-saving time changes, and difference of sunrise and sunset time between the study areas. Finally, the distribution of activity and speed of movement throughout the day were investigated in both study areas. For such analysis, data were pooled into 2-h intervals before the percentage of active locations and cumulative distances traveled was calculated for each interval within both study areas. All spatial and home-range analyses were conducted using R version 2.13.1 software (R Development Core Team 2011) and visualized in Arc View GIS 9.1 (ESRI, Redlands, California).

Statistical analyses.—Linear mixed models (Pinheiro and Bates 2000) were used to analyze relationships between dependent variables (home range size, daily range size, daily distance traveled, and duration of activity) and explanatory variables (study area, season, sex, and age of individuals). Factor “study area” incorporated differences between study areas due to combined effect of human pressure (human presence and activity, habitat structure). Mixed models were used because of the nonindependent nature of the data. When fitting models to home range size data, group ID was treated as a random factor because some individuals belonged to the same social groups (see subsection “Trapping and radiotracking). Similarly, in models used to analyze variables derived from continuous tracking sessions, individual ID was treated as a random factor because numerous animals were sampled more than once (on average, 2.9 sessions/ind.). The distribution of model residuals was checked for normality using the Shapiro–Wilk test. Data on daily range size were log-transformed to improve normality and reduce skewness. Diurnality data (proportion of active fixes during daytime) were arcsin transformed. Interactions between explanatory variables were included in each model. Akaike information criterion (AIC), with a second-order correction for small sample size (AIC_c), was used to rank models dependent on their fit (Burnham and Anderson 2002). Models with the lowest AIC_c values were considered to be the best-fit models, and models within 2 AIC_c units were considered of similar

quality to the most parsimonious model. All statistical analyses were conducted using R version 2.13.1 software (R Development Core Team 2011).

RESULTS

Spatial behavior.—Home range size estimates obtained using α -LoCoH and MCP were highly correlated, both in total home range and daily range (Spearman’s $r = 0.73$, $n = 34$, $P < 0.001$ and $r = 0.80$, $n = 71$, $P < 0.001$, respectively). Therefore, we restricted further analysis only to α -LoCoH estimates because they reflect area actually used by wild boar.

In Cracow City, average home range was less than half the size of that in BPF (Table 1). Variation in home range size of wild boar was best explained by differences between the 2 study areas (Tables 2 and 3). The model including only study area as an explanatory variable explained the variation in home range size better than other models, and accounted for 41% of the model weight (Table 2).

On a daily basis wild boar utilized areas of similar size in each study area (Table 1). However, in Cracow City daily occupied range constituted 90% of the total home range, whereas in BPF the daily range only covered 45% of the total home range. None of the explanatory variables sufficiently explained variation in daily range size. Consequently, the intercept-only model provided the best fit (Tables 2 and 3).

In Cracow City, daily distance traveled by wild boar was almost twice that of wild boar within the BPF (Table 1). On average, subadult wild boar covered 65% longer distances on a daily basis compared with adults ($\bar{X} \pm SD$; 12.6 ± 6.51 km, 8.1 ± 4.52 km, respectively). In Cracow City distances covered daily were lower during the spring compared with other seasons, whereas in BPF, contrastingly, daily movement distance was consistent during all seasons (Fig. 2a). The best model explaining variation in daily distance traveled included study area, age, season, and an interaction between study area and season (Table 2). Although overall effect of season was insignificant in the best model (Table 3), different seasonal effects between the study areas were identified, as indicated by a significant interaction term (Table 3).

Table 2.—Confidence set of mixed linear models ($\Delta\text{AIC}_c < 2$) for variables explaining home range size, daily range size, daily distance traveled, duration of activity, and diurnality of wild boar (*Sus scrofa*) in Cracow City and Białowieża Primeval Forest study areas, Poland studied in 2006–2008. Included in the table are corrected Akaike information criterion (AIC_c) values and the difference between each model and the best-fitting model (ΔAIC_c), number of model parameters (k), and model Akaike's weights (ω_i). The models were ranked dependent on the AIC_c value and in descending order (the most parsimonious models are on the top of each list).

Response variable	Variables included in a model	k	AIC_c	ΔAIC_c	ω_i
Home range size	Area	4	79.98	0	0.412
	Area + Age	5	81.23	1.25	0.220
	Area + Sex	5	81.56	1.58	0.187
	Area + Age + Sex	6	81.64	1.66	0.180
Daily range size	Intercept	3	69.80	0	0.355
	Age	4	70.69	0.89	0.227
	Sex	4	71.64	1.84	0.141
	Area + Season + Area \times Season	10	71.65	1.85	0.140
Daily distance	Area	4	71.70	1.90	0.137
	Area + Age + Season + Area \times Season	11	1,361.74	0	0.484
	Area + Age + Season	8	1,362.55	0.81	0.322
Activity duration	Area + Age + Season + Sex + Area \times Season	9	1,363.58	1.84	0.193
	Season + Area + Area \times Season	10	363.56	0	0.404
	Season + Area + Age + Area \times Season	11	364.87	1.31	0.210
	Season + Area + Sex + Area \times Season	11	364.97	1.42	0.199
Diurnality	Season	6	365.10	1.55	0.187
	Area + Season + Area \times Season	10	-383.54	0	0.342
	Area + Season + Age + Area \times Season	11	-383.40	0.13	0.320
	Area + Season + Sex + Area \times Season	11	-382.29	1.25	0.183
	Area + Season + Age + Sex + Area \times Season	12	-381.95	1.59	0.155

Table 3.—Parameters and their statistics included in the most parsimonious model from the confidential set of models ($\Delta\text{AIC}_c < 2$), and with home range size, daily range size, daily distance traveled, duration of activity, and diurnality of wild boar (*Sus scrofa*) as response variables. All explanatory variables were factors and estimates for factor levels are presented in relation to: area (Białowieża), season (winter), and age (adult).

Response variable	Parameter	Coefficient	SE	t-value	P-value
Home range size	Intercept	2.29	0.27	8.44	<0.001
	Area (Cracow)	-1.20	0.54	-2.23	0.044
Daily range size	Intercept	-0.16	0.047	-3.52	<0.001
	Intercept	4,648.3	2,491.64	1.86	0.069
Daily distance	Area (Cracow)	9,775.4	2,856.52	3.42	0.002
	Age (Subadult)	2,978.5	1,092.78	2.72	0.009
	Area (Cracow \times spring)	-8,074.1	3,380.86	-2.39	0.022
	Area (Cracow \times summer)	-5,314.7	3,642.97	-1.46	0.153
	Area (Cracow \times autumn)	-3,708.0	3,587.29	-1.03	0.307
	Season (autumn)	3,075.6	3,024.32	1.02	0.315
	Season (summer)	1,465.4	2,757.31	0.53	0.598
	Season (spring)	460.8	2,761.82	0.17	0.868
	Intercept	9.61	1.80	5.33	<0.001
	Season (autumn)	6.14	2.04	3.01	0.004
	Area (Cracow \times autumn)	-6.73	2.30	-2.91	0.005
	Area (Cracow \times spring)	-3.55	2.09	-1.70	0.096
Activity duration	Area (Cracow \times summer)	-3.81	2.25	-1.69	0.098
	Season (spring)	2.82	1.78	1.58	0.121
	Area (Cracow)	2.58	2.42	1.07	0.299
	Season (summer)	0.78	1.81	0.43	0.666
	Intercept	0.058	0.0056	10.31	<0.001
	Area (Cracow)	-0.053	0.0071	-7.45	<0.001
	Season (summer)	0.054	0.0089	6.05	<0.001
	Season (spring)	0.044	0.0088	4.91	<0.001
	Area (Cracow \times summer)	-0.043	0.0117	-3.68	<0.001
	Area (Cracow \times autumn)	-0.039	0.0115	-3.39	0.002
Diurnality	Season (autumn)	0.030	0.0097	3.05	0.004
	Area (Cracow \times spring)	-0.026	0.0108	-2.42	0.021




Fig 2.—Seasonal changes in mean values ($\pm SE$) of a) daily distance traveled, b) duration of activity, and c) diurnality (proportion of active fixes taken during daylight hours from total number of active fixes during 24-h period) of wild boar (*Sus scrofa*) in Cracow City and Białowieża Primeval Forest study areas during 2006–2008.

Activity patterns.—Variation in duration of activity was largely driven by seasonal changes and the different effects of season in the 2 study areas (Tables 2 and 3). Specifically, seasonal variation in the duration of activity was a result of the wild boar in BPF being active for longer periods (by 5.7 h on average) in autumn compared with Cracow City (Fig. 2b, Table 3). During other seasons duration of activity was comparable between study areas (Fig. 2b, Table 3). Overall, differences




Fig 3.—Distribution of a) activity and b) speed of movement during 24-h period of wild boar (*Sus scrofa*) in Cracow City and Białowieża Primeval Forest study areas during 2006–2008. Mean ($\pm SE$) values are shown for 2-h time intervals. Shaded areas represent the shifting times for sunrise and sunset throughout the year.

between study areas did not contribute significantly to explained variation in activity duration (Tables 2 and 3).

Wild boars in BPF were more active during daylight compared with Cracow City, where they were virtually inactive during daylight (Fig. 3a, Table 1). Study area, season, and an interaction between study area and season were included in the best model describing variation in diurnality (Table 2). Furthermore, seasonal changes in diurnality contributed significantly to explained variation of this parameter (Fig. 2c, Table 3). Effect of season on daytime activity was particularly strong in BPF, with the lowest values observed during winter (Fig. 2c, Table 3).

In BPF, activity was evenly distributed throughout the 24-h period (Kruskal–Wallis $H_{11,409} = 18.16, P = 0.078$; Fig. 3a), whereas in Cracow City activity was concentrated to crepuscular and dark hours (Kruskal–Wallis $H_{11,421} = 375.98, P < 0.001$; Fig. 3a). A similar pattern was observed with regard to speed of movement. Movements were uniformly distributed throughout the day in BPF, but not in Cracow City (Kruskal–Wallis $H_{11,409} = 15.13, P = 0.18$; $H_{11,421} = 106.01, P$

< 0.001, respectively; Fig. 3b). Finally, the proportion of active fixes within 2-h intervals in Cracow City was correlated with the speed of movement (Spearman's $r = 0.97$, $P < 0.001$), whereas in BPF, activity was not significantly associated with movement (Spearman's $r = 0.44$, $P = 0.15$).

DISCUSSION

This study demonstrated high behavioral plasticity in wild boar living in 2 contrasting areas differing in human pressure, both direct human presence and indirect human-induced habitat changes. Wild boar inhabiting urban areas occupied smaller home ranges, were more mobile on a daily basis, and were more nocturnal when compared with the population inhabiting a primeval forest characterized by low levels of human pressure. In contrast, daily duration of activity and daily range size were comparable between both study areas. Some behavioral parameters (daily distance traveled, activity duration, and diurnality) were affected by seasonal changes, although the effects were different in the 2 study areas. Seasonal effects were stronger in the BPF compared with the urban area of Cracow City. However, because our study was based on a relative small number of individuals that were also unevenly distributed between the study areas, our results should be interpreted with caution.

Spatial behavior in areas with low and high human pressure.—In contrast to the continuous forest habitat of BPF, the urban area of Cracow City provides patchy habitat with dispersed resources (e.g., food, resting, and breeding sites). This difference in habitat structure most likely contributed to contrasting spatiotemporal patterns observed between both study areas. Within the urban environment wild boar had to cover distance almost twice as long as those in primeval forest to meet their daily energetic requirements, as well as to locate shelter for long periods of diurnal inactivity. Interestingly, wild boar showed similar seasonal patterns in both areas, except during spring when a reduction in daily distance traveled was observed in Cracow City. Spring is the breeding season for wild boar (Fernandez-Llario and Carranza 2000; Jędrzejewska and Jędrzejewski 1998) and restricted daily movement could be a result of parturition and weaning of offspring (females yielded 65% of data for this season), as detailed in Spitz and Janeau (1990).

Wild boar is a generalist omnivore with opportunistic feeding habits, and its diet reflects local environmental conditions (Herrero et al. 2006; Schley and Roper 2003). Consequently, wild boar can take advantage of a variety of food resources offered by urban environments (e.g., crop fields, metropolitan parks and forests, garbage dumps, and orchards). In contrast to BPF, where food abundance is seasonally variable (Jędrzejewska and Jędrzejewski 1998), urban environments provide relatively high and constant food resources, which could explain the smaller home ranges observed in Cracow City. Therefore, our result may support the “food-exploitation hypothesis,” which predicts an inverse relationship between food abundance and home range size (Larter and

Gates 1994). However, in this study we did not quantify food resource abundance. Besides food abundance, habitat fragmentation could also contribute to smaller home ranges observed in the urban area.

Wild boar in BPF occupied daily ranges that constituted less than half (45%) of their total home range, whereas in Cracow City almost the entire home range (90%) was covered on a daily basis. A potential explanation could be that wild boar in the BPF shift their ranges spatially according to spatiotemporal changes in food resources, which is similar to shifts described by Keuling et al. (2009) for wild boar in Germany.

No effect of study area, season, sex, or age on daily range size was detected in our study. Other studies have had similar findings, e.g., the lack of seasonal and sexual differences in daily range size were reported in a study from Italy (Russo et al. 1997). However, variation in daily range size may also be shaped by subtle site-specific factors (e.g., daily ambient temperature—Kamler et al. 2007a), which were not included in our group of robust explanatory variables.

Activity patterns in areas with low and high human pressure.—In both study areas wild boars were active for similar periods of time (~11 h/day), which suggests that individuals had to balance comparable energy budgets. However, there was a remarkable difference between the distributions of activity throughout the day among the 2 habitats. Wild boar in the urban area exhibited crepuscular and nocturnal activity, which could have been a strategy of avoiding interference with humans. This finding is consistent with other studies investigating urban mammal populations (wild boar in Barcelona, Spain—Cahill et al. 2003; red fox [*Vulpes vulpes*] in Oxford, United Kingdom—Doncaster and Macdonald 1997). Furthermore, activity was strongly associated with movement, which suggests that wild boar visited numerous food patches and spent relatively little time foraging on one patch before finding secluded sites for daytime resting. In contrast, activity of wild boar in the primeval forest was evenly distributed throughout the 24-h period and showed a polyphasic pattern characterized by alternating phases of activity and resting. Interestingly, similar patterns have been found in other ungulates inhabiting BPF (red deer—Kamler et al. 2007b; European bison—Caboń-Raczyńska et al. 1987). However, in those species the alternating peaks of activity and resting are attributed to the foraging habits of ruminants (food intake and regurgitation), which is not the case in wild boar. High level of diurnality observed in the BPF was possibly caused by very limited hunting pressure in this area, as it had been shown that high hunting pressure increased nocturnal activity of wild boar (Keuling et al. 2008a). Thus, observed activity patterns, as well as high activity during daylight hours, are likely a consequence of low levels of human activity. Wild boar in the BPF showed mainly on-site activity, as suggested by a nonsignificant relationship between activity and speed of movement, and this can be explained by uniformly distributed and readily available resources (food, water, and resting sites) that minimize the need for extensive movement.

Wild boars from BPF were active for exceptionally long periods during the autumn when compared with other seasons, as well as to Cracow City. Two interacting factors may be responsible for this pattern. First, the severity of winter has been shown to be a main factor affecting survival and reproduction of wild boar (Geisser and Reyer 2005; Jędrzejewska and Jędrzejewski 1998). Therefore, achieving good body condition and gaining fat reserves before winter is crucial for wild boar survival. Second, the acorn crop occurring in autumn is the most efficient way to obtain sufficient fat reserves within the BPF (Jędrzejewska and Jędrzejewski 1998), explaining the higher activity observed in autumn. In contrast, no seasonal effect was observed for this parameter in Cracow City, which may indicate that urban environments provide stable, season-independent food resources. Alternatively, the milder climate within the city may reduce the requirement for gaining crucial fat reserves.

In the BPF, wild boar co-occur with 2 natural predators, wolf and lynx. However, predation constitutes a relatively small portion of natural mortality of wild boar in the area, only 19% from wolf and 1% from lynx (Jędrzejewska and Jędrzejewski 1998). Predation risk affects spatiotemporal behavior at various scales and may lead to, e.g., reduced locomotory activity and range size, and changes in habitat use (Borkowski and Owadowska 2010; Valeix et al. 2009). However, literature on spatiotemporal response of wild boar to predation risk is lacking. In our study, we could not quantify the effect of predation and separate it from other factors (low human pressure, habitat structure). Season also had strong effects on the proportion of time wild boars were active during daylight hours (diurnality). This effect was particularly pronounced in the BPF. It was apparent that seasonal changes in levels of diurnality mirrored seasonal changes in day length. Diurnality, with regard to our calculations, was a relative measure giving proportion of the total amount of activity during a 24-h period allocated to daylight hours. Therefore, it is sensitive to varying day length, especially when duration of activity is constant throughout the year, which was found to be the case in our study (except for autumn in the BPF). Overall, levels of diurnal activity within the urban area were much lower than those in the primeval forest (usually below 10% of total daily activity time). Moreover, urban populations maintained nocturnal activity almost irrespective of season, whereas forest population followed natural changes in day length. This difference may demonstrate effect of human pressure on activity patterns of wild boar.

In summary, our study showed that wild boar can adjust their spatiotemporal behavior to local conditions. The spatial behavior and activity of wild boar population inhabiting primeval forest seemed to be mainly driven by seasonal changes in food availability and abiotic conditions, whereas human activity shaped spatiotemporal patterns of the urban population.

Densities of wild boar have been increasing over the last decades within Europe (Apollonio et al. 2010; Geisser and Reyer 2004; Sáez-Royuela and Telleria 1986). Additionally, this expansion is accompanied by invasion of new habitats densely

populated by humans (Cahill et al. 2003; Fisher et al. 2004), including urban areas that may offer new unexplored niches. Several factors have been proposed to explain demographic expansion of wild boar (Bieber and Ruf 2005). Until now, however, behavioral plasticity of the species has not drawn much attention, although it may play an important role in this process.

ACKNOWLEDGMENTS

The project was funded by the Polish Ministry of Science and Higher Education (grant No. 2 P04F 073 29). We are grateful to K. Bojarska, R. Kozak, A. Maszczyk, and many students and volunteers for their help with fieldwork. We also thank the Environmental Engineering Department of Cracow Municipal Office, City Park and Zoological Garden Foundation, and “Podwawelskie” Hunting Club for permission to work in the forested areas of Cracow and help with fieldwork. Statistical analyses were improved thanks to valuable comments from K. Zub and M. Oliver. T. Borowik helped to improve graphics and H. Parris revised English. We are grateful to D. Kuijper and 2 anonymous reviewers for valuable comments on an earlier version of the manuscript.

LITERATURE CITED

- ABRAMSKY, Z., AND C. R. TRACY. 1980. Relation between home range size and regulation of population size in *Microtus ochrogaster*. *Oikos* 34:347–355.
- APOLLONIO, M., R. ANDERSEN, AND R. PUTMAN (eds.). 2010. European ungulates and their management in the 21st century. Cambridge University Press, Cambridge, United Kingdom.
- BABER, D. W., AND B. E. COBLENTZ. 1986. Density, home range, habitat use, and reproduction in feral pigs on Santa Catalina Island. *Journal of Mammalogy* 67:512–525.
- BERTOLOTTO, E. 2010. Behavioural ecology of wild boar (*Sus scrofa*) in an Apennine Environment. PhD dissertation, University of Sassari, Sassari, Italy.
- BIEBER, C., AND T. RUF. 2005. Population dynamics in wild boar *Sus scrofa*: ecology, elasticity of growth rate and implications for the management of pulsed resource consumers. *Journal of Applied Ecology* 42:1203–1213.
- BOITANI, L., L. MATTEI, D. NONIS, AND F. CORSI. 1994. Spatial and activity patterns of wild boar in Tuscany, Italy. *Journal of Mammalogy* 75:600–612.
- BÖRGER, L., ET AL. 2006. Effects of sampling regime on the mean and variance of home range size estimates. *Journal of Animal Ecology* 75:1393–1405.
- BORKOWSKI, Z., AND E. OWADOWSKA. 2010. Field vole (*Microtus agrestis*) seasonal spacing behavior: the effect of predation risk by mustelids. *Naturwissenschaften* 97:487–493.
- BURNHAM, K. P., AND D. R. ANDERSON. 2002. Model selection and multi-model inference: a practical information-theoretic approach. Springer, New York.
- CABÓN-RACZYŃSKA, K., M. KRASIŃSKA, Z. KRASIŃSKI, AND J. WÓJCICKI. 1987. Rhythm of daily activity and behavior of European bison in the Białowieża Forest in the period without snow cover. *Acta Theriologica* 32:335–372.
- CAHILL, S., F. LLIMONA, AND J. GRACIA. 2003. Spacing and nocturnal activity of wild boar *Sus scrofa* in a Mediterranean metropolitan park. *Wildlife Biology* 9:3–13.
- CAMPBELL-SMITH, G., M. CAMPBELL-SMITH, I. SINGLETON, AND M. LINKIE. 2011. Apes in space: saving an imperiled orangutan population in Sumatra. *PLOS ONE* 6(2):e17210.

- CLUTTON-BROCK, T. H., AND P. H. HARVEY. 1978. Mammals, resources and reproductive strategies. *Nature* 273:191–195.
- DAAN, S., AND J. ASCHOFF. 1982. Circadian contributions to survival. Pp. 305–321 in *Vertebrate circadian systems* (J. Aschoff, S. Daan, and G. A. Groos, eds.). Springer-Verlag, Berlin, Germany.
- DAMUTH, J. 1981. Home range, home range overlap, and species energy use among herbivorous mammals. *Biological Journal of the Linnaean Society* 15:185–193.
- DONCASTER, C. P., AND D. MACDONALD. 1997. Activity patterns and interactions of red foxes (*Vulpes vulpes*) in Oxford city. *Journal of Zoology* 241:73–87.
- FERNANDEZ-LLARIO, P., AND J. CARRANZA. 2000. Reproductive performance of the wild boar in a Mediterranean ecosystem under drought conditions. *Ethology, Ecology and Evolution* 12:335–343.
- FISHER, C., H. GOUDIN, AND M. OBERMANN. 2004. Spatial behavior of the wild boar in Geneva, Switzerland: testing the methods and first results. Pp. 149–157 in *Wild boar research 2002. A selection and edited papers from 4th International Wild Boar Symposium. Galemys*, 16 Special Issue (C. Fonseca, J. Herrero, A. Luis, and A. M. V. M. Soares, eds.). SECEM, Málaga, Spain.
- FRANK, L. G., AND R. WOODROFFE. 2001. Behaviour of carnivores in exploited and controlled populations. Pp. 419–442 in *Carnivore conservation* (J. L. Gittleman, S. M. Funk, D. Macdonald, and R. K. Wayne, eds.). Cambridge University Press, Cambridge, United Kingdom.
- GEISSER, H., AND H.-U. REYER. 2004. Efficacy of hunting, feeding, and fencing to reduce crop damage by wild boars. *Journal of Wildlife Management* 68:939–946.
- GEISSER, H., AND H.-U. REYER. 2005. The influence of food and temperature on population density of wild boar *Sus scrofa* in the Thurgau (Switzerland). *Journal of Zoology* 267:89–96.
- GETZ, W. M., S. FORTMANN-ROE, P. C. CROSS, A. J. LYONS, S. J. RYAN, AND C. C. WILMERS. 2007. LoCoH: nonparametric kernel methods for constructing home ranges and utilization distributions. *PLoS ONE* 2(2):e207.
- GETZ, W. M., AND C. C. WILMERS. 2004. A local nearest neighbour convex-hull construction of home ranges and utilization distributions. *Ecography* 27:489–505.
- HARRIS, S., W. J. CRESSWELL, P. G. FORDE, W. J. TREWHELLA, T. WOOLLARD, AND S. WRAY. 1990. Home-range analysis using radiotracking data—a review of problems and techniques particularly as applied to the study of mammals. *Mammal Review* 20:97–123.
- HERRERO, J., A. GARCÍA-SERRANO, S. COUTO, V. M. ORTUÑO, AND R. GARCÍA-GONZÁLEZ. 2006. Diet of wild boar *Sus scrofa* L. and crop damage in an intensive agroecosystem. *European Journal of Wildlife Research* 52:245–250.
- HUCK, M., J. DAVISON, AND T. J. ROPER. 2008. Comparison of two sampling protocols and four home-range estimators using radiotracking data from urban badgers *Meles meles*. *Wildlife Biology* 14:467–477.
- JĘDRZEJEWSKA, B., AND W. JĘDRZEJEWSKI. 1998. *Predation in vertebrate communities: the Białowieża Primeval Forest as a case study*. Springer-Verlag, Berlin, Germany.
- JĘDRZEJEWSKI, W., AND J. F. KAMLER. 2004. From the field: modified drop-net for capturing ungulates. *Wildlife Society Bulletin* 32:1305–1308.
- JEPPESEN, J. L. 1987. Impact of human disturbance on home range, movements and activity of red deer (*Cervus elaphus*) in a Danish environment. *Danish Review of Game Biology* 13:1–38.
- JETZ, W., C. CARBONE, J. FULFORD, AND J. H. BROWN. 2004. The scaling of animal space use. *Science* 306:266–268.
- KAMLER, J. F., B. JĘDRZEJEWSKA, AND W. JĘDRZEJEWSKI. 2007a. Factors affecting daily ranges of red deer *Cervus elaphus* in Białowieża Primeval Forest, Poland. *Acta Theriologica* 52:113–118.
- KAMLER, J. F., B. JĘDRZEJEWSKA, AND W. JĘDRZEJEWSKI. 2007b. Activity patterns of red deer in Białowieża National Park, Poland. *Journal of Mammalogy* 88:508–514.
- KENWARD, R. E. 1987. *Wildlife radio tagging*. Academic Press, London, United Kingdom.
- KEULING, O., N. STIER, AND M. ROTH. 2008a. How does hunting influence activity and spatial usage in wild boar *Sus scrofa* L. *European Journal of Wildlife Research* 54:729–737.
- KEULING, O., N. STIER, AND M. ROTH. 2008b. Annual and seasonal space use of different age classes of female wild boar *Sus scrofa* L. *European Journal of Wildlife Research* 54:403–412.
- KEULING, O., N. STIER, AND M. ROTH. 2009. Commuting, shifting or remaining? Different spatial utilisation patterns of wild boar *Sus scrofa* L. in forest and field crops during summer. *Mammalian Biology* 74:145–152.
- KITCHEN, A. M., E. M. GESE, AND E. R. SCHAUSTER. 2000. Changes in coyote activity patterns due to reduced exposure to human persecution. *Canadian Journal of Zoology* 78:853–857.
- KREEGER, T. J. 1997. *Handbook of wildlife chemical immobilization*. Wildlife Pharmaceuticals Inc., Fort Collins, Colorado.
- LARTER, N. C., AND C. C. GATES. 1994. Home range size of wood bison: effects of age, sex and forage availability. *Journal of Mammalogy* 75:142–149.
- LEMEL, J., J. TRUVÉ, AND B. SÖDERBERG. 2003. Variation in ranging and activity behaviour of European wild boar (*Sus scrofa*) in Sweden. *Wildlife Biology* 9:29–36.
- LENTH, R. V. 1981. On finding the source of a signal. *Technometrics* 23:149–154.
- MAILLARD, D., AND P. FOURNIER. 1995. Effect of shooting with hounds on home range size of wild boar (*Sus scrofa* L.) groups in Mediterranean habitat. *IBEX Journal of Mountain Ecology* 3:102–107.
- MASSEI, G., P. V. GENOV, B. STAINES, AND M. L. GORMAN. 1997. Factors influencing home range and activity of wild boar (*Sus scrofa*) in a Mediterranean coastal area. *Journal of Zoology* 242:411–423.
- MATSCHKE, G. H. 1967. Aging European wild hogs by dentition. *Journal of Wildlife Management* 31:109–113.
- MCLOUGHLIN, P. D., AND S. H. FERGUSON. 2000. A hierarchical pattern of limiting factors helps explain variation in home range size. *Ecoscience* 7(2):123–130.
- MCNAB, B. K. 1963. Bioenergetics and the determination of home range size. *American Naturalist* 97:133–140.
- MYSTERUD, A. F., F. J. PEREZ-BARBERIA, AND I. J. GORDON. 2001. The effect of season, sex, and feeding style on home range area versus body mass scaling in temperate ruminants. *Oecologia* 127:30–39.
- PINHEIRO, J. C., AND D. M. BATES. 2000. *Mixed-effects models in S and S-PLUS*. Springer-Verlag, New York.
- R DEVELOPMENT CORE TEAM. 2011. R: a language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. ISBN 3-900051-07-0, <http://www.R-project.org/>.
- RUSSO, L., G. MASSEI, AND P. V. GENOV. 1997. Daily home range and activity of wild boar in a Mediterranean area free of hunting. *Ethology, Ecology & Evolution* 9:287–294.
- SÁEZ-ROYUELA, C., AND J. L. TELLERIA. 1986. The increased population of the Wild Boar (*Sus scrofa* L.) in Europe. *Mammal Review* 16:97–101.
- SCHLEY, L., AND T. J. ROPER. 2003. Diet of wild boar *Sus scrofa* in Western Europe, with particular reference to consumption of agricultural crops. *Mammal Review* 33:43–56.

- SIKES, R. S., GANNON, W. L. and THE ANIMAL CARE AND USE COMMITTEE OF THE AMERICAN SOCIETY OF MAMMALOGISTS . 2011. Guidelines of the American Society of Mammalogists for the use of wild mammals in research. *Journal of Mammalogy* 92:235–253.
- SINGER, F. J., D. K. OTTO, A. R. TIPTON, AND C. P. HABLE. 1981. Home ranges, movements and habitat of European wild boar in Tennessee. *Journal of Wildlife Management* 45:343–353.
- SODEIKAT, G., AND K. POHLMAYER. 2001. Temporary home range modifications of wild boar family groups (*Sus scrofa* L.) caused by drive hunts in Lower Saxony (Germany). *European Journal of Wildlife Research* 48:161–166.
- SPITZ, F., AND G. JANEAU. 1990. Spatial strategies: an attempt to classify daily movements of wild boar. *Acta Theriologica* 35:129–149.
- SWIHART, R. K., AND N. A. SLADE. 1985. Testing for independence of observation in animal movements. *Ecology* 66:1176–1184.
- TAITT, M. J. 1981. The effect of extra food on small rodent populations: I. Deermice (*Peromyscus maniculatus*). *Journal of Animal Ecology* 50:111–124.
- THEUERKAUF, J., W. JĘDRZEJEWSKI, K. SCHMIDT, AND R. GULA. 2003. Spatiotemporal segregation of wolves from humans in the Białowieża Forest (Poland). *Journal of Wildlife Management* 67:706–716.
- TOLON, V., S. DRAY, A. LOISON, A. ZEILEIS, C. FISCHER, AND E. BAUBET. 2009. Responding to spatial and temporal variations in predation risk: space use of a game species in a changing landscape of fear. *Canadian Journal of Zoology* 87:1129–1137.
- TUFTO, J., R. ANDERSEN, AND J. D.C.. LINNELL. 1996. Habitat use and ecological correlates of home range size in a small cervid: the roe deer. *Journal of Animal Ecology* 65:715–724.
- VALEIX, M. ET AL. 2009. Behavioral adjustments of African herbivores to predation risk by lions: spatiotemporal variations influence habitat use. *Ecology* 90:23–30.
- VISTNES, I., C. NELLEMANN, P. JORDHY, AND O. STRAND. 2004. Effects of infrastructure on migration and range use of wild reindeer. *Journal of Wildlife Management* 68:101–108.
- WOOD, G. W., AND R. E. BRENNEMAN. 1980. Feral hog movements and habitat use in South Carolina. *Journal of Wildlife Management* 44:420–427.

Submitted 10 February 2012. Accepted 29 May 2012.

Associate Editor was Harald Beck.