

Artin's Algebra

Ashan Jayamal

2024-03-05

Contents

1 Matrices	5
1.1 Basic Operations	5
2 Group Theory	7
2.1 Laws of Compositions	7
2.2 Groups and Subgroups	7
2.3 Subgroups of the Additive Group of Integers	9
2.4 Cyclic Groups	9
2.5 Homomorphisms	9
2.6 Isomorphisms	11
2.7 Equivalence Relations and Partitions	11
2.8 Cosets	18
2.9 Modular Arithmetic	18
2.10 The Correspondence Theorem	18
2.11 Product Groups	23
2.12 Quotient Groups	29
2.13 Exercises	34
3 Liener Operators	81
4 Vector spaces.	83

Chapter 1

Matrices

1.1 Basic Operations

Let m and n be positive integers. An $m \times n$ matrix is a collection of mn numbers arranged in a rectangular array.

A handwritten diagram on a piece of paper defining an $m \times n$ matrix. It shows a bracketed rectangular arrangement of elements a_{ij} . The top row is labeled $a_{11} \dots a_{1n}$, and the bottom row is labeled $a_{m1} \dots a_{mn}$. To the right of the matrix, the text "m rows" is written vertically. Below the matrix, the text "n columns" is written.

$$\begin{bmatrix} a_{11} & \cdots & a_{1n} \\ \vdots & & \vdots \\ a_{m1} & \cdots & a_{mn} \end{bmatrix}$$

m rows

n columns

::: {.example #unnamed-chunk-1}

$$A := \begin{bmatrix} 8 & 0 & 3 \\ 78 & -5 & 2 \end{bmatrix}$$

A is 2×3 matrix.(two rows and three columns)

::: The numbers in a matrix are the matrix entries. They may be denoted by a_{ij} , where i and j are indices (integers) with $1 < i < m$ and $1 < j < n$. The index i represents the row index, and j represents the column index. So a_{ij} is the entry that appears in the i th row and j th column of the matrix.

$$i \begin{bmatrix} & & j \\ \cdots & a_{ij} & \cdots \\ & & \vdots \end{bmatrix}$$

Figure 1.1:

Chapter 2

Group Theory

2.1 Laws of Compositions

2.2 Groups and Subgroups

Definition 2.1 (Group). A group is a set G together with a law of composition that has the following properties:

- The law of composition is associative: $(ab)c = a(bc)$ for all a, b, c in G .
- G contains an identity element 1 , such that $la = a$ and $al = a$ for all a in G .
- Every element a of G has an inverse, an element b such that $ab = 1$ and $ba = 1$.

Notation: If set G with composition \cdot is a group, then we denote it by (G, \cdot)
::: {.definition #unnamed-chunk-3 name="Ableian Group"} Group G is called abliean if its law of composition is commutative. i.e.

$$\forall x \in G, xy = yx$$

:::

Example 2.1.

- $\mathbb{R}^\times := \mathbb{R} \setminus \{0\}$ is an abliean group under multiplication
- $GL_n(\mathbb{R}) := \{A \in M_n(\mathbb{R}) : A \text{ is invertible}\}$ with matrix multiplication is non-abliean group. This group is called *general linear group*.

Definition 2.2 (Order of group). The order of a group G is the number of elements that it contains. We

$$|G| := \text{number of elements of } G = \text{the order of } G$$

If the order is finite, G is said to be a finite group. If not, G is an infinite group.

Here is our notation for some familiar infinite abelian groups:

Example 2.2.

- $(\mathbb{Z}, +)$:The set of integers, with addition as its law of composition (the additive group of integers)
- $(\mathbb{R}, +)$:The set of real numbers, with addition as its law of composition (the additive group of real numbers)
- $(\mathbb{R}^{\times}, \times)$:The set of nonzero real numbers, with multiplication as its law of composition(the multiplicative group) $(\mathbb{C}, +)$:the set of complex numbers, with addition as its law of composition (the additive group of complex numbers)
- $(\mathbb{C}^{\times}, \times)$:The set of nonzero complex numbers, with multiplication as its law of composition(the multiplicative group of complex numbers)

Proposition 2.1 (Cancellation Law). *Let G be group and let $a, b, c \in G$ whose law of composition is written multiplicatively.*

- *If $ab = ac$ or if $ba = ca$, then $b = c$.*
- *If $ab = a$ or if $ba = a$, then $b = 1$*

Proof. Multiply both sides of $ab = ac$ on the left by a^{-1} to obtain $b = c$. The other proofs are analogous. \square

Remark. As you saw a^{-1} plays a major rule in above proof. So the cancellation rule does not hold when element a is not invertible.

Example 2.3 (Example that not holding the cancellation law).

$$\begin{bmatrix} 1 & 1 \\ 0 & 0 \end{bmatrix} \begin{bmatrix} 1 & 1 \\ 2 & 0 \end{bmatrix} = \begin{bmatrix} 1 & 1 \\ 0 & 0 \end{bmatrix} \begin{bmatrix} 3 & 0 \\ 0 & 1 \end{bmatrix}$$

Example 2.4. Let T be a set and $G := \{f : T \rightarrow T : f \text{ is a bijection}\}$. Then G with composition is a group. We use notation $\text{sys}(T)$ to denote the

Example 2.5. The group of permutations of the set of indices $\{1, 2, \dots, n\}$ is called the *symmetric group*, and is denoted by S_n . Then $|S_n| = n!$. So, S_n is a finite group of order $n!$.

Let's discuss some individual cases for n .

- $n = 2$

The permutations of a set $\{1, 2\}$ of two elements are the identity i and the transposition $\tau = (12)$.

$$S_2 := \{id, (12)\}$$

◦	id	(12)
id	id	(12)
(12)	(12)	id
—	—	—

- $n = 3$

S_3 has order $3! = 6$. S_3 serves as a convenient example because it is the smallest group whose law of composition isn't commutative. We will refer to it often. To describe it, we pick two particular permutations in terms of which we can write all others. We take the cyclic permutation (123) , and the transposition (12) , and label them as x and y , respectively. Then

$$x^3 = (123)^3 = (123)(123)(123) = (123)(132) = id \quad (2.1)$$

$$y^2 = (12)^2 = (12)(12) = id \quad (2.2)$$

$$yx = (12)(123) = (13) = (132)(12) = ((123)(123))(12) = (123)^2(123)$$

As a summary,

$$x^3 = 1, Y^2 = 1, yx = x^2y$$

2.3 Subgroups of the Additive Group of Integers

2.4 Cyclic Groups

2.5 Homomorphisms

Definition 2.3. Let $(G, *)$ and (G', \odot) be groups. A **homomorphism** $\phi : G \rightarrow G'$ is a map from G to G' such that for all a and b in G :

$$\phi(a * b) = \phi(a) \odot \phi(b). \quad (2.4)$$

Examples of Homomorphisms

1. **Determinant Function:** $\det : GL_n(\mathbb{R}) \rightarrow \mathbb{R}^\times$
2. **Exponential Map:** $\exp : (\mathbb{R}, +) \rightarrow (\mathbb{R}^\times, \times)$ defined by $x \mapsto e^x$

3. **Map** $\phi: (\mathbb{Z}, +) \rightarrow G$ defined by $\phi(n) = a^n$, where a is a given element of G
4. **Absolute Value Map:** $|\cdot|: (\mathbb{C}^\times, \times) \rightarrow (\mathbb{R}^\times, \times)$

Trivial Homomorphism

The trivial homomorphism $\phi: G \rightarrow G'$ between any two groups maps every element of G to the identity in G' .

Inclusion Map

If H is a subgroup of G , the inclusion map $i: H \rightarrow G$ defined by $i(x) = x$ for x in H is a homomorphism.

Proposition 2.2. *Let $\phi: G \rightarrow G'$ be a group homomorphism.*

- (a) *If a_1, \dots, a_k are elements of G , then $\phi(a_1 \dots a_k) = \phi(a_1) \dots \phi(a_k)$.*
- (b) *ϕ maps the identity to the identity: $\phi(1_G) = 1_{G'}$.*
- (c) *ϕ maps inverses to inverses: $\phi(a^{-1}) = \phi(a)^{-1}$.*

Proposition 2.3. *Let $\phi: G \rightarrow G'$ be a homomorphism of groups, and let $a, b \in G$. Let $K = \ker(\phi)$. Then following conditions are equivalent:*

- $\phi(a) = \phi(b)$,
- $a^{-1}b$ is in K ,
- b is in the coset aK ,
- The cosets bK and aK are equal.

Proof.

- (1) \implies (2). Suppose $\phi(a) = \phi(b)$. Now consider

$$\phi(a^{-1}b) = \phi(a^{-1})\phi(b) = \phi(a^{-1})\phi(b) = 1$$

Thus, $a^{-1}b \in K$.

- (2) \implies (1). If $a^{-1}b \in K$

$$1 = \phi(a^{-1}b) = \phi(a^{-1})\phi(b) = \phi(a^{-1})\phi(b)$$

- Thus, $\phi(a) = \phi(b)$.
- Suppose b is in the coset aK . Then $b = ak$ for some

□

2.6 Isomorphisms


2.7 Equivalence Relations and Partitions

Definition 2.4. Sure, here is the LaTeX version of your statement:

A partition n of a set S is a subdivision of S into nonoverlapping, nonempty subsets:

$$S = \bigcup_i A_i$$

, where A_i are disjoint nonempty subsets of S .



Example 2.6. The two sets *Even* and *Odd* partition the set of integers.

Example 2.7. With the usual notation, the sets

$$\{1\}, \{y, xy, x^2y\}, \{x, x^2\}$$

form a partition of the symmetric group S_3 .

Definition 2.5. An equivalence relation on a set S is a relation that holds between certain pairs of elements.

- Reflexive: For all a , $a \sim a$.
- Symmetric: If $a \sim b$, then $b \sim a$.
- Transitive: If $a \sim b$ and $b \sim c$, then $a \sim c$.

Example 2.8. Two triangles are said to be congruent if their sides have the same length and angles have same measure.

Congruence of triangles is an example of an equivalence relation on the set of triangles in the plane. If A, B , and C are triangles, and if A is congruent to B and B is congruent to C , then A is congruent to C . It is very easy to check three equivalence properties. I am not going to do this.

Example 2.9. Conjugacy is an equivalence relation on a group. Let G be a group. Two group elements are conjugate, $a \sim b$, if $b = gag^{-1}$ for some $g \in G$.

- *Reflexive:* Observe that $a = aaa^{-1}$ then, $a \sim a$.
- *Symmetric:* Suppose that $a \sim b$, then $b = gag^{-1}$. Thus, $a = g^{-1}bg$. Hence, $b \sim a$.
- *Transitive:* Suppose that $a \sim b$ and $b \sim c$. This means that $b = g_1ag_1^{-1}$ and $c = g_2bg_2^{-1}$ for some group elements g_1 and g_2 . Then $c = g_2(g_1ag_1^{-1})g_2^{-1} = (g_2g_1)a(g_2g_1)^{-1}$, so $a \sim c$

Proposition 2.4. An equivalence relation on a set S determines a partition of S , and conversely.

Proof.

- (\Rightarrow): Given a partition of S , the corresponding equivalence relation is defined by the rule that $a \sim b$ if a and b lie in the same subset of the partition. The axioms for an equivalence relation are obviously satisfied.
- (\Rightarrow): Given an equivalence relation (\sim), one defines a partition this way: The subset that contains a is the set of all elements b such that $a \sim b$. This subset is called the **equivalence class** of a . We'll denote it by C_a here

$$C_a = \{b \in S | a \sim b\}$$

The next lemma completes the proof of the proposition

□

Lemma 2.1. *Given an equivalence relation on a set S , the subsets of S that are equivalence classes partition S .*

Proof. This is an important point, so we will check it carefully. We must remember that the notation C_a stands for a subset defined in a certain way. The partition consists of the subsets, and several notations may describe the same subset.

- *Non-emptiness:* The reflexive axiom tells us that a is in its equivalence class. Therefore, the class C_a is nonempty.
- *Union is whole set:* Since a can be any element, the union of the equivalence classes is the whole set S .
- *Disjoint property:* To show this, we prove following claim:

– **Claim:** If C_a and C_b have an element in common, then $C_a = C_b$.
 Since we can interchange the roles of a and b , it will suffice to show that if C_a and C_b have an element, say d , in common, then $C_b \subseteq C_a$. Suppose that x is in C_b , then $b \sim x$. Since d is in both sets, $a \sim d$ and $b \sim d$, and the symmetry property tells us that $d \sim b$. So we have $a \sim d$, $d \sim b$, and $b \sim x$. Two applications of transitivity show that $a \sim x$, and therefore, x is in C_a .

□

Example 2.10. The relation on a group defined by $a \sim b$ if a and b are elements of the same order is an equivalence relation. (Trivial.)

The corresponding partition for the symmetric group S_3 are

$$\{1\}, \{y, xy, x^2y\}, \{x, x^2\}.$$

If a partition of a set S is given, we may construct a new set \bar{S} whose elements are the subsets. We imagine putting the subsets into separate piles, and we regard the piles as the elements of our new set \bar{S} . It seems advisable to have a notation to distinguish a subset from the element of the set S (the pile) that it represents. If U is a subset, we will denote by $[U]$ the corresponding element of S .

Example 2.11. If S is the set of integers and if Even and Odd denote the subsets of even and odd integers, respectively, then \bar{S} contains the two elements [Even] and [Odd].

We will use this notation more generally. When we want to regard a subset U of S as an element of a set of subsets of S , we denote it by $[U]$.

When an equivalence relation on S is given, the equivalence classes form a partition, and we obtain a new set \bar{S} whose elements are the equivalence classes $[C_a]$. We can think of the elements of this new set in another way, as the set obtained by changing what we mean by equality among elements. If a and b are in S , we interpret $a \sim b$ to mean that a and b become equal in \bar{S} , because $C_a = C_b$. With this way of looking at it, the difference between the two sets S and \bar{S} is that in \bar{S} more elements have been declared “equal,” i.e., equivalent. It seems to me that we often treat congruent triangles this way in school.

For any equivalence relation, there is a natural surjective map

$$\pi : S \rightarrow \bar{S} \quad (2.5)$$

that maps an element a of S to its equivalence class: $\pi(a) = [C_a]$. When we want to regard \bar{S} as the set obtained from S by changing the notion of equality, it will be convenient to denote the element $[C_a]$ of \bar{S} by the symbol \bar{a} . Then the map π becomes

$$\pi(a) = \bar{a} \quad (2.6)$$

We can work in \bar{S} with the symbols used for elements of S , but with bars over them to remind us of the new rule:

$$\text{If } a \text{ and } b \text{ are in } S, \text{ then } \bar{a} = \bar{b} \text{ means } a \sim b. \quad (2.7)$$

A disadvantage of this bar notation is that many symbols represent the same element of S . Sometimes this disadvantage can be overcome by choosing a particular element, a representative element, in each equivalence class. For example, the even and the odd integers are often represented by $\bar{0}$ and $\bar{1}$:

$$\{\text{[Even]}, \text{[Odd]}\} = \{\bar{0}, \bar{1}\}. \quad (2.8)$$

Though the pile picture may be easier to grasp at first, the second way of viewing S is often better because the bar notation is easier to manipulate algebraically.

2.7.1 The Equivalence Relation Defined by a Map

Any map of sets $f : S \rightarrow T$ gives us an equivalence relation on its domain S . It is defined by the rule: $a \sim b$ if $f(a) = f(b)$.

Definition 2.6 (Inverse Image and Fibers). The inverse image of an element t of T is the subset of S consisting of all elements s such that $f(s) = t$. It is denoted symbolically as

$$f^{-1}(t) = \{s \in S \mid f(s) = t\}. \quad (2.9)$$

The inverse images are also called the fibres of the map f , and the fibres that are not empty are the equivalence classes for the relation defined above.

Remark. This is symbolic notation. Please remember that **unless f is bijective, f^{-1} will not be a map.**

Here the set S of equivalence classes has another incarnation, as the image of the map. The elements of the image correspond bijectively to the nonempty fibres, which are the equivalence classes

Example 2.12. Let's consider the absolute value map from the complex numbers \mathbb{C}^\times to the positive real numbers \mathbb{R}^+ :


$$f : \mathbb{C}^\times \rightarrow \mathbb{R}^+, \quad f(z) = |z|$$

The fiber of an element t in \mathbb{R}^+ is the subset of \mathbb{C}^\times consisting of all complex numbers z such that $f(z) = |z| = t$. Symbolically, we denote the fiber as:

$$f^{-1}(t) = \{z \in \mathbb{C}^\times : |z| = t\}$$


This fiber represents all complex numbers with the same absolute value t . Thus, fibers are circles in Complex plane.

Note that the absolute value map is surjective, so each positive real number t corresponds to a unique fiber in \mathbb{C}^\times .



Some Fibres of the Absolute Value Map $\mathbb{C}^\times \rightarrow \mathbb{R}^\times$.

Example 2.13. If G is a finite group, we can define a map $f : G \rightarrow \mathbb{N}$ to the set $\{1, 2, 3, \dots\}$ of natural numbers, letting $f(a)$ be the order of the element a in G . The fibers of this map are the sets of elements with the same order (see example 2.10).




Definition 2.7 (congruence). Let G, G' be groups and $\phi : G \rightarrow G'$ be a group homomorphism. The equivalence relation on G defined by ϕ is usually denoted by \equiv , rather than by \sim , and is referred to as congruence.

$$a \equiv b \text{ if } \phi(a) = \phi(b)$$

We have seen that elements a and b of G are congruent, i.e., $\phi(a) = \phi(b)$, if and only if b is in the coset aK of the kernel K (See proposition 2.3)

Proposition 2.5. Let K be the kernel of a homomorphism $\varphi : G \rightarrow G'$. The fibre of that contains an element a of G is the coset aK of K . These cosets partition the group G , and they correspond to elements of the image of φ .



Proof. The fibre of φ over an element $g' \in G'$ is the set of all elements $a \in G$ such that $\varphi(a) = g'$. Let's denote this fibre as

$$F_{g'} = \{a \in G : \varphi(a) = g'\}$$

Now, let's take an arbitrary element $a \in G$ and consider the left coset $aK = \{ak : k \in K\}$.

We know that K is the kernel of φ , so for any $k \in K$, we have $\varphi(k) = 1_{G'}$ where $1_{G'}$ is the identity element in G' .

- Claim 1: $aK = F_{\varphi(a)}$.

- sub claim 1.1: $aK \subseteq F_{\varphi(a)}$.

For any $ak \in aK$, we have $\varphi(ak) = \varphi(a)\varphi(k) = \varphi(a)1_{G'} = \varphi(a)$. This shows that every element in the coset aK is mapped to the same element under φ as a itself. Therefore, the coset aK is a subset of the fibre $F_{\varphi(a)}$.

- sub claim 1.2 : $F_{\varphi(a)} \subseteq aK$.

If $a' \in F_{\varphi(a)}$, then $\varphi(a') = \varphi(a)$, which implies that $\varphi((a')^{-1}a) = 1_{G'}$. This means that $(a')^{-1}a \in K$, or equivalently, $a' \in aK$. (by proposition 2.3)Therefore, the fibre $F_{\varphi(a)}$ is a subset of the coset aK .

Since aK is a subset of $F_{\varphi(a)}$ and $F_{\varphi(a)}$ is a subset of aK , we conclude that $aK = F_{\varphi(a)}$. So, the fibre of φ that contains an element a of G is indeed the coset aK of K .

- Claim 2: The cosets of a subgroup K partition the group G .

A partition of a set is a collection of non-empty subsets such that every element in the set is in exactly one of these subsets.

To show that the cosets of K partition G , we need to show two things:

- sub claim 2.1: Every element of G is in at least one coset of K .
Given any $g \in G$, g is in the coset gK . So, every element of G is in at least one coset of K .
- sub claim 2.2: No element of G is in more than one coset of K .
Suppose gK and hK are two cosets of K and there is some element $x \in G$ that is in both gK and hK . This means that there exist $k_1, k_2 \in K$ such that $x = gk_1 = hk_2$ (by proposition 2.3). Then $g = hk_2k_1^{-1}$. Thus, $g \in hK$, which implies that $gK = hK$. So, no element of G is in more than one coset of K .

Therefore, the cosets of K partition the group G . □

2.8 Cosets

2.9 Modular Arithmetic

2.10 The Correspondence Theorem

Definition 2.8. Let $\phi : G \rightarrow G'$ be a group homomorphism, and let H be a subgroup of G . We may restrict ϕ to H , obtaining a homomorphism

$$\phi|_H : H \rightarrow G'$$

In other words, we take the same map but restrict its domain.

Notation: We use this notation for clarity $[\phi|_H](h)$.

Further, we can see that following observations.

- By definition, $\forall h \in H, \phi|_H(h) = \phi(h)$
- The restriction $\phi|_H$ is a homomorphism (Since ϕ is homomorphism).
- The kernel of $\phi|_H$ is the intersection of the kernel of ϕ with H :

$$\ker(\phi|_H) = (\ker\phi) \cap H$$

There is no need to prove this. This is trivial by definition of kernel.

- Image of $\phi|_H$ is the same as the image $\phi(H)$ of H under the map ϕ .

$$Im(\phi_H) = \phi(H)$$

- If $|H|$ and $|G'|$ have no common factor, $\phi(H) = \{1\}$, so H is contained in the kernel. (Since, by Artin's book corollary 2.8.13,

$$\begin{array}{c|c} |Im(\phi_H)| & |H| \\ \hline & \end{array} \quad (2.10)$$

$$\begin{array}{c|c} |Im(\phi_H)| & |G'| \\ \hline & \end{array} \quad (2.11)$$

, Thus, if $|H|$ and $|G'|$ have no common factors, $|Im(\phi_H)| = |\phi(H)| = 1$. So, $Im(\phi_H) = \phi(H) = \{1\}$

Now let's see an example.

Example 2.14. Define sign homomorphism $\sigma : S_n \rightarrow \{\pm 1\}$ by $\sigma(x) = 1$ if x is even, and $\sigma(x) = -1$ if x is odd. Then the image of the sign homomorphism is,

$$Im(\sigma) = \{\pm 1\}$$

it has order 2.

Let $H = \{x \in S_n : \text{if } x \text{ has odd order}\}$. Then H is a subgroup. So, $H \subset \ker(\sigma)$ (We can easily verify this.)

Further, The sub group of S_n with even permutations is called *Alternating group* (A_n).

2.10.1 PROBLEM

Proposition 2.6. Let $\phi : G \rightarrow G'$ be a homomorphism with kernel K , and let $H' \leq G'$. Denote the inverse image $\phi^{-1}(H')$ by H . i.e.:

$$H = \phi^{-1}(H') = \{x \in G : \phi(x) \in H'\}$$

Then,

- $\phi^{-1}(H')$ is a subgroup of G that contains K .
- If H is a normal subgroup of G' , then $\phi^{-1}(H')$ is also a normal subgroup of G .
- If ϕ is surjective and H is a normal subgroup of G , then $\phi^{-1}(H')$ is a normal subgroup of G' .

Remark. ϕ^{-1} is not a map.

Proof.

- **Claim 1:** $K \subseteq \phi^{-1}(H') = H$
Let $x \in K$. Then $\phi(x) = 1_{G'}$. Since $1_{G'} \in H'$. Thus, $x \in \phi^{-1}(H') = H$.
Therefore, $K \subseteq \phi^{-1}(H') = H$
- **Claim 2:** $\phi^{-1}(H')$ is a subgroup of G

- *Closure* : Suppose $x, y \in H = \phi^{-1}(H')$. Then $\phi(x), \phi(y) \in \phi(H')$. Since ϕ is a homomorphism $\phi(x)\phi(y) = \phi(xy)$. Since $H' \leq G'$, then $\phi(xy) = \phi(x)\phi(y) \in H'$. Thus, $xy \in \phi^{-1}(H') = H$
- *Identity* : Since $\phi(1_G) = 1_{G'}$, $1_G \in \phi^{-1}(H') = H$
- *Inverse* : Let $x \in \phi^{-1}(H') = H$. Then $\phi(x) \in H'$ and since $H' \leq G'$, then $(\phi(x))^{-1} \in H'$. Since ϕ is a homomorphism, $(\phi(x))^{-1} = \phi(x^{-1})$. Thus $(\phi(x))^{-1} = \phi(x^{-1}) \in H'$. Hence, $x^{-1} \in \phi^{-1}(H')$.

- **Claim 3:** If H' is a normal subgroup of G' , then $\phi^{-1}(H')$ is also a normal subgroup of G .

Now suppose that H' is a normal subgroup of G' . Let $x \in H$ and $g \in G$. Then, since ϕ is a homomorphism,

$$\phi(-1) = \phi(g)\phi(x)\phi(g^{-1}) = \phi(g)\phi(x)(\phi(g))^{-1}$$

So, $\phi(gxg^{-1})$ is conjugate of $\phi(x)$. Since $x \in H = \phi^{-1}(H')$, the $\phi(x) \in H'$.
Thus, $\phi(gxg^{-1}) = \phi(g)\phi(x)(\phi(g))^{-1} \in H'$. Hence, $gxg^{-1} \in \phi^{-1}(H') = H$.

- **Claim 4:** If ϕ is surjective and H is a normal subgroup of G , then $\phi^{-1}(H')$ is a normal subgroup of G' .

Suppose that ϕ is surjective and H is a normal subgroup of G . Let $a \in H'$ and $b \in G'$. Since ϕ is surjective, there are elements $x \in H$ and $y \in G$ such that $\phi(x) = a$ and $\phi(y) = b$. Since H is normal $yxy^{-1} \in H$, thus $\phi(yxy^{-1}) = bab^{-1} \in H'$.

□

Example 2.15. let denote the determinant homomorphism

$$\det : GL_n(\mathbb{R}) \rightarrow \mathbb{R}^\times$$

. Note that $\mathbb{R}^+ \trianglelefteq \mathbb{R}^\times$. (It is very clear that \$The set of positive real numbers is a subgroup of \mathbb{R}^\times , and since \mathbb{R}^\times is abelian, \mathbb{R}^+ is normal.) Now consider the inverse image of \mathbb{R}^+ ,

$$\det^{-1}(\mathbb{R}^+) = \{A \in GL_n(\mathbb{R}) : \det(A) \in \mathbb{R}^+\} = SL_n(\mathbb{R})$$

. By above proposition we can see that $SL_n(\mathbb{R})$ is a normal subgroup. (Because $\mathbb{R}^+ \trianglelefteq \mathbb{R}^\times$ and \det is a surjective map.)

Theorem 2.1 (Correspondence Theorem). *Let $\phi : G \rightarrow G'$ be a surjective group homomorphism with kernel K . There is a bijective correspondence between subgroups of G' and subgroups of G that contain K :*

$$\{\text{subgroups of } G \text{ that contain } K\} \leftrightarrow \{\text{subgroups of } G'\}$$

This correspondence is defined as follows:

$$\begin{aligned} \text{a subgroup } H \text{ of } G \text{ that contains } K &\rightsquigarrow \text{its image } \phi(H) \in G', \\ \text{a subgroup } H' \text{ of } G' &\rightsquigarrow \text{its inverse image } \phi^{-1}(H') \text{ in } G. \end{aligned}$$

- If H and H' are corresponding subgroups, then H is normal in G if and only if H' is normal in G' .
- If H and H' are corresponding subgroups, then

$$|H| = |H'||K|$$

Proof. Let H be subgroup of G that contain K . Let H' be a subgroup of G' . Now we need to check folllwings,

- $\phi(H)$ is a subgroup of G' .
- $\phi^{-1}(H')$ is a subgroup of G , and it contains K .
- H' is a normal subgroup of G' if and only if $\phi^{-1}(H')$ is a normal subgroup of G .
- (bijectivity of the correspondence) $\phi(\phi^{-1}(H')) = H'$ and $\phi^{-1}(\phi(H)) = H$
- $|(\phi^{-1}(H'))| = |H'||K|$.
- **Claim 1:** $\phi(H)$ is a subgroup of G' .
 - *Closure:* Let $x, y \in \phi(H)$. Then there is $a, b \in H$ such that $\phi(a) = x$ and $\phi(b) = y$. Since ϕ is homomorphism, $xy = \phi(a)\phi(b) = \phi(ab)$. Since $H \leq G$, $ab \in H$, $xy = \phi(ab) \in \phi(H)$.
 - *Identity:* Since $1_G \in H$, $\phi(1_G) = 1_{G'} \in \phi(H)$
 - *Inverse:* Let $x \in \phi(H)$. Then there exist $a \in H$ such that $\phi(a) = x$. Since $H \leq G$, a^{-1} exists in H . $\phi(a^{-1}) = \phi(a)^{-1} = x^{-1} \in \phi(G)$.
- **Claim 2:** $\phi^{-1}(H')$ is a subgroup of G , and it contains K .
This is true from proportion 2.6
- **Claim 3:** H' is a normal subgroup of G' if and only if $\phi^{-1}(H')$ is a normal subgroup of G .
Alreday this prooved in proportion 2.6
- **Claim 4.1:** $\phi(\phi^{-1}(H')) = H'$

- $\phi(\phi^{-1}(H')) \subset H'$
Lett $x \in \phi(\phi^{-1}(H'))$. Then there exist $y \in \phi^{-1}(H')$ such that $\phi(y) = x$. Then by definiton of the pre image $x = \phi(y) \in H'$.
- $\phi(\phi^{-1}(H')) \supset H'$
Let $a \in H'$. Since ϕ is surjective, there exists $b \in G$ such that $\phi(b) = a \in H'$. Thus $b \in \phi^{-1}(H')$. Hence $a = \phi(b) = \phi(\phi^{-1}(H'))$.

Therefore, $\phi(\phi^{-1}(H')) = H'$.

• **Claim 4.2:** $\phi^{-1}(\phi(H)) = H$

- $\phi^{-1}(\phi(H)) \supset H$
Let $x \in H$. Then $\phi(x) \in \phi(H)$. So, $x \in \phi^{-1}(\phi(H))$.
- $\phi^{-1}(\phi(H)) \subset H$
Let $y \in \phi^{-1}(\phi(H))$. By definition of inverse image, $\phi(y) \in \phi(H)$. Then there exist $z \in H$ such that $\phi(y) = \phi(z)$. Then $z^{-1}y$ is in the kernal K . (by Artins book proposition 2.5.8). Since $K \subset H$, $z^{-1}y \in H$. So, $a \in H$ and $a^{-1}z \in H$. Thus, $a(a^{-1}x) = x \in H$. Hence $\phi^{-1}(\phi(H)) \subset H$

Therefore, $\phi^{-1}(\phi(H)) = H$ - **Claim 5** : $|(\phi^{-1}(H'))| = |H'||K|$.

□

2.10.2 Problem

Example 2.16. Recall

$$\begin{aligned} S_4 &= \{id, (12), (13), (14), (23), (24), (34), (123), (124), (132), (134), (142), (143), (234), (1234)\} \\ S_3 &= \{id, (12), (13), (23), (123), (132)\} \end{aligned}$$

There are 6 such subgroups of S_3 ,

$$\langle (12) \rangle, \langle (13) \rangle, \langle (23) \rangle, \langle (123) \rangle = \langle (132) \rangle, S_3$$

. There is one proper subgroup of order 3. That is $\langle (123) \rangle$. There are 3 subgroups of order 2. They are $\langle (12) \rangle, \langle (13) \rangle, \langle (23) \rangle$

The Correspondence Theorem tells us that there are four proper subgroups of S_4 that contain K .

2.10.3 Problem

2.11 Product Groups

Let G, G' be two groups. The product set $G \times G'$, the set of pairs of elements (a, a') with a in G and a' in G' , can be made into a group by component-wise multiplication,

multiplication of pairs is defined by the rule,

$$(a, a') \cdot (b, b') = (ab, a'b') \quad \text{for } a, b \in G \text{ and for } a', b' \in G'$$

Let's prove that $G \times G'$ is a group.

Proof. Let G, G' be two groups and let $a, b, c \in G$ and $a', b', c' \in G'$

- *Closure* : $(a, a') \cdot (b, b') = (ab, a'b')$. So, since $a, b \in G$ and $a', b' \in G'$ and G and G' be a group, then $ab \in G$ and $a'b' \in G$. Thus, $(a, a') \cdot (b, b') = (ab, a'b') \in G \times G'$.
- *Asscitivity*: We can obtain following using asscitivity property of group G and G' .

$$((a, a') \cdot (b, b')) \cdot (c, c') = (ab, a'b') \cdot (c, c') = (abc, a'b'c') = (a, a') \cdot (bc, b'c') = (a, a') \cdot ((b, b') \cdot (c, c'))$$

- *identity*: $(1_G, 1_{G'})$ is the identity

$$(1_G, 1_{G'}) \cdot (b, b') = (1_G b, 1_{G'} b') = (b, b') = (b 1_G, b' 1_{G'}) = (b, b') \cdot (1_G, 1_{G'})$$


- *Inverse* : The inverse of (a, a') is $(a^{-1}, (a')^{-1})$

$$(a, a') \cdot (a^{-1}, (a')^{-1}) = (aa^{-1}, a'(a')^{-1}) = (1_G, 1_{G'}) = (a^{-1}a, (a')^{-1}a') = (a^{-1}, (a')^{-1}) \cdot (a, a')$$

□

So, The group obtained in this way is called the product of G and G' .

It is related to the two factors G and G' in a simple way that we can sum up in



terms of some homomorphisms.

The homomorphisms are defined as follows,

$$i : G \rightarrow G \times G' \quad (2.12)$$

$$x \mapsto (x, 1) \quad (2.13)$$

(2.14)

$$i' : G' \rightarrow G \times G' \quad (2.15)$$

$$x \mapsto (x', 1) \quad (2.16)$$

(2.17)

$$p : G \times G' \rightarrow G' \quad (2.18)$$

$$(x, x') \mapsto x \quad (2.19)$$

(2.20)

$$p' : G' \times G' \rightarrow G' \quad (2.21)$$

$$(x, x') \mapsto x' \quad (2.22)$$

(2.23)

(2.24)

Observe that i and i' are injective and

$$Im(i) = G \times 1'_{G'} \leq G \times G' \text{ and } Im(i') = 1_G \times G' \leq G \times G'$$

The maps p and p' are called projections and they are surjective.

$$\ker(p) = 1 \times G' \text{ and } \ker(p') = G \times 1_{G'}$$

It is obviously desirable to decompose a given group G as a product, that is, to find investigate groups H and H' such that G is isomorphic to the product $H \times H'$. The groups H and H' will be simpler, and the relation between $H \times H'$ and its factors is easily understood. It is rare that a group is a product, but it does happen occasionally

Example 2.17. Consider a cyclic group of order 6 can be decomposed. It might be suprised you. A cyclic group C_6 of order 6 is isomorphic to the product $C_2 \times C_3$ of cyclic groups of orders 2 and 3.

$$C_6 \equiv C_2 \times C_3$$

Let say

$$C_2 = \langle y \rangle = \{1, y\}, \text{ with } y^2 = 1 \quad (2.25)$$

$$C_3 = \langle z \rangle = \{1, z, z^2\}, \text{ with } y^3 = 1 \quad (2.26)$$

Let $x \in C_2 \times C_3$. Then there exist $p \in C_2$ and $q \in C_3$ such that $x = (p, q)$. Let's find order of x , that is the smallest positive integer k such that $x^k = (y^k, z^k)$ is the identity $(1, 1)$.


- Case-I: $k = 1, x^1 = (y^1, z^1) = (y, z)$
- Case-II: $k = 2, x^2 = (y^2, z^2) = (1, z^2)$
- Case-III: $k = 3, x^3 = (y^3, z^3) = (y, 1)$
- Case-IV: $k = 4, x^4 = (y^4, z^4) = (1, z)$
- Case-V: $k = 5, x^5 = (y^5, z^5) = (1, z^2)$
- Case-VI: $k = 6, x^6 = (y^6, z^6) = (1, 1)$

Thus, the smallest positive integer k such that $x^k = (1, 1)$ is 6. So, order of x is 6. Since $C_2 \times C_3$ has order 6. Furthur,

$$C_2 \times C_3 = \langle x \rangle$$

The powers of x are

$$(1, 1), (y, z), (1, z^2), (y, 1), (1, z), (y, z^2)$$



Here $i = 0, 1, 2, \dots, 5$

So let's try to see above result more generally.

Proposition 2.7. *Let r and s be relatively prime integers. A cyclic group of order rs is isomorphic to the product of a cyclic group of order r and a cyclic group of order s .*

Example 2.18 (s). On the other hand, a cyclic group of order 4 is not isomorphic to a product of two cyclic groups of order 2.

$$C_4 \not\cong C_2 \times C_2$$

Because , every element of $C_2 \times C_2$ has order 1 or 2, whereas a cyclic group of order 4 contains two elements of order 4.

The next proposition describes product groups.

Proposition 2.8. *Let G be a group and let $H, K \leq G$, and let $f : H \times K \rightarrow G$ be the multiplication map, defined by $f(h, k) = hk$. Its image is the set $HK = \{hk | h \in H, k \in K\}$.*

- a. *f is injective if and only if $H \cap K = \{1\}$.*
- b. *f is a homomorphism from the product group $H \times K$ to G if and only if elements of K commute with elements of H : $hk = kh$.*
- c. *If H is a normal subgroup of G , then HK is a subgroup of G .*
- d. *f is an isomorphism from the product group $H \times K$ to G if and only if $H \cap K = \{1\}$, $HK = G$, and also H and K are normal subgroups of G .*

It is important to note that the multiplication map may be bijective though it isn't a group homomorphism. This happens, for instance, when $G = S_3$, and with the usual notation, $H = \langle x \rangle$ and $K = \langle y \rangle$.

Proof.

a.

- (\Rightarrow) We are going to use proof by contrapositive. So, suppose that $x \in H \cap K$ such that $x \neq 1$. Since $x \in H$ then $x^{-1} \in H$.

$$f(x^{-1}, x) = x^{-1} \cdot x = 1 = 1 \cdot 1 = f(1, 1)$$

Thus, f is not injective. Thus, if f is injective then $H \cap K = \{1\}$. -(\Leftarrow) Now Suppose that $H \cap K = \{1\}$. Let $(h_1, k_1), (h_2, k_2) \in H \times K$ such that $h_1k_1 = h_2k_2$. Now multiply both sides of this equation on the left by h_1^{-1} and on the right by k_2^{-1} ,

$$h_1k_1 = h_2k_2 \tag{2.27}$$

$$h_1^{-1}(h_1k_1)k_2^{-1} = h_1^{-1}(h_2k_2)k_2^{-1} \tag{2.28}$$

$$(h_1^{-1}h_1)k_1k_2^{-1} = h_1^{-1}h_2(k_2k_2^{-1}) \tag{2.29}$$

$$k_1k_2^{-1} = h_1^{-1}h_2 \tag{2.30}$$

Note that $(k_1 k_2^{-1}) \in K$ and $h_1^{-1} h_2 \in H$. Since $H \cap K = \{1\}$,

$$k_1 k_2^{-1} = h_1^{-1} h_2 = 1$$

. Then,

$$k_1 = k_2 \quad \text{and} \quad h_1 = h_2.$$

Then

$$(h_1, k_1) = (h_2, k_2)$$

b. Let $(h_1, k_1), (h_2, k_2) \in H \times K$. Now consider,

$$f((h_1, k_1) \cdot (h_2, k_2)) = f((h_1 h_2, k_1 k_2)) = h_1 h_2 k_1 k_2 \quad (2.31)$$

$$f((h_1, k_1)) \cdot f((h_2, k_2)) = (h_1 k_1) \cdot (h_2 k_2) = h_1 k_1 h_2 k_2 \quad (2.32)$$

$$\begin{aligned} f \text{ is homomorphism} &\iff f((h_1, k_1) \cdot (h_2, k_2)) = f((h_1, k_1)) \cdot f((h_2, k_2)) \\ &\iff h_1 h_2 k_1 k_2 = h_1 k_1 h_2 k_2 \quad (2.34) \\ &\iff h_1^{-1} (h_1 h_2 k_1 k_2) k_2^{-1} = h_1^{-1} (h_1 k_1 h_2 k_2) k_2^{-1} \quad (2.35) \\ &\iff (h_1^{-1} h_1) h_2 k_1 (k_2 k_2^{-1}) = (h_1^{-1} h_1) k_1 h_2 (k_2 k_2^{-1}) \quad (2.36) \\ &\iff (h_1^{-1} h_1) h_2 k_1 (k_2 k_2^{-1}) = (h_1^{-1} h_1) k_1 h_2 (k_2 k_2^{-1}) \quad (2.37) \\ &\iff h_2 k_1 = k_1 h_2 \quad (2.38) \end{aligned}$$

c.

- Suppose that H is a normal sub group of G . Note that

$$KH = \bigcup_{k \in K} kH \quad \text{and} \quad HK = \bigcup_{k \in K} Hk$$

Since, H is normal, $kH = Hk$ for all $k \in K$. So, $HK = KH$. We are going to use sub group test,

- *non-emptyness*: Clearly, $1 = 1 \cdot 1 \in HK$ (because $1 \in H$ and $1 \in K$)
- *closure*:

$$HKHK = HHKK = HK$$

Thus, HK is closed under multiplication.

- *closed under inverse*: Let $hk \in HK$. Then

$$(hk)^{-1} = k^{-1} \cdot h^{-1} \in KH = HK$$

This proves closure of HK under inverses.

d.

- (\Leftarrow): Suppose that $H \cap K = \{1\}$, $HK = G$, and also $H, K \trianglelefteq G$. According to the (b), f is a homomorphism from $H \times K$ to G if and only if $hk = kh$ for all $h \in H$. Consider,

$$(hkh^{-1})k^{-1} = h(kh^{-1}k^{-1}).$$

Note that, Since $K \trianglelefteq G$, $hkh^{-1} \in K$. So, $(hkh^{-1})k^{-1} \in K$. Similary we can show that $h(kh^{-1}k^{-1}) \in H$. Thus,

$$(hkh^{-1})k^{-1} = h(kh^{-1}k^{-1}) \in H \cap K.$$

But from our hypothesis, $H \cap K = \{1\}$. Hence,

$$hkh^{-1}k^{-1} = 1 \quad (2.39)$$

$$hk = kh \quad (2.40)$$

Therefore, by (b) we can f is hormorphism. Now we have to proove bijectivity,

- *Injectivity:* Already prooved in 2.8 (a)
- *Surjectivity:* For any $g \in G$, we can write $g = hk$ for some $h \in H$ and $k \in K$ (since $Im(f) = HK = G$). Thus, $f(h, k) = hk = g$, so f is surjective.

Therefore f is an isomorphism.

- (\Rightarrow) Now suppose that f is isomorphism.

- **Claim i:** $H \cap K = \{1\}$.

Suppose that $x \in H \cap K$. Then,

$$f(1, x) = x = f(x, 1)$$

Since f is an isomorphism (and hence injective), we must have $x = 1$. Therefore, $H \cap K = \{1\}$.

- **Claim ii :** $HK = G$.

Since f is surjective, for any $g \in G$, there exist $h \in H$ and $k \in K$ such that $f(h, k) = hk = g$. Therefore, $HK = G$.

- **Claim iii :** H and K are normal in G .

For any $h \in H$ and $g \in G$, we can write $g = hk$ for some $k \in K$. Then

$$ghg^{-1} = (hk)h(hk)^{-1} = hk(hh^{-1})k^{-1} = h(kk^{-1}) = h \in H.$$

Thus, $H \trianglelefteq G$. Similarly, we can show that $K \trianglelefteq G$.

□

Proposition 2.9. *There are two isomorphism classes of groups of order 4, the class of the cyclic group C_4 of order 4 and the class of the Klein Four Group, which is isomorphic to the product $C_2 \times C_2$ of two groups of order 2.*

Proof. Let G be a group of order 4. The order of any element x of G divides 4, so there are two cases to consider:

- Case 1: G contains an element of order 4 ($|G|=4$). Then G is a cyclic group of order 4.
- Case 2: Every element of G except the identity has order 2.

In this case, $x = x^{-1}$ for every element x of G . Let x and y be two elements of G . Then xy has order 2, so $(xy)(x^{-1}y^{-1}) = (xy)(xy) = 1$. This shows that x and y commute,

$$xyx^{-1}y^{-1} = 1 \quad (2.41)$$

$$xyx^{-1}(y^{-1}y) = y \quad (2.42)$$

$$xyx^{-1} = y \quad (2.43)$$

$$xy(x^{-1}x) = yx \quad (2.44)$$

$$xy = yx \quad (2.45)$$

since x, y are arbitrary elements, G is abelian. So every subgroup is normal. We choose distinct elements x and y in G , and we let H and K be the cyclic groups of order 2 that they generate. Proposition 2.8 (d) shows that G is isomorphic to the product group $H \times K$.

□

2.12 Quotient Groups


Definition 2.9. The set of cosets of a normal subgroup N of a group G is often denoted by G/N .

G/N is the set of cosets of N in G

Notation: When we regard a coset C as an element of the set of cosets, the bracket notation $[C]$ may be used. If $C = aN$, we may also use the bar notation to denote the element $[C]$ by \bar{a} , and then we would denote the set of cosets by \bar{G} :

$$\bar{G} = G/N$$

Theorem 2.2. Let N be a normal subgroup of a group G , and let \bar{G} denote the set of cosets of N in G . There is a law of composition on \bar{G} that makes this set into a group, such that the map $\pi : G \rightarrow \bar{G}$ defined by $\pi(a) = \bar{a}$ is a surjective homomorphism whose kernel is N .



Remark. The map is often referred to as the canonical map from G to \bar{G} . The word “canonical” indicates that this is the only map that we might reasonably be talking about

The next corollary is very simple, but it is important enough to single out:

Corollary 2.1. Let $N \trianglelefteq G$, and let $\bar{G} = G/N$ be the canonical homomorphism. Let $a_1, \dots, a_k \in G$ such that the product $a_1 \cdots a_k \in N$. Then $\bar{a}_1 \cdots \bar{a}_k = \bar{1}$.

Proof. Let $p = a_1 \cdots a_k \in N$. This implies $\pi(p) = \bar{p} = \bar{1}$.

Since π is a homomorphism, $\pi(p) = \pi(a_1 \cdots a_k) = \pi(a_1) \cdots \pi(a_k) = \bar{a}_1 \cdots \bar{a}_k$ \square

Proof of 2.2 There are several things to be done. We must

- define a law of composition on \bar{G} ,
- prove that the law makes \bar{G} into a group,
- prove that π is a surjective homomorphism, and
- prove that the kernel of π is N .

Remark. If $A, B \subseteq G$ then AB denotes the set of products ab :

$$AB := \{x \in G : x = ab, a \in A, b \in B\}$$

- We will call this a product set, though in some other contexts the phrase “product set” refers to the set $A \times B$ of pairs of elements

Lemma 2.2. *Let N be a normal subgroup of a group G , and let aN and bN be cosets of N . The product set $(aN)(bN)$ is also a coset and*

$$(aN)(bN) = \{x \in G : x = anbm' \& n, n' \in N\} = abN$$

Proof. Since N is a subgroup, $NN = N$.

Since N is normal, left and right cosets are equal: $Nb = bN$.

$$(aN)(bN) = a(Nb)N = a(bN)N = abNN = abN$$

□

- This lemma allows us to define multiplication on the set $\overline{G} = G/N$.
- Using the bracket notation, the definition is this: If C_1 and C_2 are cosets, then $[C_1][C_2] = [C_1C_2]$, Where C_1C_2 is the product set.
- The lemma shows that this product set is another coset.
- To compute the product $[C_1][C_2]$, take any elements $a \in C_1$ and $b \in C_2$. Then $C_1 = aN$, $C_2 = bN$, and C_1C_2 is the coset abN that contains ab . So,

$$[aN][bN] = [abN] \text{ or } \overline{ab} = \overline{ab}.$$

Then by definition of the map π in theorem 2.2,

$$\pi(ab) = \overline{ab} = \overline{ab} = \pi(a)\pi(b)$$

The fact that π is a homomorphism will follow, once we show that \overline{G} is a group. Since the canonical map π is surjective, the next lemma proves this

Lemma 2.3. *Let G be a group, and let Y be a set with a law of composition (both laws written with multiplicative notation). Consider a surjective map $\varphi : G \rightarrow Y$ with the homomorphism property: $\varphi(ab) = \varphi(a)\varphi(b)$ for all a and b in G . Then Y is a group, and φ is a homomorphism.*

Proof. The group axioms that are true in G are carried over to Y by the surjective map φ .

- **Closure** : Let $y_1, y_2 \in Y$. Since φ is surjective, $y_1 = \varphi(x_1), y_2 = \varphi(x_2)$ for some $x_1, x_2 \in G$.

$$y_1 y_2 = \varphi(x_1) \varphi(x_2) = \varphi(x_1 x_2) \in Y$$

- **Associativity Property** :

Let y_1, y_2 , and y_3 be elements of Y . Since φ is surjective, $y_i = \varphi(x_i)$ for some x_i in G . Then

$$(y_1 y_2) y_3 = (\varphi(x_1) \varphi(x_2)) \varphi(x_3) = \varphi(x_1 x_2) \varphi(x_3) = \varphi((x_1 x_2) x_3) \quad (2.46)$$

$$=^* \varphi(x_1 (x_2 x_3)) = \varphi(x_1) \varphi(x_2 x_3) = y_1 (y_2 y_3) \quad (2.47)$$

The equality marked with an asterisk ($=^*$) is the associative law in G . The other equalities follow from the homomorphism property of φ .

- **Identity** Let $y \in Y$. Since φ is surjective, there exist $x \in G$ such that $y = \varphi(x)$. Let 1_G be identity of G . Then,

$$y \varphi(1_G) = \varphi(x) \varphi(1_G) = \varphi(x 1_G) = \varphi(x) = y = \varphi(1_G x) = \varphi(1_G) \varphi(x) = \varphi(1_G) y$$

Thus, $\varphi(1_G)$ is the identity of Y .

- **Inverse** : Let $y \in G$. Since φ is surjective, there exist $x \in G$ such that $y = \varphi(x)$,

$$y \varphi(x^{-1}) = \varphi(x) \varphi(x^{-1}) = \varphi(x x^{-1}) = \varphi(1_G) = \varphi(x^{-1} x) = \varphi(x^{-1}) \varphi(x) = \varphi(x^{-1}) y$$

Thus, $y^{-1} = \varphi(x)$

□

The only thing remaining to be verified is that the kernel of the homomorphism π is the subgroup N .

$$\pi(a) = \pi(1) \iff \bar{a} = \bar{1} \quad (2.48)$$

$$\iff [aN] = [1N] \quad (2.49)$$

$$\iff a \in N. \quad (2.50)$$

Thus, $\ker(\pi) = N$




Figure 2.1: A Schematic Diagram of Coset Multiplication.

Remark. Our assumption that N is a **normal** subgroup of G is crucial to lemma 2.2. If H is **not** normal, there will be left cosets C_1 and C_2 of H in G such that the product set C_1C_2 does not lie in a single left cosets.

Let's see an example for this. Going back once more to the subgroup $H = \langle y \rangle$ of S_3 . Note that the subgroup H is not normal.


The product set $(1H)(xH)$ contains four elements:

$$(1H)(xH) = \{1, y\}\{x, xy\} = \{x, xy, x^2y, x^2\}$$

. It is not a coset.

The next theorem relates the quotient group construction to a general group homomorphism, and it provides a fundamental method of identifying quotient groups.

Theorem 2.3 (First Isomorphism Theorem). *Let $\varphi : G \rightarrow G'$ be a surjective group homomorphism with kernel N . The quotient group $\bar{G} = G/N$ is isomorphic to the image G' . To be precise, let $\pi : G \rightarrow \bar{G} = G/N$ be the canonical map. There exists a unique isomorphism $\bar{\varphi} : G/N \rightarrow G'$ such that $\varphi = \psi \circ \pi$.*



Proof. The elements of $\overline{G} = G/N$ are the cosets of N , and they are also the fibres of the map φ . The map $\bar{\varphi}$ referred to in the theorem is the one that sends a non-empty fibre to its image: $\bar{\varphi}(\overline{x}) = \varphi(x)$. For any surjective map of sets $\varphi : G \rightarrow G'$, one can form the set \overline{G} of fibres, and then one obtains a diagram as above, in which $\bar{\varphi}$ is the bijective map that sends a fibre to its image. When φ is a group homomorphism, $\bar{\varphi}$ is an isomorphism because

$$\bar{\varphi}(\overline{ab}) = \varphi(ab) = \varphi(a)\varphi(b) = \bar{\varphi}(\overline{a})\bar{\varphi}(\overline{b}).$$

□

Corollary 2.2. Let $\varphi : G \rightarrow G'$ be a group homomorphism with kernel N and image H' . The quotient group $\overline{G} = G/N$ is isomorphic to the image H' .

2.13 Exercises

2.13.1 Laws of composition

Exercise 2.1. Let S be a set. Prove that the law of composition defined by $ab = a$ for all a and b in S is associative? For which sets does this law have an identity?

Solution: Let $a, b, c \in S$. Now consider following

$$(ab)c = (ac) = a = (ab) = a(bc)$$

Thus, the given law of composition is associative.

If the given law of composition has an identity element whenever every element $a \in S$ has a multiplicative inverse./ In other words, for every element $a \in S$, there exists an element $e \in S$ such that

$$ae = ea = a.$$

Thus, $ae = a = ea = e$. So, the identity element is the same as every element in S , and the law has an identity for all sets S . Thus, only singletons sets have this given law of compositions have identity. # Groups and Subgroups :: {.exercise #unnamed-chunk-61} Make a multiplication table for the symmetric group S_3 . :: **Solution**

\circ	e	(12)	(13)	(23)	(123)	(132)
e	e	(12)	(13)	(23)	(123)	(132)
(12)	(12)	e	(123)	(132)	(13)	(23)
(13)	(13)	(123)	e	(132)	(23)	(12)
(23)	(23)	(132)	(123)	e	(12)	(13)
(123)	(123)	(13)	(23)	(12)	(132)	e
(132)	(132)	(23)	(12)	(13)	e	(123)

Exercise 2.2. Let S be a set with an associative law of composition and with an identity element. Prove that the subset consisting of the invertible elements in S is a group.

Proof. Let $S^* \subseteq S$ with consisting of the invertible elements in S .

- *identity* : We are given identity element $1 \in S$ and $(1^{-1})(1) = 1 \implies 1 \in S^*$
- *Closure* : So let $a, b \in S^*$. Then a, b are invertible. Let $a^{-1}, b^{-1} \in S^*$ be inverses of a and b respectively. Now we need to check $ab \in S^*$.

$$(b^{-1}a^{-1})(ab) = b^{-1}(a^{-1}a)b = b^{-1}b = e = aa^{-1} = a(bb^{-1})a^{-1} = (ab)(b^{-1}a^{-1})$$

□

Thus, $(ab)^{-1} = b^{-1}a^{-1}$. So, $ab \in S^*$. Therefore, S^* is closed under the composition.

- *Associativity*: The associativity property inherit from the group S .

Therefore, S^* is a group.

Exercise 2.3. Let G be a group. Let $x, y, z, w \in G$.

- Solve for y , given that $xyz^{-1}w = 1$.

- (b) Suppose that $xyz = 1$. Does it follow that $yzx = 1$? Does it follow that $yxz = 1$?

Solution:

Let $x, y, z, w \in G$

$$\begin{aligned} & xyx^{-1}w = 1 \\ & x^{-1}(xyx^{-1}w) = x^{-1}(1) \quad (\text{Property of equality}) \\ & (x^{-1}x)(y x^{-1}w) = x^{-1} \quad (\text{Associativity}) \\ & 1(y x^{-1}w) = x^{-1} \quad (\text{Identity inverse}) \\ & y x^{-1}w = x^{-1} \quad (\text{Identity}) \\ & y(x^{-1}w)w^{-1} = x^{-1}w^{-1} \quad (\text{Property of equality}) \\ & y(x^{-1}w)w^{-1} = x^{-1}w^{-1} \quad (\text{Associativity}) \\ & y(x^{-1}w^{-1}) = x^{-1}w^{-1} \quad (\text{Inverse}) \\ & y(x^{-1}w^{-1}) = x^{-1}w^{-1} \quad (\text{Identity}) \\ & yz = x^{-1}w^{-1}z \quad (\text{Property of equality}) \\ & wz^{-1}z = x^{-1}w^{-1}z \quad (\text{Associativity}) \\ & yz = x^{-1}w^{-1}z \quad (\text{Inverse}) \\ & y = x^{-1}w^{-1}z \quad (\text{Identity}) \end{aligned}$$

a.

b.

$\begin{aligned} & xyz = 1 \\ & x^{-1}(xyz) = x^{-1} \\ & (x^{-1}x)(yz) = x^{-1} \\ & 1(yz) = x^{-1} \end{aligned}$	$\begin{aligned} & yz = x^{-1} \\ & (yz)(x) = x^{-1}x = 1 \\ & yzx = 1 \end{aligned}$
---	---

- The given statement is false.

$$xyz = 1 \quad (2.51)$$

$$x^{-1}(xyz) = x^{-1} \cdot 1 \quad (2.52)$$

$$(x^{-1}x)(yz) = x^{-1} \quad (2.53)$$

$$1 \cdot (yz) = x^{-1} \quad (2.54)$$

$$(yz) = x^{-1} \quad (2.55)$$

$$(yz)z^{-1} = x^{-1}z^{-1} \quad (2.56)$$

$$y(zz^{-1}) = x^{-1}z^{-1} \quad (2.57)$$

$$y \cdot 1 = x^{-1}z^{-1} \quad (2.58)$$

$$yx = x^{-1}z^{-1}x \quad (2.59)$$

$$yxz = x^{-1}z^{-1}xz \quad (2.60)$$

$$(2.61)$$

If G is not abelian or $z \neq x$. $xyz = 1 \Rightarrow yxz = 1$.

Exercise 2.4. In which of the following cases is H a subgroup of G ?

(a) $G = GL_n(\mathbb{C})$ and $H = GL_n(\mathbb{R})$.

(b) $G = \mathbb{R}^\times$ and $H = \{1, -1\}$.

(c) $G = \mathbb{Z}^+$ and H is the set of positive integers.

(d) $G = \mathbb{R}^+$ and H is the set of positive reals.

(e) $G = GL_2(\mathbb{R})$ and H is the set of matrices $\begin{bmatrix} a & 0 \\ 0 & 0 \end{bmatrix}$ with all entries equal to 0.

Solution:

(a)

- *Subset ?* : Since every matrix with real entries can be interpreted as a matrix with complex entries, we conclude that $H \subset G$.
- *Closure ?* : For every $A, B \in GL_n(\mathbb{R})$ we have $AB \in GL_n(\mathbb{R})$ since Product of invertible matrices is invertible. I proved this result in chapter 1.
- *Identity ?* : The identity in G is I_n , the identity $n \times n$ matrix. Observe that $I_n \in H = GL_n(\mathbb{R})$. (Because every entry of I_n is 1 or 0)

- *Inverse ?*: $GL_n(\mathbb{R})$ is the set of $n \times n$ invertible matrices with real entries. So, all the elements in H is invertible.

Therefore, $H \leq G$

(b)

- *Subset ?* : This is trivial $\{-1, 1\} \subset \mathbb{R}^\times$
- *Closure ?* :

$$1 \times 1 = 1 \in H \quad (2.62)$$

$$(-1) \times 1 = (-1) \in H \quad (2.63)$$

$$1 \times (-1) = (-1) \in H \quad (2.64)$$

$$(-1) \times (-1) = -1 \in H \quad (2.65)$$

$$(2.66)$$

Thus, H is closed.

- *Identity ?* : The identity in G is 1, which is also in H .
- *Inverse ?* :

$$1 \times 1 = 1 \implies (1)^{-1} = 1 \in H \quad (2.67)$$

$$(-1) \times (-1) = 1 \implies (-1)^{-1} = (-1) \in H \quad (2.68)$$

Therefore, H is sub group of G .

- (c) Note that $2 \in H$, but inverse of 2 in \mathbb{Z}^+ is $(-2) \notin H$.
Therefore, H is not a subgroup of G .

(d)

- *Subset ?* : This is trivial. $H \subset \mathbb{R}^\times$
- *Closure ?* : Product of positive real number is positive. Thus, H is closed.
- *Identity ?* : 1 is the identity of G , which is in H .
- *Inverse ?* : Let $a \in H$. Then,

$$a \cdot \frac{1}{a} = \frac{1}{a} \cdot a = 1$$

So, $\frac{1}{a} \in H$ is the inverse of a . Thus, H is closed under inverses.

- (e) Let $A = \begin{bmatrix} 2 & 0 \\ 0 & 0 \end{bmatrix}$. Then $A \in H$. But observe that $\det(A) = 0 \implies A$ is not invertible. Thus, $A \notin H$. Thus, this $H \not\subseteq G$.

Therefore, H is not a subgroup of G .

Exercise 2.5. In the definition of a subgroup, the identity element in H is required to be the identity of G . One might require only that H have an identity element, not that it need be the same as the identity in G . - Show that if H has an identity at all, then it is the identity in G . - Show that the analogous statement is true for inverses.

Exercise 2.6. Let G be a group. We define an opposite group G° with the law of composition $a * b$ as follows: - The underlying set is the same as G , - but the law of composition is $a * b = ba$.

Prove that G° is a group.

Solution:

Let G be a group. Define opposite group G° with law of composition $a * b$ as follows.

- The underline set is same as G .
- Law of composition $a * b = ba$.

Need to show ~~\underline{G}~~ G° is a group.

Let $a, b, c \in G^\circ$. Then $a, b, c \in G$.

- Closure: $a * b = ba \in G \Rightarrow a * b = ba \in G^\circ$.
Thus ~~\underline{G}~~ is closed under $*$.
- Identity: Let e be the identity of G .
Then $e \in G^\circ$.
Claim: Observe following.
 $e * a = ae = a = ea = a * e$.
Thus $e \in G^\circ$ is the identity of G° .
- Inverse: Since G is a group
 $\bar{a} \in G$ exists. Then $\bar{a} \in G^\circ$. Observe
 $\bar{a}^{-1} * a = a\bar{a} = e = \bar{a}a = a * \bar{a}$.
Thus, \bar{a}^{-1} is the inverse of a in G° .
- Associativity:

$$\begin{aligned} (a * (b * c)) &= a * (c * b) = (cb)a = c(ba) \quad (\text{Since } G \text{ is a group}) \\ &= c(a * b) = (a * b) * c \end{aligned}$$

Thus associativity holds in G° .

Therefore, G° is a group.

2.13.2 Subgroups of the Additive Group of Integers

Exercise 2.7. Let $a = 123$ and $b = 321$. Compute $d = \gcd(a, b)$, and express d as an integer combination $ra + bs$.

We use Euclidean Algorithm.

$$\begin{aligned}
 321 &= 2(123) + 75 \\
 123 &= 1(75) + 48 \\
 75 &= 1(48) + 27 \\
 48 &= 1(27) + 21 \\
 27 &= 1(21) + 6 \\
 21 &= 3(6) + 3 \\
 6 &= 3(2) + 0
 \end{aligned}$$

Thus $\gcd(123, 321) = 3$.
Now let's back-substitute it.

$$\begin{aligned}
 3 &= 21 - 3(6) \\
 &= 1(21) - 3(27 - 1(21)) \\
 &= 1(21) - 3(27) + 3(21) \\
 &= 4(21) - 3(27) \\
 &= 4(48 - 1(27)) - 3(27) \\
 &= 4(48) - 7(27) \\
 &= 4(48) - 7(75 - 1(48)) \\
 &= 11(48) - 7(75) \\
 &= 11(123 - 1(75)) - 7(75) \\
 &= 11(123) - 18(75) \\
 &= 11(123) - 18(321 - 2(123)) \\
 &= 47(123) - 18(321)
 \end{aligned}$$

Solution:

Exercise 2.8. Prove that if a and b are positive integers whose sum is a prime p , their greatest common divisor is 1.

Let $a, b \in \mathbb{Z}^+$ with
 $a+b = p$
 Let $d = \gcd(a, b)$
 $a|d$ and $b|d$
 $a = dk_1$ and $b = dk_2$
 for some $k_1, k_2 \in \mathbb{Z}$
 $a+b = dk_1 + dk_2$
 $a+b = d(k_1 + k_2)$
 Since $(a+b)$ is a prime
 the only value can get is 1.

Solution:

Exercise 2.9. Define the greatest common divisor of a set $\{a_1, \dots, a_n\}$ of n integers. Prove that it exists, and that it is an integer combination of a_1, \dots, a_n .

2.13.3 Cyclic group

Exercise 2.10. Let a and b be elements of a group G . Assume that a has order 7 and that $a^3b = ba^3$. Prove that $ab = ba$.

Solution:

$$\begin{aligned}
 & \text{Given that } a^7 = 1 \\
 & a(a^6) = 1 = (a^6)a \\
 & \text{Thus, } a^{-1} = a^6 \\
 \\
 & ba^3 = a^3b \\
 & ba^3b^{-1} = a^3 \\
 & (ba^3b^{-1})^2 = (a^3)^2 \\
 & (ba^3b^{-1})(ba^3b^{-1}) = a^6 \\
 & ba^3(b^{-1}b)a^3b^{-1} = a^{-1} \\
 & ba^6b^{-1} = a^{-1} \\
 & ba^6 = a^{-1}b \\
 & ba^{-1} = a^{-1}b \\
 & ab^{-1} = b \\
 & ab = ba
 \end{aligned}$$

Exercise 2.11. An n th root of unity is a complex number z such that $z^n = 1$.

- (a) Prove that the n th roots of unity form a cyclic subgroup of \mathbb{C}^\times of order n .
- (b) Determine the product of all the n th roots of unity.

If $z^n = 1$ then

$$z = \exp\left(\frac{2\pi i k}{n}\right), \quad k=0, 1, \dots, n-1$$

$$\begin{aligned} \text{Let } S &:= \left\{ e^{\frac{2\pi i k}{n}} \mid k=0, 1, 2, \dots, n-1 \right\} \\ &= \left\{ e^0, e^{\frac{2\pi i}{n}}, \dots, e^{\frac{2\pi i(n-1)}{n}} \right\} \end{aligned}$$

Solution: First let's check S is a subgroup of \mathbb{C}^\times

- Subset: It is trivial that $S \subseteq \mathbb{C}^\times$

- Closure: Let $x, y \in S$. Then

$$x = e^{\frac{2\pi i k_1}{n}} \text{ and}$$

$$y = e^{\frac{2\pi i k_2}{n}} \text{ for some } k_1, k_2 = 0, 1, \dots, n-1$$

$$\begin{aligned} \text{Then, } xy &= e^{\frac{(2\pi i k_1)}{n}} \cdot e^{\frac{(2\pi i k_2)}{n}} \\ &= e^{\frac{2\pi i (k_1 + k_2)}{n}} \end{aligned}$$

If $(k_1+k_2) > n$, we can find $a, b \in \mathbb{N}$
such that (by division algorithm)

$$k_1 + k_2 = an + b, \quad 0 \leq b < n$$

$$\begin{aligned} xy &= e^{\frac{2\pi i(k_1+k_2)}{n}} \\ &= e^{\frac{2\pi i}{n}(an+b)} \\ &= e^{2\pi i(a)} \cdot e^{\frac{2\pi i b}{n}} \\ &= 1 \times e^{\frac{2\pi i b}{n}} \\ &= e^{\frac{2\pi i b}{n}}, \text{ for some } b = 0, 1, 2, \dots, n-1 \end{aligned}$$

Thus, $xy \in S$

- Inverse: Let $x \in S$.

◦ If $x=1$, then $x^{-1}=1$.

◦ If $x \neq 1$,

Then $x = e^{\exp(2\pi i k_1/n)}$ for some $k_1 = 1, 2, \dots, n-1$
choose $y = e^{\exp(2\pi i (n-k_1)/n)}$

$$\text{Then, } yx = xy = e^{\frac{2\pi i k_1}{n}} \cdot e^{\frac{2\pi i (n-k_1)}{n}} = e^{\frac{2\pi i (k_1 + n - k_1)}{n}} = e^{\frac{2\pi i n}{n}} = e^{2\pi i} = 1$$

Thus, $x^{-1} = y \in S$.

By subgroup test S is a subgroup of G

Now, we are done subgrouping. Now we have to look the cyclic property.

Observe that $(e^{\frac{2\pi i}{n}})^k = e^{\frac{2\pi i k}{n}}, k=0, 1, \dots, n-1$

Therefore, $S = \langle e^{\frac{2\pi i}{n}} \rangle$

$$\begin{aligned}
 b) \sum_{k=0}^{n-1} e^{\frac{2\pi i k}{n}} &= e^{\left(\sum_{k=0}^{n-1} \frac{2\pi i k}{n}\right)} \\
 &= e^{\frac{2\pi i}{n} \left(\sum_{k=0}^{n-1} k\right)} \\
 &= e^{\frac{2\pi i}{n} \left(\frac{n(n-1)}{2}\right)} \\
 &= e^{\pi i (n-1)} \\
 &= (e^{i\pi})^{n-1} \quad (\because e^{i\pi} = -1) \\
 &= (-1)^{n-1}
 \end{aligned}$$

Exercise 2.12. Let a and b be elements of a group G . Prove that ab and ba have the same order

Proof. Let a and b be elements in a group G with identity e .

Suppose that ab has finite order n . Then $(ab)^n = e$ and n is the smallest positive integer for which this equation is true. We also have that

$$(ba)^{n+1} = (ba)(ba) \cdots (ba) = b \underbrace{(ab) \cdots (ab)}_{n \text{ times}} a = b(ab)^n a = a(ba)^m b = bea = ba.$$

Thus, since $(ba)^{n+1} = (ba)^n(ba)$ we can conclude that $(ba)^n(ba) = ba$ and then by the cancellation law, we have that $(ba)^n = e$.

Now, to show that the order of ba is n , we need to demonstrate that n is the smallest positive integer such that $(ba)^n = e$.

Suppose there exists a positive integer $m < n$ such that $(ba)^m = e$. Then,

$$(ab)^{m+1} = (ab)(ab) \cdots (ab) = a \underbrace{(ba) \cdots (ba)}_{m \text{ times}} b = a(ba)^m b = aeb = ab.$$

Thus, since $(ab)^{m+1} = (ab)^m(ab)$ we have that $(ab)^m(ab) = ab$ and then by the cancellation law, we have that $(ab)^m = e$ which contradicts the fact that n is the smallest positive integer such that $(ab)^n = e$.

Hence, n is the smallest positive integer such that $(ba)^n = e$ and therefore (ba) has finite order n . \square

2.13.4 Modular Arithmetic

Exercise 2.13. What are the possible values of a^2 modulo 4? modulo 8?

solution In modulo 4

$$\begin{aligned}\bar{0}^2 &\equiv 0 \pmod{4} \\ \bar{1}^2 &\equiv 1 \pmod{4} \\ \bar{2}^2 &\equiv 0 \pmod{4} \\ \bar{3}^2 &\equiv 1 \pmod{4}\end{aligned}$$

The possible values of $a^2 \pmod{4}$ are 0 and 1.

In modulo 8

$$\begin{aligned}\bar{0}^2 &\equiv 0 \pmod{8} \\ \bar{1}^2 &\equiv 1 \pmod{8} \\ \bar{2}^2 &\equiv 4 \pmod{8} \\ \bar{3}^2 &\equiv 1 \pmod{8} \\ \bar{4}^2 &\equiv 0 \pmod{8} \\ \bar{5}^2 &\equiv 1 \pmod{8} \\ \bar{6}^2 &\equiv 4 \pmod{8} \\ \bar{7}^2 &\equiv 1 \pmod{8}\end{aligned}$$

The possible values of $a^2 \pmod{8}$ are 0,1 and 4.

Exercise 2.14. Prove that every integer a is congruent to the sum of its decimal digits modulo 9.

Proof. Let $x \in \mathbb{Z}$. Now we can represent x as follows

$$x = a_0 10^0 + a_1 10^1 + \cdots + a_n 10^n = \sum_{i=0}^n a_i 10^i \text{ forsome } n \in \mathbb{Z}, \text{ and } a_i \in \{0, 1, \dots, 9\}$$

We need to show $x \equiv \sum_{i=0}^n a_i \pmod{9}$. So, now consider,

$$x - \sum_{i=0}^n a_i = \sum_{i=0}^n a_i 10^i - \sum_{i=0}^n a_i \quad (2.69)$$

$$= \sum_{i=0}^n (a_i 10^i - a_i) \quad (2.70)$$

$$= \sum_{i=0}^n a_i (10^i - 1) \quad (2.71)$$

By following claim we can get that,

$$x - \sum_{i=0}^n a_i \equiv 0 \pmod{9} \quad (2.72)$$

$$x \equiv \sum_{i=0}^n a_i \pmod{9} \quad (2.73)$$

□

Claim: $9|(10^k - 1)$ for any $k \in \mathbb{N}$.

Proof. We use mathematical induction.

$k = 1$

This case is trivial. Because $9|10 - 1$.

$k = n \in \mathbb{Z}$

Now assume when $n = k, 9|(10^n - 1)$.

$k = n + 1$

$$10^{n+1} - 1 = 10 \cdot 10^n - 1 = 9 \cdot 10^n + (10^n - 1)$$

Thus $9|(10^{n+1} - 1)$.

Therefore, by mathematical induction, $9|(10^n - 1)$ for any $n \in \mathbb{N}$. □

Exercise 2.15. Solve the congruence $2x \equiv 5$ modulo 9 and modulo 6.

Done Later add.

Exercise 2.16. Determine the integers n for which the pair of congruences $2x - y \equiv 1$ and $4x + 3y \equiv 2$ modulo n has a solution.

Done Later add.

Exercise 2.17. Prove the Chinese Remainder Theorem: Let a, b, u, v be integers, and assume that the greatest common divisor of a and b is 1. Then there is an integer x such that $x \equiv u$ modulo a and $X \equiv v$ modulo b .

Hint: Do the case $u = 0$ and $v = 1$ first.

Exercise 2.18. Determine the order of each of the matrices A and B when the matrix entries are interpreted modulo 3.

2.13.5 Product Group

Exercise 2.19. Let x be an element of order r of a group G , and let y be an element of G' of order s . What is the order of (x, y) in the product group $G \times G'$?

Solution: The order of (x, y) in the product group $G \times G'$ is $\text{lcm}(r, s)$. Let $n \in \mathbb{Z}^+$ such that

$$(x, y)^n = (x^n, y^n) = (1_G, 1_{G'}).$$

This implies, $x^n = 1_G$ and $y^n = 1_{G'}$. Since order of x and y are r and s respectively,

$$r|n \quad \text{and} \quad s|n$$

So, we know that the least positive integer such that above property holds is $\text{lcm}(r, s)$. Hence, The order of (x, y) in the product group $G \times G'$ is $\text{lcm}(r, s)$.

Exercise 2.20. What does Proposition 2.8 tell us when, with the usual notation for the symmetric group S_3 , K and H are the subgroups $\langle y \rangle$ and $\langle x \rangle$?

Recall: $y^2 = 1$ and $x^3 = 1$. Then

$$H = \langle x \rangle = \{1, x, x^2\} \quad \text{and} \quad K = \langle y \rangle = \{1, y\}$$

Let the multiplication map, $f : H \times K \rightarrow S_3$ defined by $f(h, k) = hk$. Then, $\text{Im}(f) = HK = \{hk : h \in \langle x \rangle, k \in \langle y \rangle\} = \{1, x, x^2, y, xy, x^2y\}$.

Claim 1: f is injective.

Observe that $H \cap K = \{1\}$. by 2.8 a) f is injective.

Claim 2: f is surjective.

Observe that $S_3 = HK = \{1, x, x^2, y, xy, x^2y\}$. Thus, f is surjective.

Claim 3 : f is not homomorphism.

$$f((x, y) \cdot (x, 1)) = f(x^2, y^2) = f(x^2, 1) = x^2 \cdot 1 = x^2 \quad (2.74)$$

$$f(x, y) \cdot f(x, 1) = (x \cdot y) \cdot (x \cdot 1) = (xy) \cdot x = y \quad (2.75)$$

Thus, f is not a homomorphism.

Exercise 2.21. Prove that the product of two infinite cyclic groups is not infinite cyclic.

Proof.

- **Claim 1 :** Infinite cyclic groups are isomorphic to \mathbb{Z} .

Let G be a infinite cyclic group and $\langle a \rangle = G$. We can define a function $f : \mathbb{Z} \rightarrow G$ by $f(n) = a^n$ for all integers n . We need to show that f is an isomorphism, which means it is bijective.

- **Subclaim 1.1 :** f is a homomorphism.

$$f(n+m) = a^{n+m} = a^n \cdot a^m = f(n) \cdot f(m)$$

Thus, f is a homomorphism.

- **Subclaim 1.2 :** f is injective

Suppose that $f(n) = f(m)$

$$f(n) = f(m) \tag{2.76}$$

$$a^n = a^m \tag{2.77}$$

$$n = m \text{ (Since order is infinite)} \tag{2.78}$$

- **Subclaim 1.3 :** f is surjective.

Let $x \in G = \langle a \rangle$. Then, $x = a^k$ for some $k \in \mathbb{Z}$. Then, observe that

$$f(k) = a^k = a$$

. Thus, f is surjective.

Therefore, f is an isomorphism. Thus, the infinite cyclic groups are isomorphic to \mathbb{Z} .

We can consider $\mathbb{Z} \times \mathbb{Z}$. Suppose that (a, b) generate the product group. But then, we see that $(2a, b)$ cannot be obtained from adding (a, b) to itself, which implies that $\mathbb{Z} \times \mathbb{Z}$ is not infinite cyclic.

□

Exercise 2.22. In each of the following cases, determine whether or not G is isomorphic to the product group $H \times K$.

- (a) $G = \mathbb{R}^\times$, $H = \{\pm 1\}$, $K = \{\text{positive real numbers}\}$.
- (b) $G = \{\text{invertible upper triangular } 2 \times 2 \text{ matrices}\}$, $H = \{\text{invertible diagonal matrices}\}$, $K = \{\text{upper triangular matrices with diagonal entries 1}\}$.
- (c) $G = e^x$, $H = \{\text{unit circle}\}$, $K = \{\text{positive real numbers}\}$.

Solution:

- (a) Observe that $G = \mathbb{R}^\times = \mathbb{R} \setminus \{0\}$ is abelian. Then $H, K \trianglelefteq G$. Further, observe that $H \cap K = \{1\}$ and $HK = \mathbb{R}^\times = G$. Thus, by proposition 2.8 $G \cong H \times K$.

- (b)

- **Claim b.1:** $K \trianglelefteq G$.

Let $g = \begin{bmatrix} a & b \\ c & \end{bmatrix}$ be an arbitrary matrix in G and $k = \begin{bmatrix} 1 & d \\ & 1 \end{bmatrix} \in K$. Then,

$$g^{-1} = \begin{bmatrix} a^{-1} & -ba^{-1}c^{-1} \\ & c^{-1} \end{bmatrix}$$

Then

$$\begin{aligned} gkg^{-1} &= \begin{bmatrix} a & b \\ c & \end{bmatrix} \begin{bmatrix} 1 & d \\ & 1 \end{bmatrix} \begin{bmatrix} a^{-1} & -ba^{-1}c^{-1} \\ & c^{-1} \end{bmatrix} \\ &= \begin{bmatrix} a & b \\ c & \end{bmatrix} \begin{bmatrix} a^{-1} & dc^{-1} - ba^{-1}c^{-1} \\ & c^{-1} \end{bmatrix} = \begin{bmatrix} 1 & adc^{-1} \\ & 1 \end{bmatrix} \in K \end{aligned}$$

Thus, $H \trianglelefteq G$.

- **Claim b.2 :** $H = Z(G)$ (H is in the center of G).

– *subclaim b.2.1* : $H \subseteq Z(G)$.

Let $h = \begin{bmatrix} x & 0 \\ & x \end{bmatrix} \in H$. Then,

$$B = \begin{bmatrix} x & 0 \\ & x \end{bmatrix} = x \begin{bmatrix} 1 & \\ & 1 \end{bmatrix} = BI$$

Then observe that, for any $A \in G$,

$$BA = (xI)A = xA = Ax = A(xI) = AB$$

– *subclaim b.2.2* : $H \supseteq Z(G)$.

Let $\begin{bmatrix} p & q \\ r & s \end{bmatrix} \in Z(G)$, then,

$$\begin{bmatrix} p & q \\ r & s \end{bmatrix} \begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix} = \begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix} \begin{bmatrix} p & q \\ r & s \end{bmatrix} \quad (2.79)$$

$$\begin{bmatrix} p & -q \\ r & -s \end{bmatrix} = \begin{bmatrix} p & q \\ -r & -s \end{bmatrix} \quad (2.80)$$

(2.81)

This yields $-q = q$ and $r = -r$ which imply $q = r = 0$. Now considering,

$$\begin{bmatrix} p & 0 \\ 0 & s \end{bmatrix} \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix} = \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix} \begin{bmatrix} p & 0 \\ 0 & s \end{bmatrix} \quad (2.82)$$

$$\begin{bmatrix} 0 & p \\ q & 0 \end{bmatrix} = \begin{bmatrix} 0 & q \\ p & 0 \end{bmatrix} \quad (2.83)$$

(2.84)

This implies that $a = d$ and since the matrix is invertible, $a \neq 0$. Thus, $H \supseteq Z(G)$.

Therefore, $H = Z(G)$. In other words, H is in the center of G .

- **Claim b.2 :** $H \trianglelefteq G$.

Since, H is in the center of G , then H is normal.

- **Claim b.3 :** $H \cap K = \{I\}$.

This is trivial.


- **Claim b.4 :** $HK = G$ Clearly, $HK \subseteq G$. Conversely, for any $g \in G$, $a \neq 0, b \neq 0$,

$$g = \begin{bmatrix} a & b \\ c & d \end{bmatrix} = \begin{bmatrix} a & 0 \\ 0 & d \end{bmatrix} \begin{bmatrix} 1 & a^{-1}b \\ 0 & 1 \end{bmatrix} \in HK$$

. Thus, $HK = G$.

Therefore, by proposition 2.8 $G \cong HK$

- (c) Since C^\times is abelian, then H and K are normal. And, $H \cap K = \{1\}$.



Consider $a + bi \in C^\times$. Then

$$a + bi = \left(\frac{a}{\sqrt{a^2 + b^2}} + \frac{b}{\sqrt{a^2 + b^2}} i \right) \sqrt{a^2 + b^2} \in HK$$

So, $HK = G$. So, by proposition 2.8 $G \cong H \times K$.

Exercise 2.23. Let G_1 and G_2 be groups, and let Z_i be the center of G_i . Prove that the center of the product group $G_1 \times G_2$ is $Z_1 \times Z_2$.

Let $Z_1 = Z(G_1) = \{x \in G_1 \mid xg = gx \ \forall g \in G_1\}$
 and $Z_2 = Z(G_2) = \{x \in G_2 \mid xg = gx \ \forall g \in G_2\}$

Need to show: $Z(G_1 \times G_2) = Z(G_1) \times Z(G_2)$

claim 1: $Z(G_1) \times Z(G_2) \subseteq Z(G_1 \times G_2)$

Let $(g_1, g_2) \in G_1 \times G_2$ and $(z_1, z_2) \in Z(G_1) \times Z(G_2)$


$(z_1, z_2) \in Z(G_1) \times Z(G_2)$

$$(g_1, g_2) \cdot (z_1, z_2) = (g_1 z_1, g_2 z_2) \neq \cancel{(z_1, z_2)}$$

$$= (z_1 g_1, z_2 g_2) \quad (\text{Since } z_1 \in Z(G_1) \text{ and } z_2 \in Z(G_2))$$


$$= (z_1, z_2)(g_1, g_2).$$

Thus, $Z(G_1) \times Z(G_2) \subseteq Z(G_1 \times G_2)$



Exercise 2.24. Let G be a group that contains normal subgroups of orders 3 and 5, respectively. Prove that G contains an element of order 15.

Exercise 2.25. Let H be a subgroup of a group G , let $\phi : G \rightarrow H$ be a homomorphism whose restriction to H is the identity map, and let N be its kernel. What can one say about the product map $H \times N \rightarrow G$?



Claim: f is injective.

Clearly $I_G \in H \cap N$.

Assume that $I_G \neq x \in H \cap N$

$$x \in H \Rightarrow f(x) = x \neq I_G \quad \text{--- (1)}$$

$$x \in N \Rightarrow f(x) = I_G \quad \text{--- (2)}$$

(1) and (2) give contradiction,

Thus, $H \cap N = \{I\}$

By proposition 2.11 (my Artins book) 2.11.4

(a), f is injective.

Claim: f is surjective

This is trivial, $\forall h \in H, h = f(h)$


claim: If G is abelian then f is homomorphism

Suppose G is abelian then $hn = nh$

$\forall h \in H \trianglelefteq G$ and $\forall n \in N \trianglelefteq G$

Thus, # by artins book proposition 2.11.4
part b) f is a homomorphism

Exercise 2.26. Let G , G' , and H be groups. Establish a bijective correspondence between homomorphisms $\phi : H \rightarrow G \times G'$ from H to the product group and pairs (φ, φ') consisting of a homomorphism $\varphi : H \rightarrow G$ and a homomorphism $\varphi' : H \rightarrow G'$.



We have to define a map.

$$f: \left\{ \begin{array}{l} \text{homomorphisms} \\ \text{from } H \text{ to } G \times G' \end{array} \right\} \xrightarrow{\quad} \left\{ \begin{array}{l} \text{pairs of} \\ \text{homomorphisms} \\ (\psi, \psi') \end{array} \right\}$$

$$\underline{\Phi} \longmapsto (\pi \circ \underline{\Phi}, \pi' \circ \underline{\Phi})$$

where π, π' are projection maps,

$$\pi: G \times G' \longrightarrow G$$

$$(g, g') \longmapsto g$$

$$\text{and } \pi': G \times G' \longrightarrow G'$$

$$(g, g') \longmapsto g'$$

Now I am going to prove

1. f is well-defined
2. f is injective
3. f is surjective.

~~First let's check c1~~

First of all Let's denote

$$A := \left\{ \underline{\Phi}: H \rightarrow G \times G' \mid \underline{\Phi} \text{ is homomorphisms} \right\}$$

Let $B := \{(\varphi, \psi) \mid \varphi: H \rightarrow G, \psi: H \rightarrow G' \text{ and } \varphi \text{ and } \psi \text{ are homomorphisms}\}$

Now let's start to check three conditions.

1 $\boxed{f(\emptyset) \in B \quad \forall \emptyset: H \rightarrow G \times G' \text{ be homomorph}}$

Let $\emptyset: H \rightarrow G \times G'$ be a homomorphism

subclaim 1.1: π is homomorphism

Let $(g_1, g'_1), (g_2, g'_2) \in G \times G'$

$$\begin{aligned} \pi((g_1, g'_1)(g_2, g'_2)) &= \pi(g_1 g_2, g'_1 g'_2) \\ &= (g_1 g_2) = \pi(g_1, g'_1) \pi(g_2, g'_2) \end{aligned}$$

Thus π is homomorphism
Now we are done the subclaim

claim: $\pi \circ \emptyset$ is homomorphism

Let $h, h' \in H$

$$\begin{aligned} [\pi \circ \emptyset](hh') &= \pi(\emptyset(chh')) = \pi(\emptyset(ch)\emptyset(ch')) \\ &= \pi(\emptyset(ch)) \pi(\emptyset(ch')) \\ &= \pi \circ \emptyset(ch) \cdot \pi \circ \emptyset(ch') \end{aligned}$$

Thus $(\pi \circ \emptyset)$ is a homomorphism

Therefore f is well-defined ■

Claim: f is injective

Let $\phi_1, \phi_2 \in A$ such that

$$\cancel{f(\phi_1)} = f(\phi_2)$$

$$(\pi \circ \phi_1, \pi' \circ \phi_1) = (\pi \circ \phi_2, \pi' \circ \phi_2)$$

Thus, for all $h \in H$,

$$\pi \circ \phi_1(h) = \pi \circ \phi_2(h) \text{ and} \quad ①$$

$$\pi' \circ \phi_1(h) = \pi' \circ \phi_2(h) \quad ②$$

Let p be an arbitrary element in H ,

Then,

$$\cancel{\phi_1(p)} = \phi_1(p) = (g_1, g'_1) \text{ for some } g_1, g'_1 \in G'$$

$$\phi_2(p) = (g_2, g'_2) \text{ for some } g_2 \in G, g'_2 \in G'$$

By equation ①

$$\pi \circ \phi_1(p) = \pi \circ \phi_2(p)$$

$$\pi(g_1, g'_1) = \pi(g_2, g'_2)$$

$$\text{thus, } g_1 = g_2 \quad \cancel{g'_1 = g'_2} \quad ③$$

By equation ②

$$\pi_0^* \phi_1(p) = \pi_0^* \phi_2(p)$$

$$\pi^*(g_1, g_1') = \pi^*(g_2, g_2')$$

$$g_1' = g_2' \quad \text{--- ④}$$

By ③ and ④,

$$\phi_1(p) = \phi_2(p)$$

Since p is arbitrary, $\phi_1 = \phi_2$

Therefore f is injective.

③ f is surjective.

Let $(\tilde{\psi}, \tilde{\psi}') \in B$

Define $\tilde{\phi}: H \rightarrow G \times G'$
 $q \mapsto (\tilde{\psi}(q), \tilde{\psi}'(q))$

sub claim 3.1: $\tilde{\phi}$ is homomorphism

Let $\tilde{h}_1, \tilde{h}_2 \in H$.

$$\begin{aligned}\tilde{\phi}(h_1, h_2) &= (\tilde{\psi}(\tilde{h}_1, \tilde{h}_2), \tilde{\psi}'(\tilde{h}_1, \tilde{h}_2)) \\ &= (\tilde{\psi}(h_1)\tilde{\psi}(h_2), \tilde{\psi}'(h_1)\tilde{\psi}'(h_2)) \\ &= (\tilde{\psi}(h_1), \tilde{\psi}'(h_1))(\tilde{\psi}(h_2), \tilde{\psi}'(h_2)) \\ &= \underbrace{\tilde{\phi}(h_1)}_{\tilde{\phi}} \cdot \underbrace{\tilde{\phi}(h_2)}_{\tilde{\phi}}\end{aligned}$$

Thus $\tilde{\phi}$ is homomorphism from H to $G \times G'$

$$\tilde{\phi} \in A$$

~~$\tilde{\phi} = R \circ \tilde{\phi}, \tilde{\phi}' = R' \circ \tilde{\phi}$~~

Now we need to check

$$\tilde{\psi} = R \circ \tilde{\phi} \text{ and } \tilde{\psi}' = R' \circ \tilde{\phi}$$

~~$\tilde{\phi}(h)$~~ Let $\forall h \in H$,
 $(\pi \circ \tilde{\phi})(h) = \pi(\tilde{\phi}(h)) = \pi(\tilde{\phi}(ch), \tilde{\phi}'(ch)) = \tilde{\phi}(h)$

thus, $\pi \circ \tilde{\phi} = \tilde{\phi}$

Similarly, $\pi' \circ \tilde{\phi} = \tilde{\phi}'$

$f(\tilde{\phi}) = (\pi \circ \tilde{\phi}, \pi' \circ \tilde{\phi}) = (\phi, \phi')$

Thus f is surjective.

Exercise 2.27. Let H and K be subgroups of a group G . Prove that the product set HK is a subgroup of G if and only if $HK = KH$.

Solution :

Proof. Let G be group and $H, K \leq G$.

- (\implies): **Claim** : if $HK \leq G$ then $HK = KH$.

Suppose $HK \leq G$.

– **sub claim:** Let $x \in KH$. Then $x = kh$ for some $h \in H$ and $k \in K$,

$$x = kh = 1kh1 = (1k)(h1) \in HK$$

(Since $1 \in H$ and $1 \in K$). Thus, $KH \subseteq HK$.

– **sub claim:** $KH \supseteq HK$.

We can not use previous method. Because we still do not know KH is sub group or not.

Let $y \in HK$. Since $HK \leq G$, $y^{-1} \in HK$. Then $y^{-1} = hk$ for some $h_0 \in H$ and $k_0 \in K$.

$$y = (y^{-1})^{-1} = (h_0 k_0)^{-1} = k_0 h_0 \in KH$$

Thus, $KH \supseteq HK$.

Therefore, $KH = HK$.

- (\Leftarrow): **Claim** : If $HK = KH$ then $HK \leq G$.

Now suppose that $HK = KH$. Suppose that $HK = KH$ is empty. Then, H, K is empty. Now we use sub group test.

- *Closure:* Let $x, y \in HK$. Then $x = h_1k_1$ and $y = h_2k_2$ for some $h_1, h_2 \in H$ and $k_1, k_2 \in K$.

$$xy = (h_1k_1)(h_2k_2) \quad (2.85)$$

$$= h_1(k_1h_2)k_2 \quad (\text{By associativity property}) \quad (2.86)$$

$$= h_1(h_3k_3)k_2 \quad (\text{for some } h_3 \in H \text{ and } k_3 \in K \text{ (Since } KH = HK)) \quad (2.87)$$

$$= (h_1h_3)(k_3k_2) \in HK \quad (2.88)$$

- *Inverse:* Let $x \in HK$. Then $x = h'k'$ for some $h' \in H$ and $k' \in K$.

$$x^{-1} = (h'k')^{-1} = k'h' \in KH = HK$$

Thus, $HK \leq G$.

Therefore, the product set HK is a subgroup of G if and only if $HK = KH$ \square

2.13.6 Quotient Groups

Exercise 2.28. Show that if a subgroup H of a group G is not normal, there are left cosets aH and bH whose product is not a coset.

Proof. Suppose that $H \leq G$ but $H \not\trianglelefteq G$. We are going to use indirect proof. So, assume that $(aH)(bH)$ is a coset for all $a, b \in G$. Then,

$$aHbH = abH$$

then for all $h_1, h_2, h_3 \in H$, So,

$$ah_1bh_2 = abh_3$$

$$ah_1bh_2 = abh_3 \quad (2.89)$$

$$a^{-1}(ah_1bh_2) = a^{-1}(abh_3) \quad (2.90)$$

$$(a^{-1}a)(h_1bh_2) = (a^{-1}a)(bh_3) \quad (2.91)$$

$$h_1bh_2 = bh_3 \quad (2.92)$$

$$b^{-1}(h_1bh_2)h_2^{-1} = b^{-1}(bh_3)h_2^{-1} \quad (2.93)$$

$$b^{-1}h_1b(h_2h_2^{-1}) = (b^{-1}b)h_3h_2^{-1} \quad (2.94)$$

$$b^{-1}h_1b = h_3h_2^{-1} \in H \quad (2.95)$$

$$(2.96)$$

Thus, $b^{-1}h_1b \in H$. and since $b \in G$ and $h_1 \in H$ is arbitrary, H is normal subgroup, which contradict the our hypothesis. Thus, there exist left cosets aH and bH whose product is not a coset. \square

Exercise 2.29. In the general linear group $GL_3(\mathbb{R})$, consider the subsets:

$$H = \left\{ \begin{pmatrix} 1 & * & * \\ 0 & 1 & * \\ 0 & 0 & 1 \end{pmatrix} \mid * \text{ represents an arbitrary real number} \right\}$$

and

$$K = \left\{ \begin{pmatrix} 1 & 0 & * \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix} \mid * \text{ represents an arbitrary real number} \right\}$$

Show that H is a subgroup of GL_3 , that K is a normal subgroup of H , and identify the quotient group H/K . Determine the center of H .

- **Claim 1:** $H \leq G$.

Certainly! Let's prove that the subset H is a subgroup of the general linear group $GL_3(\mathbb{R})$.

- *Non-emptiness:* We start by showing that H is non-empty. The identity matrix \mathbf{I}_3 belongs to H since it satisfies the conditions for H :

$$\mathbf{I}_3 = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}$$

Therefore, H is non-empty.

- *Closure under matrix multiplication:* Let \mathbf{A} and \mathbf{B} be matrices in H . We need to show that their product \mathbf{AB} is also in H . Consider:

$$\mathbf{A} = \begin{pmatrix} 1 & a & b \\ 0 & 1 & c \\ 0 & 0 & 1 \end{pmatrix} \quad \text{and} \quad \mathbf{B} = \begin{pmatrix} 1 & x & y \\ 0 & 1 & z \\ 0 & 0 & 1 \end{pmatrix}$$

The product \mathbf{AB} is:

$$\mathbf{AB} = \begin{pmatrix} 1 & a+x & by+az+b \\ 0 & 1 & c+z \\ 0 & 0 & 1 \end{pmatrix}$$

“““ Since $a+x$, $y+az+b$, and $c+z$ are arbitrary real numbers, \mathbf{AB} satisfies the conditions for H . Hence, H is closed under matrix multiplication.

- *Closure under taking inverses:* Let \mathbf{A} be a matrix in H . We need to show that its inverse \mathbf{A}^{-1} is also in H . The inverse of \mathbf{A} is:

$$\mathbf{A}^{-1} = \begin{pmatrix} 1 & -a & ac - b \\ 0 & 1 & -c \\ 0 & 0 & 1 \end{pmatrix}$$

Again, since $-a$, $ac - b$, and $-c$ are arbitrary real numbers, \mathbf{A}^{-1} satisfies the conditions for H . Therefore, H is closed under taking inverses. Hence, we have shown that H is a subgroup of $GL_3(\mathbb{R})$.

- **Claim 2 :** $K \leq H$.

I am not going to prove that this case, because this is exhausting. This is very easy.

- **Claim 3 :** K is normal sub group of H .

Let $A := \begin{bmatrix} 1 & a & b \\ 0 & 1 & c \\ 0 & 0 & 1 \end{bmatrix} \in H$ and $B := \begin{bmatrix} 1 & 0 & d \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} \in K$, where $a, b, c, d \in \mathbb{R}$.

Then,

$$ABA^{-1} = \begin{bmatrix} 1 & a & b \\ 0 & 1 & c \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} 1 & 0 & d \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} 1 & -a & ac - b \\ 0 & 1 & -c \\ 0 & 0 & 1 \end{bmatrix} \quad (2.97)$$

$$= \begin{bmatrix} 1 & a & d + b \\ 0 & 1 & c \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} 1 & -a & ac - b \\ 0 & 1 & -c \\ 0 & 0 & 1 \end{bmatrix} \quad (2.98)$$

$$= \begin{bmatrix} 1 & 0 & d \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} \in K \quad (2.99)$$

Thus, K is normal sub group of H .

- **Quotient group H/K :**

Let $h_1, h_2 \in H$ be such that

$$h_1 = \begin{bmatrix} 1 & a_1 & b_1 \\ 0 & 1 & c_1 \\ 0 & 0 & 1 \end{bmatrix} \text{ and } h_2 = \begin{bmatrix} 1 & a_2 & b_2 \\ 0 & 1 & c_2 \\ 0 & 0 & 1 \end{bmatrix}.$$

Suppose that $h_1 K = h_2 K$. By Artin's book 2.85, $h_1^{-1} h_2 \in K$.

$$h_1^{-1} h_2 = \begin{bmatrix} 1 & -a_1 & a_1 c_1 - b_1 \\ 0 & 1 & -c_1 \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} 1 & a_2 & b_2 \\ 0 & 1 & c_2 \\ 0 & 0 & 1 \end{bmatrix} = \begin{bmatrix} 1 & -a_1 + a_2 & b_2 - a_1 c_2 + a_1 c_1 - b_1 \\ 0 & 1 & -c_1 + c_2 \\ 0 & 0 & 1 \end{bmatrix}$$

Since,

$$h_1 K = h_2 K \iff h_1^{-1} h_2 \in K \quad (2.100)$$

$$\iff -a_1 + a_2 = 0 \text{ and } -c_2 - c_1 \quad (2.101)$$

$$\iff a_1 = a_2 \text{ and } c_1 = c_2 \quad (2.102)$$

Thus,

$$H/K = \left\{ hK \mid \begin{bmatrix} 1 & a & 0 \\ 0 & 1 & c \\ 0 & 0 & 1 \end{bmatrix}, a, c \in \mathbb{R} \right\}$$

- **Center of Group H**

$$Z(H) = \left\{ A \in H \mid AX = XA \text{ for all } X \in H \right\}$$

Let's find center. We need to find $a, b, c \in \mathbb{R}$ such that

$$\begin{aligned} \begin{bmatrix} 1 & a & b \\ 0 & 1 & c \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} 1 & x & y \\ 0 & 1 & z \\ 0 & 0 & 1 \end{bmatrix} &= \begin{bmatrix} 1 & x & y \\ 0 & 1 & z \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} 1 & a & b \\ 0 & 1 & c \\ 0 & 0 & 1 \end{bmatrix} \text{ for all } x, y, z \in \mathbb{R} \\ \begin{bmatrix} 1 & a+x & b+y+az \\ 0 & 1 & c+z \\ 0 & 0 & 1 \end{bmatrix} &= \begin{bmatrix} 1 & a+x & b+y+xc \\ 0 & 1 & c+z \\ 0 & 0 & 1 \end{bmatrix} \end{aligned} \quad (2.104)$$

By comparing 1×3 index, we need to find that $a, b, c \in \mathbb{R}$ $az = xc$ for all $x, y, z \in \mathbb{R}$. SO, the only possibility is $a = c = 0$. Thus,

$$Z(H) = \left\{ \begin{bmatrix} 1 & 0 & b \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} \mid b \in \mathbb{R} \right\}$$

Exercise 2.30. Let P be a partition of a group G with the property that for any pair of elements A, B of the partition, the product set AB is contained entirely within another element C of the partition. Let N be the element of P that contains 1. Prove that N is a normal subgroup of G and that P is the set of its cosets.

Proof.

- **Claim 1:** The product set $NN = N$.

Let $x \in N$, Then $x = x \cdot 1 \in NN$. Thus, $N \subseteq NN$. We know that the product set NN is contained entirely within an element of partition, but N is a partition. The only way this happen $N = NN$.

- **Claim 2:** $N \leq G$.

- *Subset*: It is very clear that $N \subseteq G$.
- *Closure*: Let $x, y \in N$. Then $(xy) \in NN = N$. So, N is closed under composition.
- *Identity*: Given that $1 \in N$.
- *Inverse*: Let $x \in N$. We have to show that x^{-1} is contained in N . We use proof by contradiction. So, Assume contrary, $x^{-1} \in M$, M is an element of partition P such that $N \neq M$.

$$xx^{-1} \in NM, xx^{-1} = 1 \in N$$

So, we would have $NM \subseteq N$ (since the product set is contained entirely within one of the sets of partition). On the other hand,

$$1x^{-1} \in NM, 1x^{-1} = x^{-1} \in N$$

So, $NM \subseteq N$ a contradiction.

Therefore, conclude that our assumption that $x^{-1} \notin N$ was wrong, that we must have $x^{-1} \in N$

- **Claim 3:** $N \trianglelefteq$

Let $a \in G$ and $n \in N$. Let $a \in A$ and $a^{-1} \in B$, where $A, B \in P$.

$$ana^{-1} = ANB$$

, But also, $1 \in N$,

$$g \cdot 1 \cdot g^{-1} = 1 \in ANB$$

Since P splits into disjoint subsets and the only one element of partition that contains 1 is N .

$$ANB = N$$

and but also $gng^{-1} \in N$. Therefore, N is normal subgroup.

-Claim 4: P is the set of its cosets.

Let A be some element of partition, and let $a \in A$.

Need to prove: $A = aN$

- Subclaim 4.1: $A \subseteq aN$.

Let $b \in A$. Let B be some element of partition. Then $b^{-1}b \in BA$, $b^{-1}b = 1 \in N$. So, $BA \subseteq N$. Specially,

$$b^{-1}a \in N$$

So there exists some $n \in N$ such that

$$b^{-1}a = n \iff a = bn \iff b = an^{-1}$$

Since $N \leq G$. $n^{-1} \in N$. Thus, $b \in aN$, so $A \subseteq aN$.

- Subclaim 4.2 : $A \supseteq aN$.

– subclaim 4.2.1: $AN \subseteq A$,

Note that $a \in A$ and $1 \in N$. Thus,

$$a \cdot 1 \in AN, \quad a \cdot 1 = a \in A$$

Since, the product set is contained entirely in one element partition. Thus, $AN \subseteq A$

Now,

$$aN = \{an | n \in N\} \subseteq \{an : a \in A, n \in N\} = AN \subseteq A$$

Thus, $ax \in A$, as required. Thus $aN \subseteq A$

Hence, we can conclude that $A = aN$.

□

Exercise 2.31. Let $H = \{\pm 1, \pm i\}$ be the subgroup of $G = C^\times$ of fourth roots of unity. Describe the cosets of H in G explicitly. Is G/H isomorphic to G ?

Cosets are in the following form

$$ZH = \{zH, z^{-1}H\}, \text{ where } z \in \mathbb{C}^{\times}$$

We are going to use 1st isomorphism, we need to find a surjective homomorphism φ such that $\ker(\varphi) = H$.

Let $\varphi: G \rightarrow G$ defined by

$$\begin{array}{ccc} & G & \longrightarrow G \\ z & \longmapsto & z^4 \end{array}$$

Claim 1: φ is homomorphism

Let $a, b \in G$

$$\varphi(ab) = (ab)^4 = a^4 b^4 = \varphi(a) \varphi(b)$$

Thus, φ is homomorphism

Claim 2: φ is surjective.

Let $c \in G$. Then let $d = \sqrt[4]{c}$. We know that d exists. Because \mathbb{C}^{\times} have all n th roots every element of \mathbb{C}^{\times} (algebraically closed). Then,

$$\varphi(d) = c$$

Thus, φ is surjective.

claim 3: $\ker(\psi) = H$

$$\begin{aligned}
 \ker(\psi) &= \{x \mid x^4 = 1\} \\
 &= \left\{ e^{\frac{2\pi i k}{4}} \mid k=0,1,2,3 \right\} \\
 &= \left\{ e^0, e^{\frac{2\pi i}{4}}, e^{\frac{4\pi i}{4}}, e^{\frac{6\pi i}{4}} \right\} \\
 &= \left\{ e^0, e^{\frac{\pi i}{2}}, e^{\pi i}, e^{\frac{3\pi i}{2}} \right\} \\
 &= \{ 1, i, -1, -i \} \\
 &= H
 \end{aligned}$$

claim 4: $\text{Im}(\psi) = G$

We know that $\text{Im}(\psi) \subseteq G$ $\xrightarrow{*}$

Let $x \in G - C^{\times}$. Then $x^{1/4}$ exists in $C^{\times} - G$

$$\psi(x^{1/4}) = x$$

Thus, $\text{Im}(\psi) \supseteq G$ $\xrightarrow{**}$

By $*$ and $**$ $\text{Im}(\psi) = G$

claim: $G/H \cong G$

By first theorem of isomorphism, we can conclude that, $G/H \cong G$

Exercise 2.32. Let G be the group of upper triangular real matrices $\begin{bmatrix} a & b \\ 0 & d \end{bmatrix}$ with a and d different from zero. For each of the following subsets, determine whether or not S is a subgroup, and whether or not S is a normal subgroup. If S is a normal subgroup, identify the quotient group G/S .

1. S is the subset defined by $b = 0$.
2. S is the subset defined by $d = 1$.
3. S is the subset defined by $a = d$.

$$\textcircled{i} \quad S_i := \left\{ \begin{bmatrix} a & 0 \\ 0 & d \end{bmatrix} : a, d \in \mathbb{R} \setminus \{0\} \right\}$$

Observe that $I = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \in S$. Thus, $S \neq \emptyset$

Let $A, B \in G$. Then

$$A := \begin{bmatrix} a_1 & 0 \\ 0 & a_2 \end{bmatrix} \quad \text{for some } a_1, a_2 \in \mathbb{R} \setminus \{0\}$$

$$B := \begin{bmatrix} b_1 & 0 \\ 0 & b_2 \end{bmatrix}$$

$$\text{Then } B^{-1} = \frac{1}{b_1 b_2} \begin{bmatrix} b_2 & 0 \\ 0 & b_1 \end{bmatrix} = \begin{bmatrix} 1/b_1 & 0 \\ 0 & 1/b_2 \end{bmatrix}$$

So, $B^{-1} \in G$. Thus,

$$\begin{aligned} AB^{-1} &= \begin{bmatrix} a_1 & 0 \\ 0 & a_2 \end{bmatrix} \begin{bmatrix} 1/b_1 & 0 \\ 0 & 1/b_2 \end{bmatrix} \\ &= \begin{bmatrix} a_1/b_1 & 0 \\ 0 & a_2/b_2 \end{bmatrix} \end{aligned}$$

Since $\frac{a_1}{b_1}, \frac{a_2}{b_2} \in \mathbb{R} \setminus \{0\}$, thus $AB^{-1} \in S$

Therefore, S is a subgroup.

Let's check S_3 is normal or not

Claim: S_3 is not normal

Let $X := \begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix} \in G$. Then

$$X^{-1} := \begin{bmatrix} 1 & -1 \\ 0 & 1 \end{bmatrix}$$

Let $Y := \begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix} \in B$. Then,

$$XYX^{-1} = \begin{bmatrix} 1 & -2 \\ 0 & -1 \end{bmatrix}. \text{ But } XYX^{-1} \notin B$$

Therefore, B is NOT a Normal subgroup

$$\textcircled{2} \quad S'' := \left\{ \begin{bmatrix} a & b \\ 0 & 1 \end{bmatrix} : a, b \in \mathbb{R} \right\}$$

claim1: $S'' \leq G$

Observe that $\begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \in S''$. Thus $S'' \neq \emptyset$.

Let $X'' = \begin{bmatrix} a & b \\ 0 & 1 \end{bmatrix}$ & $Y'' = \begin{bmatrix} e & f \\ 0 & 1 \end{bmatrix} \in S''$

$$\begin{aligned} \text{Then } (Y'')^{-1} &:= \frac{1}{e} \begin{bmatrix} 1 & -f \\ 0 & e \end{bmatrix} \\ &= \begin{bmatrix} 1/e & -f/e \\ 0 & 1 \end{bmatrix} \end{aligned}$$

$$\begin{aligned} (X'')(Y'')^{-1} &:= \begin{bmatrix} a & b \\ 0 & 1 \end{bmatrix} \begin{bmatrix} 1/e & -f/e \\ 0 & 1 \end{bmatrix} \\ &= \begin{bmatrix} ae & af+b \\ 0 & 1 \end{bmatrix} \in S'' \end{aligned}$$

Thus, $S'' \leq G$

Cosets

Let $gS, hS \in G/S$.

$$gS = hS \iff h^{-1}g \in S$$

$$\text{Let } g = \begin{bmatrix} g_1 & g_2 \\ 0 & g_3 \end{bmatrix}, h = \begin{bmatrix} h_1 & h_2 \\ 0 & h_3 \end{bmatrix}$$

where, $g_1, g_2, g_3, h_1, h_2, h_3 \in \mathbb{R}$ and $g_1, g_2, h_1, h_3 \neq 0$

$$\begin{aligned} \text{Then, } h^{-1}g &= \begin{bmatrix} 1/h_1 & -h_2/h_1h_3 \\ 0 & 1/h_3 \end{bmatrix} \begin{bmatrix} g_1 & g_2 \\ 0 & g_3 \end{bmatrix} \\ &= \begin{bmatrix} g_1/h_1 & g_2/h_1 - \frac{g_2h_2}{h_1h_3} \\ 0 & g_3/h_3 \end{bmatrix} \end{aligned}$$

$$\text{Thus } h^{-1}g \in S \iff \frac{g_3}{h_3} = 1 \iff g_3 = h_3$$

$$\text{Therefore, } gS \neq hS \iff g_3 \neq h_3$$

Note that g_1, g_2 does not help to determine whether two cosets are equal.

Thus, we can fix $a=1, b=0$,

$$G/S := \left\{ gS \mid g = \begin{bmatrix} 1 & 0 \\ 0 & c \end{bmatrix}, c \in \mathbb{R} \setminus \{0\} \right\}$$

$$\textcircled{3} \quad S''' := \left\{ \begin{bmatrix} a & b \\ 0 & a \end{bmatrix} : a, b \in \mathbb{R}, a \neq 0 \right\}$$

Observe that, $\begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \in S'''$. Thus, $S''' \neq \emptyset$

Let $A := \begin{bmatrix} a & b \\ 0 & a \end{bmatrix}$, $B := \begin{bmatrix} e & f \\ 0 & e \end{bmatrix} \in S'''$, where $a, b, e, f \in \mathbb{R} \setminus \{0\}$

$$\text{Then } B' = \frac{1}{e^2} \begin{bmatrix} e & f \\ 0 & e \end{bmatrix} = \begin{bmatrix} 1/e & f/e \\ 0 & 1/e \end{bmatrix}$$

$$AB^{-1} = \begin{bmatrix} a & b \\ 0 & a \end{bmatrix} \begin{bmatrix} 1/e & f/e \\ 0 & 1/e \end{bmatrix} = \begin{bmatrix} ae & af+be \\ 0 & ae \end{bmatrix} \in S'''$$

Thus, $S''' \leq G$

$$\text{Let } A := \begin{bmatrix} a & b \\ 0 & c \end{bmatrix} \in G. \text{ So, } A^{-1} := \begin{bmatrix} 1/a & -b \\ 0 & 1/c \end{bmatrix}$$

Let $B := \begin{bmatrix} e & f \\ 0 & e \end{bmatrix} \in S'''$. Then

$$ABA^{-1} = \begin{bmatrix} e & \cancel{(eb+af+be)} \\ 0 & e \end{bmatrix} \in S'''$$

Thus, S''' is normal.

So, we can fix that $g_2=0$

$$G/S := \left\{ gS \mid g = \begin{bmatrix} g_1 & 0 \\ 0 & g_3 \end{bmatrix}, g_1, g_3 \in \mathbb{R} \right\}$$

2.13.7 Miscellaneous Problems

Exercise 2.33 (M1). Describe the column vectors $(a, c)^T$ that occur as the first column of an integer matrix A whose inverse is also an integer matrix.

$$\text{Let } A = \begin{bmatrix} a & b \\ c & d \end{bmatrix} \in GL_2(\mathbb{Z})$$

$$A^{-1} \text{ exists} \iff \det(A) = ad - bc \neq 0$$

$$A^{-1} = \frac{1}{(ad - bc)} \begin{bmatrix} d & -b \\ -c & a \end{bmatrix}$$

Lemma: Let A be matrix with integer entries.
 A^{-1} has integer entries $\iff \det(A) = \pm 1$

We know that

$1 = \det(I) = \det(A^{-1}A) = \det(A^{-1})\det(A)$. Thus,
 A and A^{-1} have integer entries $\iff \det(A) = \pm 1$

Claim: a and c are relative prime

Assume that $g = \gcd(a, c) > 1$. Then

$$a = gk_1 \text{ and } c = gk_2 \text{ for some } k_1, k_2 \in \mathbb{Z}$$

$$\begin{aligned} \text{Thus, } \det(A) &= ad - bc = gk_1d - gk_2c \\ &= g(k_1d - k_2c) \end{aligned}$$

Thus $g \mid \det(A)$

But $\det(A) = \pm 1$. So, it contradicts $g > 1$.

Hence, $\gcd(a, c) = 1$

Therefore a and c must relative prime

Exercise 2.34 (M.2).

- (a) Prove that every group of even order contains an element of order 2.
- (b) Prove that every group of order 21 contains an element of order 3.

Claim a : Let G be a group whose identity is e . Let G be of even order. Then, $\exists x \in G : |x| = 2$

Proof. (Claim a) In any group G , the identity element e is self-inverse with the property that the identity is the only group element of order 1, and is the only such element.

That leaves an odd number of elements.

Each element in $x \in G : |x| > 2$ can be paired off with its inverse, as $|x^{-1}| = |x| > 2$ (Since order of a group element equals the order of its inverse.)

Hence there must be at least one element which has not been paired off with any of the others which is therefore self-inverse. Let's say it, 'y'.

$$\begin{array}{l} y^{-1} = y \& x y = e \& |y| = 2 \end{array}$$

Thus, every group of even order contains an element of order 2. \square

Claim b: Let G be a group whose identity is e . Let G be of order 21. ($|G|=21$) Then, $\exists x \in G : |x| = 3$

Chapter 3

Lienaeer Operators

Chapter 4

Vector spaces.