

Maple

Ashan J

2024-07-06

Contents

1	Matrices	5
1.1	Defining matrices	5
1.2	Row Opreations	11
1.3	Determinanat of Matrix	14
1.4	Exercise	16
2	System of linear equations and matrices	19
2.1	Solving Linear Systems	20
2.2	Dependent Systems:	23
2.3	Inconsistent System	24
2.4	Exercise	27
3	Intergration	29
3.1	Indefinite Integration	29
3.2	Definite Integration	31
3.3	Integration by Parts	32
3.4	Substitutions	34
3.5	Multiple Integration	35
3.6	Exercise	37
3.7	Computing the area from the integral	38
3.8	Exercise	41

Chapter 1

Matrices

There are several ways to define a matrix in Maple. Use the help command to find the general definition of a matrix. To work with Matrices first load the package `linalg` Which include most linear algebra related commands.

1.1 Defining matrices

There are several ways to define Matrix. Let's look at them through examples. You can use maple help.

```
> ? matrix
```

```
> restart;
> with(linalg):
> M1 :=matrix (3, 3, [1,-2,-3,2,-2,2,3,-3,3]);
```

```
> restart;
> with(linalg):
> M1 :=matrix (3, 3, [1,-2,-3,2,-2,2,3,-3,3]);
```

$$M1 := \begin{bmatrix} 1 & -2 & -3 \\ 2 & -2 & 2 \\ 3 & -3 & 3 \end{bmatrix}$$

```
>
```

```
> M2:= matrix(3,4,[1,1/2,-1/3,1/4,2,1/2,-3/4,5/4,3,3/5,3/7,3/8]);
```

```
> M2:= matrix(3,4,[1,1/2,-1/3,1/4,2,1/2,-3/4,5/4,3,3/5,3/7,3/8]);
```

$$M2 := \begin{bmatrix} 1 & \frac{1}{2} & -\frac{1}{3} & \frac{1}{4} \\ 2 & \frac{1}{2} & -\frac{3}{4} & \frac{5}{4} \\ 3 & \frac{3}{5} & \frac{3}{7} & \frac{3}{8} \end{bmatrix}$$

```
>
```

```
[> A:= matrix(3,3,[[1,-3,-2],[2,-3,5],[-4,2,6]]);
```

```
> A:= matrix(3,3,[[1,-3,-2],[2,-3,5],[-4,2,6]]);
```

$$A := \begin{bmatrix} 1 & -3 & -2 \\ 2 & -3 & 5 \\ -4 & 2 & 6 \end{bmatrix}$$

```
[> B:=matrix([[1,-3,-2],[2,-3,5],[-4,2,6]]);
```

```
> B:=matrix([[1,-3,-2],[2,-3,5],[-4,2,6]]);
```

$$B := \begin{bmatrix} 1 & -3 & -2 \\ 2 & -3 & 5 \\ -4 & 2 & 6 \end{bmatrix}$$

```
>
```

```
[> Matrix (2,2,fill=a);
```

```
> Matrix (2,2,fill=a);
```

```
> | A uppercase
```

$$\begin{bmatrix} a & a \\ a & a \end{bmatrix}$$

```
[> Matrix (2,2,symbol=a);
```

```
> Matrix (2,2,symbol=a);
```

$$\begin{bmatrix} a_{1,1} & a_{1,2} \\ a_{2,1} & a_{2,2} \end{bmatrix}$$

```
>
```

```
[> f:=i->x*i-1;
```

```
[> M:=Matrix(2,f);
```

```
> f:=i->x*i-1;
```

$$f:=i \rightarrow x i - 1$$

```
> M:=Matrix(2,f);
```

$$M:=\begin{bmatrix} x-1 & x-1 \\ 2x-1 & 2x-1 \end{bmatrix}$$

```
>
```

```
[> g:=(i,j)->x*(i+j-1);
```

```
[> M1:=Matrix(3,g);
```

```
> g:=(i,j)->x*(i+j-1);
```

$$g:=(i,j) \rightarrow x(i+j-1)$$

```
> M1:=Matrix(3,g);
```

$$M1:=\begin{bmatrix} x & 2x & 3x \\ 2x & 3x & 4x \\ 3x & 4x & 5x \end{bmatrix}$$

```
[> |
```

To define a diagonal matrix.

```
[> C:=diag(1,2,3);
```

```
> C:=diag(1,2,3);
```

$$C:=\begin{bmatrix} 1 & 0 & 0 \\ 0 & 2 & 0 \\ 0 & 0 & 3 \end{bmatrix}$$

```
[> diag(1,1,1);
```

```
> diag(1,1,1);
```

$$\begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

To define lower triangular matrix.

```
[> M_1:=Matrix(3,[[x],[y,y],[z,z,z]],shape=triangular[lower]);  
> M_1:=Matrix(3,[[x],[y,y],[z,z,z]],shape=triangular[lower]);
```

$$M_1 := \begin{bmatrix} x & 0 & 0 \\ y & y & 0 \\ z & z & z \end{bmatrix}$$

```
[> M_2:=Matrix(3,[[1,2,3],[4,5,6],[7,8,9]],shape=triangular[upper]);  
> M_2:=Matrix(3,[[1,2,3],[4,5,6],[7,8,9]],shape=triangular[upper]);
```

$$M_2 := \begin{bmatrix} 1 & 2 & 3 \\ 0 & 5 & 6 \\ 0 & 0 & 9 \end{bmatrix}$$

To define zero matrix

```
[> M_3:=Matrix(4,4,shape=zero);
```

```
> M_3:=Matrix(4,4,shape=zero);
```

$$M_3 := \begin{bmatrix} 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix}$$

To define identity matrix

```
[> M_3:=Matrix(3,3,shape=identity);
```

```
> M_3:=Matrix(3,3,shape=identity);
```

$$M_3 := \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

To find an entry of a matrix, follow the name of the matrix by indices inside a square bracket.


```
[> A:= matrix(3,3,[[1,-3,-2],[2,-3,5],[-4,2,6]]);
[> A[3,2];
```

```
> A:= matrix(3,3,[[1,-3,-2],[2,-3,5],[-4,2,6]]);
```

$$A := \begin{bmatrix} 1 & -3 & -2 \\ 2 & -3 & 5 \\ -4 & 2 & 6 \end{bmatrix}$$

```
> A[3,2];
```

2

```
[> B:=matrix([[1,-3,-2],[2,-3,5],[-4,2,6]]);
[> B[1,2];
```

```
> B:=matrix([[1,-3,-2],[2,-3,5],[-4,2,6]]);
```

$$B := \begin{bmatrix} 1 & -3 & -2 \\ 2 & -3 & 5 \\ -4 & 2 & 6 \end{bmatrix}$$

```
> B[1,2];
```

-3

Matrix Operation

1.1.1 Addition and Scalar Multiplication

Algebraic expressions with matrices are evaluated by the command `evalm`. scalar multiplication of matrices done by the usual symbol `*`.

```
[> evalm(A+B);
```

```
> evalm(A+B);
```

$$\begin{bmatrix} 2 & -6 & -4 \\ 4 & -6 & 10 \\ -8 & 4 & 12 \end{bmatrix}$$

```
[> evalm(2*A+3*B);
```

```
> evalm(2*A+3*B) ;
```

$$\begin{bmatrix} 5 & -15 & -10 \\ 10 & -15 & 25 \\ -20 & 10 & 30 \end{bmatrix}$$

1.1.2 Matrix Multiplication

For matrix multiplication the symbol `&*` is used.

```
[> evalm(A&*B);
```

```
> evalm(A&*B) ;
```

$$\begin{bmatrix} 3 & 2 & -29 \\ -24 & 13 & 11 \\ -24 & 18 & 54 \end{bmatrix}$$

1.1.3 Transpose

To get the transpose of a matrix the command `transpose` is used.

```
[> transpose(A);
```

```
> transpose(A) ;
```

$$\begin{bmatrix} 1 & 2 & -4 \\ -3 & -3 & 2 \\ -2 & 5 & 6 \end{bmatrix}$$

1.2 Row Operations

In your linear algebra class you will learn elementary row operations on matrices here we will use Maple to do the same thing. Let's define a new matrix A .

- **addrow(A,r1,r2,m)**

Returns a copy of a matrix A in which row $r2$ replaced with by $\text{row}(A,r2)+m*\text{row}(A,r1)$

```
[> A:=matrix([[1,4,3,10],[2,1,-1,-1],[3,-1,4,11
[> addrow(A,1,2,-2);
[> addrow(%,1,3,-3);
```

Initial matrix A :

$$A := \begin{bmatrix} 1 & 4 & 3 & 10 \\ 2 & 1 & -1 & -1 \\ 3 & -1 & 4 & 11 \end{bmatrix}$$

Command: `> addrow(A, 1, 2, -2);`

Row operation: $R_2 \rightarrow R_2 + (-2)R_1$

Resulting matrix:

$$\begin{bmatrix} 1 & 4 & 3 & 10 \\ 0 & -7 & -7 & -21 \\ 3 & -1 & 4 & 11 \end{bmatrix}$$

Command: `> addrow(%, 1, 3, -3);`

Row operation: $R_3 \rightarrow R_3 + (-3)R_1$

Resulting matrix:

$$\begin{bmatrix} 1 & 4 & 3 & 10 \\ 0 & -7 & -7 & -21 \\ 0 & -13 & -5 & -19 \end{bmatrix}$$

- **mulrow(A,row,expr)**

Returns a matrix A in which has the same entries as A with the r^{th} row multiplied by $expr$

```
[> mulrow(A,2,-1/7);
```

```
[> mulrow(%,3,-1/3);
```

$A := \begin{bmatrix} 1 & 4 & 3 & 10 \\ 2 & 1 & -1 & -1 \\ 3 & -1 & 4 & 11 \end{bmatrix}$

`> mulrow(A, 2, -1/7);`

`> mulrow(A, 2, -1/7);`

$$\begin{bmatrix} 1 & 4 & 3 & 10 \\ -2 & -1 & 1 & 1 \\ 3 & -1 & 4 & 11 \end{bmatrix}$$

`> mulrow(%, 3, -1/3);`

`> mulrow(%, 3, -1/3);`

$$\begin{bmatrix} 1 & 4 & 3 & 10 \\ -2 & -1 & 1 & 1 \\ -1 & \frac{1}{3} & -\frac{4}{3} & -\frac{11}{3} \end{bmatrix}$$

$$\left[\begin{array}{ccc|c} 1 & 4 & 3 & 10 \\ 2 & 1 & -1 & -1 \\ 3 & -1 & 4 & 11 \end{array} \right]$$

$R_2 \rightarrow (-\frac{1}{7})R_2$

$$\downarrow$$

$$\left[\begin{array}{ccc|c} 1 & 4 & 3 & 10 \\ -\frac{2}{7} & -\frac{1}{7} & \frac{1}{7} & \frac{1}{7} \\ 3 & -1 & 4 & 11 \end{array} \right]$$

$R_3 \rightarrow (-\frac{1}{3})R_3$

$$\downarrow$$

$$\left[\begin{array}{ccc|c} 1 & 4 & 3 & 10 \\ -\frac{2}{7} & -\frac{1}{7} & \frac{1}{7} & \frac{1}{7} \\ 1 & -\frac{1}{3} & \frac{4}{3} & \frac{11}{3} \end{array} \right]$$

- **swaprow(A, r1, r2)**
This command interchange row r1 and r2 of A

`[> swaprow(A, 2, 3);`

$$A := \begin{bmatrix} 1 & 4 & 3 & 10 \\ 2 & 1 & -1 & -1 \\ 3 & -1 & 4 & 11 \end{bmatrix}$$

$$> \text{swaprow}(A, 2, 3)$$

$$> \text{swaprow}(A, 2, 3);$$

$$\begin{bmatrix} 1 & 4 & 3 & 10 \\ 3 & -1 & 4 & 11 \\ 2 & 1 & -1 & -1 \end{bmatrix}$$

$$\begin{bmatrix} 1 & 4 & 3 & 10 \\ 2 & 1 & -1 & -1 \\ 3 & -1 & 4 & 11 \end{bmatrix} \xrightarrow{R_2 \leftrightarrow R_3} \begin{bmatrix} 1 & 4 & 3 & 10 \\ 3 & -1 & 4 & 11 \\ 2 & 1 & -1 & -1 \end{bmatrix}$$

Similarly, you can learn `addcol`, `mulcol`, `swapcol` commands by your self.

1.3 Determinanat of Matrix

To find the determinate of a matrix, maple has a special command `det`.

```
[> M1 :=matrix (3, 3, [1,-2,-3,2,-2,2,3,-3,3]);
[> det(M1);
```

```
> M1 :=matrix (3, 3, [1,-2,-3,2,-2,2,3,-3,3]);
                                
$$M1 := \begin{bmatrix} 1 & -2 & -3 \\ 2 & -2 & 2 \\ 3 & -3 & 3 \end{bmatrix}$$

> det(M1);
                                0
```

```
[> B:=matrix([[1,-3,-2],[2,-3,5],[-4,2,6]]);
[> det(B);
```

```
> B:=matrix([[1,-3,-2],[2,-3,5],[-4,2,6]]);
```

$$B := \begin{bmatrix} 1 & -3 & -2 \\ 2 & -3 & 5 \\ -4 & 2 & 6 \end{bmatrix}$$

```
> det(B);
```

84

1.3.1 Inverse of a matrix

The `inverse` command is used to find the inverse of a square matrix, if exists.

```
[> B:=matrix([[1,-3,-2],[2,-3,5],[-4,2,6]]);
[> inverse(B);
```

```
> B:=matrix([[1,-3,-2],[2,-3,5],[-4,2,6]]);
```

$$B := \begin{bmatrix} 1 & -3 & -2 \\ 2 & -3 & 5 \\ -4 & 2 & 6 \end{bmatrix}$$

```
> inverse(B);
```

$$\begin{bmatrix} \frac{-1}{3} & \frac{1}{6} & \frac{-1}{4} \\ \frac{-8}{21} & \frac{-1}{42} & \frac{-3}{28} \\ \frac{-2}{21} & \frac{5}{42} & \frac{1}{28} \end{bmatrix}$$

```
>
```

```
[> inverse([[2,-1],[3,2]]);
```

> **inverse**([[2,-1],[3,2]]);

$$\begin{bmatrix} \frac{2}{7} & \frac{1}{7} \\ \frac{-3}{7} & \frac{2}{7} \end{bmatrix}$$

>

1.4 Exercise

Exercise 1.1. Performs the indicated computations.

A. $\begin{bmatrix} 1 & 0 & 3 \\ 2 & -1 & 6 \end{bmatrix} + \begin{bmatrix} 2 & 0 & 4 \\ -2 & 5 & 8 \end{bmatrix}$

B. $5 \begin{bmatrix} 2 & 1 & 3 \\ -1 & 2 & 4 \\ -6 & 1 & 5 \end{bmatrix} - 3 \begin{bmatrix} -2 & 1 & 4 \\ 5 & 0 & 7 \\ 2 & -1 & 3 \end{bmatrix}$

C. $\begin{bmatrix} 2 & 3 \\ -1 & 4 \end{bmatrix} \begin{bmatrix} 5 & -1 \\ 2 & 7 \end{bmatrix}$

D. $\begin{bmatrix} 2 & 3 & 1 & 5 \\ 0 & 6 & 2 & 4 \end{bmatrix} \begin{bmatrix} 5 & 7 & 1 \\ 2 & 0 & 3 \\ 1 & 0 & 0 \\ 0 & 5 & 6 \end{bmatrix}$

E. $\begin{bmatrix} 2 & 4 & 3 \\ 1 & 3 & 5 \end{bmatrix} \begin{bmatrix} 0 & 7 & 4 \\ 2 & 3 & 0 \end{bmatrix}$

F. $3 \begin{bmatrix} -2 & 1 \\ 0 & 4 \\ 2 & 3 \end{bmatrix}$

G. $\begin{bmatrix} 1 & 0 & 3 & -1 & 5 \\ 2 & 1 & 6 & 2 & 5 \end{bmatrix} \begin{bmatrix} 7 & 1 \\ 2 & 3 \\ -1 & 0 \\ 5 & 6 \\ 2 & 3 \end{bmatrix}$

H. $\begin{bmatrix} 1 & -1 & 2 \\ 3 & 5 & 6 \\ 2 & 4 & -1 \end{bmatrix} \begin{bmatrix} 2 \\ 1 \\ 3 \end{bmatrix}$

Exercise 1.2. Determine the given matrices are invertible. If they are, compute the inverse.

A. $\begin{bmatrix} 2 & 1 \\ 3 & 2 \end{bmatrix}$

B. $\begin{bmatrix} 1 & 1 & 1 \\ 0 & 2 & 3 \\ 5 & 5 & 1 \end{bmatrix}$

C. $\begin{bmatrix} a & a \\ b & b \end{bmatrix}$

D. $\begin{bmatrix} 1 & 1 & 1 \\ 0 & 1 & 1 \\ 0 & 0 & 1 \end{bmatrix}$

E. $\begin{bmatrix} 5 & 7 & 0 \\ 2 & 1 & 3 \\ 1 & 0 & 0 \\ 0 & 4 & 3 \end{bmatrix}$

F. $\begin{bmatrix} 1 & 1 & 1 & 1 \\ 1 & 2 & -1 & 2 \\ 1 & -1 & 2 & 1 \\ 1 & 3 & 3 & 2 \end{bmatrix}$

G. $\begin{bmatrix} 1 & -3 & 0 & 2 \\ 3 & -12 & -2 & -6 \\ -2 & 10 & 2 & 5 \\ -1 & 6 & 1 & 3 \end{bmatrix}$

Exercise 1.3. Compute the 3×3 matrix whose entries are given by the function y^{ij} , Where $i = j = 1, 2, 3$.

Exercise 1.4. Compute the matrix, $\begin{bmatrix} x & x^2 \\ x^2 & x^3 \end{bmatrix}$ by defining a suitable function as in the exercise 1.3

Exercise 1.5. Which of the following matrices are skew-symmetric? (A square matrix is symmetric if $A^T = A$, where A^T is the transpose of A .)

A. $\begin{bmatrix} 1 & -6 \\ 6 & 0 \end{bmatrix}$

B. $\begin{bmatrix} 0 & -6 \\ 6 & 0 \end{bmatrix}$

C. $\begin{bmatrix} 2 & -2 & 2 \\ 2 & 2 & -2 \\ 2 & 2 & 2 \end{bmatrix}$

D. $\begin{bmatrix} 0 & 1 & -1 \\ -1 & 0 & 2 \\ 1 & -2 & 0 \end{bmatrix}$

Exercise 1.6. Convert the following matrix in to an upper triangular matrix.

$$\begin{bmatrix} 2 & 3 & -1 \\ -3 & -2 & 1 \\ 0 & 1 & 1 \end{bmatrix}$$

Exercise 1.7. Convert the following matrices into a identity matrix.

A. $\begin{bmatrix} 2 & 7 & 3 \\ 1 & 3 & 2 \\ 3 & 7 & 9 \end{bmatrix}$

B. $\begin{bmatrix} 2 & -3 \\ 3 & 4 \end{bmatrix}$

Chapter 2

System of linear equations and matrices

A linear system is a collection of first degree equations. A solution to a system consists of one or more sets of specific values that are common solutions to each of the individual equations. Here is a simple example which we can solve quite easily using the solve command. ## Augment of a matrix

```
[> with(linalg):  
[> A:=matrix([[1,2],[2,3]]);
```

Output:

```
[> with(linalg):  
[> matrix([[1,2],[2,3]]);
```

$$\begin{bmatrix} 1 & 2 \\ 2 & 3 \end{bmatrix}$$

```
[> B:=matrix([[3,4,5],[6,7,8]]);
```

Output:

```
> B:=matrix([[3,4,5],[6,7,8]]);
```

$$B := \begin{bmatrix} 3 & 4 & 5 \\ 6 & 7 & 8 \end{bmatrix}$$

```
augment(A,B);
```

Output:

```
> augment(A,B);
```

$$\begin{bmatrix} 1 & 2 & 3 & 4 & 5 \\ 2 & 3 & 6 & 7 & 8 \end{bmatrix}$$

The function `augment` joins two or more matrices together horizontally. The matrices and vectors must have the same number of rows.

2.1 Solving Linear Systems

$$\begin{array}{rrcrcl} 2x & + & 5y & - & 4z & = & 9 \\ 3x & + & 5y & + & 2z & = & 12 \\ 4x & - & y & + & 5z & = & -3 \end{array}$$

$$\underbrace{\begin{bmatrix} 2 & 5 & -4 \\ 3 & 5 & 2 \\ 4 & -1 & 5 \end{bmatrix}}_A \underbrace{\begin{bmatrix} x \\ y \\ z \end{bmatrix}}_X = \underbrace{\begin{bmatrix} 9 \\ 12 \\ -3 \end{bmatrix}}_B$$

2.1.1 Method I

A linear system is a collection of first degree equations. A solution to a system consists of one or more set of specific values that our common solution to each of the individual equations. Here is a simple example which we can solve quite easily using the `solve` command.

```
[> sys:={2*x+5*y-4*z=9,3*x+5*y+2*z=12,4*x-y+5*z=-3};
[> solve(sys,{x,y,z});
```

Output:

```
> sys:={2*x+5*y-4*z=9,3*x+5*y+2*z=12,4*x-y+5*z=-3};
      sys := {2 x + 5 y - 4 z = 9, 3 x + 5 y + 2 z = 12, 4 x - y + 5 z = -3}
> solve(sys, {x,y,z});
      {x = -33/37, y = 99/37, z = 24/37}
```

Maple will automatically use fractions, however you can force decimal answers using the *evalf* command.

```
[> evalf(%);
```

Output:

```
> evalf(%);
      {x = -0.8918918919, y = 2.675675676, z = 0.6486486486}
```

2.1.2 Method II

We can also convert the above system of equations to a matrix system.

```
[> A:=genmatrix(sys,[x,y,z],b);
[> evalm(b);
```

Output:

```
> A:=genmatrix(sys,[x,y,z],b);
      A :=  $\begin{bmatrix} 2 & 5 & -4 \\ 3 & 5 & 2 \\ 4 & -1 & 5 \end{bmatrix}$ 
> evalm(b);
       $[9, 12, -3]$ 
```

In this case ,

- A is the coefficient matrix, and

- b is vector representing the constant values.

The command `evalm(b)` evaluated b as a matrix (a vector is a $n \times 1$ matrix). In other words, this command simply writes out what the vector b looks like.

```
[> linsolve(A,b);
```

Output:

```
> linsolve(A,b);
```

$$\begin{bmatrix} \frac{-33}{37} & \frac{99}{37} & \frac{24}{37} \end{bmatrix}$$

2.1.3 Method III

Another way to solve a matrix equation $Ax = b$ is to left multiply both sides by the inverse matrix A^{-1} , if it exists, to get the solution $x = A^{-1}b$.

```
[> inverse(A);
[> evalm(inverse(A)*b);
```

Output:

```
> inverse(A); |
```

$$\begin{bmatrix} \frac{9}{37} & \frac{-7}{37} & \frac{10}{37} \\ \frac{-7}{111} & \frac{26}{111} & \frac{-16}{111} \\ \frac{-23}{111} & \frac{22}{111} & \frac{-5}{111} \end{bmatrix}$$

```
> evalm(inverse(A)*b);
```

$$\begin{bmatrix} \frac{-33}{37} & \frac{99}{37} & \frac{24}{37} \end{bmatrix}$$

The `evalm` command forces a matrix computation and express the result.

2.2 Dependent Systems:

Some systems are dependent which means there are an infinite number of solutions. The form of these solutions will entail the use of parameters. Let's look an example using two different methods; `solve` and `linsolve`.

Example 2.1.

$$\begin{array}{rrrrrr} x & - & 2y & + & z & = & 3 \\ x & + & y & - & 2z & = & -4 \\ 2x & - & y & - & z & = & -1 \end{array}$$

```
[> restart;
[> with(linalg):
[> sys:={x-2*y+z=3,x+y-2*z=-4,2*x-y-z=-1};
[> solve(sys,{x,y,z});
```

```
> restart;
> with(linalg):
> sys:={x-2*y+z=3,x+y-2*z=-4,2*x-y-z=-1};
                                     sys := {x - 2 y + z = 3, x + y - 2 z = -4, 2 x - y - z = -1}
> solve(sys,{x,y,z});
```

$$\{x = -\frac{5}{3} + z, y = -\frac{7}{3} + z, z = z\}$$

output:

```
[> A:=genmatrix(sys,[x,y,z],b);
[> evalm(b);
[> linsolve(A,b);
```

output:

```
> A:=genmatrix(sys,[x,y,z],b);
```

$$A := \begin{bmatrix} 1 & -2 & 1 \\ 1 & 1 & -2 \\ 2 & -1 & -1 \end{bmatrix}$$

```
> evalm(b);
```

$$[3, -4, -1]$$

```
> linsolve(A,b);
```

$$\left[-\frac{5}{3} + t_1, -\frac{7}{3} + t_1, t_1 \right]$$

In both cases, the solution contains a parameter. The `solve` command express it in terms actual variable used, and the `linsolve` command use the funny `t` character to distinguish it from a variable you might have defined yourself.

2.3 Inconsistent System

An Inconsistent system has no solution. Maple generally, refers to answer questions which have no answer.

Example 2.2.

$$\begin{array}{rrcrcl} x & + & y & - & 3z & = & 10 \\ x & + & y & - & z & = & 1 \\ x & + & y & + & z & = & 8 \end{array}$$

Let's try to solve this system.

```
[> restart;
[> with(linalg):
[> sys:={x+y-3*z=10,x+y-z=1,x+y+z=8};
[> solve(sys,{x,y,z});
```

Output:

```
> restart;
> with(linalg):
> sys:={x+y-3*z=10,x+y-z=1,x+y+z=8};
                                sys := {x + y - 3 z = 10, x + y - z = 1, x + y + z = 8}
> solve(sys,{x,y,z});
> |
```

This method does not work. It does not return anything.

Let's try `linsolve` method.

```
[> A:=genmatrix(sys,[x,y,z],b);
[> linsolve(A,b);
```

Output:


```

> A:=genmatrix(sys,[x,y,z],b);

```

$$A := \begin{bmatrix} 1 & 1 & -3 \\ 1 & 1 & -1 \\ 1 & 1 & 1 \end{bmatrix}$$

```

> linsolve(A,b);
>
>

```

This method also does not work.

```
[> evalm(inverse(A)&*b);
```

Output:

```

> evalm(inverse(A) &*b);
Error, (in linalg:-inverse) singular matrix
>

```

2.3.0.1 Automatic Reduction

```

> sys:={3*x+5*y+2*z=12,2*x+5*y-4*z=9,4*x-y+5*z=-3};
> evalm(b);
> C:=augment(A,b);

```

Output:

```

> sys:={3*x+5*y+2*z=12,2*x+5*y-4*z=9,4*x-y+5*z=-3};

```

$$\text{sys} := \{2x + 5y - 4z = 9, 3x + 5y + 2z = 12, 4x - y + 5z = -3\}$$

```

> evalm(b);

```

$$[10, 1, 8]$$

```

> C:=augment(A,b);

```

$$C := \begin{bmatrix} 1 & 1 & -3 & 10 \\ 1 & 1 & -1 & 1 \\ 1 & 1 & 1 & 8 \end{bmatrix}$$

2.3.1 Method IV

An even faster method is simplifying let maple to do all the work for us. The `gausselim` command will perform all of the steps of Gaussian eliminations and reduce an augmented matrix to **row echelon form**.

```
[> sys:={3*x+5*y-2*z=12,2*x+5*y-4*z=9,4*x-y+5*z=-3};
[> A:=genmatrix(sys,[x,y,z],b);
[> evalm(b);
[> C:=augment(A,b);
[> gausselim(C);
```



```
> sys:={3*x+5*y-2*z=12,2*x+5*y-4*z=9,4*x-y+5*z=-3};
sys := {2 x + 5 y - 4 z = 9, 3 x + 5 y - 2 z = 12, 4 x - y + 5 z = -3}
> A:=genmatrix(sys,[x,y,z],b);
```

$$A := \begin{bmatrix} 2 & 5 & -4 \\ 3 & 5 & -2 \\ 4 & -1 & 5 \end{bmatrix}$$

```
> evalm(b);
```

$$[9, 12, -3]$$

```
> C:=augment(A,b);
```

$$C := \begin{bmatrix} 2 & 5 & -4 & 9 \\ 3 & 5 & -2 & 12 \\ 4 & -1 & 5 & -3 \end{bmatrix}$$

```
> gausselim(C);
```

$$\begin{bmatrix} 2 & 5 & -4 & 9 \\ 0 & -11 & 13 & -21 \\ 0 & 0 & \frac{23}{22} & \frac{36}{11} \end{bmatrix}$$

```
> |
```

2.3.2 Method V

The command `gaussjordan` does the same thing. It performs all the steps of Gauss Jordan elimination and reduces and reduces an augmented matrix into reduced row echelon form.

```
[> gaussjordan(C);
```

```
> gaussjord(C) ;
```

$$\begin{bmatrix} 1 & 0 & 0 & \frac{-75}{23} \\ 0 & 1 & 0 & \frac{129}{23} \\ 0 & 0 & 1 & \frac{72}{23} \end{bmatrix}$$

```
>
```

2.4 Exercise

Exercise 2.1. Solve the following system of equations.

$$\begin{array}{rrcrcl} x_1 & + & 2x_2 & + & 3x_3 & = & 9 \\ 2x_1 & - & x_2 & + & x_3 & = & 8 \\ 3x_1 & & & - & x_3 & = & 3 \end{array}$$

Exercise 2.2. A small manufacturing plant makes three types of inflatable boats: one-person, two-person, and four-person models. Each boat requires the services of three departments, as listed in the table. The cutting, assembly, and packaging departments have available a maximum of 380, 330, and 120 labor hours per week, respectively.

Department	One Person Boat	Two-Person Boat	Four-Person Boat
Cutting	0.5 hr	1.0 hr	1.5 hr
Assembling	0.6 hr	0.9 hr	1.2 hr
Packaging	0.2 hr	0.3 hr	0.5 hr

- (A) How many boats of each type must be produced each week for the plant to operate at full capacity?
- (B) How is the production schedule in part A affected if the packaging department is no longer used?
- (C) How is the production schedule in part A affected if the four-person boat is no longer produced?

Chapter 3

Integration

The `int` command is used to compute both definite and indefinite integrals of Maple expressions as shown by the following examples.

3.1 Indefinite Integration

```
[> int(x^2,x);
```

```
> int (x^2,x) ;
```

$$\frac{x^3}{3}$$

```
[> int(sin(2*t),t);
```

> int(sin(2*t), t) ;

$$-\frac{1}{2} \cos(2 t)$$

```
[> f:=x->(3*x-6)/(x^2-4);
[> int(f(x),x);
```

> f:=x->(3*x-6)/(x^2-4) ;

$$f:=x \rightarrow \frac{3x-6}{x^2-4}$$

> int(f(x), x) ;

$$3 \ln(x+2)$$

```
[> Int(sin(x),x);
```

> Int(sin(x), x) ;

$$\int \sin(x) dx$$

```
[> Int(3*x^2+2)^(5/3),x);
```

> **Int(x*(3*x^2+2)^(5/3),x);**

$$\int x (3x^2 + 2)^{(5/3)} dx$$

[> Int(x*(3*x^2+2)^(5/3),x)=int(x*(3*x^2+2)^(5/3),x);

> **Int(x*(3*x^2+2)^(5/3),x)=int(x*(3*x^2+2)^(5/3),x);**

$$\int x (3x^2 + 2)^{(5/3)} dx = \frac{(3x^2 + 2)^{(8/3)}}{16}$$

Notice that Maple doesn't include a constant of integration for indefinite integrals. Where there are constants, Parameters other variable around, Maple assume that you mean to take the integral as the variable you specify changes, and that all other letters in the expression represent constants.

3.2 Definite Integration

[> Int(g(x),x=a..b);

> **Int(g(x),x=a..b);**

$$\int_a^b g(x) dx$$

[> Int(x^2*exp(x),x=0..2)=int(x^2*exp(x),x=0..2);

> **Int(x^2*exp(x),x=0..2)=int(x^2*exp(x),x=0..2);**

$$\int_0^2 x^2 e^x dx = -2 + 2e^2$$

[> f := x -> x*sin(x);

[> int(f(x),x=0..Pi);

```
> f := x -> x*sin(x) ;
```

$$f := x \rightarrow x \sin(x)$$

```
> int(f(x), x=0..Pi) ;
```

$$\pi$$

Some functions can not be integrated analytically, but the definite integrals of such functions still have meaning and MAPLE can determine them.

```
[> int(exp(-x^2)*ln(x), x) ;
```

```
> int(exp(-x^2)*ln(x), x) ;
```

$$\int e^{(-x^2)} \ln(x) dx$$

```
[> int(exp(-x^2)*ln(x), x=0..infinity) ;
```

```
[> evalf(%);
```

```
> int(exp(-x^2)*ln(x), x=0..infinity) ;
```

$$-\frac{\sqrt{\pi} \gamma}{4} - \frac{1}{2} \sqrt{\pi} \ln(2)$$

```
> evalf(%);
```

$$-0.8700577270$$

3.3 Integration by Parts

The `intparts` command is used to integrate by parts, and the command exist in `student` package.

The `intparts` command has two arguments: the first is the expression to be anti-differentiated, and the second is the choice for, the piece which is to be differentiated.

$$\int uD(v)dx = uv - \int vD(u)dx$$

Example 3.1. Method of integration by parts to compute $\int x \cos(x)$

```
[> restart;
[> with(student):
[> p1:= Int(x*cos(x),x);
[> p2:= intparts(p1,x);
[> p3:= value(p2);
[> p1 = p3 + C;
```

```
> restart;
> with(student):
> p1:= Int(x*cos(x),x);
```

$$p1 := \int x \cos(x) dx$$

```
> p2:= intparts(p1,x);
```

$$p2 := x \sin(x) - \int \sin(x) dx$$

```
> p3:= value(p2);
```

$$p3 := x \sin(x) + \cos(x)$$

```
> p1 = p3 + C;
```

$$\int x \cos(x) dx = x \sin(x) + \cos(x) + C$$

```
[> restart;
[> with(student):
[> Int(sqrt(x)*ln(x),x)=intparts(Int(sqrt(x)*ln(x),x),ln(x));
```

```

> restart;
> with(student):
> Int(sqrt(x)*ln(x),x)=intparts(Int(sqrt(x)*ln(x),x),ln(x));

```

$$\int \sqrt{x} \ln(x) dx = \frac{2}{3} \ln(x) x^{(3/2)} - \int \frac{2\sqrt{x}}{3} dx$$

```

[> Int(sqrt(x)*ln(x),x)=value(intparts(Int(sqrt(x)*ln(x),x),ln(x)));

```

```

> Int(sqrt(x)*ln(x),x)=value(intparts(Int(sqrt(x)*ln(x),x),ln(x)));

```

$$\int \sqrt{x} \ln(x) dx = \frac{2}{3} \ln(x) x^{(3/2)} - \frac{4x^{(3/2)}}{9}$$

3.4 Substitutions

These are done using the `changevar` command. Look at the following example.

Example 3.2. Evaluate $\int 4 \cos(\sin(x)) dx$ using the substitution, $u = \cos(x)$.

```

[> Int(cos(x)^4*sin(x),x)=changevar(u=cos(x),Int(cos(x)^4*sin(x),x),u);

```

```

> Int(cos(x)^4*sin(x),x)=changevar(u=cos(x),Int(cos(x)^4*sin(x),x),u);

```

$$\int \cos(x)^4 \sin(x) dx = \int -u^4 du$$

```

[> v1:=value(changevar(u=cos(x),Int(cos(x)^4*sin(x),x),u));

```

```

> v1:=value(changevar(u=cos(x),Int(cos(x)^4*sin(x),x),u));

```

$$v1 := -\frac{u^5}{5}$$

To back-substitute, we use the `subs` command,

```

[> subs(u=cos(x),v1);

```

```

> subs(u=cos(x), v1);

```

$$-\frac{1}{5} \cos(x)^5$$

3.5 Multiple Integration

3.5.1 Double integration (Area calculation)

```
[> Int(Int(x^2+2*x,x),x);
```

```
> Int(Int(x^2+2*x,x),x);
```

$$\iint x^2 + 2x \, dx \, dx$$

```
[> int(int(x^2+2*x,x),x);
```

```
> int(int(x^2+2*x,x),x);
```

$$\frac{1}{12}x^4 + \frac{1}{3}x^3$$

```
[> Int(Int(x+y^2,y=0..x),x=1..2);
```

```
> Int(Int(x+y^2,y=0..x),x=1..2);
```

$$\int_1^2 \int_0^x x + y^2 \, dy \, dx$$

```
[> a:=Int(x+y^2,y=0..x);
```

```
> a:=Int(x+y^2,y=0..x);
```

$$a := \int_0^x x + y^2 \, dy$$

```
[> b:=Int(a,x=1..2);
```

```
> b:=Int(a,x=1..2);
```

$$b := \int_1^2 \int_0^x x + y^2 \, dy \, dx$$

```
[> Int(Int(x+y^2,y=0..x),x=1..2)=int(int(x+y^2,y=0..x),x=1..2);
```

```
> Int(Int(x+y^2,y=0..x),x=1..2)=int(int(x+y^2,y=0..x),x=1..2);
```

$$\int_1^2 \int_0^x x + y^2 \, dy \, dx = \frac{43}{12}$$

```
[> A:=Doubleint(x+y^2,y=0..x,x=1..2);
```

```
[> value(A);
```

```
> A:=Doubleint(x+y^2,y=0..x,x=1..2);
```

$$A := \int_1^2 \int_0^x x + y^2 \, dy \, dx$$

```
> value(A);
```

$$\frac{43}{12}$$

3.5.2 Triple integration (Volume Calculation)

```
[> Int(Int(Int(x^3*exp(y)*sin(z),x=0..2),z=-Pi..2*Pi),y=0..1)=int(int(int(x^3*exp(y)*s
```

```
> A:=Doubleint(x+y^2,y=0..x,x=1..2);
```

$$A := \int_1^2 \int_0^x x + y^2 \, dy \, dx$$

```
> value(A);
```

$$\frac{43}{12}$$

```
[> V:=Tripleint(x^3*exp(y)*sin(z),x=0..2,z=-Pi..2*Pi,y=0..1);
[> value(V);

> Int(Int(Int(x^3*exp(y)*sin(z),x=0..2),z=-Pi..2*Pi),y=0..1)
=Int(int(int(x^3*exp(y)*sin(z),x=0..2),z=-Pi..2*Pi),y=0..1);
```

$$\int_0^1 \int_{-\pi}^{2\pi} \int_0^2 x^3 e^y \sin(z) \, dx \, dz \, dy = 8 - 8e$$

Note: Since there is often more than one way to find an indefinite integral, it may happen that the answer you obtain by doing the techniques seen in class is different than the one obtained with Maple.

3.6 Exercise

Exercise 3.1. Integrate the following expressions,

- i. $2x^2 + \frac{x}{2}$
- ii. $x^3(3x^2 + 2)^{\frac{5}{3}}$
- iii. e^{-x^2}

Exercise 3.2. Find the definite integral of the following over the given intervals,

- i. $\int_2^{10} x^3(3x^2 + 2)^{\frac{5}{3}} \, dx$
- ii. $\int_0^{\frac{\pi}{2}} \cos(x) \sin(x^2) \, dx$
- iii. $\int_0^{\pi} e^x \cos(x) \, dx$
- iv. $\int_0^{\pi} e^{2\sin(x)} \cos(x) \, dx$

Exercise 3.3. Use of the method of integration by parts to compute,

- i. $\int x e^{-2x} \, dx$
- ii. $\int x \ln(x) \, dx$

Exercise 3.4. Evaluate the following expressions

- i. $\int_{-3}^3 \int_{-3}^3 3x^2 + 5y^2 + 4 \, dx \, dy$
- ii. $\int_{-3}^3 \int_{-3}^3 9 \ln(x) - 5y^2 \, dx \, dy$

Note: Since there is often more than one way to find an indefinite integral, it may happen that the answer you obtain by doing the techniques seen in class is different than the one obtained with Maple. Compare the answers obtained by Maple and the answer you obtained manually for the integral, $\int \cos(3x) \, dx$

There are two main ways to use the definite integral. - The easiest one to understand is as a means for computing areas (and volumes). - The second way the definite integral is used is as a sum. That is, we use the definite integral to “add things up”.

3.7 Computing the area from the integral

Example 3.3. Find the area under the curve $x^2 \sin(x)$

```
[> int(x^2*sin(x),x=-Pi..Pi);
```

```
[> plot(x^2*sin(x),x=-Pi..Pi);
```

```
[> a1:=int(x^2*sin(x),x=-Pi..0);
```

```
‘[> a2:=int(x^2*sin(x),x=0..Pi);
```

```
[> abs(a1)+a2;
```

Example 3.4. Find the area bounded by the curves $-x^2 + 4x + 6$ and $\frac{x}{3} + 2$

```
[> f := x-> -x^2+4*x+6;
```

```
[> g := x-> x/3+2;
```

```
[> plot({f(x),g(x)},x=-2..6);
```


```
> f := x-> -x^2+4*x+6;
```

```
g := x-> x/3+2;
```

```
plot({f(x),g(x)},x=-2..6);
```

$$f := x \rightarrow -x^2 + 4x + 6$$

$$g := x \rightarrow \frac{1}{3}x + 2$$



```
[> a := fsolve(f(x)=g(x),x=-2..0);  
[> b := fsolve(f(x)=g(x),x=4..6);  
[> Area:=int(f(x)-g(x),x=a..b);
```

```

> a := fsolve(f(x)=g(x), x=-2..0);
                                     a := -0.8798034327
> b := fsolve(f(x)=g(x), x=4..6);
                                     b := 4.546470099
>
> Area:=int(f(x)-g(x), x=a..b);
                                     Area := 26.62893493

```

```
[> evalf(solve(-x^2+4*x+6-x/3-2,x));
```

```
[> Area:=int(f(x)-g(x), x=-.8798034327..0)+int(f(x)-g(x), x=0..4.546470099);
```

Example 3.5. Find the area of the region enclosed between the two curves $f(x) = 2x + 5$ and $g(x) = x^2 + 2$ from $x = 0$ to $x = 6$.

```


[> g:=x->x^2+2;f:=x->2*x+5;
[> plot({f(x),g(x)},x=-2..7,color=[red,blue]);

```

```

> g:=x->x^2+2;f:=x->2*x+5;
                                     g := x → x2 + 2
                                     f := x → 2x + 5
> plot({f(x),g(x)},x=-2..7,color=[red,blue]);

```

```
[> solve(x^2+2-2*x-5,x);
```

```
[> Area:=int(f(x)-g(x),x=0..3)+int(g(x)-f(x),x=3..6);
```

3.8 Exercise

Exercise 3.5. Area under the curve $10 - x^2$ in the interval $[0, 4]$.

Exercise 3.6. Area bounded by the curves $f(x) = \cos(x)$ and $y = \frac{1}{2}$ in the interval 0 to 8.

Exercise 3.7. Area bounded by the curves $f(x) = \frac{1}{4}x^2 - 4$ and $g(x) = \frac{1}{4}x + 1$.