



**BLACK HOLES**

OBSERVATORIO  
ASTRONÓMICO  
NACIONAL

# Classical Black Holes

---

## 10. Viscous Torques

Edward Larrañaga

# Outline for Part 1

## 1. Viscous Torques

### 1.1 Differential Rotation

### 1.2 Roche Lobe Overflow

## 2. Formation of the Accretion Disk

# Viscous Torques

## Accretion Disk

- If the only interaction between fluid elements is gravity, the angular momentum is conserved.

## Accretion Disk

- If the only interaction between fluid elements is gravity, the angular momentum is conserved.
- In the description of the accretion disk it is needed a force responsible for the redistribution of angular momentum.

# Differential Rotation

# Differential Rotation

- Keplerian Velocity

$$\Omega_k = \Omega_k(r)$$

# Differential Rotation

- Keplerian Velocity

$$\Omega_k = \Omega_k(r)$$

- Neighboring material at different radii moves with different velocity



# Differential Rotation

- Keplerian Velocity

$$\Omega_k = \Omega_k(r)$$

- Neighboring material at different radii moves with different velocity
- The disk is not a solid body. It moves with *differential rotation*.

## Differential Rotation

- Keplerian Velocity

$$\Omega_k = \Omega_k(r)$$

- Neighboring material at different radii moves with different velocity
- The disk is not a solid body. It moves with *differential rotation*.
- The thermal motion of the fluid molecules and the turbulent motion of the fluid produce *viscous stresses*.

## Differential Rotation

- In our simple description we will consider only momentum transport in the radial direction, produced by the process known as *shear viscosity*.

## Differential Rotation

- In our simple description we will consider only momentum transport in the radial direction, produced by the process known as *shear viscosity*.
- It appears when there are internal distortions (usually local stresses that are proportional to the local rate of strain)

## Differential Rotation

- In our simple description we will consider only momentum transport in the radial direction, produced by the process known as *shear viscosity*.
- It appears when there are internal distortions (usually local stresses that are proportional to the local rate of strain)
- Although the following is the simplest description, it can be used to describe other mechanisms of angular momentum transport such as magnetic loops that couple fluid elements at macroscopic distances across the disk.

# Modeling Viscosity

- $\lambda$ : Typical scale in the accretion disk

## Modeling Viscosity


- $\lambda$ : Typical scale in the accretion disk
- $\tilde{v}$ : typical speed in the accretion disk

## Modeling Viscosity

- $\lambda$ : Typical scale in the accretion disk
- $\tilde{v}$ : typical speed in the accretion disk
- As a first model, consider a uniform gas moving only in the tangential direction with velocity  $v_\phi(r)$ .



## Modeling Viscosity



## Modeling Viscosity

The only non-vanishing component of the stress tensor is the  $\phi$ -component of the force per unit surface of the  $r = \text{constant}$  surface,

$$\sigma_{r\phi} = -\eta \left. \frac{\partial v_\phi}{\partial r} \right|_R$$

## Modeling Viscosity

The only non-vanishing component of the stress tensor is the  $\phi$ -component of the force per unit surface of the  $r = \text{constant}$  surface,

$$\sigma_{r\phi} = -\eta \left. \frac{\partial v_\phi}{\partial r} \right|_R$$

$\eta$ : dynamical viscosity

$$v_\phi(r) = \Omega_k(r)r$$

$\frac{\partial v_\phi}{\partial r}$ : velocity gradient

## Modeling Viscosity

$$\sigma_{r\phi} = -\eta R \left. \frac{\partial \Omega_k}{\partial r} \right|_R$$

$\phi$ -component of the force per unit surface of the  $r = \text{constant}$  surface.

## Modeling Viscosity

The dynamical viscosity is usually written in terms of the *kinematical viscosity*,  $\nu$ , as  $\eta = \rho\nu$

## Modeling Viscosity

The dynamical viscosity is usually written in terms of the *kinematical viscosity*,  $\nu$ , as  $\eta = \rho\nu$

The kinematical viscosity is modeled as

$$\nu \sim \lambda \tilde{\nu}$$

## Modeling Viscosity

The dynamical viscosity is usually written in terms of the *kinematical viscosity*,  $\nu$ , as  $\eta = \rho\nu$

The kinematical viscosity is modeled as

$$\nu \sim \lambda \tilde{v}$$

Molecular transport:

$\lambda$ : mean free path

$\tilde{v}$ : thermal speed

## Modeling Viscosity

The dynamical viscosity is usually written in terms of the *kinematical viscosity*,  $\nu$ , as  $\eta = \rho\nu$

The kinematical viscosity is modeled as

$$\nu \sim \lambda \tilde{v}$$

Molecular transport:

$\lambda$ : mean free path

$\tilde{v}$ : thermal speed

Turbulent motion:

$\lambda$ : spatial scale (or characteristic wavelength) of the turbulence

$\tilde{v}$ : typical velocity of the eddies



## Modeling Viscosity

$$\sigma_{r\phi} = -\rho\nu R \left. \frac{\partial \Omega_k}{\partial r} \right|_R$$

$\phi$ -component of the force per unit surface of the  $r = \text{constant}$  surface.

## Modeling Viscosity

$$\sigma_{r\phi} = -\rho\nu R \left. \frac{\partial \Omega_k}{\partial r} \right|_R$$

$\phi$ -component of the force per unit surface of the  $r = \text{constant}$  surface.

The net torque ( $\vec{R} \times \vec{F}$ ) is calculated by multiplying  $\sigma_{r\phi}$  by the area of the  $r = R = \text{constant}$  surface,  $2\pi RH$ .

## Modeling Viscosity

$$\sigma_{r\phi} = -\rho\nu R \left. \frac{\partial \Omega_k}{\partial r} \right|_R$$

$\phi$ -component of the force per unit surface of the  $r = \text{constant}$  surface.

The net torque ( $\vec{R} \times \vec{F}$ ) is calculated by multiplying  $\sigma_{r\phi}$  by the area of the  $r = R = \text{constant}$  surface,  $2\pi RH$ .

$H$ : Height of the disk

## Modeling Viscosity

Net torque on the outer ring due to the inner one .

## Modeling Viscosity

Net torque on the outer ring due to the inner one .

$$\text{Torque} = R \times 2\pi RH \times \sigma_{r\phi}$$

## Modeling Viscosity

Net torque on the outer ring due to the inner one .

$$\text{Torque} = R \times 2\pi R H \times \sigma_{r\phi}$$

$$\text{Torque} = -2\pi R^3 H \rho \nu \left. \frac{\partial \Omega_k}{\partial r} \right|_R$$

# Modeling Viscosity

Using the surface mass density,

## Modeling Viscosity

Using the surface mass density,

$$\Sigma = \int_0^H \rho(z) dz = \rho H$$



## Modeling Viscosity

Using the surface mass density,

$$\Sigma = \int_0^H \rho(r) dz = \rho H$$

$$\text{Torque} = -2\pi R^3 \Sigma \nu \left. \frac{\partial \Omega_k}{\partial r} \right|_R$$

## Modeling Viscosity

We define the torque at the coordinate  $r$  as the function,

## Modeling Viscosity

We define the torque at the coordinate  $r$  as the function,

$$G(r) = 2\pi r^3 \Sigma \nu \frac{\partial \Omega_k}{\partial r}$$

- If  $\frac{\partial \Omega_k}{\partial r} = 0$

## Modeling Viscosity

We define the torque at the coordinate  $r$  as the function,

$$G(r) = 2\pi r^3 \Sigma \nu \frac{\partial \Omega_k}{\partial r}$$

- If  $\frac{\partial \Omega_k}{\partial r} = 0$   
→ Rigid body rotation

## Modeling Viscosity

We define the torque at the coordinate  $r$  as the function,

$$G(r) = 2\pi r^3 \Sigma \nu \frac{\partial \Omega_k}{\partial r}$$

- If  $\frac{\partial \Omega_k}{\partial r} = 0$ 
  - Rigid body rotation
  - No torque

## Modeling Viscosity

We define the torque at the coordinate  $r$  as the function,

$$G(r) = 2\pi r^3 \Sigma \nu \frac{\partial \Omega_k}{\partial r}$$

- If  $\frac{\partial \Omega_k}{\partial r} = 0$ 
  - Rigid body rotation
  - No torque
- If  $\Omega_k(r)$  decreases outwards,  $\frac{\partial \Omega_k}{\partial r} < 0$ ,

## Modeling Viscosity

We define the torque at the coordinate  $r$  as the function,

$$G(r) = 2\pi r^3 \Sigma \nu \frac{\partial \Omega_k}{\partial r}$$

- If  $\frac{\partial \Omega_k}{\partial r} = 0$ 
  - Rigid body rotation
  - No torque
- If  $\Omega_k(r)$  decreases outwards,  $\frac{\partial \Omega_k}{\partial r} < 0$ , and  $G(r) < 0$ ,

## Modeling Viscosity

We define the torque at the coordinate  $r$  as the function,

$$G(r) = 2\pi r^3 \Sigma \nu \frac{\partial \Omega_k}{\partial r}$$

- If  $\frac{\partial \Omega_k}{\partial r} = 0$ 
  - Rigid body rotation
  - No torque
- If  $\Omega_k(r)$  decreases outwards,  $\frac{\partial \Omega_k}{\partial r} < 0$ , and  $G(r) < 0$ ,
  - angular momentum goes from inner circles to outer circles.



## Modeling Viscosity

We define the torque at the coordinate  $r$  as the function,

$$G(r) = 2\pi r^3 \Sigma \nu \frac{\partial \Omega_k}{\partial r}$$

- If  $\frac{\partial \Omega_k}{\partial r} = 0$ 
  - Rigid body rotation
  - No torque
- If  $\Omega_k(r)$  decreases outwards,  $\frac{\partial \Omega_k}{\partial r} < 0$ , and  $G(r) < 0$ ,
  - angular momentum goes from inner circles to outer circles.
  - The gas slowly spirals in!

## Roche Lobe Overflow

A star will fill its Roche lobe when it has a radius  $R_{*R}$  such that a sphere with this radius has the same volume as the Roche lobe.

## Roche Lobe Overflow

A star will fill its Roche lobe when it has a radius  $R_{*R}$  such that a sphere with this radius has the same volume as the Roche lobe. A numerical computation gives the value (Eggleton, 1983)

$$\frac{R_{*R}}{a} \approx \frac{0.49q^{2/3}}{0.6q^{2/3} + \ln(1 + q^{1/3})}$$

## Roche Lobe Overflow

A star will fill its Roche lobe when it has a radius  $R_{*R}$  such that a sphere with this radius has the same volume as the Roche lobe. A numerical computation gives the value (Eggleton, 1983)

$$\frac{R_{*R}}{a} \approx \frac{0.49q^{2/3}}{0.6q^{2/3} + \ln(1 + q^{1/3})}$$

For  $0.1 \leq q \leq 0.8$  there is the Paczynski approximation

$$\frac{R_{*R}}{a} \approx \frac{2}{3^{4/3} \left( \frac{q}{1+q} \right)^{1/3}}$$

$$\frac{R_{*R}}{a} \approx 0.462 \left( \frac{M_*}{M + M_*} \right)^{1/3}$$

## Self-sustainability of the Roche Overflow

Roche overflow  $\longrightarrow$  change  $q$   
 $\longrightarrow$  change  $a, \tau$   
 $\longrightarrow$  Roche lobe ?

## Self-sustainability of the Roche Overflow

Roche overflow  $\longrightarrow$  change  $q$   
 $\longrightarrow$  change  $a, \tau$   
 $\longrightarrow$  Roche lobe ?

If Roche lobe grows  $\longrightarrow$  overflow stops

## Self-sustainability of the Roche Overflow

Roche overflow  $\longrightarrow$  change  $q$   
 $\longrightarrow$  change  $a, \tau$   
 $\longrightarrow$  Roche lobe ?

If Roche lobe grows  $\longrightarrow$  overflow stops

If Roche lobe shrinks  $\longrightarrow$  overflow continues!

# Angular Momentum

Total Angular Momentum

$$J = \left( M r_{BH}^2 + M_* r_*^2 \right) \frac{2\pi}{\tau}$$



# Angular Momentum

Total Angular Momentum

$$J = \left( M r_{BH}^2 + M_* r_*^2 \right) \frac{2\pi}{\tau}$$

Using the center of mass definition,

# Angular Momentum

Total Angular Momentum

$$J = \left( Mr_{BH}^2 + M_* r_*^2 \right) \frac{2\pi}{\tau}$$

Using the center of mass definition,

$$J = MM_* \sqrt{\frac{Ga}{M + M_*}}$$

# Angular Momentum

Total Angular Momentum

$$J = \left( M r_{BH}^2 + M_* r_*^2 \right) \frac{2\pi}{\tau}$$

Using the center of mass definition,

$$J = M M_* \sqrt{\frac{G a}{M + M_*}}$$

$$a = \frac{J^2}{G} \frac{M + M_*}{M^2 M_*^2}$$

## Angular Momentum

Considering that all the mass lost by the star,  $\dot{M}_* < 0$ , is accreted by the BH,

$$\dot{M} + \dot{M}_* = 0$$

## Angular Momentum

Considering that all the mass lost by the star,  $\dot{M}_* < 0$ , is accreted by the BH,

$$\dot{M} + \dot{M}_* = 0$$

and differentiating the expression for the angular momentum,

## Angular Momentum

Considering that all the mass lost by the star,  $\dot{M}_* < 0$ , is accreted by the BH,

$$\dot{M} + \dot{M}_* = 0$$

and differentiating the expression for the angular momentum,

$$\frac{\dot{a}}{a} = \frac{2\dot{J}}{J} + \frac{2(-\dot{M}_*)}{M_*} \left[ 1 - \frac{M_*}{M} \right]$$

## Angular Momentum

$$\frac{\dot{a}}{a} = \frac{2\dot{J}}{J} + \frac{2(-\dot{M}_*)}{M_*} \left[ 1 - \frac{M_*}{M} \right]$$

For conservative systems:  $\dot{J} = 0$ , and since  $\dot{M}_* < 0$ , then

## Angular Momentum

$$\frac{\dot{a}}{a} = \frac{2\dot{J}}{J} + \frac{2(-\dot{M}_*)}{M_*} \left[ 1 - \frac{M_*}{M} \right]$$

For conservative systems:  $\dot{J} = 0$ , and since  $\dot{M}_* < 0$ , then

$$\frac{\dot{a}}{a} = \frac{2(-\dot{M}_*)}{M_*} \left[ 1 - \frac{M_*}{M} \right] > 0$$

for  $q < 1$ .



## Angular Momentum

$$\frac{\dot{a}}{a} = \frac{2\dot{J}}{J} + \frac{2(-\dot{M}_*)}{M_*} \left[ 1 - \frac{M_*}{M} \right]$$

For conservative systems:  $\dot{J} = 0$ , and since  $\dot{M}_* < 0$ , then

$$\frac{\dot{a}}{a} = \frac{2(-\dot{M}_*)}{M_*} \left[ 1 - \frac{M_*}{M} \right] > 0$$

for  $q < 1$ .

For conservative systems: BH grows putting more mass near the CM and the star moves in a wider orbit, increasing  $a$ , in order to conserve  $J$ .

## Self-sustainability of the Roche Overflow

$$\frac{R_{*R}}{a} = 0.462 \left( \frac{M_*}{M + M_*} \right)^{1/3}$$

## Self-sustainability of the Roche Overflow

$$\frac{R_{*R}}{a} = 0.462 \left( \frac{M_*}{M + M_*} \right)^{1/3}$$

$$\frac{\dot{R}_{*R}}{R_{*R}} = \frac{\dot{a}}{a} + \frac{1}{3} \frac{\dot{M}_*}{M_*}$$

## Self-sustainability of the Roche Overflow

$$\frac{R_{*R}}{a} = 0.462 \left( \frac{M_*}{M + M_*} \right)^{1/3}$$

$$\frac{\dot{R}_{*R}}{R_{*R}} = \frac{\dot{a}}{a} + \frac{1}{3} \frac{\dot{M}_*}{M_*}$$

$$\frac{\dot{R}_{*R}}{R_{*R}} = \frac{2\dot{J}}{J} + \frac{2(-\dot{M}_*)}{M_*} \left[ \frac{5}{6} - \frac{M_*}{M} \right]$$

## Self-sustainability of the Roche Overflow

$$\frac{\dot{R}_{*R}}{R_{*R}} = \frac{2\dot{J}}{J} + \frac{2(-\dot{M}_*)}{M_*} \left[ \frac{5}{6} - \frac{M_*}{M} \right]$$

## Self-sustainability of the Roche Overflow

$$\frac{\dot{R}_{*R}}{R_{*R}} = \frac{2\dot{J}}{J} + \frac{2(-\dot{M}_*)}{M_*} \left[ \frac{5}{6} - \frac{M_*}{M} \right]$$

- Conservative systems ( $\dot{J} = 0$ ) with  $q < \frac{5}{6}$ :  
Roche lobe of the star grows and the overflow stops.

## Self-sustainability of the Roche Overflow

$$\frac{\dot{R}_{*R}}{R_{*R}} = \frac{2\dot{J}}{J} + \frac{2(-\dot{M}_*)}{M_*} \left[ \frac{5}{6} - \frac{M_*}{M} \right]$$

## Self-sustainability of the Roche Overflow

$$\frac{\dot{R}_{*R}}{R_{*R}} = \frac{2\dot{J}}{J} + \frac{2(-\dot{M}_*)}{M_*} \left[ \frac{5}{6} - \frac{M_*}{M} \right]$$

- Conservative systems ( $\dot{J} = 0$ ) with  $q > \frac{5}{6}$ :  
Roche lobe of the star shrinks and the overflow continues.



## Self-sustainability of the Roche Overflow

$$\frac{\dot{R}_{*R}}{R_{*R}} = \frac{2\dot{J}}{J} + \frac{2(-\dot{M}_*)}{M_*} \left[ \frac{5}{6} - \frac{M_*}{M} \right]$$

## Self-sustainability of the Roche Overflow

$$\frac{\dot{R}_{*R}}{R_{*R}} = \frac{2\dot{J}}{J} + \frac{2(-\dot{M}_*)}{M_*} \left[ \frac{5}{6} - \frac{M_*}{M} \right]$$

- If the angular momentum diminishes ( $\dot{J} < 0$ ) it accentuates the diminution of the Roche lobe of the star.

## Self-sustainability of the Roche Overflow

$$\frac{\dot{R}_{*R}}{R_{*R}} = \frac{2\dot{J}}{J} + \frac{2(-\dot{M}_*)}{M_*} \left[ \frac{5}{6} - \frac{M_*}{M} \right]$$

- If the angular momentum diminishes ( $\dot{J} < 0$ ) it accentuates the diminution of the Roche lobe of the star.

The Roche overflow is rapid and violent but stops when  $q$  is smaller than  $\frac{5}{6}$ .

## Self-sustainability of the Roche Overflow

The transfer of mass continues if

- The star expands (stellar evolution). Here the Roche lobe must be large enough to accomodate the star, hence this must be a *long-period* system

## Self-sustainability of the Roche Overflow

The transfer of mass continues if

- The star expands (stellar evolution). Here the Roche lobe must be large enough to accomodate the star, hence this must be a *long-period* system
- The binary system loses angular momentum. There are many mechanisms for losing  $J$ :

## Self-sustainability of the Roche Overflow

The transfer of mass continues if

- The star expands (stellar evolution). Here the Roche lobe must be large enough to accomodate the star, hence this must be a *long-period* system
- The binary system loses angular momentum. There are many mechanisms for losing  $J$ :
  - Gravitational radiation

## Self-sustainability of the Roche Overflow

The transfer of mass continues if

- The star expands (stellar evolution). Here the Roche lobe must be large enough to accomodate the star, hence this must be a *long-period* system
- The binary system loses angular momentum. There are many mechanisms for losing  $J$ :
  - Gravitational radiation
  - Tidal forces on the star

## Self-sustainability of the Roche Overflow

The transfer of mass continues if

- The star expands (stellar evolution). Here the Roche lobe must be large enough to accomodate the star, hence this must be a *long-period* system
- The binary system loses angular momentum. There are many mechanisms for losing  $J$ :
  - Gravitational radiation
  - Tidal forces on the star
  - Wind (magnetically linked to the star)



# Outline for Part 2

## 1. Viscous Torques

### 1.1 Differential Rotation

### 1.2 Roche Lobe Overflow

## 2. Formation of the Accretion Disk

# Formation of the Accretion Disk

# Formation of the Accretion Disk

The transferred material has a high specific angular momentum.  
Therefore, accretion is not a direct process.

# Formation of the Accretion Disk

The transferred material has a high specific angular momentum.

Therefore, accretion is not a direct process.

As seen from the BH, the matter spreads as from a nozzle rotating around the center of mass.

# Formation of the Accretion Disk

The transferred material has a high specific angular momentum.


Therefore, accretion is not a direct process.

As seen from the BH, the matter spreads as from a nozzle rotating around the center of mass.

$v_{\parallel}$  : parallel component of the velocity with respect to the line of centers.

$v_{\perp}$  : perpendicular component of the velocity with respect to the line of centers.

## Formation of the Accretion Disk



# Formation of the Accretion Disk

$$v_{\parallel} \lesssim c_s$$

# Formation of the Accretion Disk

$$v_{\parallel} \lesssim c_s$$

$c_s$  : speed of sound in the envelope of the star.



## Formation of the Accretion Disk

$$v_{\parallel} \lesssim c_s$$

$c_s$  : speed of sound in the envelope of the star.

For normal stellar envelope temperatures, ( $< 10^5$  K);

## Formation of the Accretion Disk

$$v_{\parallel} \lesssim c_s$$

$c_s$  : speed of sound in the envelope of the star.

For normal stellar envelope temperatures, ( $< 10^5$  K);  $c_s \lesssim 10$  km/s

## Formation of the Accretion Disk

$$v_{\perp} \sim b\omega = \frac{2\pi b}{\tau}$$

## Formation of the Accretion Disk

$$v_{\perp} \sim b\omega = \frac{2\pi b}{\tau}$$

Using

$$b \approx a [0.500 - 0.227 \log_{10}(q)]$$

$$a = 3.5 \times 10^{10} \left( \frac{M}{M_{\odot}} \right)^{1/3} (1+q)^{1/3} \tau_{hr}^{2/3} [\text{cm}]$$

## Formation of the Accretion Disk

$$v_{\perp} \sim b\omega = \frac{2\pi b}{\tau}$$

Using

$$b \approx a [0.500 - 0.227 \log_{10}(q)]$$

$$a = 3.5 \times 10^{10} \left( \frac{M}{M_{\odot}} \right)^{1/3} (1+q)^{1/3} \tau_{hr}^{2/3} \text{ [cm]}$$

we get

$$v_{\perp} \gtrsim 305.6 \left( \frac{M}{M_{\odot}} \right)^{1/3} (1+q)^{1/3} \tau_{hr}^{-1/3} \text{ [km/s]}$$

## Formation of the Accretion Disk

$$v_{\parallel} \lesssim c_s \lesssim 10 \text{ km/s}$$

$$v_{\perp} \gtrsim 305.6 \left( \frac{M}{M_{\odot}} \right)^{1/3} (1+q)^{1/3} \tau_{hr}^{-1/3} [\text{km/s}]$$

## Formation of the Accretion Disk

$$v_{\parallel} \lesssim c_s \lesssim 10 \text{ km/s}$$

$$v_{\perp} \gtrsim 305.6 \left( \frac{M}{M_{\odot}} \right)^{1/3} (1+q)^{1/3} \tau_{hr}^{-1/3} [\text{km/s}]$$

$$v_{\perp} \gtrsim 104.2 \left( \frac{M}{M_{\odot}} \right)^{1/3} (1+q)^{1/3} \tau_{days}^{-1/3} [\text{km/s}]$$

## Formation of the Accretion Disk

$$v_{\parallel} \lesssim c_s \lesssim 10 \text{ km/s}$$

$$v_{\perp} \gtrsim 305.6 \left( \frac{M}{M_{\odot}} \right)^{1/3} (1+q)^{1/3} \tau_{hr}^{-1/3} [\text{km/s}]$$

$$v_{\perp} \gtrsim 104.2 \left( \frac{M}{M_{\odot}} \right)^{1/3} (1+q)^{1/3} \tau_{days}^{-1/3} [\text{km/s}]$$

$$v_{\perp} \gg v_{\parallel}$$



# Formation of the Accretion Disk

- A particle (or parcel of gas) is released from rest at the point  $L_1$ , i.e. pressure effects are neglected.

# Formation of the Accretion Disk

- A particle (or parcel of gas) is released from rest at the point  $L_1$ , i.e. pressure effects are neglected.
- Because of the motion of the star, this particle is moving with velocity  $v_{\perp}$  as seen from the BH.

# Formation of the Accretion Disk

- A particle (or parcel of gas) is released from rest at the point  $L_1$ , i.e. pressure effects are neglected.
- Because of the motion of the star, this particle is moving with velocity  $v_{\perp}$  as seen from the BH.
- The particle will describe an elliptical trajectory around the BH but the presence of the star will make this ellipse to precess slowly.

# Formation of the Accretion Disk

- In the stream of accreting particles, their orbits will intersect, resulting in dissipation of energy via collisions.

# Formation of the Accretion Disk

- In the stream of accreting particles, their orbits will intersect, resulting in dissipation of energy via collisions.
- However, angular momentum will be conserved in this collisions.

# Formation of the Accretion Disk

- In the stream of accreting particles, their orbits will intersect, resulting in dissipation of energy via collisions.
- However, angular momentum will be conserved in this collisions.
- Hence, particles will go into the trajectories with minimum energy for a given angular momentum,

# Formation of the Accretion Disk

- In the stream of accreting particles, their orbits will intersect, resulting in dissipation of energy via collisions.
- However, angular momentum will be conserved in this collisions.
- Hence, particles will go into the trajectories with minimum energy for a given angular momentum, i.e. circular orbits!

# Formation of the Accretion Disk

- In the stream of accreting particles, their orbits will intersect, resulting in dissipation of energy via collisions.
- However, angular momentum will be conserved in this collisions.
- Hence, particles will go into the trajectories with minimum energy for a given angular momentum, i.e. circular orbits!
- This process is called *circularization*.



# Formation of the Accretion Disk

- In the stream of accreting particles, their orbits will intersect, resulting in dissipation of energy via collisions.
- However, angular momentum will be conserved in this collisions.
- Hence, particles will go into the trajectories with minimum energy for a given angular momentum, i.e. circular orbits!
- This process is called *circularization*.
- The radius of the resulting circular orbit is called *circularization radius*,  $r_{\text{circ}}$ .

# Formation of the Accretion Disk

$\Omega_k$ : Keplerian angular velocity in the circular orbit

# Formation of the Accretion Disk

$\Omega_k$ : Keplerian angular velocity in the circular orbit

$v_\phi$ : Tangential velocity in the circular orbit

# Formation of the Accretion Disk

$\Omega_k$ : Keplerian angular velocity in the circular orbit

$v_\phi$ : Tangential velocity in the circular orbit

$$r_{\text{circ}} \Omega_k^2(r_{\text{circ}}) = \frac{v_\phi^2(r_{\text{circ}})}{r_{\text{circ}}} = \frac{GM}{r_{\text{circ}}^2}$$

# Formation of the Accretion Disk

$\Omega_k$ : Keplerian angular velocity in the circular orbit

$v_\phi$ : Tangential velocity in the circular orbit

$$r_{\text{circ}} \Omega_k^2(r_{\text{circ}}) = \frac{v_\phi^2(r_{\text{circ}})}{r_{\text{circ}}} = \frac{GM}{r_{\text{circ}}^2}$$

$$v_\phi(r_{\text{circ}}) = \sqrt{\frac{GM}{r_{\text{circ}}}}$$

# Formation of the Accretion Disk

Conservation of angular momentum

# Formation of the Accretion Disk

Conservation of angular momentum

$$r_{\text{circ}} v_{\phi} = b^2 \omega$$

# Formation of the Accretion Disk

Conservation of angular momentum

$$r_{\text{circ}} v_{\phi} = b^2 \omega$$

$$\frac{r_{\text{circ}}}{a} = \frac{4\pi^2}{GM\tau} a^3 \left( \frac{b}{a} \right)^4$$



# Formation of the Accretion Disk

Conservation of angular momentum

$$r_{\text{circ}} v_{\phi} = b^2 \omega$$

$$\frac{r_{\text{circ}}}{a} = \frac{4\pi^2}{GM\tau} a^3 \left( \frac{b}{a} \right)^4$$

and using Kepler's third law,

$$\frac{r_{\text{circ}}}{a} = (1 + q) \left( \frac{b}{a} \right)^4$$

# Formation of the Accretion Disk

Conservation of angular momentum

$$r_{\text{circ}} v_{\phi} = b^2 \omega$$

$$\frac{r_{\text{circ}}}{a} = \frac{4\pi^2}{GM\tau} a^3 \left( \frac{b}{a} \right)^4$$

and using Kepler's third law,

$$\frac{r_{\text{circ}}}{a} = (1 + q) \left( \frac{b}{a} \right)^4$$

$$\frac{r_{\text{circ}}}{a} = (1 + q) [0.500 - 0.227 \log_{10} q]^4$$

## Formation of the Accretion Disk

$$r_{\text{circ}} = (1 + q)^{4/3} [0.500 - 0.227 \log_{10} q]^4 \left( \frac{M}{M_{\odot}} \right)^{1/3} \tau_{\text{days}}^{2/3} [R_{\odot}]$$

## Formation of the Accretion Disk

- The stream of gas moves in a ring with  $r = r_{\text{circ}}$ .

## Formation of the Accretion Disk

- The stream of gas moves in a ring with  $r = r_{\text{circ}}$ .
- In the ring the particles have collisions, shocks, viscous dissipation and other processes that transform some of the potential energy into heat (producing radiation).

## Formation of the Accretion Disk

- The stream of gas moves in a ring with  $r = r_{circ}$ .
- In the ring the particles have collisions, shocks, viscous dissipation and other processes that transform some of the potential energy into heat (producing radiation).
- However, this release of energy needs the losing of angular momentum.

# Formation of the Accretion Disk

- The stream of gas moves in a ring with  $r = r_{\text{circ}}$ .
- In the ring the particles have collisions, shocks, viscous dissipation and other processes that transform some of the potential energy into heat (producing radiation).
- However, this release of energy needs the losing of angular momentum.
- In the absence of external torques, the only possible process is a *transfer of angular momentum* from inner regions outwards by internal torques.

# Formation of the Accretion Disk

- The stream of gas moves in a ring with  $r = r_{\text{circ}}$ .
- In the ring the particles have collisions, shocks, viscous dissipation and other processes that transform some of the potential energy into heat (producing radiation).
- However, this release of energy needs the losing of angular momentum.
- In the absence of external torques, the only possible process is a *transfer of angular momentum* from inner regions outwards by internal torques.
- The redistribution of angular momentum makes particles in the outer parts move outwards (gaining angular momentum) and the particles in the inner parts spiral inwards.



## Formation of the Accretion Disk

- The accretion disk will extend from  $r_{in} \geq r_{ISCO}$  up to  $r_{out} \leq b$

## Formation of the Accretion Disk

- The accretion disk will extend from  $r_{in} \geq r_{ISCO}$  up to  $r_{out} \leq b$
- Viscous torques may be modeled using different processes:

## Formation of the Accretion Disk

- The accretion disk will extend from  $r_{in} \geq r_{ISCO}$  up to  $r_{out} \leq b$
- Viscous torques may be modeled using different processes:
  - Viscous torques due to differential rotation in the accretion disk (produced by the thermal motion of the fluid molecules). This is a local mechanism for angular momentum transport

# Formation of the Accretion Disk

- The accretion disk will extend from  $r_{in} \geq r_{ISCO}$  up to  $r_{out} \leq b$
- Viscous torques may be modeled using different processes:
  - Viscous torques due to differential rotation in the accretion disk (produced by the thermal motion of the fluid molecules). This is a local mechanism for angular momentum transport
  - Magnetic loops that couple fluid elements located at macroscopic distances across the disk. This is a non-local mechanism for angular momentum transport.

# Formation of the Accretion Disk

- The accretion disk will extend from  $r_{in} \geq r_{ISCO}$  up to  $r_{out} \leq b$
- Viscous torques may be modeled using different processes:
  - Viscous torques due to differential rotation in the accretion disk (produced by the thermal motion of the fluid molecules). This is a local mechanism for angular momentum transport
  - Magnetic loops that couple fluid elements located at macroscopic distances across the disk. This is a non-local mechanism for angular momentum transport.
  - Turbulence in the fluid may be an origin of angular momentum transport. Turbulence may be produced by mechanisms as Thermally driven convection, Pure hydrodynamic instabilities or Magnetohydrodynamic (MHD) turbulence.

# Formation of the Accretion Disk

- The accretion disk will extend from  $r_{in} \geq r_{ISCO}$  up to  $r_{out} \leq b$
- Viscous torques may be modeled using different processes:
  - Viscous torques due to differential rotation in the accretion disk (produced by the thermal motion of the fluid molecules). This is a local mechanism for angular momentum transport
  - Magnetic loops that couple fluid elements located at macroscopic distances across the disk. This is a non-local mechanism for angular momentum transport.
  - Turbulence in the fluid may be an origin of angular momentum transport. Turbulence may be produced by mechanisms as Thermally driven convection, Pure hydrodynamic instabilities or Magnetohydrodynamic (MHD) turbulence.
- However, this release of energy needs the losing of angular momentum.

# Formation of the Accretion Disk


- The accretion disk will extend from  $r_{in} \geq r_{ISCO}$  up to  $r_{out} \leq b$
- Viscous torques may be modeled using different processes:
  - Viscous torques due to differential rotation in the accretion disk (produced by the thermal motion of the fluid molecules). This is a local mechanism for angular momentum transport
  - Magnetic loops that couple fluid elements located at macroscopic distances across the disk. This is a non-local mechanism for angular momentum transport.
  - Turbulence in the fluid may be an origin of angular momentum transport. Turbulence may be produced by mechanisms as Thermally driven convection, Pure hydrodynamic instabilities or Magnetohydrodynamic (MHD) turbulence.
- However, this release of energy needs the losing of angular momentum.
- In the absence of external torques, the only possible process is *a transfer of angular momentum* from inner regions outwards

# Formation of the Accretion Disk

- The accretion disk will extend from  $r_{in} \geq r_{ISCO}$  up to  $r_{out} \leq b$
- Viscous torques may be modeled using different processes:
  - Viscous torques due to differential rotation in the accretion disk (produced by the thermal motion of the fluid molecules). This is a local mechanism for angular momentum transport
  - Magnetic loops that couple fluid elements located at macroscopic distances across the disk. This is a non-local mechanism for angular momentum transport.
  - Turbulence in the fluid may be an origin of angular momentum transport. Turbulence may be produced by mechanisms as Thermally driven convection, Pure hydrodynamic instabilities or Magnetohydrodynamic (MHD) turbulence.
- However, this release of energy needs the losing of angular momentum.
- In the absence of external torques, the only possible process is *a transfer of angular momentum* from inner regions outwards



# Temperature of the gas in the accretion structure



Next Lecture

## 10. Accretion Disks. Detailed Description