



BLACK HOLES

OBSERVATORIO
ASTRONÓMICO
NACIONAL

Classical Black Holes

11. Novikov-Thorne Thin Disks

Edward Larrañaga

Outline for Part 1

- 1. Novikov-Thorne Thin Disk
 - 1.1 Thin Disk General Description
 - 1.2 Kerr Spacetime
- 2. Transfer Function and Spectrum for Thin Disks

Novikov-Thorne Thin Disks

Novikov-Thorne Thin Disks

- Geometrically thin and optically thick accretion disk.

Novikov-Thorne Thin Disks

- Geometrically thin and optically thick accretion disk.
- Has four parameters: BH mass, BH spin, mass accretion rate and viscosity parameter.

Novikov-Thorne Thin Disks

- The spacetime is stationary, axisymmetric, asymptotically flat, and reflection- symmetric with respect to the equatorial plane.

Novikov-Thorne Thin Disks

- The spacetime is stationary, axisymmetric, asymptotically flat, and reflection-symmetric with respect to the equatorial plane.
- The accretion disk is non-self-gravitating; that is, the impact of the disk's mass on the background metric is ignored.

Novikov-Thorne Thin Disks

- The spacetime is stationary, axisymmetric, asymptotically flat, and reflection-symmetric with respect to the equatorial plane.
- The accretion disk is non-self-gravitating; that is, the impact of the disk's mass on the background metric is ignored.
- The accretion disk is in the equatorial plane; that is, the disk is perpendicular to the black hole spin.

Novikov-Thorne Thin Disks

- The spacetime is stationary, axisymmetric, asymptotically flat, and reflection-symmetric with respect to the equatorial plane.
- The accretion disk is non-self-gravitating; that is, the impact of the disk's mass on the background metric is ignored.
- The accretion disk is in the equatorial plane; that is, the disk is perpendicular to the black hole spin.
- The inner edge of the disk is at the ISCO radius.

Modeling Viscosity




Fig. 6.1 *Top panel* Accretion disk luminosity in Eddington units versus time for 766 spectra of LMC X-3. The shaded region does not satisfy the thin disk selection criterion $L/L_{\text{Edd}} < 0.3$, as well as the data below the dotted line, which marks $L/L_{\text{Edd}} = 0.05$. *Bottom panel* fitted value of the inner disk radius of the 411 spectra in the *top panel* that can meet the thin disk selection criterion. See the text for more details. From [51]. © AAS. Reproduced with permission

Novikov-Thorne Thin Disks

- The accretion disk is geometrically thin, namely the disk opening angle is $h/r \ll 1$, where $h(r)$ is the semi-thickness of the disk at the radial coordinate r .

Novikov-Thorne Thin Disks

- The accretion disk is geometrically thin, namely the disk opening angle is $h/r \ll 1$, where $h(r)$ is the semi-thickness of the disk at the radial coordinate r .
- We suppose to average over time scales Δt that are short enough to assume that the spacetime is stationary (for instance, the mass accreted by the central object does not appreciable change the background metric) and large enough to neglect possible inhomogeneities in the accretion fluid.

Novikov-Thorne Thin Disks

- The accretion disk is geometrically thin, namely the disk opening angle is $h/r \ll 1$, where $h(r)$ is the semi-thickness of the disk at the radial coordinate r .
- We suppose to average over time scales Δt that are short enough to assume that the spacetime is stationary (for instance, the mass accreted by the central object does not appreciable change the background metric) and large enough to neglect possible inhomogeneities in the accretion fluid.

Any physical quantity $\Psi(t, r, \theta, \phi)$ will be averaged over t and ϕ ,

$$\Psi(r, \theta) = \frac{1}{2\pi\Delta t} \int_0^{\Delta t} \int_0^{2\pi} \Psi(t, r, \theta, \phi) d\phi dt$$

Novikov-Thorne Thin Disks

- The particles of the gas follow nearly-geodesic circular orbits in the equatorial plane.

Novikov-Thorne Thin Disks

- The particles of the gas follow nearly-geodesic circular orbits in the equatorial plane.
- Radial heat transport is ignored, and energy and angular momentum are radiated from the disk surface.

Novikov-Thorne Thin Disks

- The particles of the gas follow nearly-geodesic circular orbits in the equatorial plane.
- Radial heat transport is ignored, and energy and angular momentum are radiated from the disk surface.
This assumption requires a thin accretion disk.

Novikov-Thorne Thin Disks

- The particles of the gas follow nearly-geodesic circular orbits in the equatorial plane.
- Radial heat transport is ignored, and energy and angular momentum are radiated from the disk surface.
This assumption requires a thin accretion disk.
- Magnetic fields are ignored.

Novikov-Thorne Thin Disks

- Energy and angular momentum from the disk's surface are only carried away by photons with wavelength $\lambda \ll M$.

Novikov-Thorne Thin Disks

- Energy and angular momentum from the disk's surface are only carried away by photons with wavelength $\lambda \ll M$.
This assumption let us neglect possible coherent superpositions of radiation reaction in nearby regions of the disk.

Novikov-Thorne Thin Disks

- Energy and angular momentum from the disk's surface are only carried away by photons with wavelength $\lambda \ll M$.
This assumption let us neglect possible coherent superpositions of radiation reaction in nearby regions of the disk.
- The effect of energy and angular momentum transport by photons emitted from the disk and returning to the disk due to strong light bending in the vicinity of the black hole (returning radiation) is neglected.

Thin Disk General Description

Thin Disk General Description

- The time-averaged radial structure of the Novikov-Thorn disk is obtained from the conservation laws (energy, angular momentum and rest mass)

Thin Disk General Description

- The time-averaged radial structure of the Novikov-Thorn disk is obtained from the conservation laws (energy, angular momentum and rest mass)
- The description will be independent of the specific properties of the accretion.

Mass Accretion Rate

Conservation of the rest mass:

Mass Accretion Rate

Conservation of the rest mass:

$$\nabla_\mu(\rho u^\mu) = 0$$

Mass Accretion Rate

Conservation of the rest mass:

$$\nabla_\mu(\rho u^\mu) = 0$$

ρ : Time-averaged rest mass density

u^μ : Time-averaged 4-velocity of the fluid

Thin Disk General Description

Integrating over the 3-volume of the disk (between r and $r + \Delta r$) and using Gauss' theorem,

Thin Disk General Description

Integrating over the 3-volume of the disk (between r and $r + \Delta r$) and using Gauss' theorem,

$$\dot{M} = -2\pi\sqrt{-\gamma}\Sigma u^r = \text{constant}$$

Thin Disk General Description

Integrating over the 3-volume of the disk (between r and $r + \Delta r$) and using Gauss' theorem,

$$\dot{M} = -2\pi\sqrt{-\gamma}\Sigma u^r = \text{constant}$$

$$\gamma = - \left(\frac{g_{t\phi}^2}{g_{\phi\phi}} - g_{tt} \right) g_{rr} g_{\phi\phi}$$

$$\Sigma(r) = \int_{-h}^h \rho dz$$

Energy Flux and Torque

From the conservation of energy,

$$\nabla_{\mu} T^{t\mu} = 0$$

and angular momentum,

$$\nabla_{\mu} T^{\phi\mu} = 0$$

Energy Flux and Torque

we obtain the time-averaged energy flux emitted from the surface of the disk,

$$\mathcal{F}(r) = \frac{\dot{M}}{4\pi M^2} F(r)$$

Energy Flux and Torque

we obtain the time-averaged energy flux emitted from the surface of the disk,

$$\mathcal{F}(r) = \frac{\dot{M}}{4\pi M^2} F(r)$$

and the time-averaged torque,

$$W_{\phi}^r(r) = \frac{\dot{M}}{2\pi M^2} \frac{(\Omega \ell_z - \varepsilon)}{\partial_r \Omega} F(r)$$

Energy Flux and Torque

we obtain the time-averaged energy flux emitted from the surface of the disk,

$$\mathcal{F}(r) = \frac{\dot{M}}{4\pi M^2} F(r)$$

and the time-averaged torque,

$$W_{\phi}^r(r) = \frac{\dot{M}}{2\pi M^2} \frac{(\Omega \ell_z - \varepsilon)}{\partial_r \Omega} F(r)$$

ε : specific energy

ℓ_z : specific axial component of the angular momentum

Ω : Angular velocity for equatorial circular geodesics

Energy Flux and Torque

$$F(r) = -\frac{\partial_r \Omega}{(\varepsilon - \Omega l_z)^2} \frac{M^2}{\sqrt{-\gamma}} \int_{r_{in}}^r [(\varepsilon - \Omega l_z) \partial_x l_z] dx$$

Energy Flux and Torque

$$F(r) = -\frac{\partial_r \Omega}{(\varepsilon - \Omega l_z)^2} \frac{M^2}{\sqrt{-\gamma}} \int_{r_{in}}^r [(\varepsilon - \Omega l_z) \partial_x l_z] dx$$

r_{in} : inner edge of the disk (presumably the ISCO radius)

Efficiency

$$L_{acc} = \eta \dot{M}$$

Efficiency

$$L_{acc} = \eta \dot{M}$$

η : Total efficiency

Efficiency

$$L_{acc} = \eta \dot{M}$$

η : Total efficiency

$$\eta = \eta_r + \eta_k$$

Efficiency

$$L_{acc} = \eta \dot{M}$$

η : Total efficiency

$$\eta = \eta_r + \eta_k$$

η_r : Radiative Efficiency (can be measured from bolometric luminosity)

Efficiency

$$L_{acc} = \eta \dot{M}$$

η : Total efficiency

$$\eta = \eta_r + \eta_k$$

η_r : Radiative Efficiency (can be measured from bolometric luminosity)

$$L_{bol} = \eta_r \dot{M}$$

Efficiency

$$L_{acc} = \eta \dot{M}$$

η : Total efficiency

$$\eta = \eta_r + \eta_k$$

Efficiency

$$L_{acc} = \eta \dot{M}$$

η : Total efficiency

$$\eta = \eta_r + \eta_k$$

η_k : gravitational energy transformed into kinetic energy of jets or outflows

Efficiency

$$L_{acc} = \eta \dot{M}$$

η : Total efficiency

$$\eta = \eta_r + \eta_k$$

η_k : gravitational energy transformed into kinetic energy of jets or outflows

For Novikov-Thorn disks: $\eta_k = 0$

Efficiency

$$L_{acc} = \eta \dot{M}$$

η : Total efficiency

$$\eta = \eta_r + \eta_k$$

η_k : gravitational energy transformed into kinetic energy of jets or outflows

For Novikov-Thorn disks: $\eta_k = 0$

$$\eta_{NT} = \eta_r$$

Efficiency

- The particles of the accreting gas slowly fall onto the BH

Efficiency

- The particles of the accreting gas slowly fall onto the BH
- When particles reach the ISCO radius, they quickly plunge onto the BH without emitting additional radiation

Efficiency

- The particles of the accreting gas slowly fall onto the BH
- When particles reach the ISCO radius, they quickly plunge onto the BH without emitting additional radiation
- A fraction of the emitted radiation is captured by the BH

Efficiency

- The particles of the accreting gas slowly fall onto the BH
- When particles reach the ISCO radius, they quickly plunge onto the BH without emitting additional radiation
- A fraction of the emitted radiation is captured by the BH

$$\eta_N T = 1 - \epsilon_{ISCO} - \zeta$$

Efficiency

- The particles of the accreting gas slowly fall onto the BH
- When particles reach the ISCO radius, they quickly plunge onto the BH without emitting additional radiation
- A fraction of the emitted radiation is captured by the BH

$$\eta_N T = 1 - \epsilon_{ISCO} - \zeta$$

ϵ_{ISCO} : Specific energy of a test particle at the ISCO

ζ : Fraction of energy captured by the BH

Efficiency

$$\zeta = \frac{1}{\dot{M}} \int_{r_{\text{ISCO}}}^{\infty} \left[\int_0^{\pi/2} \int_0^{2\pi} c\Upsilon(-n_t) \cos \theta \sin \theta d\phi d\theta \right] \mathcal{F}(r) 4r dr$$

Efficiency

$$\zeta = \frac{1}{\dot{M}} \int_{r_{\text{ISCO}}}^{\infty} \left[\int_0^{\pi/2} \int_0^{2\pi} c \Upsilon(-n_t) \cos \theta \sin \theta d\phi d\theta \right] \mathcal{F}(r) 4r dr$$

$$c = \begin{cases} 0 & \text{radiation escapes} \\ 1 & \text{radiation is captured} \end{cases}$$

Efficiency

$$\zeta = \frac{1}{\dot{M}} \int_{r_{\text{ISCO}}}^{\infty} \left[\int_0^{\pi/2} \int_0^{2\pi} c\Upsilon(-n_t) \cos \theta \sin \theta d\phi d\theta \right] \mathcal{F}(r) 4r dr$$

Efficiency

$$\zeta = \frac{1}{\dot{M}} \int_{r_{\text{ISCO}}}^{\infty} \left[\int_0^{\pi/2} \int_0^{2\pi} c\Upsilon(-n_t) \cos \theta \sin \theta d\phi d\theta \right] \mathcal{F}(r) 4r dr$$

Υ : takes into account possible angular dependence of the emission

$$\begin{cases} \Upsilon = 1 & \text{isotropic emission} \\ \Upsilon \propto 1 + 2 \cos \theta & \text{limb-darkened emission} \\ \dots & \end{cases}$$

Efficiency

$$\zeta = \frac{1}{\dot{M}} \int_{r_{\text{ISCO}}}^{\infty} \left[\int_0^{\pi/2} \int_0^{2\pi} c\Upsilon(-n_t) \cos \theta \sin \theta d\phi d\theta \right] \mathcal{F}(r) 4r dr$$

Efficiency

$$\zeta = \frac{1}{\dot{M}} \int_{r_{\text{ISCO}}}^{\infty} \left[\int_0^{\pi/2} \int_0^{2\pi} c \Upsilon(-n_t) \cos \theta \sin \theta d\phi d\theta \right] \mathcal{F}(r) 4r dr$$

n^μ : normalized photon 4-momentum

$$n^\mu = \frac{k^\mu}{k_{(t)}}$$

k^μ : photon 4-momentum

$k_{(t)}$: photon energy in the rest frame of the emitter

Kerr Spacetime

Kerr Spacetime

$$\varepsilon = \frac{r^{3/2} - 2Mr^{1/2} \pm aM^{1/2}}{r^{3/4} \sqrt{r^{3/2} - 3Mr^{1/2} \pm 2aM^{1/2}}}$$

Kerr Spacetime

$$\varepsilon = \frac{r^{3/2} - 2Mr^{1/2} \pm aM^{1/2}}{r^{3/4} \sqrt{r^{3/2} - 3Mr^{1/2} \pm 2aM^{1/2}}}$$

$$\ell_z = \pm \frac{M^{1/2} (r^2 \mp 2aM^{1/2}r^{1/2} + a^2)}{r^{3/4} \sqrt{r^{3/2} - 3Mr^{1/2} \pm 2aM^{1/2}}}$$

Kerr Spacetime

$$\varepsilon = \frac{r^{3/2} - 2Mr^{1/2} \pm aM^{1/2}}{r^{3/4} \sqrt{r^{3/2} - 3Mr^{1/2} \pm 2aM^{1/2}}}$$

$$\ell_z = \pm \frac{M^{1/2} (r^2 \mp 2aM^{1/2}r^{1/2} + a^2)}{r^{3/4} \sqrt{r^{3/2} - 3Mr^{1/2} \pm 2aM^{1/2}}}$$

$$\Omega_{\pm} = \pm \frac{M^{1/2}}{r^{3/2} \pm aM^{1/2}}$$

Kerr Spacetime

$$\varepsilon = \frac{r^{3/2} - 2Mr^{1/2} \pm aM^{1/2}}{r^{3/4} \sqrt{r^{3/2} - 3Mr^{1/2} \pm 2aM^{1/2}}}$$

$$\ell_z = \pm \frac{M^{1/2} (r^2 \mp 2aM^{1/2}r^{1/2} + a^2)}{r^{3/4} \sqrt{r^{3/2} - 3Mr^{1/2} \pm 2aM^{1/2}}}$$

$$\Omega_{\pm} = \pm \frac{M^{1/2}}{r^{3/2} \pm aM^{1/2}}$$

$$\sqrt{-\gamma} = r$$

Kerr Spacetime

$$F(x) = \frac{3}{2} \frac{1}{x^4(x^3 - 3x + 2a)} \left[x - x_0 - \frac{3}{2}a \ln \left(\frac{x}{x_0} \right) \right. \\ \left. - \frac{3(x_1 - a)^2}{x_1(x_1 - x_2)(x_1 - x_3)} \ln \left(\frac{x - x_1}{x_0 - x_1} \right) \right. \\ \left. - \frac{3(x_2 - a)^2}{x_2(x_2 - x_1)(x_2 - x_3)} \ln \left(\frac{x - x_2}{x_0 - x_2} \right) \right. \\ \left. - \frac{3(x_3 - a)^2}{x_3(x_3 - x_1)(x_3 - x_2)} \ln \left(\frac{x - x_3}{x_0 - x_3} \right) \right]$$

Radial Structure Evolution

$$x = \sqrt{\frac{r}{M}}$$

$$x_0 = \sqrt{\frac{r_{\text{ISCO}}}{M}}$$

x_1, x_2 and x_3 are the roots of $x^3 - 3x + 2a = 0$,

Radial Structure Evolution

$$x = \sqrt{\frac{r}{M}}$$

$$x_0 = \sqrt{\frac{r_{\text{ISCO}}}{M}}$$

x_1, x_2 and x_3 are the roots of $x^3 - 3x + 2a = 0$,

$$x_1 = 2 \cos \left(\frac{1}{3} \arccos a - \frac{\pi}{3} \right)$$

$$x_2 = 2 \cos \left(\frac{1}{3} \arccos a + \frac{\pi}{3} \right)$$

$$x_3 = -2 \cos \left(\frac{1}{3} \arccos a \right)$$

Radiative Efficiency in Kerr spactime

Neglecting the radiation captured by the BH, $\zeta = 0$ the NT efficiency is $\eta_{NT} = 1 - \epsilon_{ISCO}$

Radiative Efficiency in Kerr spactime

Neglecting the radiation captured by the BH, $\zeta = 0$ the NT efficiency is $\eta_{NT} = 1 - \epsilon_{ISCO}$
where ϵ_{ISCO} is

$$\epsilon = \frac{r^{3/2} - 2Mr^{1/2} \pm aM^{1/2}}{r^{3/4} \sqrt{r^{3/2} - 3Mr^{1/2} \pm 2aM^{1/2}}}$$

Radiative Efficiency in Kerr spacetime

Neglecting the radiation captured by the BH, $\zeta = 0$ the NT efficiency is $\eta_{NT} = 1 - \epsilon_{ISCO}$

where ϵ_{ISCO} is

$$\epsilon = \frac{r^{3/2} - 2Mr^{1/2} \pm aM^{1/2}}{r^{3/4} \sqrt{r^{3/2} - 3Mr^{1/2} \pm 2aM^{1/2}}}$$

evaluated at

$$r_{ISCO} = 3M + Z_2 \mp \sqrt{(3M - Z_1)(3M + Z_1 + 2Z_2)}$$

Radiative Efficiency in Kerr spacetime

Neglecting the radiation captured by the BH, $\zeta = 0$ the NT efficiency is $\eta_{NT} = 1 - \epsilon_{ISCO}$

where ϵ_{ISCO} is

$$\epsilon = \frac{r^{3/2} - 2Mr^{1/2} \pm aM^{1/2}}{r^{3/4} \sqrt{r^{3/2} - 3Mr^{1/2} \pm 2aM^{1/2}}}$$

evaluated at

$$r_{ISCO} = 3M + Z_2 \mp \sqrt{(3M - Z_1)(3M + Z_1 + 2Z_2)}$$

$$Z_1 = M + (M^2 - a^2)^{1/3} [(M + a)^{1/3} + (M - a)^{1/3}]$$

Radiative Efficiency in Kerr spacetime

Neglecting the radiation captured by the BH, $\zeta = 0$ the NT efficiency is $\eta_{NT} = 1 - \epsilon_{ISCO}$

where ϵ_{ISCO} is

$$\epsilon = \frac{r^{3/2} - 2Mr^{1/2} \pm aM^{1/2}}{r^{3/4} \sqrt{r^{3/2} - 3Mr^{1/2} \pm 2aM^{1/2}}}$$

evaluated at

$$r_{ISCO} = 3M + Z_2 \mp \sqrt{(3M - Z_1)(3M + Z_1 + 2Z_2)}$$

$$Z_1 = M + (M^2 - a^2)^{1/3} [(M + a)^{1/3} + (M - a)^{1/3}]$$

$$Z_2 = \sqrt{3a^2 + Z_1^2}$$

Radiative Efficiency in Kerr spactime

$$\eta_{NT} = \eta_{NT}(a)$$

Radiative Efficiency in Kerr spactime

$$\eta_{NT} = \eta_{NT}(a)$$

Particular cases:

Radiative Efficiency in Kerr spactime

$$\eta_{NT} = \eta_{NT}(a)$$

Particular cases:

$$\eta_{NT}(a = 0) = 0.057$$

Radiative Efficiency in Kerr spacetime

$$\eta_{NT} = \eta_{NT}(a)$$

Particular cases:

$$\eta_{NT}(a = 0) = 0.057$$

$$\eta_{NT}(a = 1) = 0.423 \text{ (co-rotating disk)}$$

Radiative Efficiency in Kerr spacetime

$$\eta_{NT} = \eta_{NT}(a)$$

Particular cases:

$$\eta_{NT}(a = 0) = 0.057$$

$$\eta_{NT}(a = 1) = 0.423 \text{ (co-rotating disk)}$$

$$\eta_{NT}(a = 1) = 0.038 \text{ (counter-rotating disk)}$$

Formation of the Accretion Disk




Fig. 6.2 Novikov-Thorne radiative efficiency $\eta_{NT} = 1 - E_{ISCO}$ as a function of the spin a in the Kerr metric. The *upper curve* is for corotating orbits, the *lower curve* for counterrotating orbits. The *dotted horizontal lines* correspond (from *top to bottom*) to the Novikov-Thorne radiative efficiency for $a/M = 1$ ($\eta_{NT} \approx 0.423$), $a/M = 0.998$ ($\eta_{NT} \approx 0.321$), $a/M = 0$ ($\eta_{NT} \approx 0.057$), and $a/M = 1$ in the case of a counterrotating disk ($\eta_{NT} \approx 0.038$)

Outline for Part 2

- 1. Novikov-Thorne Thin Disk
 - 1.1 Thin Disk General Description
 - 1.2 Kerr Spacetime
- 2. Transfer Function and Spectrum for Thin Disks

Transfer Function

Spectrum of the Thin Accretion Disk

The spectrum produced by a thin accretion disk is calculated by using the *transfer function* and considering two parts in the calculation,

Spectrum of the Thin Accretion Disk

The spectrum produced by a thin accretion disk is calculated by using the *transfer function* and considering two parts in the calculation,

- The first part corresponds to find the local spectrum (radiation at the surface of the accretion disk)

Spectrum of the Thin Accretion Disk

The spectrum produced by a thin accretion disk is calculated by using the *transfer function* and considering two parts in the calculation,

- The first part corresponds to find the local spectrum (radiation at the surface of the accretion disk)
- The second part corresponds to describe the propagation of the radiation from the disk to observer

Local Spectrum

- The local spectrum of the radiation depends only on the astrophysical model (it does not depend on the spacetime geometry)

Local Spectrum

- The local spectrum of the radiation depends only on the astrophysical model (it does not depend on the spacetime geometry)
- The local spectrum will be given by the specific intensity $I_e(\nu_e, r_e, \vartheta_e)$

Local Spectrum

- The local spectrum of the radiation depends only on the astrophysical model (it does not depend on the spacetime geometry)
- The local spectrum will be given by the specific intensity $I_e(\nu_e, r_e, \vartheta_e)$

ν_e : emitted photon frequency (in the rest frame of the disk)

r_e : emission radius

ϑ_e : emission angle with respect to the normal to the disk

Formation of the Accretion Disk




Fig. 6.3 The viewing angle i is the inclination angle of the disk, namely the angle between the axis normal to the disk and the line of sight of the distant observer; it is the same in Newtonian gravity and in general relativity. The emission angle ϑ_e is the angle, measured in the rest-frame of the gas and at a certain emission point, between the normal to the disk and the photon propagation direction. In Newtonian gravity, $i = \vartheta_e$, but in a curved spacetime the two angles are different in general

Observed Spectrum

The observed flux (in units of $\text{erg s}^{-1} \text{cm}^{-2} \text{Hz}^{-1}$)

$$F_{obs}(\nu_{obs}) = \int I_{obs}(\nu_{obs}, X, Y) d\tilde{\Omega}$$

Observed Spectrum

The observed flux (in units of $\text{erg s}^{-1} \text{cm}^{-2} \text{Hz}^{-1}$)

$$F_{obs}(\nu_{obs}) = \int I_{obs}(\nu_{obs}, X, Y) d\tilde{\Omega}$$

I_{obs} : Specific intensity detected by the observer

$d\tilde{\Omega} = \frac{dX dY}{D^2}$: Element of solid angle subtended by the image of the disk in the observer's sky

D : Distance observer-source

Observed Spectrum

The observed flux (in units of $\text{erg s}^{-1} \text{cm}^{-2} \text{Hz}^{-1}$)

$$F_{obs}(\nu_{obs}) = \int I_{obs}(\nu_{obs}, X, Y) d\tilde{\Omega}$$

I_{obs} : Specific intensity detected by the observer

$d\tilde{\Omega} = \frac{dXdY}{D^2}$: Element of solid angle subtended by the image of the disk in the observer's sky

D : Distance observer-source

$$F_{obs}(\nu_{obs}) = \int g^3 I_e(\nu_e, r_e, \vartheta_e) d\tilde{\Omega}$$

Observed Spectrum

The observed flux (in units of $\text{erg s}^{-1} \text{cm}^{-2} \text{Hz}^{-1}$)

$$F_{obs}(\nu_{obs}) = \int I_{obs}(\nu_{obs}, X, Y) d\tilde{\Omega}$$

I_{obs} : Specific intensity detected by the observer

$d\tilde{\Omega} = \frac{dX dY}{D^2}$: Element of solid angle subtended by the image of the disk in the observer's sky

D : Distance observer-source

$$F_{obs}(\nu_{obs}) = \int g^3 I_e(\nu_e, r_e, \vartheta_e) d\tilde{\Omega}$$

$g = \frac{\nu_{obs}}{\nu_e}$: redshift factor

Observed Spectrum

The observed flux (in units of $\text{erg s}^{-1} \text{cm}^{-2} \text{Hz}^{-1}$)

$$F_{obs}(\nu_{obs}) = \int I_{obs}(\nu_{obs}, X, Y) d\tilde{\Omega}$$

I_{obs} : Specific intensity detected by the observer

$d\tilde{\Omega} = \frac{dX dY}{D^2}$: Element of solid angle subtended by the image of the disk in the observer's sky

D : Distance observer-source

$$F_{obs}(\nu_{obs}) = \int g^3 I_e(\nu_e, r_e, \vartheta_e) d\tilde{\Omega}$$

$g = \frac{\nu_{obs}}{\nu_e}$: redshift factor

$$I_{obs} = g^3 I_e$$

The transfer function

- In order to describe the spacetime geometry information and the position of the observer, we introduce the *transfer function*, $f(g^*, r_e, i)$.

The transfer function

- In order to describe the spacetime geometry information and the position of the observer, we introduce the *transfer function*, $f(g^*, r_e, i)$.
- This function includes all the relativistic effects such as gravitational and Doppler redshift, light bending, etc.

The transfer function

- In order to describe the spacetime geometry information and the position of the observer, we introduce the *transfer function*, $f(g^*, r_e, i)$.
- This function includes all the relativistic effects such as gravitational and Doppler redshift, light bending, etc.

$$g^* = \frac{g - g_{min}}{g_{max} - g_{min}}$$

The transfer function

- In order to describe the spacetime geometry information and the position of the observer, we introduce the *transfer function*, $f(g^*, r_e, i)$.
- This function includes all the relativistic effects such as gravitational and Doppler redshift, light bending, etc.

$$g^* = \frac{g - g_{min}}{g_{max} - g_{min}}$$

$g_{max} = g_{max}(r_e, i)$: maximum value of the redshift factor

$g_{min} = g_{min}(r_e, i)$: minimum value of the redshift factor

The transfer function

$$F_{obs}(\nu_{obs}) = \int g^3 l_e(\nu_e, r_e, \vartheta_e) d\tilde{\Omega}$$

The transfer function

$$F_{obs}(\nu_{obs}) = \int g^3 l_e(\nu_e, r_e, \vartheta_e) d\tilde{\Omega}$$

$$F_{obs}(\nu_{obs}) = \frac{1}{D^2} \int \int g^3 l_e(\nu_e, r_e, \vartheta_e) dX dY$$

The transfer function

$$F_{obs}(\nu_{obs}) = \int g^3 l_e(\nu_e, r_e, \vartheta_e) d\tilde{\Omega}$$

$$F_{obs}(\nu_{obs}) = \frac{1}{D^2} \int \int g^3 l_e(\nu_e, r_e, \vartheta_e) dX dY$$

$$F_{obs}(\nu_{obs}) = \frac{1}{D^2} \int_{r_{ISCO}}^{\infty} \int_0^1 \frac{\pi r_e g^2}{\sqrt{g^* (1 - g^*)}} f(g^*, r_e, i) l_e(\nu_e, r_e, \vartheta_e) dg^* dr_e$$

The transfer function

$$F_{obs}(\nu_{obs}) = \int g^3 l_e(\nu_e, r_e, \vartheta_e) d\tilde{\Omega}$$

$$F_{obs}(\nu_{obs}) = \frac{1}{D^2} \int \int g^3 l_e(\nu_e, r_e, \vartheta_e) dX dY$$

$$F_{obs}(\nu_{obs}) = \frac{1}{D^2} \int_{r_{ISCO}}^{\infty} \int_0^1 \frac{\pi r_e g^2}{\sqrt{g^*(1-g^*)}} f(g^*, r_e, i) l_e(\nu_e, r_e, \vartheta_e) dg^* dr_e$$

$$f(g^*, r_e, i) = \frac{1}{\pi r_e} g \sqrt{g^*(1-g^*)} \left| \frac{\partial(X, Y)}{\partial(g^*, r_e)} \right|$$

Explicit form of the transfer function

- Consider a stationary and axisymmetric spacetime.

Explicit form of the transfer function

- Consider a stationary and axisymmetric spacetime.
- We need to relate the position of the received photon, (X, Y) with the emission point, r_e , and with the redshift factor g , and the emission angle, ϑ .

Explicit form of the transfer function

- Consider a stationary and axisymmetric spacetime.
- We need to relate the position of the received photon, (X, Y) with the emission point, r_e , and with the redshift factor g , and the emission angle, ϑ .
- This relation is obtained by evolving (numerically) the geodesic equations from the observation point to the emission point.

Explicit form of the transfer function. Redshift

- Numerically solving the geodesic equations give the position of the emitted photon in the accretion disk, r_e

Explicit form of the transfer function. Redshift

- Numerically solving the geodesic equations give the position of the emitted photon in the accretion disk, r_e
- The redshift factor is defined as the relation between the measured frequency (in the observer's frame) and the emitted frequency (in the rest frame of the gas)

Explicit form of the transfer function. Redshift

- Numerically solving the geodesic equations give the position of the emitted photon in the accretion disk, r_e
- The redshift factor is defined as the relation between the measured frequency (in the observer's frame) and the emitted frequency (in the rest frame of the gas)

$$g = \frac{\nu_{obs}}{\nu_e} = \frac{-u_{obs}^{\mu} k_{\mu}}{-u_e^{\nu} k_{\nu}}$$

Explicit form of the transfer function. Redshift

- Numerically solving the geodesic equations give the position of the emitted photon in the accretion disk, r_e
- The redshift factor is defined as the relation between the measured frequency (in the observer's frame) and the emitted frequency (in the rest frame of the gas)

$$g = \frac{\nu_{obs}}{\nu_e} = \frac{-u_{obs}^{\mu} k_{\mu}}{-u_e^{\nu} k_{\nu}}$$

$u_{obs}^{\mu} = u_e^t(1, 0, 0, \Omega)$: 4-velocity of the observer

$u_e^{\nu} = (1, 0, 0, 0)$: 4-velocity of the gas particles

Explicit form of the transfer function. Redshift

- Using the normalization condition $g_{\mu\nu}u_e^\mu u_e^\nu = -1$,

Explicit form of the transfer function. Redshift

- Using the normalization condition $g_{\mu\nu}u_e^\mu u_e^\nu = -1$,

$$g = \frac{\sqrt{-g_{tt} - 2g_{t\phi}\Omega - g_{\phi\phi}\Omega^2}}{1 - \lambda\Omega}$$

Explicit form of the transfer function. Redshift

- Using the normalization condition $g_{\mu\nu}u_e^\mu u_e^\nu = -1$,

$$g = \frac{\sqrt{-g_{tt} - 2g_{t\phi}\Omega - g_{\phi\phi}\Omega^2}}{1 - \lambda\Omega}$$

$$\lambda = -\frac{k_\phi}{k_t}$$

$k_t = -\varepsilon$: energy of the photon (constant of motion)

Explicit form of the transfer function. Emission angle

- Normal to the disk

$$n^\mu = \left(0, 0, \sqrt{g^{\theta\theta}}, 0 \right) \Big|_{r_e, \theta_e = \pi/2}$$

Explicit form of the transfer function. Emission angle

- Normal to the disk

$$n^\mu = \left(0, 0, \sqrt{g^{\theta\theta}}, 0 \right) \Big|_{r_e, \theta_e = \pi/2}$$

- The emission angle is given by

$$\cos \vartheta_e = \pm \frac{n^\mu k_\mu}{u_e^\nu k_\nu} \Big|_e$$

Explicit form of the transfer function. Emission angle

- Normal to the disk

$$n^\mu = \left(0, 0, \sqrt{g^{\theta\theta}}, 0 \right) \Big|_{r_e, \theta_e = \pi/2}$$

- The emission angle is given by

$$\cos \vartheta_e = \pm \frac{n^\mu k_\mu}{u_e^\nu k_\nu} \Big|_e$$

$$\cos \vartheta_e = \pm \sqrt{g^{\theta\theta}} \frac{\sqrt{-g_{tt} - 2g_{t\phi}\Omega - g_{\phi\phi}\Omega^2}}{1 - \lambda\Omega} \frac{k_\theta}{k_t}$$

Explicit form of the transfer function. Jacobian

- At the end of the numerical integration of the geodesic equations one has

Explicit form of the transfer function. Jacobian

- At the end of the numerical integration of the geodesic equations one has

$$r_e = r_e(X, Y)$$

$$g = g(X, Y)$$

$$\vartheta = \vartheta(X, Y)$$

Explicit form of the transfer function. Jacobian

- At the end of the numerical integration of the geodesic equations one has

$$r_e = r_e(X, Y)$$


$$g = g(X, Y)$$

$$\vartheta = \vartheta(X, Y)$$

- From the first two relations we may calculate the Jacobian as

$$\left| \frac{\partial(X, Y)}{\partial(g^*, r_e)} \right| = (g_{max} - g_{min}) \left| \frac{\partial X}{\partial g} \frac{\partial Y}{\partial r_e} - \frac{\partial X}{\partial r_e} \frac{\partial Y}{\partial g} \right|$$

Temperature of the gas in the accretion structure



Next Lecture

10. Accretion Disks. Detailed Description