



BLACK HOLES

OBSERVATORIO
ASTRONÓMICO
NACIONAL

Classical Black Holes

06. Astrophysics of Supermassive Black Holes

Edward Larrañaga

Outline for Part 1

1. Supermassive Black Holes in the Astrophysical Scenario

1.1 Supermassive Black Holes

1.2 Origin of Supermassive Black Holes

1.3 Intermediate-Mass Black Holes

2. Mathematical Description of the Collapse

2.1 Spherically Symmetric Collapse

2.2 Dust Collapse

2.3 Homogeneous Dust Collapse

2.4 Inhomogeneous Dust Collapse

Supermassive Black Holes

Astrophysical observations have discovered evidence of the existence of black holes, with masses in the range $M \sim 10^5 - 10^{10} M_{\odot}$.

Supermassive Black Holes

Astrophysical observations have discovered evidence of the existence of black holes, with masses in the range $M \sim 10^5 - 10^{10} M_{\odot}$.

Dynamical evidence supports that supermassive black holes are located at the center of almost all galaxies (there are some exceptions such as the galaxy A2261-BCG).

Supermassive Black Holes

Astrophysical observations have discovered evidence of the existence of black holes, with masses in the range $M \sim 10^5 - 10^{10} M_{\odot}$.

Dynamical evidence supports that supermassive black holes are located at the center of almost all galaxies (there are some exceptions such as the galaxy A2261-BCG).

In the Milky Way, the central black hole, called Sagittarius A* has a mass $\sim 4 \times 10^6 M_{\odot}$.

Origin of Supermassive Black Holes

There are many hypotheses on the origin of supermassive black holes. For example:

Origin of Supermassive Black Holes

There are many hypotheses on the origin of supermassive black holes. For example:

- Gravitational collapse of the first generation of stars (Population III).

Origin of Supermassive Black Holes

There are many hypotheses on the origin of supermassive black holes. For example:

- Gravitational collapse of the first generation of stars (Population III).
- Gas-dynamical instabilities.

Origin of Supermassive Black Holes

There are many hypotheses on the origin of supermassive black holes. For example:

- Gravitational collapse of the first generation of stars (Population III).
- Gas-dynamical instabilities.
- Stellar-dynamical stabilities.

Origin of Supermassive Black Holes

There are many hypotheses on the origin of supermassive black holes. For example:

- Gravitational collapse of the first generation of stars (Population III).
- Gas-dynamical instabilities.
- Stellar-dynamical stabilities.
- Dark Matter collapse

Origin of Supermassive Black Holes

There are many hypotheses on the origin of supermassive black holes. For example:

- Gravitational collapse of the first generation of stars (Population III).
- Gas-dynamical instabilities.
- Stellar-dynamical stabilities.
- Dark Matter collapse
- Primordial fluctuations

Origin of Supermassive Black Holes

The first hypothesis on the origin of supermassive black holes states that,

Population III stars are formed out of the collapse of zero-metallicity gas.

Origin of Supermassive Black Holes

The first hypothesis on the origin of supermassive black holes states that,

Population III stars are formed out of the collapse of zero-metallicity gas.

These stars are expected to be very massive, $M_* > 100M_{\odot}$

Origin of Supermassive Black Holes

The first hypothesis on the origin of supermassive black holes states that,

Population III stars are formed out of the collapse of zero-metallicity gas.

These stars are expected to be very massive, $M_* > 100M_{\odot}$

Then, they will die with a gravitational collapse, leaving behind black holes with masses between 40 and $1000M_{\odot}$

Origin of Supermassive Black Holes

The first hypothesis on the origin of supermassive black holes states that,

Population III stars are formed out of the collapse of zero-metallicity gas.

These stars are expected to be very massive, $M_* > 100M_{\odot}$

Then, they will die with a gravitational collapse, leaving behind black holes with masses between 40 and $1000M_{\odot}$

After that, Galaxies formed around these early black holes.

Origin of Supermassive Black Holes

The mechanism of gas-dynamical instabilities states that

Origin of Supermassive Black Holes

The mechanism of gas-dynamical instabilities states that

If fragmentation is inhibited in early massive cloud by, for example, turbulences, and cooling proceeds gradually, the gas will contract until rotation stops the collapse.

Origin of Supermassive Black Holes

The mechanism of gas-dynamical instabilities states that

If fragmentation is inhibited in early massive cloud by, for example, turbulences, and cooling proceeds gradually, the gas will contract until rotation stops the collapse.

Then, global dynamical instabilities (like bar-instabilities) can transport angular momentum outwards, allowing the core collapse to continue.

Origin of Supermassive Black Holes

The mechanism of gas-dynamical instabilities states that

If fragmentation is inhibited in early massive cloud by, for example, turbulences, and cooling proceeds gradually, the gas will contract until rotation stops the collapse.

Then, global dynamical instabilities (like bar-instabilities) can transport angular momentum outwards, allowing the core collapse to continue.

The gas accumulated in the center produce a very massive central object producing a black hole with a mass of $\sim 10^4 M_{\odot}$ or higher.

Origin of Supermassive Black Holes

The mechanism of stellar-dynamical stabilities states that

Origin of Supermassive Black Holes

The mechanism of stellar-dynamical stabilities states that

If fragmentation occurs, star formation in a collapsing cloud of gas produces a compact nuclear stellar cluster.

Origin of Supermassive Black Holes

The mechanism of stellar-dynamical stabilities states that

If fragmentation occurs, star formation in a collapsing cloud of gas produces a compact nuclear stellar cluster.

Collisions of stars in the cluster are frequent and they can produce a black hole of $\sim 10^2 - 10^4 M_{\odot}$

Origin of Supermassive Black Holes

The mechanism of Dark Matter collapse states that

Origin of Supermassive Black Holes

The mechanism of Dark Matter collapse states that

Dark matter collapse proceeds as described for a cloud of dust.

Origin of Supermassive Black Holes

The mechanism of Dark Matter collapse states that

Dark matter collapse proceeds as described for a cloud of dust.

No heating in the collapse until the rate of annihilation of dark particles becomes significant.

Origin of Supermassive Black Holes

The mechanism of Dark Matter collapse states that

Dark matter collapse proceeds as described for a cloud of dust.

No heating in the collapse until the rate of annihilation of dark particles becomes significant.

This process create an object called a *Dark Matter Star* for which the annihilation radiation supports the gravitational collapse.

Origin of Supermassive Black Holes

The temperature of this Dark Matter star is estimated between 4000 and 10000 K and its radius of approximately 10^{12} m.

Origin of Supermassive Black Holes

The temperature of this Dark Matter star is estimated between 4000 and 10000 K and its radius of approximately 10^{12} m.

The luminosity of these objects is about $10^6 L_{\odot}$ and their mass is estimated in $1000 M_{\odot}$

Origin of Supermassive Black Holes

The temperature of this Dark Matter star is estimated between 4000 and 10000 K and its radius of approximately 10^{12} m.

The luminosity of these objects is about $10^6 L_{\odot}$ and their mass is estimated in $1000 M_{\odot}$

When the dark matter is completely annihilated, the object will collapse into a massive black hole.

Origin of Supermassive Black Holes

The last hypothesis for the origin of supermassive black holes is from primordial fluctuations.

Origin of Supermassive Black Holes

The last hypothesis for the origin of supermassive black holes is from primordial fluctuations.

At the regions where the density fluctuations are large enough, the gravitational force may overcome the pressure, giving as a result a complete collapse.

Origin of Supermassive Black Holes

The last hypothesis for the origin of supermassive black holes is from primordial fluctuations.

At the regions where the density fluctuations are large enough, the gravitational force may overcome the pressure, giving as a result a complete collapse.

This process create primordial black holes with masses up to $\sim 1000M_{\odot}$.

Intermediate-Mass Black Holes

Intermediate-mass black holes have masses in the range
 $10^2 - 10^4 M_{\odot}$

Intermediate-Mass Black Holes

Intermediate-mass black holes have masses in the range
 $10^2 - 10^4 M_{\odot}$

They can be created from Population III stars collapse.

Intermediate-Mass Black Holes

Intermediate-mass black holes have masses in the range $10^2 - 10^4 M_{\odot}$

They can be created from Population III stars collapse.

They can also be created at the center of stellar clusters by merger of different kinds of objects, or they can be a supermassive black hole in the process of growth.

Intermediate-Mass Black Holes

Up to today, there are no dynamical measurements of their masses but only some luminosity observations that evidence black holes with masses beyond the stellar limit.

Intermediate-Mass Black Holes

Up to today, there are no dynamical measurements of their masses but only some luminosity observations that evidence black holes with masses beyond the stellar limit.

Thus, the conclusion from these observations is that there are intermediate-mass black holes or some stellar black holes having accretion luminosity exceeding the Eddington limit.

Chandrasekhar's Limit (1931)

$$\frac{M^{4/3}}{r^5} \propto \frac{GM^2}{r^5}$$

Chandrasekhar's Limit (1931)

$$\frac{M^{4/3}}{r^5} \propto \frac{GM^2}{r^5}$$

This relations is satisfied by the unique mass

$$M_C = 0.197 \left[\left(\frac{hc}{G} \right)^3 \frac{1}{m_p^2} \right] \frac{1}{\mu_e^2}$$

Chandrasekhar's Limit (1931)

$$\frac{M^{4/3}}{r^5} \propto \frac{GM^2}{r^5}$$

This relations is satisfied by the unique mass

$$M_C = 0.197 \left[\left(\frac{hc}{G} \right)^3 \frac{1}{m_p^2} \right] \frac{1}{\mu_e^2}$$

μ_e : mean molecular wight of the electrons

Chandrasekhar's Limit (1931)

$$\frac{M^{4/3}}{r^5} \propto \frac{GM^2}{r^5}$$

This relations is satisfied by the unique mass

$$M_C = 0.197 \left[\left(\frac{hc}{G} \right)^3 \frac{1}{m_p^2} \right] \frac{1}{\mu_e^2}$$

μ_e : mean molecular wight of the electrons

$$M_C \approx 1.4M_\odot$$

Mass of a completely degenerated star.

Neutron Stars. Degenerate Gas of Neutrons

Chandrasekhar (1939)

Neutron Stars. Degenerate Gas of Neutrons

Chandrasekhar (1939)

If the degenerate star attains sufficiently high densities, the protons and electrons will combine to form neutrons.

Neutron Stars. Degenerate Gas of Neutrons

Chandrasekhar (1939)

If the degenerate star attains sufficiently high densities, the protons and electrons will combine to form neutrons.

Baade and Zwicky (1934)

Neutron Stars. Degenerate Gas of Neutrons

Chandrasekhar (1939)

If the degenerate star attains sufficiently high densities, the protons and electrons will combine to form neutrons.

Baade and Zwicky (1934)

"...supernovae represent the transitions from ordinary stars into neutron stars which in their final stages consist of extremely closely packed neutrons."

Neutron Stars. Degenerate Gas of Neutrons

Oppenheimer and Volkoff (1939)

Oppenheimer and Snyder (1939)

Neutron Stars. Degenerate Gas of Neutrons

Oppenheimer and Volkoff (1939)

Oppenheimer and Snyder (1939)

“...when all thermonuclear sources of energy are exhausted a sufficiently heavy star will collapse. This contraction will continue indefinitely till the radius of the star approaches asymptotically its gravitational radius. Light from the surface of the star will be progressively reddened and can escape over a progressively narrower range of angles till eventually the star tends to close itself off from any communication with a distant observer”.

Black Holes

If the collapsing core is too massive to be supported by the degenerate pressure of neutrons, there is no known mechanism capable of finding a new equilibrium configuration, and the body should undergo a complete collapse.

In this case, the final product is a black hole.

Outline for Part 2

1. Supermassive Black Holes in the Astrophysical Scenario

1.1 Supermassive Black Holes

1.2 Origin of Supermassive Black Holes

1.3 Intermediate-Mass Black Holes

2. Mathematical Description of the Collapse

2.1 Spherically Symmetric Collapse

2.2 Dust Collapse

2.3 Homogeneous Dust Collapse

2.4 Inhomogeneous Dust Collapse

Spherically Symmetric Collapse

- Spherical Symmetry:

$$ds^2 = -e^{2\alpha(t,r)} dt^2 + e^{2\beta(t,r)} dr^2 + R^2(t,r) d\Omega^2$$

Spherically Symmetric Collapse

- Spherical Symmetry:

$$ds^2 = -e^{2\alpha(t,r)} dt^2 + e^{2\beta(t,r)} dr^2 + R^2(t,r) d\Omega^2$$

- Suppose that the collapsing body can be described as a perfect fluid.

Then, we consider that coordinates t and r are attached to every collapsing particle (co-moving coordinates).

Spherically Symmetric Collapse

- In the co-moving frame, the 4-velocity of the fluid is just

$$u^\mu = (e^{-\alpha}, 0, 0, 0)$$

and thus $u^2 = -1$.

Spherically Symmetric Collapse

- In the co-moving frame, the 4-velocity of the fluid is just

$$u^\mu = (e^{-\alpha}, 0, 0, 0)$$

and thus $u^2 = -1$.

- The energy-momentum tensor in the co-moving frame is

$$T^\mu_\nu = \text{diag}(\rho, P, P, P)$$

Spherically Symmetric Collapse

The Einstein equations for this metric give

$$\begin{aligned} G_t^t = 8\pi T_t^t &\Rightarrow \frac{F'}{R^2 R'} = 8\pi\rho \\ G_r^r = 8\pi T_r^r &\Rightarrow \frac{F}{R^2 \dot{R}} = -8\pi P \\ G_r^t = 0 &\Rightarrow \dot{R}' - \dot{R}\alpha' - \dot{\beta}R' = 0 \end{aligned}$$

Spherically Symmetric Collapse

The Einstein equations for this metric give

$$\begin{aligned} G_t^t = 8\pi T_t^t &\Rightarrow \frac{F'}{R^2 R'} = 8\pi\rho \\ G_r^r = 8\pi T_r^r &\Rightarrow \frac{F}{R^2 \dot{R}} = -8\pi P \\ G_r^t = 0 &\Rightarrow \dot{R}' - \dot{R}\alpha' - \dot{\beta}R' = 0 \end{aligned}$$

$$F = R \left(1 - e^{-2\beta} R'^2 + e^{-2\alpha} \dot{R}^2 \right)$$

Misner-Sharp mass

Spherically Symmetric Collapse

The Misner-Sharp mass

$$F = R \left(1 - e^{-2\beta} R'^2 + e^{-2\alpha} \dot{R}^2 \right)$$

is defined by the relation

$$1 - \frac{F}{R} = g_{\mu\nu} (\partial^\mu R) (\partial^\nu R)$$

Spherically Symmetric Collapse

The Misner-Sharp mass

$$F = R \left(1 - e^{-2\beta} R'^2 + e^{-2\alpha} \dot{R}^2 \right)$$

is defined by the relation

$$1 - \frac{F}{R} = g_{\mu\nu} (\partial^\mu R) (\partial^\nu R)$$

* Note that $n^\mu = \partial^\mu R$ is normal to the surfaces $R = \text{constant}$.
Thus, when $1 - \frac{F}{R} = 0$, the corresponding surface is null.

Spherically Symmetric Collapse

From (t, t) -component of the Field equations we obtain the Misner-Sharp mass as

$$G_t^t = 8\pi T_t^t \Rightarrow \frac{F'}{R^2 R'} = 8\pi\rho$$

Spherically Symmetric Collapse

From (t, t) -component of the Field equations we obtain the Misner-Sharp mass as

$$G_t^t = 8\pi T_t^t \Rightarrow \frac{F'}{R^2 R'} = 8\pi\rho$$

$$F(r) = \int_0^r F' d\tilde{r} = 8\pi \int_0^r \rho R^2 R' d\tilde{r} = 2M(r)$$

Spherically Symmetric Collapse

Finally, the conservation of the energy-momentum tensor gives the equation

$$\nabla_{\mu} T^{\mu}_{\nu} = 0 \Rightarrow \alpha' = -\frac{p'}{\rho + p}$$

Dust Collapse

Dust Collapse

Dust is characterized by $P = 0$.

Dust Collapse

Dust is characterized by $P = 0$.

$$\begin{aligned} G_t^t = 8\pi T_t^t &\Rightarrow \frac{F'}{R^2 R'} = 8\pi\rho \\ G_r^r = 8\pi T_r^r &\Rightarrow \frac{\dot{F}}{R^2 \dot{R}} = 0 \\ G_r^t = 0 &\Rightarrow \dot{R}' - \dot{R}\alpha' - \dot{\beta}R' = 0 \\ \nabla_\mu T_\nu^\mu = 0 &\Rightarrow \alpha' = 0 \end{aligned}$$

Dust Collapse

Dust is characterized by $P = 0$.

$$\begin{aligned}G_t^t = 8\pi T_t^t &\Rightarrow \frac{F'}{R^2 R'} = 8\pi\rho \\G_r^r = 8\pi T_r^r &\Rightarrow \frac{\dot{F}}{R^2 \dot{R}} = 0 \\G_r^t = 0 &\Rightarrow \dot{R}' - \dot{R}\alpha' - \dot{\beta}R' = 0 \\\nabla_\mu T_\nu^\mu = 0 &\Rightarrow \alpha' = 0\end{aligned}$$

Then

$$F = F(r)$$

$$\alpha = \alpha(t)$$

Dust Collapse

$F = F(r)$: No inflow or outflow of energy through spherical shells.
Thus, the exterior metric is Schwarzschild.

Dust Collapse

$F = F(r)$: No inflow or outflow of energy through spherical shells.
Thus, the exterior metric is Schwarzschild.

If the boundary of the cloud of dust is located at the co-moving coordinate $r = r_b$, we have $F(r_b) = 2M$ with M the Schwarzschild's mass of the exterior metric.

Dust Collapse

$F = F(r)$: No inflow or outflow of energy through spherical shells.
Thus, the exterior metric is Schwarzschild.

If the boundary of the cloud of dust is located at the co-moving coordinate $r = r_b$, we have $F(r_b) = 2M$ with M the Schwarzschild's mass of the exterior metric.

* If $P \neq 0$ the exterior metric must be a non-vacuum Vaidya spacetime.

Dust Collapse

$\alpha = \alpha(t)$: One can re-define the time coordinate such that

$$e^{\alpha(t)} dt \Rightarrow dt$$

and therefore $g_{tt} = -1$

Dust Collapse

$$\dot{R}' - \dot{R}\alpha' - \dot{\beta}R' = 0$$

Dust Collapse

$$\dot{R}' - \dot{R}\alpha' - \dot{\beta}R' = 0$$

$$\dot{R}' - \dot{\beta}R' = 0$$

Dust Collapse

$$\dot{R}' - \dot{R}\alpha' - \dot{\beta}R' = 0$$

$$\dot{R}' - \dot{\beta}R' = 0$$

$$\frac{1}{R'} \frac{d(R')}{dt} = \frac{d\beta}{dt}$$

Dust Collapse

$$\dot{R}' - \dot{R}\alpha' - \dot{\beta}R' = 0$$

$$\dot{R}' - \dot{\beta}R' = 0$$

$$\frac{1}{R'} \frac{d(R')}{dt} = \frac{d\beta}{dt}$$

$$\frac{d(\log R')}{dt} = \frac{d\beta}{dt}$$

Dust Collapse

$$\dot{R}' - \dot{R}\alpha' - \dot{\beta}R' = 0$$

$$\dot{R}' - \dot{\beta}R' = 0$$

$$\frac{1}{R'} \frac{d(R')}{dt} = \frac{d\beta}{dt}$$

$$\frac{d(\log R')}{dt} = \frac{d\beta}{dt}$$

$$R' = e^{\beta+h(r)}$$

Dust Collapse

$$R' = e^{\beta+h(r)}$$

Dust Collapse

$$R' = e^{\beta+h(r)}$$

Introducing $f(r) = e^{2h(r)} - 1$ we have

$$e^{2\beta} = \frac{R'^2}{1+f}$$

Dust Collapse

$$ds^2 = -dt^2 + \frac{R'^2(t, r)}{1 + f(r)} dr^2 + R^2(t, r) d\Omega^2$$

Lemaitre-Tolman-Bondi Metric

Dust Collapse

Kretschmann Scalar:

$$K = 12 \frac{F'^2}{R^4 R'^2} - 32 \frac{FF'}{R^5 R'} + 48 \frac{F^2}{R^6}$$

Dust Collapse

Kretschmann Scalar:

$$K = 12 \frac{F'^2}{R^4 R'^2} - 32 \frac{FF'}{R^5 R'} + 48 \frac{F^2}{R^6}$$

Divergence at $R = 0$

Dust Collapse

One can re-scale to set $R(t, r)$ to equal the co-moving radius r at the time $t = 0$, i.e. we impose $R(0, r) = r$ and introduce a scale factor a such that

$$R(t, r) = ra(t, r)$$

$$a(0, r) = 1$$

Dust Collapse

One can re-scale to set $R(t, r)$ to equal the co-moving radius r at the time $t = 0$, i.e. we impose $R(0, r) = r$ and introduce a scale factor a such that

$$R(t, r) = ra(t, r)$$

$$a(0, r) = 1$$

$$a(t_s, r) = 0 \text{ (at the time of the formation of the singularity, } t = t_s)$$

Dust Collapse

One can re-scale to set $R(t, r)$ to equal the co-moving radius r at the time $t = 0$, i.e. we impose $R(0, r) = r$ and introduce a scale factor a such that

$$R(t, r) = ra(t, r)$$

$$a(0, r) = 1$$

$$a(t_s, r) = 0 \text{ (at the time of the formation of the singularity, } t = t_s)$$

The collapse also impose the condition $\dot{a} < 0$

Dust Collapse

One of the field equations,

$$\frac{F'}{R^2 R'} = 8\pi\rho$$

implies that a regular value of the density at $t = 0$ is obtained if the Misner-Sharp mass has the form

$$F(r) = r^3 m(r)$$

where $m(r)$ is a (sufficiently) regular function in the region $[0, r_b]$.

Dust Collapse

We have $F' = 3r^2m + r^3m'$ and therefore

$$\frac{F'}{R^2R'} = 8\pi\rho$$

Dust Collapse

We have $F' = 3r^2m + r^3m'$ and therefore

$$\frac{F'}{R^2R'} = 8\pi\rho$$

$$\frac{3r^2m + r^3m'}{(ra)^2(a + ra')} = 8\pi\rho$$

Dust Collapse

We have $F' = 3r^2m + r^3m'$ and therefore

$$\frac{F'}{R^2R'} = 8\pi\rho$$

$$\frac{3r^2m + r^3m'}{(ra)^2(a + ra')} = 8\pi\rho$$

$$\frac{3m + rm'}{a^2(a + ra')} = 8\pi\rho$$

Dust Collapse

It is usual to consider $m(r)$ as a polynomial around $r = 0$,

$$m(r) = \sum_{k=0}^{\infty} m_k r^k$$

where m_k are constants

Dust Collapse

From the definition of the Misner-Sharp mass we have

$$\dot{R}^2 = \frac{F}{R} + f$$

Dust Collapse

From the definition of the Misner-Sharp mass we have

$$\dot{R}^2 = \frac{F}{R} + f$$

Using the proposed forms for F and R we obtain

$$r\dot{a} = -\sqrt{\frac{mr^2}{ra} + f}$$

Dust Collapse

From the definition of the Misner-Sharp mass we have

$$\dot{R}^2 = \frac{F}{R} + f$$

Using the proposed forms for F and R we obtain

$$r\dot{a} = -\sqrt{\frac{mr^2}{ra} + f}$$

$$\dot{a} = -\sqrt{\frac{m}{a} + \frac{f}{r^2}}$$

Dust Collapse

Evaluating at the time $t = 0$ we have $a(0, r) = 1$ and thus

$$\dot{a}(0, r) = -\sqrt{m + \frac{f}{r^2}}$$

Dust Collapse

Evaluating at the time $t = 0$ we have $a(0, r) = 1$ and thus

$$\dot{a}(0, r) = -\sqrt{m + \frac{f}{r^2}}$$

Note that the function f defines the initial velocity of the particles in the cloud.

Dust Collapse

Evaluating at the time $t = 0$ we have $a(0, r) = 1$ and thus

$$\dot{a}(0, r) = -\sqrt{m + \frac{f}{r^2}}$$

Note that the function f defines the initial velocity of the particles in the cloud. It is usual to write this function as an expansion around $r = 0$ as

$$f(r) = r^2 b(r)$$

with

$$b(r) = \sum_{k=0}^{\infty} b_k r^k$$

Homogeneous Dust Collapse

Homogeneous Dust Collapse

Simplest Model of Gravitational Collapse. Oppenheimer-Snyder
(1939)

Homogeneous Dust Collapse

Simplest Model of Gravitational Collapse. Oppenheimer-Snyder (1939)

Homogeneous: $\rho = \rho(t)$

Homogeneous Dust Collapse

Simplest Model of Gravitational Collapse. Oppenheimer-Snyder (1939)

Homogeneous: $\rho = \rho(t)$

This implies that

$$m' = 0 \longrightarrow m = m_0$$

Homogeneous Dust Collapse

Simplest Model of Gravitational Collapse. Oppenheimer-Snyder
(1939)

Homogeneous Dust Collapse

Simplest Model of Gravitational Collapse. Oppenheimer-Snyder (1939)

It also implies that

$$a' = 0 \longrightarrow \frac{f}{r^2} = \text{constant}$$

Homogeneous Dust Collapse

Simplest Model of Gravitational Collapse. Oppenheimer-Snyder (1939)

It also implies that

$$a' = 0 \longrightarrow \frac{f}{r^2} = \text{constant}$$
$$f(r) = r^2 b(r) = r^2 \text{constant}$$

Homogeneous Dust Collapse

Simplest Model of Gravitational Collapse. Oppenheimer-Snyder (1939)

It also implies that

$$a' = 0 \longrightarrow \frac{f}{r^2} = \text{constant}$$

$$f(r) = r^2 b(r) = r^2 \text{constant}$$

which is accomplished by choosing $b = b_0$

Homogeneous Dust Collapse

Simplest Model of Gravitational Collapse. Oppenheimer-Snyder (1939)

It also implies that

$$a' = 0 \longrightarrow \frac{f}{r^2} = \text{constant}$$

$$f(r) = r^2 b(r) = r^2 \text{constant}$$

which is accomplished by choosing $b = b_0$

Then

$$\dot{a} = -\sqrt{\frac{m_0}{a} + b_0}$$

Homogeneous Dust Collapse

The Lemaitre-Tolman-Bondi line element becomes

$$ds^2 = -dt^2 + \frac{R'^2(t, r)}{1 + f(r)} dr^2 + R^2(t, r) d\Omega^2$$

Homogeneous Dust Collapse

The Lemaitre-Tolman-Bondi line element becomes

$$ds^2 = -dt^2 + \frac{R'^2(t, r)}{1 + f(r)} dr^2 + R^2(t, r) d\Omega^2$$

$$ds^2 = -dt^2 + \frac{a^2(t)}{1 + r^2 b_0} dr^2 + r^2 a^2(t) d\Omega^2$$

Homogeneous Dust Collapse

The Lemaitre-Tolman-Bondi line element becomes

$$ds^2 = -dt^2 + \frac{R'^2(t, r)}{1 + f(r)} dr^2 + R^2(t, r) d\Omega^2$$

$$ds^2 = -dt^2 + \frac{a^2(t)}{1 + r^2 b_0} dr^2 + r^2 a^2(t) d\Omega^2$$

$$ds^2 = -dt^2 + a^2(t) \left[\frac{dr^2}{1 + b_0 r^2} + r^2 d\Omega^2 \right]$$

Homogeneous Dust Collapse

$$ds^2 = -dt^2 + a^2(t) \left[\frac{dr^2}{1 + b_0 r^2} + r^2 d\Omega^2 \right]$$

For $b_0 = 0$ this line element becomes

$$ds^2 = -dt^2 + a^2(t) [dr^2 + r^2 d\Omega^2]$$

and describes the counterpart of a flat spacetime, in which the collapse is marginally bound (i.e. the falling particles have vanishing velocity at infinity).

Homogeneous Dust Collapse

For $b_0 = 0$ the scale factor is

$$\dot{a} = -\sqrt{\frac{m_0}{a} + b_0}$$

Homogeneous Dust Collapse

For $b_0 = 0$ the scale factor is

$$\dot{a} = -\sqrt{\frac{m_0}{a} + b_0}$$

$$\frac{da}{dt} = -\sqrt{\frac{m_0}{a}}$$

Homogeneous Dust Collapse

For $b_0 = 0$ the scale factor is

$$\dot{a} = -\sqrt{\frac{m_0}{a} + b_0}$$

$$\frac{da}{dt} = -\sqrt{\frac{m_0}{a}}$$

$$\sqrt{a} da = -\sqrt{m_0} dt$$

Homogeneous Dust Collapse

For $b_0 = 0$ the scale factor is

$$\dot{a} = -\sqrt{\frac{m_0}{a} + b_0}$$

$$\frac{da}{dt} = -\sqrt{\frac{m_0}{a}}$$

$$\sqrt{a} da = -\sqrt{m_0} dt$$

$$\frac{2}{3}a^{3/2} - \frac{2}{3}a^{3/2}(0) = -\sqrt{m_0}t$$

Homogeneous Dust Collapse

The value of the initial condition was supposed as $a(0) = 1$. Thus

$$a(t) = \left[1 - \frac{\sqrt[3]{m_0}}{2} t \right]^{2/3}$$

Homogeneous Dust Collapse

The value of the initial condition was supposed as $a(0) = 1$. Thus

$$a(t) = \left[1 - \frac{3\sqrt{m_0}}{2} t \right]^{2/3}$$

Therefore, the singularity $R = 0$ occurs at the time t_s for which $a(t_s) = 0$. This is

$$t_s = \frac{2}{3\sqrt{m_0}}$$

Homogeneous Dust Collapse

The curve $t_H(r)$ describing the time at which the shell r crosses the horizon can be obtained from the condition

$$1 - \frac{F}{R} = 0$$

Homogeneous Dust Collapse

The curve $t_H(r)$ describing the time at which the shell r crosses the horizon can be obtained from the condition

$$1 - \frac{F}{R} = 0$$

$$1 - \frac{m_0 r^3}{ra(t_H)} = 0$$

Homogeneous Dust Collapse

The curve $t_H(r)$ describing the time at which the shell r crosses the horizon can be obtained from the condition

$$1 - \frac{F}{R} = 0$$

$$1 - \frac{m_0 r^3}{ra(t_H)} = 0$$

$$a(t_H) = m_0 r^2$$

Homogeneous Dust Collapse

$$a(t_H) = m_0 r^2$$

Homogeneous Dust Collapse

$$a(t_H) = m_0 r^2$$

For $b_0 = 0$ we have

$$\left[1 - \frac{3\sqrt{m_0}}{2} t_H \right]^{2/3} = m_0 r^2$$

Homogeneous Dust Collapse

$$a(t_H) = m_0 r^2$$


For $b_0 = 0$ we have

$$\left[1 - \frac{3\sqrt{m_0}}{2} t_H \right]^{2/3} = m_0 r^2$$

$$t_H(r) = \frac{2}{3\sqrt{m_0}} - \frac{2}{3} m_0 r^3$$


Gravitational Collapse of a Homogeneous Cloud of Dust

Eddington-Finkelstein Diagram



Gravitational Collapse of a Homogeneous Cloud of Dust

Eddington-Finkelstein Diagram



Inhomogeneous Dust Collapse

Inhomogeneous Dust Collapse

ρ depends on both t and r , and thus

Inhomogeneous Dust Collapse

ρ depends on both t and r , and thus

$$\rho = \rho(t, r)$$

$$m = m(r)$$

$$b = b(r)$$

$$a = a(t, r)$$

Inhomogeneous Dust Collapse

The simplest form for the function $m(r)$ is

$$m(r) = m_0 + m_2 r^2$$

Inhomogeneous Dust Collapse

The simplest form for the function $m(r)$ is

$$m(r) = m_0 + m_2 r^2$$

This function defines the density in terms of the two parameters m_0 and m_2 .

Inhomogeneous Dust Collapse

The simplest form for the function $m(r)$ is

$$m(r) = m_0 + m_2 r^2$$

This function defines the density in terms of the two parameters m_0 and m_2 .

The condition that ρ must be a decreasing function of r implies that $m_2 < 0$.

Inhomogeneous Dust Collapse

The simplest form for the function $m(r)$ is

$$m(r) = m_0 + m_2 r^2$$

This function defines the density in terms of the two parameters m_0 and m_2 .

The condition that ρ must be a decreasing function of r implies that $m_2 < 0$.

* $m_1 = 0$ in order to have a non divergent density.

Inhomogeneous Dust Collapse

The equation

$$\dot{R}^2 = \frac{F}{R} + f$$

gives this time

$$ra(\dot{t}, r) = -\sqrt{\frac{mr^2}{ra} + br^2}$$

Inhomogeneous Dust Collapse

The equation

$$\dot{R}^2 = \frac{F}{R} + f$$

gives this time

$$ra(\dot{t}, r) = -\sqrt{\frac{mr^2}{ra} + br^2}$$

$$a(\dot{t}, r) = -\sqrt{\frac{m}{a} + b}$$

Inhomogeneous Dust Collapse

In the marginally bound case, $b = 0$, it gives

Inhomogeneous Dust Collapse

In the marginally bound case, $b = 0$, it gives

$$\dot{a}(t, r) = -\sqrt{\frac{m(r)}{a}}$$

Inhomogeneous Dust Collapse

In the marginally bound case, $b = 0$, it gives

$$\dot{a}(t, r) = -\sqrt{\frac{m(r)}{a}}$$

$$\sqrt{a(t, r)} da = -\sqrt{m(r)} dt$$

Inhomogeneous Dust Collapse

In the marginally bound case, $b = 0$, it gives

$$\dot{a}(t, r) = -\sqrt{\frac{m(r)}{a}}$$

$$\sqrt{a(t, r)} da = -\sqrt{m(r)} dt$$

$$\frac{2}{3}a^{3/2}(t, r) - \frac{2}{3}a^{3/2}(0, r) = -\sqrt{m(r)}t$$

Inhomogeneous Dust Collapse

The value of the initial condition was supposed as $a(0) = 1$. Thus

$$a(t, r) = \left[1 - \frac{3\sqrt{m(r)}}{2} t \right]^{2/3}$$

Inhomogeneous Dust Collapse

The value of the initial condition was supposed as $a(0) = 1$. Thus

$$a(t, r) = \left[1 - \frac{3\sqrt{m(r)}}{2} t \right]^{2/3}$$

This equation shows how each shell of the distribution (each r) collapses with a different scale factor and with a different velocity.

Inhomogeneous Dust Collapse

The time to reach the singularity and the horizon are, respectively,

Inhomogeneous Dust Collapse

The time to reach the singularity and the horizon are, respectively,

$$t_s(r) = \frac{2}{3\sqrt{m(r)}}$$

Inhomogeneous Dust Collapse

The time to reach the singularity and the horizon are, respectively,

$$t_s(r) = \frac{2}{3\sqrt{m(r)}}$$

$$t_H(r) = \frac{2}{3\sqrt{m(r)}} - \frac{2}{3}m(r)r^3$$

Inhomogeneous Dust Collapse


Using $m(r) = m_0 + m_2 r^2$ gives the functions

$$t_s(r) = \frac{2}{3\sqrt{m_0 + m_2 r^2}}$$

$$t_H(r) = \frac{2}{3\sqrt{m_0 + m_2 r^2}} - \frac{2}{3}r^3 (m_0 + m_2 r^2)$$

Gravitational Collapse of a Inhomogeneous Cloud of Dust

Eddington-Finkelstein Diagram



Next Lecture

06. Black Holes Astrophysics