



BLACK HOLES

OBSERVATORIO
ASTRONÓMICO
NACIONAL

Classical Black Holes

09. Accretion in Binary Systems

Edward Larrañaga

Outline for Part 1

1. Roche Lobe Overflow

1.1 Binary Systems

1.2 Roche Lobe Overflow

2. Formation of the Accretion Disk

Accretion by Roche Lobe Overflow

Binary Systems

Binary System Geometry

- System of a star and a compact object (NS or BH)

Binary System Geometry

- System of a star and a compact object (NS or BH)
- Gravitational tidal forces deform the outer layers of the star into a tear drop shape

Binary System Geometry

- System of a star and a compact object (NS or BH)
- Gravitational tidal forces deform the outer layers of the star into a tear drop shape
- As the star rotates, tidal forces forced it to pulsate (change its form into the direction of the compact object)


Binary System Geometry

- System of a star and a compact object (NS or BH)
- Gravitational tidal forces deform the outer layers of the star into a tear drop shape
- As the star rotates, tidal forces forced it to pulsate (change its form into the direction of the compact object)
- Thermodynamical processes in the star's interior may damp (or amplify) the pulsations

Binary System Geometry

- System of a star and a compact object (NS or BH)
- Gravitational tidal forces deform the outer layers of the star into a tear drop shape
- As the star rotates, tidal forces forced it to pulsate (change its form into the direction of the compact object)
- Thermodynamical processes in the star's interior may damp (or amplify) the pulsations
- This process may dissipate orbital and rotational energy until the system reaches the state of minimum energy for constant angular momentum: *synchronous rotation in circular orbits*

Binary System Geometry



Binary System Geometry

Center of mass:

$$r_{BH} + r_* = a$$

$$Mr_{BH} = M_* r_*$$

Binary System Geometry

Center of mass:

$$r_{BH} + r_* = a$$

$$Mr_{BH} = M_* r_*$$

Angular velocity of the system:

$$\omega = \frac{v_*}{r_*} = \frac{v_{BH}}{r_{BH}}$$

Binary System Geometry

In the co-rotating frame (centered at the CM), gravitational force is balanced by centrifugal force.

Binary System Geometry

In the co-rotating frame (centered at the CM), gravitational force is balanced by centrifugal force.

$$\vec{f}_c = m\omega^2 r \hat{r}$$

Binary System Geometry

In the co-rotating frame (centered at the CM), gravitational force is balanced by centrifugal force.

$$\vec{f}_c = m\omega^2 r \hat{r}$$

$$U_c = -\frac{1}{2}m\omega^2 r^2$$

Binary System Geometry

In the co-rotating frame (centered at the CM), gravitational force is balanced by centrifugal force.

$$\vec{f}_c = m\omega^2 r \hat{r}$$

$$U_c = -\frac{1}{2}m\omega^2 r^2$$

$$U_g = -\frac{GMm}{s} - \frac{GM_* m}{s_*}$$

Binary System Geometry

Total potential:

Binary System Geometry

Total potential:

$$\Phi = \frac{U_g + U_c}{m}$$

Binary System Geometry

Total potential:

$$\Phi = \frac{U_g + U_c}{m}$$

$$\Phi = -\frac{GM}{s} - \frac{GM_*}{s_*} - \frac{1}{2}\omega^2 r^2$$

Binary System Geometry

Using:

Binary System Geometry

Using: Law of cosines

$$s^2 = r^2 + r_{BH}^2 + 2rr_{BH} \cos \theta$$

$$s_*^2 = r^2 + r_*^2 - 2rr_* \cos \theta$$

Binary System Geometry

Using: Law of cosines

$$s^2 = r^2 + r_{BH}^2 + 2rr_{BH} \cos \theta$$

$$s_*^2 = r^2 + r_*^2 - 2rr_* \cos \theta$$

Kepler's third law

$$\omega^2 = \left(\frac{2\pi}{\tau} \right)^2 = \frac{G(M + M_*)}{a^3} = \frac{G(M + M_*)}{(r_{BH} + r_*)^3}$$

Lagrange Points

Lagrange Points:

$$\begin{cases} \frac{\partial \Phi}{\partial x} = 0 \\ \frac{\partial \Phi}{\partial y} = 0 \end{cases}$$

Lagrange Points

The location of the point L_1 from M_{BH} and from M_* are given by the approximations

$$b \approx a [0.500 - 0.227 \log_{10}(q)]$$

$$b_* \approx a [0.500 + 0.227 \log_{10}(q)]$$

B. Peterson. *An Introduction to Active galactic Nuclei*. Cambridge University Press. (1997)

B. W. Carroll and D. A. Ostlie. *An Introduction to Modern Astrophysics*. Addison-Wesley (1996)

Equipotential Surfaces

- Equipotential surfaces are perpendicular to the effective force $\vec{f} = -m\vec{\nabla}\Phi$.


Equipotential Surfaces

- Equipotential surfaces are perpendicular to the effective force $\vec{f} = -m\vec{\nabla}\Phi$.
- Hydrostatic equilibrium guarantees that the pressure is constant along equipotentials (there is no parallel force to the equipotential surface)

Equipotential Surfaces

- Equipotential surfaces are perpendicular to the effective force $\vec{f} = -m\vec{\nabla}\Phi$.
- Hydrostatic equilibrium guarantees that the pressure is constant along equipotentials (there is no parallel force to the equipotential surface)
- Since pressure is due to the weight of the overlaying layer of material, density has a constant value along the equipotential.

Binary System Geometry



Binary System Geometry

Kepler's third law

$$\omega^2 = \left(\frac{2\pi}{\tau} \right)^2 = \frac{G(M + M_*)}{a^3}$$

Binary System Geometry

Kepler's third law

$$\omega^2 = \left(\frac{2\pi}{\tau} \right)^2 = \frac{G(M + M_*)}{a^3}$$

$$a = \sqrt[3]{\frac{GM_\odot}{4\pi^2}} \left(\frac{M}{M_\odot} \right)^{1/3} (1 + q)^{1/3} \tau^{2/3}$$

Binary System Geometry

Kepler's third law

$$\omega^2 = \left(\frac{2\pi}{\tau} \right)^2 = \frac{G(M + M_*)}{a^3}$$

$$a = \sqrt[3]{\frac{GM_\odot}{4\pi^2}} \left(\frac{M}{M_\odot} \right)^{1/3} (1 + q)^{1/3} \tau^{2/3}$$

$$q = \frac{M_*}{M}$$

Binary System Geometry

Binary System Geometry

$$a = 1.5 \times 10^8 \left(\frac{M}{M_{\odot}} \right)^{1/3} (1+q)^{1/3} \tau_{\text{seg}}^{2/3} [\text{cm}]$$

Binary System Geometry

$$a = 1.5 \times 10^8 \left(\frac{M}{M_{\odot}} \right)^{1/3} (1+q)^{1/3} \tau_{seg}^{2/3} [\text{cm}]$$

$$a = 2.3 \times 10^9 \left(\frac{M}{M_{\odot}} \right)^{1/3} (1+q)^{1/3} \tau_{min}^{2/3} [\text{cm}]$$

Binary System Geometry

$$a = 1.5 \times 10^8 \left(\frac{M}{M_{\odot}} \right)^{1/3} (1+q)^{1/3} \tau_{\text{seg}}^{2/3} [\text{cm}]$$

$$a = 2.3 \times 10^9 \left(\frac{M}{M_{\odot}} \right)^{1/3} (1+q)^{1/3} \tau_{\text{min}}^{2/3} [\text{cm}]$$

$$a = 3.5 \times 10^{10} \left(\frac{M}{M_{\odot}} \right)^{1/3} (1+q)^{1/3} \tau_{\text{hr}}^{2/3} [\text{cm}]$$

Binary System Geometry

$$a = 1.5 \times 10^8 \left(\frac{M}{M_{\odot}} \right)^{1/3} (1+q)^{1/3} \tau_{\text{seg}}^{2/3} [\text{cm}]$$

$$a = 2.3 \times 10^9 \left(\frac{M}{M_{\odot}} \right)^{1/3} (1+q)^{1/3} \tau_{\text{min}}^{2/3} [\text{cm}]$$

$$a = 3.5 \times 10^{10} \left(\frac{M}{M_{\odot}} \right)^{1/3} (1+q)^{1/3} \tau_{\text{hr}}^{2/3} [\text{cm}]$$

$$a = 2.9 \times 10^{11} \left(\frac{M}{M_{\odot}} \right)^{1/3} (1+q)^{1/3} \tau_{\text{days}}^{2/3} [\text{cm}]$$

Binary System Geometry

$$a = 1.5 \times 10^8 \left(\frac{M}{M_{\odot}} \right)^{1/3} (1+q)^{1/3} \tau_{\text{seg}}^{2/3} [\text{cm}]$$

$$a = 2.3 \times 10^9 \left(\frac{M}{M_{\odot}} \right)^{1/3} (1+q)^{1/3} \tau_{\text{min}}^{2/3} [\text{cm}]$$

$$a = 3.5 \times 10^{10} \left(\frac{M}{M_{\odot}} \right)^{1/3} (1+q)^{1/3} \tau_{\text{hr}}^{2/3} [\text{cm}]$$

$$a = 2.9 \times 10^{11} \left(\frac{M}{M_{\odot}} \right)^{1/3} (1+q)^{1/3} \tau_{\text{days}}^{2/3} [\text{cm}]$$

$$a = 1.5 \times 10^{13} \left(\frac{M}{M_{\odot}} \right)^{1/3} (1+q)^{1/3} \tau_{\text{yr}}^{2/3} [\text{cm}]$$

Roche Lobe Overflow

A star will fill its Roche lobe when it has a radius R_{*R} such that a sphere with this radius has the same volume as the Roche lobe.

Roche Lobe Overflow

A star will fill its Roche lobe when it has a radius R_{*R} such that a sphere with this radius has the same volume as the Roche lobe. A numerical computation gives the value (Eggleton, 1983)

$$\frac{R_{*R}}{a} \approx \frac{0.49q^{2/3}}{0.6q^{2/3} + \ln(1 + q^{1/3})}$$

Roche Lobe Overflow

A star will fill its Roche lobe when it has a radius R_{*R} such that a sphere with this radius has the same volume as the Roche lobe. A numerical computation gives the value (Eggleton, 1983)

$$\frac{R_{*R}}{a} \approx \frac{0.49q^{2/3}}{0.6q^{2/3} + \ln(1 + q^{1/3})}$$

For $0.1 \leq q \leq 0.8$ there is the Paczynski approximation

$$\frac{R_{*R}}{a} \approx \frac{2}{3^{4/3} \left(\frac{q}{1+q} \right)^{1/3}}$$

$$\frac{R_{*R}}{a} \approx 0.462 \left(\frac{M_*}{M + M_*} \right)^{1/3}$$

Self-sustainability of the Roche Overflow

Roche overflow \longrightarrow change q
 \longrightarrow change a, τ
 \longrightarrow Roche lobe ?

Self-sustainability of the Roche Overflow

Roche overflow \longrightarrow change q
 \longrightarrow change a, τ
 \longrightarrow Roche lobe ?

If Roche lobe grows \longrightarrow overflow stops

Self-sustainability of the Roche Overflow

Roche overflow \longrightarrow change q
 \longrightarrow change a, τ
 \longrightarrow Roche lobe ?

If Roche lobe grows \longrightarrow overflow stops

If Roche lobe shrinks \longrightarrow overflow continues!

Angular Momentum

Total Angular Momentum

$$J = \left(M r_{BH}^2 + M_* r_*^2 \right) \frac{2\pi}{\tau}$$

Angular Momentum

Total Angular Momentum

$$J = \left(M r_{BH}^2 + M_* r_*^2 \right) \frac{2\pi}{\tau}$$

Using the center of mass definition,

Angular Momentum

Total Angular Momentum

$$J = \left(M r_{BH}^2 + M_* r_*^2 \right) \frac{2\pi}{\tau}$$

Using the center of mass definition,

$$J = M M_* \sqrt{\frac{G a}{M + M_*}}$$

Angular Momentum

Total Angular Momentum

$$J = \left(M r_{BH}^2 + M_* r_*^2 \right) \frac{2\pi}{\tau}$$

Using the center of mass definition,

$$J = M M_* \sqrt{\frac{G a}{M + M_*}}$$

$$a = \frac{J^2}{G} \frac{M + M_*}{M^2 M_*^2}$$

Angular Momentum

Considering that all the mass lost by the star, $\dot{M}_* < 0$, is accreted by the BH,

$$\dot{M} + \dot{M}_* = 0$$

Angular Momentum

Considering that all the mass lost by the star, $\dot{M}_* < 0$, is accreted by the BH,

$$\dot{M} + \dot{M}_* = 0$$

and differentiating the expression for the angular momentum,

Angular Momentum

Considering that all the mass lost by the star, $\dot{M}_* < 0$, is accreted by the BH,

$$\dot{M} + \dot{M}_* = 0$$

and differentiating the expression for the angular momentum,

$$\frac{\dot{a}}{a} = \frac{2\dot{J}}{J} + \frac{2(-\dot{M}_*)}{M_*} \left[1 - \frac{M_*}{M} \right]$$

Angular Momentum

$$\frac{\dot{a}}{a} = \frac{2\dot{J}}{J} + \frac{2(-\dot{M}_*)}{M_*} \left[1 - \frac{M_*}{M} \right]$$

For conservative systems: $\dot{J} = 0$, and since $\dot{M}_* < 0$, then

Angular Momentum

$$\frac{\dot{a}}{a} = \frac{2\dot{J}}{J} + \frac{2(-\dot{M}_*)}{M_*} \left[1 - \frac{M_*}{M} \right]$$

For conservative systems: $\dot{J} = 0$, and since $\dot{M}_* < 0$, then

$$\frac{\dot{a}}{a} = \frac{2(-\dot{M}_*)}{M_*} \left[1 - \frac{M_*}{M} \right] > 0$$

for $q < 1$.

Angular Momentum

$$\frac{\dot{a}}{a} = \frac{2\dot{J}}{J} + \frac{2(-\dot{M}_*)}{M_*} \left[1 - \frac{M_*}{M} \right]$$

For conservative systems: $\dot{J} = 0$, and since $\dot{M}_* < 0$, then

$$\frac{\dot{a}}{a} = \frac{2(-\dot{M}_*)}{M_*} \left[1 - \frac{M_*}{M} \right] > 0$$

for $q < 1$.

For conservative systems: BH grows putting more mass near the CM and the star moves in a wider orbit, increasing a , in order to conserve J .

Self-sustainability of the Roche Overflow

$$\frac{R_{*R}}{a} = 0.462 \left(\frac{M_*}{M + M_*} \right)^{1/3}$$

Self-sustainability of the Roche Overflow

$$\frac{R_{*R}}{a} = 0.462 \left(\frac{M_*}{M + M_*} \right)^{1/3}$$

$$\frac{\dot{R}_{*R}}{R_{*R}} = \frac{\dot{a}}{a} + \frac{1}{3} \frac{\dot{M}_*}{M_*}$$

Self-sustainability of the Roche Overflow

$$\frac{R_{*R}}{a} = 0.462 \left(\frac{M_*}{M + M_*} \right)^{1/3}$$

$$\frac{\dot{R}_{*R}}{R_{*R}} = \frac{\dot{a}}{a} + \frac{1}{3} \frac{\dot{M}_*}{M_*}$$

$$\frac{\dot{R}_{*R}}{R_{*R}} = \frac{2\dot{J}}{J} + \frac{2(-\dot{M}_*)}{M_*} \left[\frac{5}{6} - \frac{M_*}{M} \right]$$

Self-sustainability of the Roche Overflow

$$\frac{\dot{R}_{*R}}{R_{*R}} = \frac{2\dot{J}}{J} + \frac{2(-\dot{M}_*)}{M_*} \left[\frac{5}{6} - \frac{M_*}{M} \right]$$

Self-sustainability of the Roche Overflow

$$\frac{\dot{R}_{*R}}{R_{*R}} = \frac{2\dot{J}}{J} + \frac{2(-\dot{M}_*)}{M_*} \left[\frac{5}{6} - \frac{M_*}{M} \right]$$

- Conservative systems ($\dot{J} = 0$) with $q < \frac{5}{6}$:
Roche lobe of the star grows and the overflow stops.

Self-sustainability of the Roche Overflow

$$\frac{\dot{R}_{*R}}{R_{*R}} = \frac{2\dot{J}}{J} + \frac{2(-\dot{M}_*)}{M_*} \left[\frac{5}{6} - \frac{M_*}{M} \right]$$

Self-sustainability of the Roche Overflow

$$\frac{\dot{R}_{*R}}{R_{*R}} = \frac{2\dot{J}}{J} + \frac{2(-\dot{M}_*)}{M_*} \left[\frac{5}{6} - \frac{M_*}{M} \right]$$

- Conservative systems ($\dot{J} = 0$) with $q > \frac{5}{6}$:
Roche lobe of the star shrinks and the overflow continues.

Self-sustainability of the Roche Overflow

$$\frac{\dot{R}_{*R}}{R_{*R}} = \frac{2\dot{J}}{J} + \frac{2(-\dot{M}_*)}{M_*} \left[\frac{5}{6} - \frac{M_*}{M} \right]$$

Self-sustainability of the Roche Overflow

$$\frac{\dot{R}_{*R}}{R_{*R}} = \frac{2\dot{J}}{J} + \frac{2(-\dot{M}_*)}{M_*} \left[\frac{5}{6} - \frac{M_*}{M} \right]$$

- If the angular momentum diminishes ($\dot{J} < 0$) it accentuates the diminution of the Roche lobe of the star.

Self-sustainability of the Roche Overflow

$$\frac{\dot{R}_{*R}}{R_{*R}} = \frac{2\dot{J}}{J} + \frac{2(-\dot{M}_*)}{M_*} \left[\frac{5}{6} - \frac{M_*}{M} \right]$$

- If the angular momentum diminishes ($\dot{J} < 0$) it accentuates the diminution of the Roche lobe of the star.

The Roche overflow is rapid and violent but stops when q is smaller than $\frac{5}{6}$.

Self-sustainability of the Roche Overflow

The transfer of mass continues if

- The star expands (stellar evolution). Here the Roche lobe must be large enough to accomodate the star, hence this must be a *long-period* system

Self-sustainability of the Roche Overflow

The transfer of mass continues if

- The star expands (stellar evolution). Here the Roche lobe must be large enough to accomodate the star, hence this must be a *long-period* system
- The binary system loses angular momentum. There are many mechanisms for losing J :

Self-sustainability of the Roche Overflow

The transfer of mass continues if

- The star expands (stellar evolution). Here the Roche lobe must be large enough to accomodate the star, hence this must be a *long-period* system
- The binary system loses angular momentum. There are many mechanisms for losing J :
 - Gravitational radiation

Self-sustainability of the Roche Overflow

The transfer of mass continues if

- The star expands (stellar evolution). Here the Roche lobe must be large enough to accomodate the star, hence this must be a *long-period* system
- The binary system loses angular momentum. There are many mechanisms for losing J :
 - Gravitational radiation
 - Tidal forces on the star

Self-sustainability of the Roche Overflow

The transfer of mass continues if

- The star expands (stellar evolution). Here the Roche lobe must be large enough to accomodate the star, hence this must be a *long-period* system
- The binary system loses angular momentum. There are many mechanisms for losing J :
 - Gravitational radiation
 - Tidal forces on the star
 - Wind (magnetically linked to the star)

Outline for Part 2

1. Roche Lobe Overflow

1.1 Binary Systems

1.2 Roche Lobe Overflow

2. Formation of the Accretion Disk

Formation of the Accretion Disk

Formation of the Accretion Disk

The transferred material has a high specific angular momentum.
Therefore, accretion is not a direct process.

Formation of the Accretion Disk

The transferred material has a high specific angular momentum.

Therefore, accretion is not a direct process.

As seen from the BH, the matter spreads as from a nozzle rotating around the center of mass.

Formation of the Accretion Disk

The transferred material has a high specific angular momentum.


Therefore, accretion is not a direct process.

As seen from the BH, the matter spreads as from a nozzle rotating around the center of mass.

v_{\parallel} : parallel component of the velocity with respect to the line of centers.

v_{\perp} : perpendicular component of the velocity with respect to the line of centers.

Formation of the Accretion Disk



Formation of the Accretion Disk

$$v_{\parallel} \lesssim c_s$$

Formation of the Accretion Disk

$$v_{\parallel} \lesssim c_s$$

c_s : speed of sound in the envelope of the star.

Formation of the Accretion Disk

$$v_{\parallel} \lesssim c_s$$

c_s : speed of sound in the envelope of the star.

For normal stellar envelope temperatures, ($< 10^5$ K);

Formation of the Accretion Disk

$$v_{\parallel} \lesssim c_s$$

c_s : speed of sound in the envelope of the star.

For normal stellar envelope temperatures, ($< 10^5$ K); $c_s \lesssim 10$ km/s

Formation of the Accretion Disk

$$v_{\perp} \sim b\omega = \frac{2\pi b}{\tau}$$

Formation of the Accretion Disk

$$v_{\perp} \sim b\omega = \frac{2\pi b}{\tau}$$

Using

$$b \approx a [0.500 - 0.227 \log_{10}(q)]$$

$$a = 3.5 \times 10^{10} \left(\frac{M}{M_{\odot}} \right)^{1/3} (1+q)^{1/3} \tau_{hr}^{2/3} [\text{cm}]$$

Formation of the Accretion Disk

$$v_{\perp} \sim b\omega = \frac{2\pi b}{\tau}$$

Using

$$b \approx a [0.500 - 0.227 \log_{10}(q)]$$

$$a = 3.5 \times 10^{10} \left(\frac{M}{M_{\odot}} \right)^{1/3} (1+q)^{1/3} \tau_{hr}^{2/3} [\text{cm}]$$

we get

$$v_{\perp} \gtrsim 305.6 \left(\frac{M}{M_{\odot}} \right)^{1/3} (1+q)^{1/3} \tau_{hr}^{-1/3} [\text{km/s}]$$

Formation of the Accretion Disk

$$v_{\parallel} \lesssim c_s \lesssim 10 \text{ km/s}$$

$$v_{\perp} \gtrsim 305.6 \left(\frac{M}{M_{\odot}} \right)^{1/3} (1+q)^{1/3} \tau_{hr}^{-1/3} [\text{km/s}]$$

Formation of the Accretion Disk

$$v_{\parallel} \lesssim c_s \lesssim 10 \text{ km/s}$$

$$v_{\perp} \gtrsim 305.6 \left(\frac{M}{M_{\odot}} \right)^{1/3} (1+q)^{1/3} \tau_{hr}^{-1/3} [\text{km/s}]$$

$$v_{\perp} \gtrsim 104.2 \left(\frac{M}{M_{\odot}} \right)^{1/3} (1+q)^{1/3} \tau_{days}^{-1/3} [\text{km/s}]$$

Formation of the Accretion Disk

$$v_{\parallel} \lesssim c_s \lesssim 10 \text{ km/s}$$

$$v_{\perp} \gtrsim 305.6 \left(\frac{M}{M_{\odot}} \right)^{1/3} (1+q)^{1/3} \tau_{hr}^{-1/3} [\text{km/s}]$$

$$v_{\perp} \gtrsim 104.2 \left(\frac{M}{M_{\odot}} \right)^{1/3} (1+q)^{1/3} \tau_{days}^{-1/3} [\text{km/s}]$$

$$v_{\perp} \gg v_{\parallel}$$

Formation of the Accretion Disk

- A particle (or parcel of gas) is released from rest at the point L_1 , i.e. pressure effects are neglected.

Formation of the Accretion Disk

- A particle (or parcel of gas) is released from rest at the point L_1 , i.e. pressure effects are neglected.
- Because of the motion of the star, this particle is moving with velocity v_{\perp} as seen from the BH.

Formation of the Accretion Disk

- A particle (or parcel of gas) is released from rest at the point L_1 , i.e. pressure effects are neglected.
- Because of the motion of the star, this particle is moving with velocity v_{\perp} as seen from the BH.
- The particle will describe an elliptical trajectory around the BH but the presence of the star will make this ellipse to precess slowly.

Formation of the Accretion Disk

- In the stream of accreting particles, their orbits will intersect, resulting in dissipation of energy via collisions.

Formation of the Accretion Disk

- In the stream of accreting particles, their orbits will intersect, resulting in dissipation of energy via collisions.
- However, angular momentum will be conserved in this collisions.

Formation of the Accretion Disk

- In the stream of accreting particles, their orbits will intersect, resulting in dissipation of energy via collisions.
- However, angular momentum will be conserved in this collisions.
- Hence, particles will go into the trajectories with minimum energy for a given angular momentum,

Formation of the Accretion Disk

- In the stream of accreting particles, their orbits will intersect, resulting in dissipation of energy via collisions.
- However, angular momentum will be conserved in this collisions.
- Hence, particles will go into the trajectories with minimum energy for a given angular momentum, i.e. circular orbits!

Formation of the Accretion Disk

- In the stream of accreting particles, their orbits will intersect, resulting in dissipation of energy via collisions.
- However, angular momentum will be conserved in this collisions.
- Hence, particles will go into the trajectories with minimum energy for a given angular momentum, i.e. circular orbits!
- This process is called *circularization*.

Formation of the Accretion Disk

- In the stream of accreting particles, their orbits will intersect, resulting in dissipation of energy via collisions.
- However, angular momentum will be conserved in this collisions.
- Hence, particles will go into the trajectories with minimum energy for a given angular momentum, i.e. circular orbits!
- This process is called *circularization*.
- The radius of the resulting circular orbit is called *circularization radius*, r_{circ} .

Formation of the Accretion Disk

Ω_k : Keplerian angular velocity in the circular orbit

Formation of the Accretion Disk

Ω_k : Keplerian angular velocity in the circular orbit

v_ϕ : Tangential velocity in the circular orbit

Formation of the Accretion Disk

Ω_k : Keplerian angular velocity in the circular orbit

v_ϕ : Tangential velocity in the circular orbit

$$r_{\text{circ}} \Omega_k^2(r_{\text{circ}}) = \frac{v_\phi^2(r_{\text{circ}})}{r_{\text{circ}}} = \frac{GM}{r_{\text{circ}}^2}$$

Formation of the Accretion Disk

Ω_k : Keplerian angular velocity in the circular orbit

v_ϕ : Tangential velocity in the circular orbit

$$r_{\text{circ}} \Omega_k^2(r_{\text{circ}}) = \frac{v_\phi^2(r_{\text{circ}})}{r_{\text{circ}}} = \frac{GM}{r_{\text{circ}}^2}$$

$$v_\phi(r_{\text{circ}}) = \sqrt{\frac{GM}{r_{\text{circ}}}}$$

Formation of the Accretion Disk

Conservation of angular momentum

Formation of the Accretion Disk

Conservation of angular momentum

$$r_{\text{circ}} v_{\phi} = b^2 \omega$$

Formation of the Accretion Disk

Conservation of angular momentum

$$r_{\text{circ}} v_{\phi} = b^2 \omega$$

$$\frac{r_{\text{circ}}}{a} = \frac{4\pi^2}{GM\tau} a^3 \left(\frac{b}{a} \right)^4$$

Formation of the Accretion Disk

Conservation of angular momentum

$$r_{\text{circ}} v_{\phi} = b^2 \omega$$

$$\frac{r_{\text{circ}}}{a} = \frac{4\pi^2}{GM\tau} a^3 \left(\frac{b}{a} \right)^4$$

and using Kepler's third law,

$$\frac{r_{\text{circ}}}{a} = (1 + q) \left(\frac{b}{a} \right)^4$$

Formation of the Accretion Disk

Conservation of angular momentum

$$r_{\text{circ}} v_{\phi} = b^2 \omega$$

$$\frac{r_{\text{circ}}}{a} = \frac{4\pi^2}{GM\tau} a^3 \left(\frac{b}{a} \right)^4$$

and using Kepler's third law,

$$\frac{r_{\text{circ}}}{a} = (1 + q) \left(\frac{b}{a} \right)^4$$

$$\frac{r_{\text{circ}}}{a} = (1 + q) [0.500 - 0.227 \log_{10} q]^4$$

Formation of the Accretion Disk

$$r_{\text{circ}} = (1 + q)^{4/3} [0.500 - 0.227 \log_{10} q]^4 \left(\frac{M}{M_{\odot}} \right)^{1/3} \tau_{\text{days}}^{2/3} [R_{\odot}]$$

Formation of the Accretion Disk

- The stream of gas moves in a ring with $r = r_{\text{circ}}$.

Formation of the Accretion Disk

- The stream of gas moves in a ring with $r = r_{\text{circ}}$.
- In the ring the particles have collisions, shocks, viscous dissipation and other processes that transform some of the potential energy into heat (producing radiation).

Formation of the Accretion Disk

- The stream of gas moves in a ring with $r = r_{circ}$.
- In the ring the particles have collisions, shocks, viscous dissipation and other processes that transform some of the potential energy into heat (producing radiation).
- However, this release of energy needs the losing of angular momentum.

Formation of the Accretion Disk

- The stream of gas moves in a ring with $r = r_{\text{circ}}$.
- In the ring the particles have collisions, shocks, viscous dissipation and other processes that transform some of the potential energy into heat (producing radiation).
- However, this release of energy needs the losing of angular momentum.
- In the absence of external torques, the only possible process is a *transfer of angular momentum* from inner regions outwards by internal torques.

Formation of the Accretion Disk

- The stream of gas moves in a ring with $r = r_{\text{circ}}$.
- In the ring the particles have collisions, shocks, viscous dissipation and other processes that transform some of the potential energy into heat (producing radiation).
- However, this release of energy needs the losing of angular momentum.
- In the absence of external torques, the only possible process is a *transfer of angular momentum* from inner regions outwards by internal torques.
- The redistribution of angular momentum makes particles in the outer parts move outwards (gaining angular momentum) and the particles in the inner parts spiral inwards.

Formation of the Accretion Disk

- The accretion disk will extend from $r_{in} \geq r_{ISCO}$ up to $r_{out} \leq b$

Formation of the Accretion Disk

- The accretion disk will extend from $r_{in} \geq r_{ISCO}$ up to $r_{out} \leq b$
- Viscous torques may be modeled using different processes:

Formation of the Accretion Disk

- The accretion disk will extend from $r_{in} \geq r_{ISCO}$ up to $r_{out} \leq b$
- Viscous torques may be modeled using different processes:
 - Viscous torques due to differential rotation in the accretion disk (produced by the thermal motion of the fluid molecules). This is a local mechanism for angular momentum transport

Formation of the Accretion Disk

- The accretion disk will extend from $r_{in} \geq r_{ISCO}$ up to $r_{out} \leq b$
- Viscous torques may be modeled using different processes:
 - Viscous torques due to differential rotation in the accretion disk (produced by the thermal motion of the fluid molecules). This is a local mechanism for angular momentum transport
 - Magnetic loops that couple fluid elements located at macroscopic distances across the disk. This is a non-local mechanism for angular momentum transport.

Formation of the Accretion Disk

- The accretion disk will extend from $r_{in} \geq r_{ISCO}$ up to $r_{out} \leq b$
- Viscous torques may be modeled using different processes:
 - Viscous torques due to differential rotation in the accretion disk (produced by the thermal motion of the fluid molecules). This is a local mechanism for angular momentum transport
 - Magnetic loops that couple fluid elements located at macroscopic distances across the disk. This is a non-local mechanism for angular momentum transport.
 - Turbulence in the fluid may be an origin of angular momentum transport. Turbulence may be produced by mechanisms as Thermally driven convection, Pure hydrodynamic instabilities or Magnetohydrodynamic (MHD) turbulence.

Formation of the Accretion Disk

- The accretion disk will extend from $r_{in} \geq r_{ISCO}$ up to $r_{out} \leq b$
- Viscous torques may be modeled using different processes:
 - Viscous torques due to differential rotation in the accretion disk (produced by the thermal motion of the fluid molecules). This is a local mechanism for angular momentum transport
 - Magnetic loops that couple fluid elements located at macroscopic distances across the disk. This is a non-local mechanism for angular momentum transport.
 - Turbulence in the fluid may be an origin of angular momentum transport. Turbulence may be produced by mechanisms as Thermally driven convection, Pure hydrodynamic instabilities or Magnetohydrodynamic (MHD) turbulence.
- However, this release of energy needs the losing of angular momentum.


Formation of the Accretion Disk

- The accretion disk will extend from $r_{in} \geq r_{ISCO}$ up to $r_{out} \leq b$
- Viscous torques may be modeled using different processes:
 - Viscous torques due to differential rotation in the accretion disk (produced by the thermal motion of the fluid molecules). This is a local mechanism for angular momentum transport
 - Magnetic loops that couple fluid elements located at macroscopic distances across the disk. This is a non-local mechanism for angular momentum transport.
 - Turbulence in the fluid may be an origin of angular momentum transport. Turbulence may be produced by mechanisms as Thermally driven convection, Pure hydrodynamic instabilities or Magnetohydrodynamic (MHD) turbulence.
- However, this release of energy needs the losing of angular momentum.
- In the absence of external torques, the only possible process is *a transfer of angular momentum* from inner regions outwards

Formation of the Accretion Disk

- The accretion disk will extend from $r_{in} \geq r_{ISCO}$ up to $r_{out} \leq b$
- Viscous torques may be modeled using different processes:
 - Viscous torques due to differential rotation in the accretion disk (produced by the thermal motion of the fluid molecules). This is a local mechanism for angular momentum transport
 - Magnetic loops that couple fluid elements located at macroscopic distances across the disk. This is a non-local mechanism for angular momentum transport.
 - Turbulence in the fluid may be an origin of angular momentum transport. Turbulence may be produced by mechanisms as Thermally driven convection, Pure hydrodynamic instabilities or Magnetohydrodynamic (MHD) turbulence.
- However, this release of energy needs the losing of angular momentum.
- In the absence of external torques, the only possible process is *a transfer of angular momentum* from inner regions outwards

Temperature of the gas in the accretion structure



Next Lecture

10. Accretion Disks. Detailed Description