



BLACK HOLES

OBSERVATORIO
ASTRONÓMICO
NACIONAL

Classical Black Holes

10. Viscous Torques

Edward Larrañaga

Outline for Part 1

1. Viscous Torques

1.1 Differential Rotation

1.2 Radial Structure Evolution

1.3 The α -Prescription for Viscosity

2. The *alpha*-Prescription for Viscosity

Viscous Torques

Accretion Disk

- If the only interaction between fluid elements is gravity, the angular momentum is conserved.

Accretion Disk

- If the only interaction between fluid elements is gravity, the angular momentum is conserved.
- In the description of the accretion disk it is needed a force responsible for the redistribution of angular momentum.

Differential Rotation

Differential Rotation

- Keplerian Velocity

$$\Omega = \Omega(r)$$

Differential Rotation

- Keplerian Velocity

$$\Omega = \Omega(r)$$

- Neighboring material at different radii moves with different velocity

Differential Rotation

- Keplerian Velocity

$$\Omega = \Omega(r)$$

- Neighboring material at different radii moves with different velocity
- The disk is not a solid body. It moves with *differential rotation*.

Differential Rotation

- Keplerian Velocity

$$\Omega = \Omega(r)$$

- Neighboring material at different radii moves with different velocity
- The disk is not a solid body. It moves with *differential rotation*.
- The thermal motion of the fluid molecules and the turbulent motion of the fluid produce *viscous stresses*.

Differential Rotation

- In our simple description we will consider only momentum transport in the radial direction, produced by the process known as *shear viscosity*.

Differential Rotation

- In our simple description we will consider only momentum transport in the radial direction, produced by the process known as *shear viscosity*.
- It appears when there are internal distortions (usually local stresses that are proportional to the local rate of strain)

Differential Rotation

- In our simple description we will consider only momentum transport in the radial direction, produced by the process known as *shear viscosity*.
- It appears when there are internal distortions (usually local stresses that are proportional to the local rate of strain)
- Although the following is the simplest description, it can be used to describe other mechanisms of angular momentum transport such as magnetic loops that couple fluid elements at macroscopic distances across the disk.

Modeling Viscosity

- λ : Typical scale in the accretion disk


Modeling Viscosity

- λ : Typical scale in the accretion disk
- \tilde{v} : typical speed in the accretion disk

Modeling Viscosity

- λ : Typical scale in the accretion disk
- \tilde{v} : typical speed in the accretion disk
- As a first model, consider a uniform gas moving only in the tangential direction with velocity $v_\phi(r)$.

Modeling Viscosity



Modeling Viscosity

The only non-vanishing component of the stress tensor is the ϕ -component of the force per unit surface of the $r = \text{constant}$ surface,

$$\sigma_{r\phi} = -\eta \left. \frac{\partial v_\phi}{\partial r} \right|_R$$

Modeling Viscosity

The only non-vanishing component of the stress tensor is the ϕ -component of the force per unit surface of the $r = \text{constant}$ surface,

$$\sigma_{r\phi} = -\eta \left. \frac{\partial v_\phi}{\partial r} \right|_R$$

η : dynamical viscosity

$$v_\phi(r) = \Omega(r)r$$

$\frac{\partial v_\phi}{\partial r}$: velocity gradient

Modeling Viscosity

$$\sigma_{r\phi} = -\eta R \left. \frac{\partial \Omega}{\partial r} \right|_R$$

ϕ -component of the force per unit surface of the $r = \text{constant}$ surface.

Modeling Viscosity

The dynamical viscosity is usually written in terms of the *kinematical viscosity*, ν , as $\eta = \rho\nu$

Modeling Viscosity

The dynamical viscosity is usually written in terms of the *kinematical viscosity*, ν , as $\eta = \rho\nu$

The kinematical viscosity is modeled as

$$\nu \sim \lambda \tilde{\nu}$$

Modeling Viscosity

The dynamical viscosity is usually written in terms of the *kinematical viscosity*, ν , as $\eta = \rho\nu$

The kinematical viscosity is modeled as

$$\nu \sim \lambda \tilde{v}$$

Molecular transport:

λ : mean free path

\tilde{v} : thermal speed

Modeling Viscosity

The dynamical viscosity is usually written in terms of the *kinematical viscosity*, ν , as $\eta = \rho\nu$

The kinematical viscosity is modeled as

$$\nu \sim \lambda \tilde{v}$$

Molecular transport:

λ : mean free path

\tilde{v} : thermal speed

Turbulent motion:

λ : spatial scale (or characteristic wavelength) of the turbulence

\tilde{v} : typical velocity of the eddies

Modeling Viscosity

$$\sigma_{r\phi} = -\rho\nu R \left. \frac{\partial\Omega}{\partial r} \right|_R$$

ϕ -component of the force per unit surface of the $r = \text{constant}$ surface.

Modeling Viscosity

$$\sigma_{r\phi} = -\rho\nu R \left. \frac{\partial\Omega}{\partial r} \right|_R$$

ϕ -component of the force per unit surface of the $r = \text{constant}$ surface.

The net torque ($\vec{R} \times \vec{F}$) is calculated by multiplying $\sigma_{r\phi}$ by the area of the $r = R = \text{constant}$ surface, $2\pi RH$.

Modeling Viscosity

$$\sigma_{r\phi} = -\rho\nu R \left. \frac{\partial\Omega}{\partial r} \right|_R$$

ϕ -component of the force per unit surface of the $r = \text{constant}$ surface.

The net torque ($\vec{R} \times \vec{F}$) is calculated by multiplying $\sigma_{r\phi}$ by the area of the $r = R = \text{constant}$ surface, $2\pi RH$.

H : Height of the disk

Modeling Viscosity

Net torque on the outer ring due to the inner one .

Modeling Viscosity

Net torque on the outer ring due to the inner one .

$$\text{Torque} = R \times 2\pi RH \times \sigma_{r\phi}$$

Modeling Viscosity

Net torque on the outer ring due to the inner one .

$$\text{Torque} = R \times 2\pi R H \times \sigma_{r\phi}$$

$$\text{Torque} = -2\pi R^3 H \rho \nu \left. \frac{\partial \Omega}{\partial r} \right|_R$$

Modeling Viscosity

Using the surface mass density,

Modeling Viscosity

Using the surface mass density,

$$\Sigma = \int_0^H \rho(z) dz = \rho H$$

Modeling Viscosity

Using the surface mass density,

$$\Sigma = \int_0^H \rho(r) dz = \rho H$$

$$\text{Torque} = -2\pi R^3 \Sigma \nu \left. \frac{\partial \Omega}{\partial r} \right|_R$$

Modeling Viscosity

We define the torque on the inner ring by the outer one at the coordinate r as the function,

Modeling Viscosity

We define the torque on the inner ring by the outer one at the coordinate r as the function,

$$G(r) = 2\pi r^3 \Sigma \nu \frac{\partial \Omega}{\partial r}$$

- If $\frac{\partial \Omega}{\partial r} = 0$

Modeling Viscosity

We define the torque on the inner ring by the outer one at the coordinate r as the function,

$$G(r) = 2\pi r^3 \Sigma \nu \frac{\partial \Omega}{\partial r}$$

- If $\frac{\partial \Omega}{\partial r} = 0$
→ Rigid body rotation

Modeling Viscosity

We define the torque on the inner ring by the outer one at the coordinate r as the function,

$$G(r) = 2\pi r^3 \Sigma \nu \frac{\partial \Omega}{\partial r}$$

- If $\frac{\partial \Omega}{\partial r} = 0$
 - Rigid body rotation
 - No torque

Modeling Viscosity

We define the torque on the inner ring by the outer one at the coordinate r as the function,

$$G(r) = 2\pi r^3 \Sigma \nu \frac{\partial \Omega}{\partial r}$$

- If $\frac{\partial \Omega}{\partial r} = 0$
 - Rigid body rotation
 - No torque
- If $\Omega(r)$ decreases outwards, $\frac{\partial \Omega}{\partial r} < 0$,

Modeling Viscosity

We define the torque on the inner ring by the outer one at the coordinate r as the function,

$$G(r) = 2\pi r^3 \Sigma \nu \frac{\partial \Omega}{\partial r}$$

- If $\frac{\partial \Omega}{\partial r} = 0$
 - Rigid body rotation
 - No torque
- If $\Omega(r)$ decreases outwards, $\frac{\partial \Omega}{\partial r} < 0$,
 - $G(r) < 0$

Modeling Viscosity

We define the torque on the inner ring by the outer one at the coordinate r as the function,

$$G(r) = 2\pi r^3 \Sigma \nu \frac{\partial \Omega}{\partial r}$$

- If $\frac{\partial \Omega}{\partial r} = 0$
 - Rigid body rotation
 - No torque
- If $\Omega(r)$ decreases outwards, $\frac{\partial \Omega}{\partial r} < 0$,
 - $G(r) < 0$
 - angular momentum goes from inner circles to outer circles.


Modeling Viscosity

We define the torque on the inner ring by the outer one at the coordinate r as the function,

$$G(r) = 2\pi r^3 \Sigma \nu \frac{\partial \Omega}{\partial r}$$

- If $\frac{\partial \Omega}{\partial r} = 0$
 - Rigid body rotation
 - No torque
- If $\Omega(r)$ decreases outwards, $\frac{\partial \Omega}{\partial r} < 0$,
 - $G(r) < 0$
 - angular momentum goes from inner circles to outer circles.
 - The gas slowly spirals in!

Differential viscous torque



Modeling Viscosity

The net torque on the ring between r and $r + dr$ is

$$G(r + dr) - G(r) = \frac{\partial G}{\partial r} dr$$

Modeling Viscosity

The net torque on the ring between r and $r + dr$ is

$$G(r + dr) - G(r) = \frac{\partial G}{\partial r} dr$$

The work done by this torque is

Modeling Viscosity

The net torque on the ring between r and $r + dr$ is

$$G(r + dr) - G(r) = \frac{\partial G}{\partial r} dr$$

The work done by this torque is

$$dW = \frac{\partial G}{\partial r} dr d\phi$$

Modeling Viscosity

The net torque on the ring between r and $r + dr$ is

$$G(r + dr) - G(r) = \frac{\partial G}{\partial r} dr$$

The work done by this torque is

$$dW = \frac{\partial G}{\partial r} dr d\phi$$

and the associated power is

$$P = \frac{\partial G}{\partial r} \Omega dr = \left[\frac{\partial}{\partial r} (G(r)\Omega(r)) - G(r) \frac{\partial \Omega}{\partial r} \right] dr$$

Modeling Viscosity

$$P = \left[\frac{\partial}{\partial r} (G(r)\Omega(r)) - G(r) \frac{\partial \Omega}{\partial r} \right] dr$$

$\frac{\partial}{\partial r} (G(r)\Omega(r)) dr$: Rate of *convection* of the rotational energy through the gas by the torques.

Modeling Viscosity

$$P = \left[\frac{\partial}{\partial r} (G(r)\Omega(r)) - G(r) \frac{\partial \Omega}{\partial r} \right] dr$$

$\frac{\partial}{\partial r} (G(r)\Omega(r)) dr$: Rate of *convection* of the rotational energy through the gas by the torques.

Integration over the whole disk depends only on boundary conditions,

Modeling Viscosity

$$P = \left[\frac{\partial}{\partial r} (G(r)\Omega(r)) - G(r) \frac{\partial \Omega}{\partial r} \right] dr$$

$\frac{\partial}{\partial r} (G(r)\Omega(r)) dr$: Rate of *convection* of the rotational energy through the gas by the torques.

Integration over the whole disk depends only on boundary conditions,

$$\int_{r_{in}}^{r_{out}} \frac{\partial}{\partial r} (G(r)\Omega(r)) dr = \Omega(r)G(r) \Big|_{r_{in}}^{r_{out}}$$

Modeling Viscosity

$$P = \left[\frac{\partial}{\partial r} (G(r)\Omega(r)) - G(r) \frac{\partial \Omega}{\partial r} \right] dr$$

$-G(r) \frac{\partial \Omega}{\partial r} dr$: Local rate of loss of mechanical energy, which is transformed into heat.

Modeling Viscosity

$$P = \left[\frac{\partial}{\partial r} (G(r)\Omega(r)) - G(r)\frac{\partial \Omega}{\partial r} \right] dr$$

$-G(r)\frac{\partial \Omega}{\partial r}dr$: Local rate of loss of mechanical energy, which is transformed into heat.

This is the viscous dissipation rate per ring of width dr .

Modeling Viscosity

$$P = \left[\frac{\partial}{\partial r} (G(r)\Omega(r)) - G(r) \frac{\partial \Omega}{\partial r} \right] dr$$

$-G(r) \frac{\partial \Omega}{\partial r} dr$: Local rate of loss of mechanical energy, which is transformed into heat.

This is the viscous dissipation rate per ring of width dr .

$$D(r) = G(r) \frac{\partial \Omega}{\partial r} dr$$

Modeling Viscosity

Each ring of width dr has two sides, with area $\sim 2\pi r dr$.

Modeling Viscosity

Each ring of width dr has two sides, with area $\sim 2\pi r dr$.

We define the dissipation rate per unit plane surface as

$$D(r) = \frac{G(r) \frac{\partial \Omega}{\partial r} dr}{2(2\pi r dr)}$$

Modeling Viscosity

Each ring of width dr has two sides, with area $\sim 2\pi r dr$.

We define the dissipation rate per unit plane surface as

$$D(r) = \frac{G(r) \frac{\partial \Omega}{\partial r} dr}{2(2\pi r dr)}$$

$$D(r) = \frac{1}{2} \nu \Sigma \left(r \frac{\partial \Omega}{\partial r} \right)^2$$

Modeling Viscosity

$$D(r) = \frac{1}{2} \nu \Sigma \left(r \frac{\partial \Omega}{\partial r} \right)^2$$

Modeling Viscosity

$$D(r) = \frac{1}{2} \nu \Sigma \left(r \frac{\partial \Omega}{\partial r} \right)^2$$

$D(r) \geq 0$ always \rightarrow it always dissipate.

Modeling Viscosity

$$D(r) = \frac{1}{2} \nu \Sigma \left(r \frac{\partial \Omega}{\partial r} \right)^2$$

$D(r) \geq 0$ always \rightarrow it always dissipate.

The zero corresponds to the rigid body, $\left(\frac{\partial \Omega}{\partial r} = 0 \right)$

Modeling Viscosity

$$D(r) = \frac{1}{2} \nu \Sigma \left(r \frac{\partial \Omega}{\partial r} \right)^2$$

Modeling Viscosity

$$D(r) = \frac{1}{2} \nu \Sigma \left(r \frac{\partial \Omega}{\partial r} \right)^2$$

Using the Keplerian angular velocity,

$$\Omega = \Omega_k(r) = \sqrt{\frac{GM}{r^3}}$$

Modeling Viscosity

$$D(r) = \frac{1}{2} \nu \Sigma \left(r \frac{\partial \Omega}{\partial r} \right)^2$$

Using the Keplerian angular velocity,

$$\Omega = \Omega_k(r) = \sqrt{\frac{GM}{r^3}}$$

$$D(r) = \frac{9}{8} \nu \Sigma \frac{GM}{r^3}$$

Radial Structure Evolution

Radial Structure Evolution

In order to describe the radial structure of disk and its evolution we will use cylindrical coordinates (r, ϕ, z) .

Radial Structure Evolution

In order to describe the radial structure of disk and its evolution we will use cylindrical coordinates (r, ϕ, z) .

Angular velocity:

$$\Omega = \Omega_k(r) = \sqrt{\frac{GM}{r^3}}$$

Radial Structure Evolution

In order to describe the radial structure of disk and its evolution we will use cylindrical coordinates (r, ϕ, z) .

Angular velocity:

$$\Omega = \Omega_k(r) = \sqrt{\frac{GM}{r^3}}$$

Tangential velocity:

$$v_\phi = r\Omega_k(r)$$

Radial Structure Evolution

In order to describe the radial structure of disk and its evolution we will use cylindrical coordinates (r, ϕ, z) .

Angular velocity:

$$\Omega = \Omega_k(r) = \sqrt{\frac{GM}{r^3}}$$

Tangential velocity:

$$v_\phi = r\Omega_k(r)$$

Small radial *drift* velocity: $v_r = v_r(t, r) < 0$ (negative for accretion)

Radial Structure Evolution

In order to describe the radial structure of disk and its evolution we will use cylindrical coordinates (r, ϕ, z) .

Angular velocity:

$$\Omega = \Omega_k(r) = \sqrt{\frac{GM}{r^3}}$$

Tangential velocity:

$$v_\phi = r\Omega_k(r)$$

Small radial *drift* velocity: $v_r = v_r(t, r) < 0$ (negative for accretion)

Surface mass density: $\Sigma = \Sigma(t, r)$

Radial Structure Evolution

Consider m .

Radial Structure Evolution

Consider m.

Mass: $2\pi r dr \Sigma$

Radial Structure Evolution

Consider m.

Mass: $2\pi r dr \Sigma$

Total angular momentum: $(2\pi r dr \Sigma) r^2 \Omega$

Radial Structure Evolution

Consider m.

Mass: $2\pi r dr \Sigma$

Total angular momentum: $(2\pi r dr \Sigma) r^2 \Omega$

Rate of change of mass:

$$\frac{\partial}{\partial t} (2\pi r dr \Sigma) \cong -2\pi dr \frac{\partial}{\partial r} (r \Sigma v_r)$$

Radial Structure Evolution

Consider m.

Mass: $2\pi r dr \Sigma$

Total angular momentum: $(2\pi r dr \Sigma) r^2 \Omega$

Rate of change of mass:

$$\frac{\partial}{\partial t} (2\pi r dr \Sigma) \cong -2\pi dr \frac{\partial}{\partial r} (r \Sigma v_r)$$

In the limit $dr \rightarrow 0$ this gives the mass conservation equation

$$r \frac{\partial \Sigma}{\partial t} + \frac{\partial}{\partial r} (r \Sigma v_r) = 0$$

Radial Structure Evolution

Rate of change of the total angular momentum:

$$\frac{\partial}{\partial t} (2\pi r dr \Sigma r^2 \Omega) \cong -2\pi dr \frac{\partial}{\partial r} (r \Sigma v_r r^2 \Omega) + \frac{\partial G}{\partial r} dr$$

Radial Structure Evolution

Rate of change of the total angular momentum:

$$\frac{\partial}{\partial t} (2\pi r dr \Sigma r^2 \Omega) \cong -2\pi dr \frac{\partial}{\partial r} (r \Sigma v_r r^2 \Omega) + \frac{\partial G}{\partial r} dr$$

In the limit $dr \rightarrow 0$ this gives the momentum conservation equation

$$r \frac{\partial}{\partial t} (\Sigma r^2 \Omega) + \frac{\partial}{\partial r} (r \Sigma v_r r^2 \Omega) = \frac{1}{2\pi} \frac{\partial G}{\partial r}$$

Radial Structure Evolution

Equations determining the radial structure of the accretion disk:

$$r \frac{\partial \Sigma}{\partial t} + \frac{\partial}{\partial r} (r \Sigma v_r) = 0$$

$$r \frac{\partial}{\partial t} (\Sigma r^2 \Omega) + \frac{\partial}{\partial r} (r \Sigma v_r r^2 \Omega) = \frac{1}{2\pi} \frac{\partial G}{\partial r}$$

$$G(r, t) = 2\pi r \nu \Sigma r^2 \frac{\partial \Omega}{\partial r}$$

Equation for the kinematical viscosity ν

Radial Structure Evolution

Combining the equations determining the radial structure of the accretion disk and assuming that $\frac{\partial \Omega}{\partial t} = 0$, we obtain

$$r\Sigma v_r \frac{\partial}{\partial r} (r^2 \Omega) = \frac{1}{2\pi} \frac{\partial G}{\partial r}$$

Radial Structure Evolution

Combining the equations determining the radial structure of the accretion disk and assuming that $\frac{\partial \Omega}{\partial t} = 0$, we obtain

$$r\Sigma v_r \frac{\partial}{\partial r} (r^2 \Omega) = \frac{1}{2\pi} \frac{\partial G}{\partial r}$$

Eliminating the velocity, v_r , we have

$$r \frac{\partial \Sigma}{\partial t} = - \frac{\partial}{\partial r} \left[\frac{1}{2\pi \frac{\partial(r^2 \Omega)}{\partial r}} \frac{\partial G}{\partial r} \right]$$

Radial Structure Evolution

Using the Keplerian angular velocity we obtain the equation for the time evolution of the surface mass density

Radial Structure Evolution

Using the Keplerian angular velocity we obtain the equation for the time evolution of the surface mass density

$$\frac{\partial \Sigma}{\partial t} = -\frac{3}{r} \frac{\partial}{\partial r} \left[r^{1/2} \frac{\partial}{\partial r} (\nu \Sigma r^{1/2}) \right]$$

Radial Structure Evolution

Using the Keplerian angular velocity we obtain the equation for the time evolution of the surface mass density

$$\frac{\partial \Sigma}{\partial t} = \frac{3}{r} \frac{\partial}{\partial r} \left[r^{1/2} \frac{\partial}{\partial r} (\nu \Sigma r^{1/2}) \right]$$

This is a *nonlinear diffusion equation* for Σ and one needs to know the function ν to solve it.

Radial Structure Evolution

Using the Keplerian angular velocity we obtain the equation for the time evolution of the surface mass density

$$\frac{\partial \Sigma}{\partial t} = \frac{3}{r} \frac{\partial}{\partial r} \left[r^{1/2} \frac{\partial}{\partial r} (\nu \Sigma r^{1/2}) \right]$$

This is a *nonlinear diffusion equation* for Σ and one needs to know the function ν to solve it.

The radial velocity becomes

$$v_r = -\frac{3}{\Sigma r^{1/2}} \frac{\partial}{\partial r} [\nu \Sigma r^{1/2}]$$

Example of Viscosity

Consider the particular case in which $\nu = \text{constant}$.

Example of Viscosity

Consider the particular case in which $\nu = \text{constant}$.

$$\frac{\partial}{\partial t} (\Sigma r^{1/2}) = \frac{3\nu}{r} \left(r^{1/2} \frac{\partial}{\partial r} \right)^2 (\Sigma r^{1/2})$$

Example of Viscosity

Change $s = 2r^{1/2}$

Example of Viscosity

Change $s = 2r^{1/2}$

$$\frac{\partial}{\partial t} (\Sigma r^{1/2}) = \frac{12\nu}{s^2} \frac{\partial^2}{\partial s^2} (\Sigma r^{1/2})$$

Example of Viscosity

Change $s = 2r^{1/2}$

$$\frac{\partial}{\partial t} (\Sigma r^{1/2}) = \frac{12\nu}{s^2} \frac{\partial^2}{\partial s^2} (\Sigma r^{1/2})$$

Now consider $\Sigma r^{1/2} = T(t)S(s)$,

Example of Viscosity

Change $s = 2r^{1/2}$

$$\frac{\partial}{\partial t} (\Sigma r^{1/2}) = \frac{12\nu}{s^2} \frac{\partial^2}{\partial s^2} (\Sigma r^{1/2})$$

Now consider $\Sigma r^{1/2} = T(t)S(s)$,

$$\frac{1}{T} \frac{dT}{dt} = \frac{12\nu}{s''} \frac{d^2 S}{ds^2} = -\lambda^2 = \text{constant}$$

Example of Viscosity

Change $s = 2r^{1/2}$

$$\frac{\partial}{\partial t} (\Sigma r^{1/2}) = \frac{12\nu}{s^2} \frac{\partial^2}{\partial s^2} (\Sigma r^{1/2})$$

Now consider $\Sigma r^{1/2} = T(t)S(s)$,

$$\frac{1}{T} \frac{dT}{dt} = \frac{12\nu}{s''} \frac{d^2 S}{ds^2} = -\lambda^2 = \text{constant}$$

T : Exponential function

S : Bessel function

Example of Viscosity

The solution of this equation considering the initial distribution

$$\Sigma(r, t = 0) = \frac{m}{2\pi r_0} \delta(r - r_0)$$

Example of Viscosity

The solution of this equation considering the initial distribution

$$\Sigma(r, t = 0) = \frac{m}{2\pi r_0} \delta(r - r_0)$$

is the function

$$\Sigma(x, \tau) = \frac{m}{\pi r_0^2} \tau^{-1} x^{-1/4} \exp \left[-\frac{1 + x^2}{\tau} \right] I_{1/4} \left(\frac{2x}{\tau} \right)$$

Example of Viscosity

The solution of this equation considering the initial distribution

$$\Sigma(r, t = 0) = \frac{m}{2\pi r_0} \delta(r - r_0)$$

is the function


$$\Sigma(x, \tau) = \frac{m}{\pi r_0^2} \tau^{-1} x^{-1/4} \exp \left[-\frac{1 + x^2}{\tau} \right] I_{1/4} \left(\frac{2x}{\tau} \right)$$

$I_{1/4}(z)$: Modified Bessel function

$$x = \frac{r}{r_0}$$

$$\tau = \frac{12\nu t}{r_0^2}$$

Formation of the Accretion Disk



The α -Prescription for Viscosity

The α -Prescription for Viscosity

Outline for Part 2

1. Viscous Torques
 - 1.1 Differential Rotation
 - 1.2 Radial Structure Evolution
 - 1.3 The α -Prescription for Viscosity
2. The *alpha*-Prescription for Viscosity

The *alpha*-Prescription for Viscosity

The *alpha*-Prescription for Viscosity

The transferred material has a high specific angular momentum.
Therefore, accretion is not a direct process.

The *alpha*-Prescription for Viscosity

The transferred material has a high specific angular momentum.

Therefore, accretion is not a direct process.

As seen from the BH, the matter spreads as from a nozzle rotating around the center of mass.

The *alpha*-Prescription for Viscosity

The transferred material has a high specific angular momentum.


Therefore, accretion is not a direct process.

As seen from the BH, the matter spreads as from a nozzle rotating around the center of mass.

v_{\parallel} : parallel component of the velocity with respect to the line of centers.

v_{\perp} : perpendicular component of the velocity with respect to the line of centers.

Formation of the Accretion Disk



Formation of the Accretion Disk

$$v_{\parallel} \lesssim c_s$$

Formation of the Accretion Disk

$$v_{\parallel} \lesssim c_s$$

c_s : speed of sound in the envelope of the star.

Formation of the Accretion Disk

$$v_{\parallel} \lesssim c_s$$

c_s : speed of sound in the envelope of the star.

For normal stellar envelope temperatures, ($< 10^5$ K);

Formation of the Accretion Disk

$$v_{\parallel} \lesssim c_s$$

c_s : speed of sound in the envelope of the star.

For normal stellar envelope temperatures, ($< 10^5$ K); $c_s \lesssim 10$ km/s

Formation of the Accretion Disk

$$v_{\perp} \sim b\omega = \frac{2\pi b}{\tau}$$

Formation of the Accretion Disk

$$v_{\perp} \sim b\omega = \frac{2\pi b}{\tau}$$

Using

$$b \approx a [0.500 - 0.227 \log_{10}(q)]$$

$$a = 3.5 \times 10^{10} \left(\frac{M}{M_{\odot}} \right)^{1/3} (1+q)^{1/3} \tau_{hr}^{2/3} [\text{cm}]$$

Formation of the Accretion Disk

$$v_{\perp} \sim b\omega = \frac{2\pi b}{\tau}$$

Using

$$b \approx a [0.500 - 0.227 \log_{10}(q)]$$

$$a = 3.5 \times 10^{10} \left(\frac{M}{M_{\odot}} \right)^{1/3} (1+q)^{1/3} \tau_{hr}^{2/3} [\text{cm}]$$

we get

$$v_{\perp} \gtrsim 305.6 \left(\frac{M}{M_{\odot}} \right)^{1/3} (1+q)^{1/3} \tau_{hr}^{-1/3} [\text{km/s}]$$

Formation of the Accretion Disk

$$v_{\parallel} \lesssim c_s \lesssim 10 \text{ km/s}$$

$$v_{\perp} \gtrsim 305.6 \left(\frac{M}{M_{\odot}} \right)^{1/3} (1+q)^{1/3} \tau_{hr}^{-1/3} \text{ [km/s]}$$

Formation of the Accretion Disk

$$v_{\parallel} \lesssim c_s \lesssim 10 \text{ km/s}$$

$$v_{\perp} \gtrsim 305.6 \left(\frac{M}{M_{\odot}} \right)^{1/3} (1+q)^{1/3} \tau_{hr}^{-1/3} \text{ [km/s]}$$

$$v_{\perp} \gtrsim 104.2 \left(\frac{M}{M_{\odot}} \right)^{1/3} (1+q)^{1/3} \tau_{days}^{-1/3} \text{ [km/s]}$$

Formation of the Accretion Disk

$$v_{\parallel} \lesssim c_s \lesssim 10 \text{ km/s}$$

$$v_{\perp} \gtrsim 305.6 \left(\frac{M}{M_{\odot}} \right)^{1/3} (1+q)^{1/3} \tau_{hr}^{-1/3} [\text{km/s}]$$

$$v_{\perp} \gtrsim 104.2 \left(\frac{M}{M_{\odot}} \right)^{1/3} (1+q)^{1/3} \tau_{days}^{-1/3} [\text{km/s}]$$

$$v_{\perp} \gg v_{\parallel}$$

Formation of the Accretion Disk

- A particle (or parcel of gas) is released from rest at the point L_1 , i.e. pressure effects are neglected.

Formation of the Accretion Disk

- A particle (or parcel of gas) is released from rest at the point L_1 , i.e. pressure effects are neglected.
- Because of the motion of the star, this particle is moving with velocity v_{\perp} as seen from the BH.

Formation of the Accretion Disk

- A particle (or parcel of gas) is released from rest at the point L_1 , i.e. pressure effects are neglected.
- Because of the motion of the star, this particle is moving with velocity v_{\perp} as seen from the BH.
- The particle will describe an elliptical trajectory around the BH but the presence of the star will make this ellipse to precess slowly.

Formation of the Accretion Disk

- In the stream of accreting particles, their orbits will intersect, resulting in dissipation of energy via collisions.

Formation of the Accretion Disk

- In the stream of accreting particles, their orbits will intersect, resulting in dissipation of energy via collisions.
- However, angular momentum will be conserved in this collisions.

Formation of the Accretion Disk

- In the stream of accreting particles, their orbits will intersect, resulting in dissipation of energy via collisions.
- However, angular momentum will be conserved in this collisions.
- Hence, particles will go into the trajectories with minimum energy for a given angular momentum,

Formation of the Accretion Disk

- In the stream of accreting particles, their orbits will intersect, resulting in dissipation of energy via collisions.
- However, angular momentum will be conserved in this collisions.
- Hence, particles will go into the trajectories with minimum energy for a given angular momentum, i.e. circular orbits!

Formation of the Accretion Disk

- In the stream of accreting particles, their orbits will intersect, resulting in dissipation of energy via collisions.
- However, angular momentum will be conserved in this collisions.
- Hence, particles will go into the trajectories with minimum energy for a given angular momentum, i.e. circular orbits!
- This process is called *circularization*.

Formation of the Accretion Disk

- In the stream of accreting particles, their orbits will intersect, resulting in dissipation of energy via collisions.
- However, angular momentum will be conserved in this collisions.
- Hence, particles will go into the trajectories with minimum energy for a given angular momentum, i.e. circular orbits!
- This process is called *circularization*.
- The radius of the resulting circular orbit is called *circularization radius*, r_{circ} .

Formation of the Accretion Disk

Ω_k : Keplerian angular velocity in the circular orbit

Formation of the Accretion Disk

Ω_k : Keplerian angular velocity in the circular orbit

v_ϕ : Tangential velocity in the circular orbit

Formation of the Accretion Disk

Ω_k : Keplerian angular velocity in the circular orbit

v_ϕ : Tangential velocity in the circular orbit

$$r_{\text{circ}} \Omega_k^2(r_{\text{circ}}) = \frac{v_\phi^2(r_{\text{circ}})}{r_{\text{circ}}} = \frac{GM}{r_{\text{circ}}^2}$$

Formation of the Accretion Disk

Ω_k : Keplerian angular velocity in the circular orbit

v_ϕ : Tangential velocity in the circular orbit

$$r_{\text{circ}} \Omega_k^2(r_{\text{circ}}) = \frac{v_\phi^2(r_{\text{circ}})}{r_{\text{circ}}} = \frac{GM}{r_{\text{circ}}^2}$$

$$v_\phi(r_{\text{circ}}) = \sqrt{\frac{GM}{r_{\text{circ}}}}$$

Formation of the Accretion Disk

Conservation of angular momentum

Formation of the Accretion Disk

Conservation of angular momentum

$$r_{\text{circ}} v_{\phi} = b^2 \omega$$

Formation of the Accretion Disk

Conservation of angular momentum

$$r_{\text{circ}} v_{\phi} = b^2 \omega$$

$$\frac{r_{\text{circ}}}{a} = \frac{4\pi^2}{GM\tau} a^3 \left(\frac{b}{a} \right)^4$$

Formation of the Accretion Disk

Conservation of angular momentum

$$r_{\text{circ}} v_{\phi} = b^2 \omega$$

$$\frac{r_{\text{circ}}}{a} = \frac{4\pi^2}{GM\tau} a^3 \left(\frac{b}{a} \right)^4$$

and using Kepler's third law,

$$\frac{r_{\text{circ}}}{a} = (1 + q) \left(\frac{b}{a} \right)^4$$

Formation of the Accretion Disk

Conservation of angular momentum

$$r_{\text{circ}} v_{\phi} = b^2 \omega$$

$$\frac{r_{\text{circ}}}{a} = \frac{4\pi^2}{GM\tau} a^3 \left(\frac{b}{a} \right)^4$$

and using Kepler's third law,

$$\frac{r_{\text{circ}}}{a} = (1 + q) \left(\frac{b}{a} \right)^4$$

$$\frac{r_{\text{circ}}}{a} = (1 + q) [0.500 - 0.227 \log_{10} q]^4$$

Formation of the Accretion Disk

$$r_{\text{circ}} = (1 + q)^{4/3} [0.500 - 0.227 \log_{10} q]^4 \left(\frac{M}{M_{\odot}} \right)^{1/3} \tau_{\text{days}}^{2/3} [R_{\odot}]$$

Formation of the Accretion Disk

- The stream of gas moves in a ring with $r = r_{\text{circ}}$.

Formation of the Accretion Disk

- The stream of gas moves in a ring with $r = r_{circ}$.
- In the ring the particles have collisions, shocks, viscous dissipation and other processes that transform some of the potential energy into heat (producing radiation).

Formation of the Accretion Disk

- The stream of gas moves in a ring with $r = r_{\text{circ}}$.
- In the ring the particles have collisions, shocks, viscous dissipation and other processes that transform some of the potential energy into heat (producing radiation).
- However, this release of energy needs the losing of angular momentum.

Formation of the Accretion Disk

- The stream of gas moves in a ring with $r = r_{\text{circ}}$.
- In the ring the particles have collisions, shocks, viscous dissipation and other processes that transform some of the potential energy into heat (producing radiation).
- However, this release of energy needs the losing of angular momentum.
- In the absence of external torques, the only possible process is a *transfer of angular momentum* from inner regions outwards by internal torques.

Formation of the Accretion Disk

- The stream of gas moves in a ring with $r = r_{\text{circ}}$.
- In the ring the particles have collisions, shocks, viscous dissipation and other processes that transform some of the potential energy into heat (producing radiation).
- However, this release of energy needs the losing of angular momentum.
- In the absence of external torques, the only possible process is a *transfer of angular momentum* from inner regions outwards by internal torques.
- The redistribution of angular momentum makes particles in the outer parts move outwards (gaining angular momentum) and the particles in the inner particles spiral inwards.

Formation of the Accretion Disk

- The accretion disk will extend from $r_{in} \geq r_{ISCO}$ up to $r_{out} \leq b$

Formation of the Accretion Disk

- The accretion disk will extend from $r_{in} \geq r_{ISCO}$ up to $r_{out} \leq b$
- Viscous torques may be modeled using different processes:

Formation of the Accretion Disk

- The accretion disk will extend from $r_{in} \geq r_{ISCO}$ up to $r_{out} \leq b$
- Viscous torques may be modeled using different processes:
 - Viscous torques due to differential rotation in the accretion disk (produced by the thermal motion of the fluid molecules). This is a local mechanism for angular momentum transport

Formation of the Accretion Disk

- The accretion disk will extend from $r_{in} \geq r_{ISCO}$ up to $r_{out} \leq b$
- Viscous torques may be modeled using different processes:
 - Viscous torques due to differential rotation in the accretion disk (produced by the thermal motion of the fluid molecules). This is a local mechanism for angular momentum transport
 - Magnetic loops that couple fluid elements located at macroscopic distances across the disk. This is a non-local mechanism for angular momentum transport.

Formation of the Accretion Disk

- The accretion disk will extend from $r_{in} \geq r_{ISCO}$ up to $r_{out} \leq b$
- Viscous torques may be modeled using different processes:
 - Viscous torques due to differential rotation in the accretion disk (produced by the thermal motion of the fluid molecules). This is a local mechanism for angular momentum transport
 - Magnetic loops that couple fluid elements located at macroscopic distances across the disk. This is a non-local mechanism for angular momentum transport.
 - Turbulence in the fluid may be an origin of angular momentum transport. Turbulence may be produced by mechanisms as Thermally driven convection, Pure hydrodynamic instabilities or Magnetohydrodynamic (MHD) turbulence.

Formation of the Accretion Disk

- The accretion disk will extend from $r_{in} \geq r_{ISCO}$ up to $r_{out} \leq b$
- Viscous torques may be modeled using different processes:
 - Viscous torques due to differential rotation in the accretion disk (produced by the thermal motion of the fluid molecules). This is a local mechanism for angular momentum transport
 - Magnetic loops that couple fluid elements located at macroscopic distances across the disk. This is a non-local mechanism for angular momentum transport.
 - Turbulence in the fluid may be an origin of angular momentum transport. Turbulence may be produced by mechanisms as Thermally driven convection, Pure hydrodynamic instabilities or Magnetohydrodynamic (MHD) turbulence.
- However, this release of energy needs the losing of angular momentum.


Formation of the Accretion Disk

- The accretion disk will extend from $r_{in} \geq r_{ISCO}$ up to $r_{out} \leq b$
- Viscous torques may be modeled using different processes:
 - Viscous torques due to differential rotation in the accretion disk (produced by the thermal motion of the fluid molecules). This is a local mechanism for angular momentum transport
 - Magnetic loops that couple fluid elements located at macroscopic distances across the disk. This is a non-local mechanism for angular momentum transport.
 - Turbulence in the fluid may be an origin of angular momentum transport. Turbulence may be produced by mechanisms as Thermally driven convection, Pure hydrodynamic instabilities or Magnetohydrodynamic (MHD) turbulence.
- However, this release of energy needs the losing of angular momentum.
- In the absence of external torques, the only possible process is *a transfer of angular momentum* from inner regions outwards

Formation of the Accretion Disk

- The accretion disk will extend from $r_{in} \geq r_{ISCO}$ up to $r_{out} \leq b$
- Viscous torques may be modeled using different processes:
 - Viscous torques due to differential rotation in the accretion disk (produced by the thermal motion of the fluid molecules). This is a local mechanism for angular momentum transport
 - Magnetic loops that couple fluid elements located at macroscopic distances across the disk. This is a non-local mechanism for angular momentum transport.
 - Turbulence in the fluid may be an origin of angular momentum transport. Turbulence may be produced by mechanisms as Thermally driven convection, Pure hydrodynamic instabilities or Magnetohydrodynamic (MHD) turbulence.
- However, this release of energy needs the losing of angular momentum.
- In the absence of external torques, the only possible process is *a transfer of angular momentum* from inner regions outwards

Temperature of the gas in the accretion structure



Next Lecture

10. Accretion Disks. Detailed Description