



BLACK HOLES

OBSERVATORIO
ASTRONÓMICO
NACIONAL

Classical Black Holes

08. Accretion

Edward Larrañaga

Outline for Part 1

1. Accretion Basics

- 1.1 Spherical Accretion
- 1.2 Eddington Luminosity
- 1.3 Estimation of the Central Mass
- 1.4 Eddington Accretion Rate
- 1.5 Growth Time
- 1.6 Temperatures
- 1.7 Compactness

2. Geodesics in Kerr Spacetime

- 2.1 Hamilton-Jacobi Formulation
- 2.2 Equations of Motion
- 2.3 Imaging a Black Hole

Accretion Basics

Accretion Basics

Process of matter falling into the potential well of a gravitating object.

Accretion Regimes

1. Spherical Accretion
2. Cylindrical Accretion
3. Accretion Disk
4. Two-Stream Accretion

Spherical Accretion

- No (significant) angular momentum

Spherical Accretion

- No (significant) angular momentum
- Determined by the relation between
 c_s : speed of sound in matter
 v_{rel} : relative velocity between accretor and matter

Spherical Accretion

- No (significant) angular momentum
- Determined by the relation between
 c_s : speed of sound in matter
 v_{rel} : relative velocity between accretor and matter
- $v_{rel} \ll c_s$

Spherical Accretion

- No (significant) angular momentum
- Determined by the relation between
 c_s : speed of sound in matter
 v_{rel} : relative velocity between accretor and matter
- $v_{rel} \ll c_s$
- If the accretor is a BH,

$$v_{rel} = v$$

v : velocity of accreting matter (i.e. the BH doesn't move!)

Cylindrical Accretion

- Small angular momentum

Cylindrical Accretion

- Small angular momentum
- $v_{rel} \geq c_s$

Accretion Disk

- Angular momentum is high enough to form an accretion disk

Accretion Disk

- Angular momentum is high enough to form an accretion disk
- Matter spirals down into the accretor

Two-Stream Accretion

- Quasi-spherically symmetric inflow coexist with an accretion disk

Spherical Accretion

Spherical Accretion

We assume a completely ionized hydrogen gas cloud as the accretion structure.

Spherical Accretion

We assume a completely ionized hydrogen gas cloud as the accretion structure.

To avoid the disintegration of the accretion structure, the outward force due to radiation pressure must be counterbalanced by the gravitational force.

Radiation Pressure

Outward energy flux at distance r from the center

$$F = \frac{L}{4\pi r^2}$$

Radiation Pressure

Outward energy flux at distance r from the center

$$F = \frac{L}{4\pi r^2}$$

L : bolometric luminosity [$\text{erg} \cdot \text{s}^{-1}$]

Radiation Pressure

Outward energy flux at distance r from the center

$$F = \frac{L}{4\pi r^2}$$

L : bolometric luminosity [$\text{erg} \cdot \text{s}^{-1}$] For photons:

$$p^\mu = \left(\frac{E}{c}, \vec{p} \right) \quad p^2 = \frac{E^2}{c^2} - |\vec{p}|^2 = 0$$

Radiation Pressure

Outward energy flux at distance r from the center

$$F = \frac{L}{4\pi r^2}$$

L : bolometric luminosity [$\text{erg} \cdot \text{s}^{-1}$] For photons:

$$p^\mu = \left(\frac{E}{c}, \vec{p} \right) \quad p^2 = \frac{E^2}{c^2} - |\vec{p}|^2 = 0$$

Then, the outwards momentum flux (or pressure) is

$$P_{\text{rad}} = \frac{F}{c} = \frac{L}{4\pi r^2 c}$$

Radiation Pressure

The radiation force on a single electron is

$$\vec{f}_{rad} = (\sigma_e P_{rad}) \hat{r} = \sigma_e \frac{L}{4\pi r^2 c} \hat{r}$$

Radiation Pressure

The radiation force on a single electron is

$$\vec{f}_{rad} = (\sigma_e P_{rad}) \hat{r} = \sigma_e \frac{L}{4\pi r^2 c} \hat{r}$$

σ_e : Thomson cross-section

Radiation Pressure

The radiation force on a single electron is

$$\vec{f}_{rad} = (\sigma_e P_{rad}) \hat{r} = \sigma_e \frac{L}{4\pi r^2 c} \hat{r}$$

σ_e : Thomson cross-section

$$\sigma_e = \frac{8\pi}{3} \left(\frac{e^2}{m_e c^2} \right)^2 = 6.65 \times 10^{-25} \text{ cm}^2$$

Interaction with protons is negligible because it is lower by a factor of $\left(\frac{m_p}{m_e} \right)^2 \sim 3 \times 10^6$

Gravitational Force

The gravitational force between the central object M and one electron-proton pair is

Gravitational Force

The gravitational force between the central object M and one electron-proton pair is

$$\vec{f}_g = -\frac{GM(m_e + m_p)}{r^2}\hat{r} \sim -\frac{GMm_p}{r^2}\hat{r}$$

Spherical Accretion

To avoid the disintegration of the accretion structure,

Spherical Accretion

To avoid the disintegration of the accretion structure,

$$|\vec{f}_{rad}| \leq |\vec{f}_g|$$

Spherical Accretion

To avoid the disintegration of the accretion structure,

$$|\vec{f}_{rad}| \leq |\vec{f}_g|$$

$$\sigma_e \frac{L}{4\pi r^2 c} \leq \frac{GMm_p}{r^2}$$

Spherical Accretion

To avoid the disintegration of the accretion structure,

$$|\vec{f}_{rad}| \leq |\vec{f}_g|$$

$$\sigma_e \frac{L}{4\pi r^2 c} \leq \frac{GMm_p}{r^2}$$

$$L \leq \frac{4\pi GMm_p c}{\sigma_e}$$

Eddington Luminosity

$$L \leq L_E$$

Eddington Luminosity

$$L \leq L_E$$

$$L_E = \frac{4\pi GMm_p c}{\sigma_e}$$

Eddington Luminosity

$$L \leq L_E$$

$$L_E = \frac{4\pi GMm_p c}{\sigma_e}$$

$$L_E = 6.31 \times 10^4 M [\text{erg} \cdot \text{s}^{-1} \cdot \text{kg}^{-1}]$$

Eddington Luminosity

$$L \leq L_E$$

$$L_E = \frac{4\pi GMm_p c}{\sigma_e}$$

$$L_E = 6.31 \times 10^4 M \text{ [erg} \cdot \text{s}^{-1} \cdot \text{kg}^{-1}\text{]}$$

$$L_E = 1.26 \times 10^{38} \left(\frac{M}{M_\odot} \right) \text{ [erg} \cdot \text{s}^{-1}\text{]}$$

Eddington Luminosity

$$L \leq L_E$$

$$L_E = \frac{4\pi G M m_p c}{\sigma_e}$$

$$L_E = 6.31 \times 10^4 M \text{ [erg} \cdot \text{s}^{-1} \cdot \text{kg}^{-1}\text{]}$$

$$L_E = 1.26 \times 10^{38} \left(\frac{M}{M_\odot} \right) \text{ [erg} \cdot \text{s}^{-1}\text{]}$$

Eddington Luminosity: Maximum luminosity of a source M powered by spherical accretion.

Estimation of the Central Mass from its Luminosity

$$M \geq \frac{L\sigma_e}{4\pi Gm_p c}$$

Estimation of the Central Mass from its Luminosity

$$M \geq \frac{L\sigma_e}{4\pi Gm_p c}$$

$$M \geq \frac{L}{1.26 \times 10^{38}} [M_{\odot}]$$

Estimation of the Central Mass from its Luminosity

$$M \geq \frac{L\sigma_e}{4\pi Gm_p c}$$

$$M \geq \frac{L}{1.26 \times 10^{38}} [M_\odot]$$

$$M \geq 8 \times 10^5 L_{44} [M_\odot]$$

Estimation of the Central Mass from its Luminosity

$$M \geq \frac{L\sigma_e}{4\pi Gm_p c}$$

$$M \geq \frac{L}{1.26 \times 10^{38}} [M_\odot]$$

$$M \geq 8 \times 10^5 L_{44} [M_\odot]$$

L_{44} : Central Source Luminosity in units of $10^{44} \text{ erg} \cdot \text{s}^{-1}$ (typical value for AGN's)

Estimation of the Central Mass from its Luminosity

$$M \geq \frac{L\sigma_e}{4\pi G m_p c}$$

$$M \geq \frac{L}{1.26 \times 10^{38}} [M_\odot]$$

$$M \geq 8 \times 10^5 L_{44} [M_\odot]$$

L_{44} : Central Source Luminosity in units of $10^{44} \text{ erg} \cdot \text{s}^{-1}$ (typical value for AGN's)

$M_E = 8 \times 10^5 L_{44} M_\odot$: Eddington's Mass. Minimum mass for a given luminosity

Estimation of the Central Mass from its Luminosity

AGN's have typically $L \sim 10^{43} - 10^{47} \text{ erg} \cdot \text{s}^{-1}$

Estimation of the Central Mass from its Luminosity

AGN's have typically $L \sim 10^{43} - 10^{47} \text{ erg} \cdot \text{s}^{-1}$

Black holes with $M \sim 10^5 - 10^9 M_{\odot}$

Estimation of the Central Mass from its Luminosity

Name	z	L_{bol}	M_{BH}	ref.	Type
3C 120	0.033	45.34	I	7.42	1 SY1
3C 390.3	0.056	44.88	I	8.55	1 SY1
Akn 120	0.032	44.91	I	8.27	1 SY1
F 9	0.047	45.23	F	7.91	1 SY1
IC 4329A	0.016	44.78	I	6.77	1 SY1
Mrk 79	0.022	44.57	I	7.86	1 SY1
Mrk 110	0.035	44.71	F	6.82	1 SY1
Mrk 335	0.026	44.69	I	6.69	1 SY1
Mrk 509	0.034	45.03	I	7.86	1 SY1
Mrk 590	0.026	44.63	I	7.20	1 SY1
Mrk 817	0.032	44.99	I	7.60	1 SY1
NGC 3227	0.004	43.86	I	7.64	1 SY1
NGC 3516	0.009	44.29	I	7.36	3 SY1
NGC 3783	0.010	44.41	I	6.94	2 SY1
NGC 4051	0.002	43.56	I	6.13	1 SY1
NGC 4151	0.003	43.73	I	7.13	1 SY1
NGC 4593	0.009	44.09	I	6.91	3 SY1
NGC 5548	0.017	44.83	I	8.03	1 SY1
NGC 7469	0.016	45.28	I	6.84	1 SY1
PG 0026+129	0.142	45.39	I	7.58	1 RQQ
PG 0052+251	0.155	45.93	F	8.41	1 RQQ
PG 0804+761	0.100	45.93	F	8.24	1 RQQ
PG 0844+349	0.064	45.36	F	7.38	1 RQQ
PG 0953+414	0.239	46.16	F	8.24	1 RQQ
PG 1211+143	0.085	45.81	F	7.49	1 RQQ
PG 1229+204	0.064	45.01	I	8.56	1 RQQ
PG 1307+085	0.155	45.83	F	7.90	1 RQQ
PG 1351+640	0.087	45.50	I	8.48	1 RQQ
PG 1411+442	0.089	45.58	F	7.57	1 RQQ
PG 1426+015	0.086	45.19	I	7.92	1 RQQ
PG 1613+658	0.129	45.66	I	8.62	1 RQQ

Name	z	L_{bol}	M_{BH}	ref.	Type
PG 1617+175	0.114	45.52	F	7.88	1 RQQ
PG 1700+518	0.292	46.56	F	8.31	1 RQQ
PG 2130+099	0.061	45.47	I	7.74	1 RQQ
PG 1226+023	0.158	47.35	I	7.22	1 RLQ
PG 1704+608	0.371	46.33	I	8.23	1 RLQ

^a Column (1) Name, (2) redshift, (3) log of the bolometric luminosity (ergs s^{-1}), (4) method for bolometric luminosity estimation (I: flux integration; F: SED fitting), (5) black hole mass estimate from reverberation mapping (for Kaspi et al. (2000) sample, where black hole mass is log mean of rms FWHM and mean FWHM mass, in solar masses), (6) reference for black hole mass estimation, and (7) AGN type.

References. — (1) Kaspi et al. (2000), (2) Onken & Peterson (2002), (3) Ho (1999).

Accretion Rate

The luminosity is just a fraction of the relativistic energy of the accreting mass, $E = mc^2$. The other fraction goes into the BH making it grow.

Accretion Rate

The luminosity is just a fraction of the relativistic energy of the accreting mass, $E = mc^2$. The other fraction goes into the BH making it grow.

$$L \propto \frac{dE}{dt} = \frac{dm}{dt}c^2 = \frac{dM}{dt}c^2$$

Accretion Rate

The luminosity is just a fraction of the relativistic energy of the accreting mass, $E = mc^2$. The other fraction goes into the BH making it grow.

$$L \propto \frac{dE}{dt} = \frac{dm}{dt}c^2 = \frac{dM}{dt}c^2$$

$$L = \eta \dot{M} c^2$$

Accretion Rate

The luminosity is just a fraction of the relativistic energy of the accreting mass, $E = mc^2$. The other fraction goes into the BH making it grow.

$$L \propto \frac{dE}{dt} = \frac{dm}{dt}c^2 = \frac{dM}{dt}c^2$$

$$L = \eta \dot{M} c^2$$

η : Efficiency of the process

Accretion Rate

Accretion produces radiation by conversion of gravitational potential.

$$U = \frac{GMm}{r}$$

Accretion Rate

Accretion produces radiation by conversion of gravitational potential.

$$U = \frac{GMm}{r}$$

$$L \sim \frac{dU}{dt} = \frac{GM}{r} \frac{dm}{dt} = \frac{GM}{r} \dot{M}$$

Accretion Rate

Accretion produces radiation by conversion of gravitational potential.

$$U = \frac{GMm}{r}$$

$$L \sim \frac{dU}{dt} = \frac{GM}{r} \frac{dm}{dt} = \frac{GM}{r} \dot{M}$$

$$\eta = \frac{GM}{rc^2}$$

Accretion Rate

In order to estimate the efficiency, consider the ISCO for Schwarzschild,

$$r_{ISCO} = 3r_s = \frac{6GM}{c^2}$$

Accretion Rate

In order to estimate the efficiency, consider the ISCO for Schwarzschild,

$$r_{ISCO} = 3r_s = \frac{6GM}{c^2}$$

A particle falling from this orbit into the BH loses the energy

Accretion Rate

In order to estimate the efficiency, consider the ISCO for Schwarzschild,

$$r_{ISCO} = 3r_s = \frac{6GM}{c^2}$$

A particle falling from this orbit into the BH loses the energy Hence

Accretion Rate

In order to estimate the efficiency, consider the ISCO for Schwarzschild,

$$r_{ISCO} = 3r_s = \frac{6GM}{c^2}$$

A particle falling from this orbit into the BH loses the energy Hence

$$\eta \sim 0.1 - 0.2$$

Accretion Rate

In some books consider the accretion of particles falling from $r = 5r_S$, because it gives most of the optical/UV continuum radiation.

Accretion Rate

In some books consider the accretion of particles falling from $r = 5r_S$, because it gives most of the optical/UV continuum radiation.

Using this point, one obtains an efficiency of $\eta \sim 0.1$

Accretion Rate

For a typical AGN, $L \sim 10^{47} \text{ erg} \cdot \text{s}^{-1}$.

Accretion Rate

For a typical AGN, $L \sim 10^{47} \text{ erg} \cdot \text{s}^{-1}$.

- If $\eta \sim 0.007$ as in the Hydrogen burning process (Nuclear fusion), it gives

Accretion Rate

For a typical AGN, $L \sim 10^{47} \text{ erg} \cdot \text{s}^{-1}$.

- If $\eta \sim 0.007$ as in the Hydrogen burning process (Nuclear fusion), it gives

$$\dot{M} = \frac{L}{\eta c^2} \sim 250 M_{\odot} \cdot \text{yr}^{-1}$$

Accretion Rate

For a typical AGN, $L \sim 10^{47} \text{ erg} \cdot \text{s}^{-1}$.

- If $\eta \sim 0.007$ as in the Hydrogen burning process (Nuclear fusion), it gives

$$\dot{M} = \frac{L}{\eta c^2} \sim 250 M_{\odot} \cdot \text{yr}^{-1}$$

- If $\eta \sim 0.1$ we obtain a more realistic value of

Accretion Rate

For a typical AGN, $L \sim 10^{47} \text{ erg} \cdot \text{s}^{-1}$.

- If $\eta \sim 0.007$ as in the Hydrogen burning process (Nuclear fusion), it gives

$$\dot{M} = \frac{L}{\eta c^2} \sim 250 M_{\odot} \cdot \text{yr}^{-1}$$

- If $\eta \sim 0.1$ we obtain a more realistic value of

$$\dot{M} = \frac{L}{\eta c^2} \sim 250 M_{\odot} \cdot \text{yr}^{-1}$$

Accretion Rate

For a typical AGN, $L \sim 10^{47} \text{ erg} \cdot \text{s}^{-1}$.

- If $\eta \sim 0.007$ as in the Hydrogen burning process (Nuclear fusion), it gives

$$\dot{M} = \frac{L}{\eta c^2} \sim 250 M_{\odot} \cdot \text{yr}^{-1}$$

- If $\eta \sim 0.1$ we obtain a more realistic value of

$$\dot{M} = \frac{L}{\eta c^2} \sim 250 M_{\odot} \cdot \text{yr}^{-1}$$

Accretion Rate

$$\dot{M} = \frac{L}{\eta c^2} = 1.11 \times 10^{23} \frac{L_{44}}{\eta} \left[\frac{\text{gr}}{\text{s}} \right]$$

Accretion Rate

$$\dot{M} = \frac{L}{\eta c^2} = 1.11 \times 10^{23} \frac{L_{44}}{\eta} \left[\frac{\text{gr}}{\text{s}} \right]$$

$$\dot{M} = 1.77 \times 10^{-2} L_{44} \left(\frac{\eta}{0.1} \right)^{-1} \left[\frac{M_{\odot}}{\text{yr}} \right]$$

Eddington Accretion Rate

$$\dot{M}_E = \frac{L_E}{\eta c^2} = \frac{4\pi GMm_p}{\sigma_e \eta c}$$

Eddington Accretion Rate

$$\dot{M}_E = \frac{L_E}{\eta c^2} = \frac{4\pi G M m_p}{\sigma_e \eta c}$$

$$\dot{M}_E = 2.67 \times 10^{-8} \left(\frac{\eta}{0.1} \right)^{-1} \frac{M}{M_\odot} \left[\frac{M_\odot}{\text{yr}} \right]$$

Eddington Accretion Rate

$$\dot{M}_E = \frac{L_E}{\eta c^2} = \frac{4\pi GMm_p}{\sigma_e \eta c}$$

$$\dot{M}_E = 2.67 \times 10^{-8} \left(\frac{\eta}{0.1} \right)^{-1} \frac{M}{M_\odot} \left[\frac{M_\odot}{\text{yr}} \right]$$

$$\dot{M}_E = 3 \left(\frac{\eta}{0.1} \right)^{-1} M_8 \left[\frac{M_\odot}{\text{yr}} \right]$$

Eddington Accretion Rate

$$\dot{M}_E = \frac{L_E}{\eta c^2} = \frac{4\pi G M m_p}{\sigma_e \eta c}$$

$$\dot{M}_E = 2.67 \times 10^{-8} \left(\frac{\eta}{0.1} \right)^{-1} \frac{M}{M_\odot} \left[\frac{M_\odot}{\text{yr}} \right]$$

$$\dot{M}_E = 3 \left(\frac{\eta}{0.1} \right)^{-1} M_8 \left[\frac{M_\odot}{\text{yr}} \right]$$

\dot{M}_E : Maximum possible accretion rate for a mass $M_8 = \frac{M}{10^8 M_\odot}$.

Eddington Accretion Rate

May $\dot{M} > \dot{M}_E$?

Eddington Accretion Rate

May $\dot{M} > \dot{M}_E$?

1. It depends on a careful determination of η . E.g. if $\eta < 0.1$, the outwards flux is diminished.

Eddington Accretion Rate

May $\dot{M} > \dot{M}_E$?

1. It depends on a careful determination of η . E.g. if $\eta < 0.1$, the outwards flux is diminished.
2. \dot{M}_E can be exceeded with non-spherical models.

Growth Time

$$\dot{M} = \frac{dM}{dt} = \frac{L}{\eta c^2}$$

Growth Time

$$\dot{M} = \frac{dM}{dt} = \frac{L}{\eta c^2}$$

$$\dot{M} = \frac{dM}{dt} = \frac{L}{L_E} \frac{4\pi G M m_p}{\sigma_e \eta c}$$

Growth Time

$$\dot{M} = \frac{dM}{dt} = \frac{L}{\eta c^2}$$

$$\dot{M} = \frac{dM}{dt} = \frac{L}{L_E} \frac{4\pi G M m_p}{\sigma_e \eta c}$$

$$\int \frac{dM}{M} = \frac{L}{L_E} \frac{4\pi G m_p}{\sigma_e \eta c} \int dt$$

Growth Time

$$\dot{M} = \frac{dM}{dt} = \frac{L}{\eta c^2}$$

$$\dot{M} = \frac{dM}{dt} = \frac{L}{L_E} \frac{4\pi G M m_p}{\sigma_e \eta c}$$

$$\int \frac{dM}{M} = \frac{L}{L_E} \frac{4\pi G m_p}{\sigma_e \eta c} \int dt$$

$$M(t) = M_0 \exp \left(\frac{t}{t_{\text{growth}}} \right)$$

Growth Time

$$\dot{M} = \frac{dM}{dt} = \frac{L}{\eta c^2}$$

$$\dot{M} = \frac{dM}{dt} = \frac{L}{L_E} \frac{4\pi G M m_p}{\sigma_e \eta c}$$

$$\int \frac{dM}{M} = \frac{L}{L_E} \frac{4\pi G m_p}{\sigma_e \eta c} \int dt$$

$$M(t) = M_0 \exp \left(\frac{t}{t_{\text{growth}}} \right)$$

$$t_{\text{growth}} = \frac{\sigma_e \eta c}{4\pi G m_p} \left(\frac{L_E}{L} \right)$$

Growth Time

$$t_{\text{growth}} = \frac{\sigma_e \eta c}{4\pi G m_p} \left(\frac{L_E}{L} \right)$$

Growth Time

$$t_{\text{growth}} = \frac{\sigma_e \eta c}{4\pi G m_p} \left(\frac{L_E}{L} \right)$$

$$t_{\text{growth}} = 3.7 \times 10^8 \eta \left(\frac{L_E}{L} \right) [\text{yr}]$$

Growth Time

$$t_{\text{growth}} = \frac{\sigma_e \eta c}{4\pi G m_p} \left(\frac{L_E}{L} \right)$$

$$t_{\text{growth}} = 3.7 \times 10^8 \eta \left(\frac{L_E}{L} \right) [\text{yr}]$$

For $L \sim L_E$, the BH grows exponentially on time scales of the order $\sim 10^8$ yr.

Radiation Temperature

The *continuum spectrum* of the emitted radiation is characterized by a temperature

$$T_{rad} = \frac{h\bar{\nu}}{k_B}$$

Radiation Temperature

The *continuum spectrum* of the emitted radiation is characterized by a temperature

$$T_{rad} = \frac{h\bar{\nu}}{k_B}$$

$\bar{\nu}$: frequency of a typical (average) photon

Black Body Temperature

For a source with accretion luminosity L and radius r , it is defined the *blackbody temperature* through

$$F = \sigma T_{eff}^4$$

Black Body Temperature

For a source with accretion luminosity L and radius r , it is defined the *blackbody temperature* through

$$F = \sigma T_{eff}^4$$

$$\sigma = 5.6 \times 10^{-5} \frac{\text{erg}}{\text{cm}^2 \cdot \text{s} \cdot \text{K}^4 \cdot \text{sr}}$$

Steffan-Boltzman Constant

Black Body Temperature

For a source with accretion luminosity L and radius r , it is defined the *blackbody temperature* through

$$F = \sigma T_{eff}^4$$

$$\sigma = 5.6 \times 10^{-5} \frac{\text{erg}}{\text{cm}^2 \cdot \text{s} \cdot \text{K}^4 \cdot \text{sr}}$$

Steffan-Boltzman Constant

$$T_{eff} = \left(\frac{L}{4\pi r^2 \sigma} \right)^{1/4}$$

Black Body Temperature

Using $L = \frac{GM}{r} \dot{M}$,

Black Body Temperature

Using $L = \frac{GM}{r} \dot{M}$,

$$T_{eff} = \left(\frac{GM\dot{M}}{4\pi r^3 \sigma} \right)^{1/4}$$

Black Body Temperature

Using $L = \frac{GM}{r} \dot{M}$,

$$T_{eff} = \left(\frac{GM\dot{M}}{4\pi r^3 \sigma} \right)^{1/4}$$

$$T_{eff} = 1.01 \times 10^6 M_8^{-1/4} \left(\frac{\dot{M}}{\dot{M}_E} \right)^{1/4} \left(\frac{r}{r_S} \right)^{-3/4}$$

compactness

One way to estimate the compactness of a source is using the luminosity and the effective surface temperature.

$$r_{BB} = \sqrt{\frac{L}{4\pi\sigma T_{eff}^4}}$$

Example

Consider a system in our galaxy with $L = 10^{37} \text{ erg} \cdot \text{s}^{-1}$

Example

Consider a system in our galaxy with $L = 10^{37} \text{ erg} \cdot \text{s}^{-1}$ From the Eddington limit, the mass of the central object must be

$$M \geq \frac{L}{1.26 \times 10^{38}} M_{\odot} \sim \frac{10^{37}}{10^{38}} M_{\odot} \sim 0.1 M_{\odot}$$

Example

If the radiation is in the optical-UV,

<https://rechneronline.de/spectrum>

Example

If the radiation is in the optical-UV,

$$\nu_{max} \sim 10^{15} \text{ Hz}$$

<https://rechneronline.de/spectrum>

Example

If the radiation is in the optical-UV,

$$\nu_{max} \sim 10^{15} \text{ Hz}$$

$$T_{eff} \sim T_c = \frac{10^{15}}{5.88 \times 10^{10}} \sim 10^5 \text{ K}$$

<https://rechneronline.de/spectrum>

Example

If the radiation is in the optical-UV,

$$\nu_{max} \sim 10^{15} \text{Hz}$$

$$T_{eff} \sim T_c = \frac{10^{15}}{5.88 \times 10^{10}} \sim 10^5 \text{K}$$

$$r_{bb} \sim 10^{12} \text{cm} \sim 10^7 \text{km}$$

<https://rechneronline.de/spectrum>

Example

If the radiation is in the optical-UV,

$$\nu_{max} \sim 10^{15} \text{ Hz}$$

$$T_{eff} \sim T_c = \frac{10^{15}}{5.88 \times 10^{10}} \sim 10^5 \text{ K}$$

$$r_{bb} \sim 10^{12} \text{ cm} \sim 10^7 \text{ km}$$

Typical size of a Star!

<https://rechneronline.de/spectrum>

Example

If the radiation is in soft X-rays at 1 keV,

<https://rechneronline.de/spectrum>

Example

If the radiation is in soft X-rays at 1 keV,

$$\nu_{max} \sim 10^{17} \text{Hz}$$

<https://rechneronline.de/spectrum>

Example

If the radiation is in soft X-rays at 1 keV,

$$\nu_{max} \sim 10^{17} \text{Hz}$$

$$T_{eff} \sim T_c = \frac{10^{17}}{5.88 \times 10^{10}} \sim 10^7 \text{K}$$

<https://rechneronline.de/spectrum>

Example

If the radiation is in soft X-rays at 1 keV,

$$\nu_{max} \sim 10^{17} \text{Hz}$$

$$T_{eff} \sim T_c = \frac{10^{17}}{5.88 \times 10^{10}} \sim 10^7 \text{K}$$

$$r_{bb} \sim 10^6 \text{cm} \sim 10 \text{km}$$

<https://rechneronline.de/spectrum>

Example

If the radiation is in soft X-rays at 1 keV,

$$\nu_{max} \sim 10^{17} \text{Hz}$$

$$T_{eff} \sim T_c = \frac{10^{17}}{5.88 \times 10^{10}} \sim 10^7 \text{K}$$

$$r_{bb} \sim 10^6 \text{cm} \sim 10 \text{km}$$

Typical size of a neutron star or a BH! <https://rechneronline.de/spectrum>

Virial Temperature

T_{th} : Temperature reached by the accreted material if all the gravitational energy is transformed into thermal energy.

Virial Temperature

T_{th} : Temperature reached by the accreted material if all the gravitational energy is transformed into thermal energy.

$2 \langle K \rangle + \langle U \rangle = 0$: Virial Theorem

Virial Temperature

T_{th} : Temperature reached by the accreted material if all the gravitational energy is transformed into thermal energy.

$2\langle K\rangle + \langle U\rangle = 0$: Virial Theorem

$$\langle U\rangle = \frac{GM(m_p + m_e)}{r} \sim \frac{GMm_p}{r}$$

Virial Temperature

T_{th} : Temperature reached by the accreted material if all the gravitational energy is transformed into thermal energy.

$2 \langle K \rangle + \langle U \rangle = 0$: Virial Theorem

$$\langle U \rangle = \frac{GM(m_p + m_e)}{r} \sim \frac{GMm_p}{r}$$

$$2 \langle K \rangle \sim 2 \times \frac{3}{2} k_B T_{th}$$

Virial Temperature

T_{th} : Temperature reached by the accreted material if all the gravitational energy is transformed into thermal energy.

$2 \langle K \rangle + \langle U \rangle = 0$: Virial Theorem

$$\langle U \rangle = \frac{GM(m_p + m_e)}{r} \sim \frac{GMm_p}{r}$$

$$2 \langle K \rangle \sim 2 \times \frac{3}{2} k_B T_{th}$$

$$T_{th} = \frac{GMm_p}{3k_B r} = T_{vir}$$

Temperatures

If the accretion energy is converted directly into radiation escaping without interaction,

Temperatures

If the accretion energy is converted directly into radiation escaping without interaction,

$$T_{rad} \sim T_{th}$$

Temperatures

If the accretion energy is converted directly into radiation escaping without interaction,

$$T_{rad} \sim T_{th}$$

If the accretion flow is optically thick, the radiation reaches thermal equilibrium with the accreted material before escaping out to the observer,

Temperatures

If the accretion energy is converted directly into radiation escaping without interaction,

$$T_{rad} \sim T_{th}$$

If the accretion flow is optically thick, the radiation reaches thermal equilibrium with the accreted material before escaping out to the observer,

$$T_{rad} \sim T_{eff}$$

Temperatures

If the accretion energy is converted directly into radiation escaping without interaction,

$$T_{rad} \sim T_{th}$$

If the accretion flow is optically thick, the radiation reaches thermal equilibrium with the accreted material before escaping out to the observer,

$$T_{rad} \sim T_{eff}$$

In general,

$$T_{eff} \lesssim T_{rad} \lesssim T_{th}$$

Outline for Part 2

1. Accretion Basics

- 1.1 Spherical Accretion
- 1.2 Eddington Luminosity
- 1.3 Estimation of the Central Mass
- 1.4 Eddington Accretion Rate
- 1.5 Growth Time
- 1.6 Temperatures
- 1.7 Compactness

2. Geodesics in Kerr Spacetime

- 2.1 Hamilton-Jacobi Formulation
- 2.2 Equations of Motion
- 2.3 Imaging a Black Hole

Hamilton-Jacobi Formulation

Hamilton-Jacobi Formulation

$$\mathcal{L} = \frac{1}{2} g_{\mu\nu} \dot{x}^\mu \dot{x}^\nu$$

Hamilton-Jacobi Formulation

$$\mathcal{L} = \frac{1}{2} g_{\mu\nu} \dot{x}^\mu \dot{x}^\nu$$

$$p_\mu = \frac{\partial \mathcal{L}}{\partial \dot{x}^\mu} = g_{\mu\nu} \dot{x}^\nu$$

Hamilton-Jacobi Formulation

$$\mathcal{L} = \frac{1}{2} g_{\mu\nu} \dot{x}^\mu \dot{x}^\nu$$

$$p_\mu = \frac{\partial \mathcal{L}}{\partial \dot{x}^\mu} = g_{\mu\nu} \dot{x}^\nu$$

$$\mathcal{H} = \frac{1}{2} g^{\mu\nu} p_\mu p_\nu$$

Hamilton-Jacobi Formulation

Hamilton's principal function

$$S = S(x^\mu; \lambda)$$

Hamilton-Jacobi Formulation

Hamilton's principal function

$$S = S(x^\mu; \lambda)$$

$$p_\mu = \frac{\partial S}{\partial x^\mu}$$

Hamilton-Jacobi Formulation

Hamilton's principal function

$$S = S(x^\mu; \lambda)$$

$$p_\mu = \frac{\partial S}{\partial x^\mu}$$

Hamilton-Jacobi Equation

Hamilton-Jacobi Formulation

Hamilton's principal function

$$S = S(x^\mu; \lambda)$$

$$p_\mu = \frac{\partial S}{\partial x^\mu}$$

Hamilton-Jacobi Equation

$$\frac{1}{2} g^{\mu\nu} \frac{\partial S}{\partial x^\mu} \frac{\partial S}{\partial x^\nu} - \frac{\partial S}{\partial \lambda} = 0$$

Kerr's Solution

Boyer-Lindquist coordinates: (t, r, θ, ϕ)

$$ds^2 = -\frac{\Delta - a^2 \sin^2 \theta}{\varrho} dt^2 - \left(\frac{r^2 + a^2 - \Delta}{\varrho} \right) 2a \sin^2 \theta dt d\phi \\ + \frac{\varrho}{\Delta} dr^2 + \varrho d\theta^2 + \left(\frac{(r^2 + a^2)^2 - \Delta a^2 \sin^2 \theta}{\varrho} \right) \sin^2 \theta d\phi^2.$$

Kerr's Solution

Boyer-Lindquist coordinates: (t, r, θ, ϕ)

$$ds^2 = -\frac{\Delta - a^2 \sin^2 \theta}{\varrho} dt^2 - \left(\frac{r^2 + a^2 - \Delta}{\varrho} \right) 2a \sin^2 \theta dt d\phi \\ + \frac{\varrho}{\Delta} dr^2 + \varrho d\theta^2 + \left(\frac{(r^2 + a^2)^2 - \Delta a^2 \sin^2 \theta}{\varrho} \right) \sin^2 \theta d\phi^2.$$

$$\varrho = r^2 + a^2 \cos^2 \theta$$

$$\Delta = r^2 - 2Mr + a^2$$

Kerr's Solution

$$\begin{aligned}\left(\frac{\partial}{\partial s}\right)^2 = & -\frac{A}{\varrho\Delta}\left(\frac{\partial}{\partial t}\right)^2 - \frac{4aMr}{\varrho\Delta}\left(\frac{\partial}{\partial t}\right)\left(\frac{\partial}{\partial\varphi}\right) + \frac{\Delta}{\varrho}\left(\frac{\partial}{\partial r}\right)^2 \\ & + \frac{1}{\varrho}\left(\frac{\partial}{\partial\theta}\right)^2 + \frac{\Delta - a^2\sin^2\theta}{\varrho\Delta\sin^2\theta}\left(\frac{\partial}{\partial\varphi}\right)^2\end{aligned}$$

Kerr's Solution

$$\left(\frac{\partial}{\partial s}\right)^2 = -\frac{A}{\varrho\Delta}\left(\frac{\partial}{\partial t}\right)^2 - \frac{4aMr}{\varrho\Delta}\left(\frac{\partial}{\partial t}\right)\left(\frac{\partial}{\partial\varphi}\right) + \frac{\Delta}{\varrho}\left(\frac{\partial}{\partial r}\right)^2 \\ + \frac{1}{\varrho}\left(\frac{\partial}{\partial\theta}\right)^2 + \frac{\Delta - a^2\sin^2\theta}{\varrho\Delta\sin^2\theta}\left(\frac{\partial}{\partial\varphi}\right)^2$$

$$A = (r^2 + a^2)^2 - a^2\Delta\sin^2\theta$$

$$\varrho = r^2 + a^2\cos^2\theta$$

$$\Delta = r^2 - 2Mr + a^2$$

Hamilton-Jacobi Formulation

Hamilton-Jacobi Equation

$$2\frac{\partial S}{\partial \lambda} = g^{\mu\nu} \frac{\partial S}{\partial x^\mu} \frac{\partial S}{\partial x^\nu}$$

Hamilton-Jacobi Formulation

Hamilton-Jacobi Equation

$$2\frac{\partial S}{\partial \lambda} = g^{\mu\nu} \frac{\partial S}{\partial x^\mu} \frac{\partial S}{\partial x^\nu}$$

$$\begin{aligned} 2\frac{\partial S}{\partial \lambda} = & -\frac{A}{\rho\Delta} \left(\frac{\partial S}{\partial t} \right)^2 - \frac{4aMr}{\rho\Delta} \left(\frac{\partial S}{\partial t} \right) \left(\frac{\partial S}{\partial \varphi} \right) + \frac{\Delta}{\rho} \left(\frac{\partial S}{\partial r} \right)^2 \\ & + \frac{1}{\rho} \left(\frac{\partial S}{\partial \theta} \right)^2 + \frac{\Delta - a^2 \sin^2 \theta}{\rho\Delta \sin^2 \theta} \left(\frac{\partial S}{\partial \varphi} \right)^2 \end{aligned}$$

Hamilton-Jacobi Formulation

Hamilton Principal Function

$$S = \frac{1}{2}\lambda\delta - \varepsilon t + \ell_z\varphi + S_r(\theta) + S_\theta(\theta)$$

Separation of the Hamilton-Jacobi Equation. Carter Constant

$$\Delta \left(\frac{dS_r}{dr} \right)^2 - \frac{1}{\Delta} [(r^2 + a^2)\varepsilon - a\ell_z]^2 + (\ell_z - a\varepsilon)^2 + \delta r^2 =$$
$$- \left(\frac{dS_\theta}{d\theta} \right)^2 - \left(\frac{\ell_z^2}{\sin^2 \theta} - a^2\varepsilon^2 + \delta a^2 \right) \cos^2 \theta = \mathcal{C}$$

Hamilton-Jacobi Formulation

Separation of the Hamilton-Jacobi Equation. Carter Constant

$$\Delta \left(\frac{dS_r}{dr} \right)^2 = \frac{1}{\Delta} [(r^2 + a^2)\varepsilon - al_z]^2 - [\mathcal{C} + (\ell_z - a\varepsilon)^2 + \delta r^2]$$
$$\left(\frac{dS_\theta}{d\theta} \right)^2 = \mathcal{C} - \left(\frac{\ell_z^2}{\sin^2 \theta} - a^2 \varepsilon^2 + \delta a^2 \right) \cos^2 \theta$$

Hamilton-Jacobi Formulation

$$S_r = \int \frac{\sqrt{R(r')}}{\Delta} dr'$$
$$S_\theta = \int \sqrt{\Theta(\theta')} d\theta'$$

Hamilton-Jacobi Formulation

$$S_r = \int \frac{\sqrt{R(r')}}{\Delta} dr'$$
$$S_\theta = \int \sqrt{\Theta(\theta')} d\theta'$$

$$R(r) = [(r^2 + a^2)\varepsilon - a\ell_z]^2 - \Delta [\mathcal{C} + (\ell_z - a\varepsilon)^2 + \delta r^2]$$

$$\Theta(\theta) = \mathcal{C} - \left[\frac{\ell_z^2}{\sin^2 \theta} + a^2 (\delta - \varepsilon^2) \right] \cos^2 \theta$$

Carter Constant

$$\left(\frac{dS_\theta}{d\theta}\right)^2 = \mathcal{C} - \left(\frac{\ell_z^2}{\sin^2 \theta} - a^2 \varepsilon^2 + \delta a^2\right) \cos^2 \theta$$

Carter Constant

$$\left(\frac{dS_\theta}{d\theta}\right)^2 = \mathcal{C} - \left(\frac{\ell_z^2}{\sin^2 \theta} - a^2 \varepsilon^2 + \delta a^2\right) \cos^2 \theta$$

$$\mathcal{C} = p_\theta^2 + p_\varphi^2 \cot^2 \theta + a^2(\delta - \varepsilon^2) \cos^2 \theta$$

Carter Constant

Schwarzschild:

$$\mathcal{C} = \left(p_\theta^2 + \frac{p_\phi^2}{\sin^2 \theta} \right) - p_\phi^2 = \ell^2 - \ell_z^2$$

where $\ell = p_\theta^2 + \frac{p_\phi^2}{\sin^2 \theta}$ is the total angular momentum.

Carter Constant

Kerr:

Carter Constant

Kerr:

- \mathcal{C} has not a direct physical interpretation.
- $\mathcal{C} = 0$ implies that the motion is in the equatorial plane.

Equations of Motion

Equations of Motion

Hamilton Canonical Equations

Equations of Motion

Hamilton Canonical Equations

$$\dot{x}^\mu = p^\mu = g^{\mu\nu} p_\nu = g^{\mu\nu} \frac{\partial S}{\partial x^\nu}$$

Equations of Motion

$$\varrho^2 \dot{r}^2 = R$$

$$\varrho^2 \dot{\theta}^2 = \Theta$$

$$\varrho \dot{\phi} = \frac{1}{\Delta} \left[2aMr\varepsilon + (\varrho - 2Mr) \frac{\ell_z}{\sin^2 \theta} \right]$$

$$\varrho \dot{t} = \frac{1}{\Delta} [A\varepsilon + 2aMr\ell_z]$$

Equations of Motion

$$\varrho^2 \dot{r}^2 = R$$

$$\varrho^2 \dot{\theta}^2 = \Theta$$

$$\varrho \dot{\phi} = \frac{1}{\Delta} \left[2aMr\varepsilon + (\varrho - 2Mr) \frac{\ell_z}{\sin^2 \theta} \right]$$

$$\varrho \dot{t} = \frac{1}{\Delta} [A\varepsilon + 2aMr\ell_z]$$

$$R(r) = [(r^2 + a^2)\varepsilon - a\ell_z]^2 - \Delta [\mathcal{C} + (\ell_z - a\varepsilon)^2 + \delta r^2]$$

$$\Theta(\theta) = \mathcal{C} - \left[\frac{\ell_z^2}{\sin^2 \theta} + a^2 (\delta - \varepsilon^2) \right] \cos^2 \theta$$

$$A = (r^2 + a^2)^2 - a^2 \Delta \sin^2 \theta$$

$$\varrho = r^2 + a^2 \cos^2 \theta$$

$$\Delta = r^2 - 2Mr + a^2$$

Imaging a Black Hole

Image Plane of a Distant Observer

Image Plane of a Distant Observer


- A distant observer receives the electromagnetic radiation from the accretion disk, around the black hole.

Image Plane of a Distant Observer

- A distant observer receives the electromagnetic radiation from the accretion disk, around the black hole.
- It is usual to define a plane of observation and consider the photons with momentum orthogonal to the plane. These photons' trajectories are integrated backwards in time to find the position of the emission point in the disk.

Image Plane of a Distant Observer

Fig. 3.5 The Cartesian coordinates (x, y, z) are centered at the black hole, while the Cartesian coordinates (X, Y, Z) are for the image plane of the distant observer, who is located at the distant D from the *black hole* and with the inclination angle i . From [1]



Coordinate Systems

- (X, Y, Z) : Cartesian coordinates in the image plane

Coordinate Systems

- (X, Y, Z) : Cartesian coordinates in the image plane
- (x, y, z) : Cartesian coordinates centered at the black hole.

Coordinate Systems

- (X, Y, Z) : Cartesian coordinates in the image plane
- (x, y, z) : Cartesian coordinates centered at the black hole.
- i : Inclination angle of the observer with respect to the z direction.

Coordinate Systems

- (X, Y, Z) : Cartesian coordinates in the image plane
- (x, y, z) : Cartesian coordinates centered at the black hole.
- i : Inclination angle of the observer with respect to the z direction.
- D : Distance observer-black hole.

Coordinate transformations

$$x = D \sin i - Y \cos i + Z \sin i$$

$$y = X$$

$$z = D \cos i + Y \sin i + Z \cos i$$

$$r = \sqrt{x^2 + y^2 + z^2}$$

$$\theta = \arccos \left(\frac{z}{r} \right)$$

$$\phi = \arctan \left(\frac{z}{r} \right)$$

Photon Trajectory Tracing

Photon Trajectory Tracing

Consider a photon received at $(X_0, Y_0, 0)$ with 3-momentum $\mathbf{k}_0 = -k_0 \hat{\mathbf{Z}}$, i.e. perpendicular to the observer plane.

The initial conditions for the position of the photon (to trace back the trajectory), as seen from the black hole and in spherical coordinates, are

$$t_0 = 0$$

$$r_0 = \sqrt{X_0^2 + Y_0^2 + D^2}$$

$$\theta_0 = \arccos \left(\frac{Y_0 \sin i + D \cos i}{r_0} \right)$$

$$\varphi_0 = \arctan \left(\frac{X_0}{D \sin i - Y_0 \cos i} \right)$$

Photon Trajectory Tracing

The initial conditions for the 4-momentum of the photon k^μ (to trace back the trajectory), as seen from the black hole and in spherical coordinates, are calculated with the transformation law,

$$k^\mu = \frac{\partial x^\mu}{\partial \bar{x}^\alpha} \bar{k}^\alpha$$

Photon Trajectory Tracing

The initial conditions for the 4-momentum of the photon k^μ (to trace back the trajectory), as seen from the black hole and in spherical coordinates, are calculated with the transformation law,

$$k^\mu = \frac{\partial x^\mu}{\partial \bar{x}^\alpha} \bar{k}^\alpha$$

$$k_0^r = -\frac{D}{r} k_0$$

$$k_0^\theta = \frac{\cos i - (Y_0 \sin i + D \cos i) \frac{D}{r_0^2}}{\sqrt{X_0^2 + (D \sin i - Y_0 \cos i)^2}} k_0$$

$$k_0^\phi = \frac{X_0 \sin i}{X_0^2 + (D \sin i - Y_0 \cos i)^2} k_0$$

Photon Trajectory Tracing

The k_0^t component of the initial 4-momentum is calculated by the condition $g_{\mu\nu}g^{\mu}k'^{\nu} = 0$,

$$k_0^t = \sqrt{(k_0^r)^2 + r_0^2(k_0^\theta)^2 + r_0^2 \sin^2 \theta_0 (k_0^\phi)^2}$$

Photon Trajectory Tracing

Given the initial conditions for position and momentum, it is possible to trace the trajectory of any photon in the observer plane back to the accretion disk.

Non-Coordinate Basis

Tetrads

Introduce a non-coordinate basis or orthonormal tetrad,

Tetrads

Introduce a non-coordinate basis or orthonormal tetrad,

$$\begin{aligned}\mathbf{E}_{(a)} &= E_{(a)}^{\mu} \partial_{\mu} \\ \mathbf{E}^{(a)} &= E_{\mu}^{(a)} dx^{\mu},\end{aligned}$$

subject to the conditions

$$\begin{aligned}\eta_{(a)(b)} &= E_{(a)}^{\mu} E_{(b)}^{\nu} g_{\mu\nu} \\ \eta^{(a)(b)} &= E_{\mu}^{(a)} E_{\nu}^{(b)} g^{\mu\nu}\end{aligned}$$

and $\det|E_{(a)}^{\mu}| > 0$ (to preserve the orientation).

Tetrads

Components of a vector in the orthonormal tetrad basis,

Tetrads

Components of a vector in the orthonormal tetrad basis,

$$V^{(a)} = E_{\mu}^{(a)} V^{\mu}$$

$$V_{(a)} = E_{(a)}^{\mu} V_{\mu}$$

General Metric

Consider a general stationary, axisymmetric, asymptotically flat metric

$$ds^2 = -e^{2\alpha(r)} dt^2 + e^{2\beta(r)} dr^2 + e^{2\gamma(r,\theta)} d\theta^2 + e^{2\epsilon(r,\theta)} (d\phi - \omega dt)^2$$

General Metric

Consider a general stationary, axisymmetric, asymptotically flat metric

$$ds^2 = -e^{2\alpha(r)} dt^2 + e^{2\beta(r)} dr^2 + e^{2\gamma(r,\theta)} d\theta^2 + e^{2\epsilon(r,\theta)} (d\phi - \omega dt)^2$$

We identify the *locally non-rotating observers* as those whose world-lines have

General Metric

Consider a general stationary, axisymmetric, asymptotically flat metric

$$ds^2 = -e^{2\alpha(r)} dt^2 + e^{2\beta(r)} dr^2 + e^{2\gamma(r,\theta)} d\theta^2 + e^{2\epsilon(r,\theta)} (d\phi - \omega dt)^2$$

We identify the *locally non-rotating observers* as those whose world-lines have

$$r = \text{constant}$$

$$\theta = \text{constant}$$

$$\phi = \omega t + \text{constant}$$

General Metric

The non-coordinate basis of the locally non-rotating observers is given by the tetrad

$$E_{(t)}^\mu = (e^{-\beta}, 0, 0, \omega e^{-\beta})$$

$$E_{(r)}^\mu = (0, e^{-\alpha}, 0, 0)$$

$$E_{(\theta)}^\mu = (0, 0, e^{-\gamma}, 0)$$

$$E_{(\phi)}^\mu = (0, 0, 0, e^{-\epsilon})$$

General Metric

and the dual basis,

$$E_{\mu}^{(t)} = (e^{\beta}, 0, 0, 0)$$

$$E_{\mu}^{(r)} = (0, e^{\alpha}, 0, 0)$$

$$E_{\mu}^{(\theta)} = (0, 0, e^{\gamma}, 0)$$

$$E_{\mu}^{(\varphi)} = (-\omega e^{\epsilon}, 0, 0, e^{\epsilon})$$

Kerr Metric

For the particular case of the Kerr metric the tetrad describing the non-coordinate basis of the locally non-rotating observers is

$$E_{(t)}^{\mu} = \left(\sqrt{\frac{A}{\varrho\Delta}}, 0, 0, \frac{2Mar}{\sqrt{A\varrho\Delta}} \right)$$

$$E_{(r)}^{\mu} = \left(0, \sqrt{\frac{\Delta}{\varrho}}, 0, 0 \right)$$

$$E_{(\theta)}^{\mu} = \left(0, 0, \frac{1}{\sqrt{\varrho}}, 0 \right)$$

$$E_{(\varphi)}^{\mu} = \left(0, 0, 0, \frac{1}{\sin\theta} \sqrt{\frac{\varrho}{A}} \right)$$

Kerr Metric

and the dual basis is

$$E_{\mu}^{(t)} = \left(\sqrt{\frac{\varrho \Delta}{A}}, 0, 0, 0 \right)$$

$$E_{\mu}^{(r)} = \left(0, \sqrt{\frac{\varrho}{\Delta}}, 0, 0 \right)$$

$$E_{\mu}^{(\theta)} = (0, 0, \sqrt{\varrho}, 0)$$

$$E_{\mu}^{(\varphi)} = \left(-\frac{2Mar \sin \theta}{\sqrt{\varrho A}}, 0, 0, \sqrt{\frac{A}{\varrho}} \sin \theta \right)$$

Momentum Components in the Non-Coordinate Basis

The momentum components of a particle moving in Kerr's spacetime are

$$p_\mu = \frac{\partial S}{\partial x^\mu}$$

Momentum Components in the Non-Coordinate Basis

The momentum components of a particle moving in Kerr's spacetime are

$$p_\mu = \frac{\partial S}{\partial x^\mu}$$

$$p_t = -\varepsilon$$

$$p_r = \frac{\sqrt{R}}{\Delta}$$

$$p_\theta = \sqrt{\Theta}$$

$$p_\phi = \ell_z$$

Momentum Components in the Non-Coordinate Basis

In the non-coordinate basis, the momentum components of a particle are

$$p^{(a)} = E_{\mu}^{(a)} p^{\mu} = \eta^{(a)(b)} E_{(b)}^{\mu} p_{\mu}$$

Momentum Components in the Non-Coordinate Basis

In the non-coordinate basis, the momentum components of a particle are

$$p^{(a)} = E_{\mu}^{(a)} p^{\mu} = \eta^{(a)(b)} E_{(b)}^{\mu} p_{\mu}$$

$$p^{(t)} = -E_{(t)}^{\mu} p_{\mu}$$

$$p^r = E_{(r)}^{\mu} p_{\mu}$$

$$p^{\theta} = E_{(\theta)}^{\mu} p_{\mu}$$

$$p^{\varphi} = E_{(\varphi)}^{\mu} p_{\mu}$$

Celestial Coordinates

Celestial Coordinates

The position of a photon in the image plane of the distant observer is given by the coordinates

$$\begin{cases} X_0 &= \alpha = \lim_{r \rightarrow \infty} \left(\frac{rp^{(\varphi)}}{p^{(t)}} \right) \\ Y_0 &= \beta = \lim_{r \rightarrow \infty} \left(\frac{rp^{(\theta)}}{p^{(t)}} \right) \end{cases}$$

Celestial Coordinates

$$\begin{cases} \alpha &= -\xi \csc i \\ \beta &= \pm \sqrt{\eta + a^2 \cos^2 i - \xi^2 \cot^2 i} \end{cases}$$

Celestial Coordinates

$$\begin{cases} \alpha &= -\xi \csc i \\ \beta &= \pm \sqrt{\eta + a^2 \cos^2 i - \xi^2 \cot^2 i} \end{cases}$$

$$\begin{cases} \xi &= \frac{l_z}{\varepsilon} \\ \eta &= \frac{\mathcal{C}}{\varepsilon^2} \end{cases}$$

Black Hole's Shadow

Effective Potential for Photons

$$R(r) = [(r^2 + a^2)\varepsilon - al_z]^2 - \Delta [\mathcal{C} + (l_z - a\varepsilon)^2 + \delta r^2]$$
$$\Theta(\theta) = \mathcal{C} - \left[\frac{\ell_z^2}{\sin^2 \theta} + a^2 (\delta - \varepsilon^2) \right] \cos^2 \theta$$

Effective Potential for Photons

$$R(r) = [(r^2 + a^2)\varepsilon - al_z]^2 - \Delta [\mathcal{C} + (\ell_z - a\varepsilon)^2 + \delta r^2]$$
$$\Theta(\theta) = \mathcal{C} - \left[\frac{\ell_z^2}{\sin^2 \theta} + a^2 (\delta - \varepsilon^2) \right] \cos^2 \theta$$

Consider these expressions for photons, $\delta = 0$, and using the quantities

$$\begin{cases} \xi &= \frac{\ell_z}{\varepsilon} \\ \eta &= \frac{\mathcal{C}}{\varepsilon^2} \end{cases}$$

Effective Potential for Photons

$$R(r) = [r^2 + a^2 - a\xi]^2 \varepsilon^2 - \Delta [\eta + (\xi - a)^2] \varepsilon^2$$
$$\Theta(\theta) = \eta \varepsilon^2 - \left[\frac{\xi^2}{\sin^2 \theta} - a^2 \right] \varepsilon^2 \cos^2 \theta$$

Effective Potential for Photons

$$\mathcal{R}(r) = \frac{R(r)}{\xi^2} = [r^2 + a^2 - a\xi]^2 - \Delta [\eta + (\xi - a)^2]$$
$$\mathcal{Y}(\theta) = \frac{\Theta(\theta)}{\xi^2} = [\eta + (\xi - a)^2] - [a \sin \theta - \xi \csc \theta]^2$$

Effective Potential for Photons

$$\mathcal{R}(r) = \frac{R(r)}{\xi^2} = [r^2 + a^2 - a\xi]^2 - \Delta [\eta + (\xi - a)^2]$$
$$\mathcal{Y}(\theta) = \frac{\Theta(\theta)}{\xi^2} = [\eta + (\xi - a)^2] - [a \sin \theta - \xi \csc \theta]^2$$

$$\Delta = r^2 - 2Mr + a^2$$

Equations of Motion for Photons

$$\varrho^2 \dot{r}^2 = \mathcal{R}$$

$$\varrho^2 \dot{\theta}^2 = \mathcal{G}$$

$$\varrho \dot{\phi} = \frac{1}{\Delta} \left[2aMr + \frac{\xi(\varrho - 2Mr)}{\sin^2 \theta} \right]$$

$$\varrho \dot{t} = \frac{1}{\Delta} [A + 2aMr\xi]$$

Photon Sphere

Circular motion of photons: $\dot{r} = 0$

Photon Sphere

Circular motion of photons: $\dot{r} = 0$

$$\begin{cases} \mathcal{R} &= 0 \\ \partial_r \mathcal{R} &= 0 \end{cases}$$

Photon Sphere

Solving for ξ and η we obtain these quantities for the circular orbit as functions of the parameter r ,

Photon Sphere

Solving for ξ and η we obtain these quantities for the circular orbit as functions of the parameter r ,

$$\xi_c = \frac{M(r^2 - a^2) - r\Delta}{a(r - M)}$$
$$\eta_c = \frac{r^3 [4M\Delta - r(r - M)^2]}{a^2(r - M)^2}$$

Photon Sphere

There are three possible cases regarding the stability of circular orbits of photons

Photon Sphere

There are three possible cases regarding the stability of circular orbits of photons

1. If $\partial_r^2 \mathcal{R} > 0$: Stable circular orbits

Photon Sphere

There are three possible cases regarding the stability of circular orbits of photons

1. If $\partial_r^2 \mathcal{R} > 0$: Stable circular orbits
2. If $\partial_r^2 \mathcal{R} < 0$: Unstable circular orbits. The photon straddles the boundary between two regions: $\partial_r \mathcal{R} = 0$; if perturbed one way it falls into the horizon, if perturbed the other way it flies outwards.

Photon Sphere

There are three possible cases regarding the stability of circular orbits of photons

1. If $\partial_r^2 \mathcal{R} > 0$: Stable circular orbits
2. If $\partial_r^2 \mathcal{R} < 0$: Unstable circular orbits. The photon straddles the boundary between two regions: $\partial_r \mathcal{R} = 0$; if perturbed one way it falls into the horizon, if perturbed the other way it flies outwards.
3. If $\partial_r^2 \mathcal{R} = 0$: Marginally stable circular orbit (Photon Sphere).
 $r = r_{ps}$.

Black Hole's Shadow

$$\alpha = -\xi_c \csc i$$

$$\beta = \sqrt{\eta_c + a^2 \cos^2 i - \xi_c^2 \cot^2 i}$$

Black Hole's Shadow

$$\alpha = -\xi_c \csc i$$

$$\beta = \sqrt{\eta_c + a^2 \cos^2 i - \xi_c^2 \cot^2 i}$$

$$\xi_c = \frac{M(r^2 - a^2) - r\Delta}{a(r - M)}$$

$$\eta_c = \frac{r^3 [4M\Delta - r(r - M)^2]}{a^2(r - M)^2}$$

Black Hole's Shadow

Schwarzschild's Black Hole:

$$\alpha^2 + \beta^2 = R_{shadow}^2$$

Black Hole's Shadow

Schwarzschild's Black Hole:

$$\alpha^2 + \beta^2 = R_{shadow}^2$$

$$R_{shadow} = \sqrt{27}M$$

Shadow of Sagittarius A^{*}

$$M_{\text{SgrA}^*} = 4 \times 10^6 M_{\odot}$$

Shadow of Sagittarius A^{*}

$$M_{\text{SgrA}^*} = 4 \times 10^6 M_{\odot}$$

$$r_H = 2M = \frac{2GM}{c^2}$$

Shadow of Sagittarius A^{*}

$$M_{\text{SgrA}^*} = 4 \times 10^6 M_{\odot}$$

$$r_H = 2M = \frac{2GM}{c^2}$$

$$r_H = 1.2 \times 10^{10} \text{ m.} = 1.2 \times 10^7 \text{ km.}$$

Shadow of Sagittarius A^{*}

$$M_{\text{SgrA}^*} = 4 \times 10^6 M_{\odot}$$

$$r_H = 2M = \frac{2GM}{c^2}$$

$$r_H = 1.2 \times 10^{10} \text{ m.} = 1.2 \times 10^7 \text{ km.}$$

$$r_H = 0.1 \text{ AU}$$

Shadow of Sagittarius A^{*}

$$M_{\text{SgrA}^*} = 4 \times 10^6 M_{\odot}$$

$$r_H = 2M = \frac{2GM}{c^2}$$

$$r_H = 1.2 \times 10^{10} \text{ m.} = 1.2 \times 10^7 \text{ km.}$$

$$r_H = 0.1 \text{ AU}$$

$$R_{\text{shadow}} = \sqrt{27M^2} = 3\sqrt{3}M$$

Shadow of Sagittarius A^{*}

$$M_{\text{SgrA}^*} = 4 \times 10^6 M_{\odot}$$

$$r_H = 2M = \frac{2GM}{c^2}$$

$$r_H = 1.2 \times 10^{10} \text{ m.} = 1.2 \times 10^7 \text{ km.}$$

$$r_H = 0.1 \text{ AU}$$

$$R_{\text{shadow}} = \sqrt{27M^2} = 3\sqrt{3}M$$

$$R_{\text{shadow}} = 3\sqrt{3}\frac{GM}{c^2} = 3 \times 10^{10} \text{ m.}$$

Shadow of Sagittarius A^{*}

$$M_{\text{SgrA}^*} = 4 \times 10^6 M_{\odot}$$

$$r_H = 2M = \frac{2GM}{c^2}$$

$$r_H = 1.2 \times 10^{10} \text{ m.} = 1.2 \times 10^7 \text{ km.}$$

$$r_H = 0.1 \text{ AU}$$

$$R_{\text{shadow}} = \sqrt{27M^2} = 3\sqrt{3}M$$

$$R_{\text{shadow}} = 3\sqrt{3}\frac{GM}{c^2} = 3 \times 10^{10} \text{ m.}$$

$$R_{\text{shadow}} = 3 \times 10^7 \text{ km.} = 0.2 \text{ AU}$$

Shadow of Sagittarius A^{*}

$$R_{shadow} = 0.2 \text{ AU} = 9.7 \times 10^{-10} \text{ kpc}$$

Shadow of Sagittarius A^{*}

$$R_{shadow} = 0.2 \text{ AU} = 9.7 \times 10^{-10} \text{ kpc}$$

$$D = 8 \text{ kpc}$$

$$\theta_{shadow} = \frac{R_{shadow}}{D}$$

Shadow of Sagittarius A^{*}

$$R_{shadow} = 0.2 \text{ AU} = 9.7 \times 10^{-10} \text{ kpc}$$

$$D = 8 \text{ kpc}$$

$$\theta_{shadow} = \frac{R_{shadow}}{D}$$

$$\theta_{shadow} = 1.2 \times 10^{-11} \text{ rad}$$

Shadow of Sagittarius A^{*}

$$R_{shadow} = 0.2 \text{ AU} = 9.7 \times 10^{-10} \text{ kpc}$$

$$D = 8 \text{ kpc}$$

$$\theta_{shadow} = \frac{R_{shadow}}{D}$$

$$\theta_{shadow} = 1.2 \times 10^{-11} \text{ rad}$$

$$\theta_{shadow} = 2.5 \mu\text{arcsec}$$

Shadow of Sagittarius A^{*}

Very Long Baseline Interferometry (VLBI)

Shadow of Sagittarius A^{*}

Very Long Baseline Interferometry (VLBI)

$$\theta_{res} \sim \frac{\lambda}{d}$$

Shadow of Sagittarius A^{*}

Very Long Baseline Interferometry (VLBI)

$$\theta_{res} \sim \frac{\lambda}{d}$$

$$\lambda \sim 1 \text{ mm}$$

Shadow of Sagittarius A^{*}

Very Long Baseline Interferometry (VLBI)

$$\theta_{res} \sim \frac{\lambda}{d}$$

$$\lambda \sim 1 \text{ mm}$$

$$d \sim 10^3 \text{ km}$$

Shadow of Sagittarius A^{*}

Very Long Baseline Interferometry (VLBI)

$$\theta_{res} \sim \frac{\lambda}{d}$$

$$\lambda \sim 1 \text{ mm}$$

$$d \sim 10^3 \text{ km}$$

$$\theta_{res} \sim 10 \mu\text{arcsec}$$

Next Lecture

08. Accretion

Alternative derivation of the Celestial Coordinates

Celestial Coordinates

- A distant observer receives the electromagnetic radiation from the surroundings of the black hole.

Celestial Coordinates

- A distant observer receives the electromagnetic radiation from the surroundings of the black hole.
- Let the observer be at $r \rightarrow \infty$ with inclination angle i (between the rotation axis of the black hole and the observer's line of sight).

Celestial Coordinates

- A distant observer receives the electromagnetic radiation from the surroundings of the black hole.
- Let the observer be at $r \rightarrow \infty$ with inclination angle i (between the rotation axis of the black hole and the observer's line of sight).
- The celestial coordinates (α, β) are the apparent angular distances of the image on the celestial sphere measured by the observer.


Celestial Coordinates

- The distant observer may set up a euclidean coordinate system (x, y, z) with the black hole at the origin and its rotation axis along z .

Celestial Coordinates

- The distant observer may set up a euclidean coordinate system (x, y, z) with the black hole at the origin and its rotation axis along z .
- Using the Boyer-Lindquist coordinates describing the black hole, the observer will be located at some coordinates $(r_0, \theta_0, \varphi_0)$, with r_0 very large.

Celestial Coordinates



Celestial Coordinates

- Note that the inclination angle is precisely $\theta_0 = i$. Hence the position of the observer is (r_0, i, φ_0)

Celestial Coordinates

- Note that the inclination angle is precisely $\theta_0 = i$. Hence the position of the observer is (r_0, i, φ_0)
- Without loosing generality, we rotate the (x, y, z) system so that the angular position of the observer in the Boyer-Lindquist coordinates is $\varphi_0 = 0$.


Celestial Coordinates

- Note that the inclination angle is precisely $\theta_0 = i$. Hence the position of the observer is (r_0, i, φ_0)
- Without loosing generality, we rotate the (x, y, z) system so that the angular position of the observer in the Boyer-Lindquist coordinates is $\varphi_0 = 0$.
- The position of the observer becomes $(r_0, i, 0)$

Celestial Coordinates

- Note that the inclination angle is precisely $\theta_0 = i$. Hence the position of the observer is (r_0, i, φ_0)
- Without loosing generality, we rotate the (x, y, z) system so that the angular position of the observer in the Boyer-Lindquist coordinates is $\varphi_0 = 0$.
- The position of the observer becomes $(r_0, i, 0)$
- Then, the observer lies in the $x - z$ plane while the y -axis lies in the (α, β) plane (remember that the observer plane is perpendicular to the line of sight).

Celestial Coordinates



Trajectory of a Light Ray

In the observer's reference frame, an incoming light ray trajectory may be described by a parametric curve

$$\begin{cases} X = X(r) \\ Y = Y(r) \\ Z = Z(r) \end{cases}$$

such that

$$r^2 = X^2(r) + Y^2(r) + Z^2(r)$$

Trajectory of a Light Ray

The tangent vector to this parametric curve at the observer's location is

$$\vec{\mu} = (\mu_1, \mu_2, \mu_3) = \left(\left. \frac{dX}{dr} \right|_{r_0}, \left. \frac{dY}{dr} \right|_{r_0}, \left. \frac{dZ}{dr} \right|_{r_0} \right)$$

Trajectory of a Light Ray

From the point of view of the observer, this tangent vector defines the trajectory of the photon as a straight line which intersects the observer's celestial plane at the coordinates (α_i, β_i) and can be written parametrically as

$$\frac{x - x_0}{\mu_1} = \frac{y - y_0}{\mu_2} = \frac{z - z_0}{\mu_3}$$

Trajectory of a Light Ray

(x_0, y_0, z_0) are the coordinates of the observer's position.

Trajectory of a Light Ray

(x_0, y_0, z_0) are the coordinates of the observer's position.
This can be written as

$$(x_0, y_0, z_0) = (r_0 \sin i, 0, r_0 \cos i)$$

Trajectory of a Light Ray

(x_0, y_0, z_0) are the coordinates of the observer's position.
This can be written as

$$(x_0, y_0, z_0) = (r_0 \sin i, 0, r_0 \cos i)$$

The celestial coordinates (α_i, β_i) can be written as

$$(\alpha_i, \beta_i) = (x_i, y_i, z_i) = (-\beta_i \cos i, \alpha_i, \beta_i \sin i)$$

Trajectory of a Light Ray

$$\frac{x - x_0}{\mu_1} = \frac{y - y_0}{\mu_2} = \frac{z - z_0}{\mu_3}$$

Trajectory of a Light Ray

$$\frac{x - x_0}{\mu_1} = \frac{y - y_0}{\mu_2} = \frac{z - z_0}{\mu_3}$$

$$\frac{-\beta_i \cos i - r_0 \sin i}{\mu_1} = \frac{\alpha_i}{\mu_2} = \frac{\beta_i \sin i - r_0 \cos i}{\mu_3}$$

Trajectory of a Light Ray

Using the transformation between cartesian and spherical coordinates,

$$X(r) = r \sin \theta \cos \varphi$$

$$Y(r) = r \sin \theta \sin \varphi$$

$$Z(r) = r \cos \theta,$$

we obtain the components of $\vec{\mu}$,

$$\mu_1 = \left. \frac{dX}{dr} \right|_{r_0} = \sin i + r_0 \cos i \left. \frac{d\theta}{dr} \right|_{r_0}$$

$$\mu_2 = \left. \frac{dY}{dr} \right|_{r_0} = r_0 \sin i \left. \frac{d\varphi}{dr} \right|_{r_0}$$

$$\mu_3 = \left. \frac{dZ}{dr} \right|_{r_0} = \cos i - r_0 \sin i \left. \frac{d\theta}{dr} \right|_{r_0}$$

Trajectory of a Light Ray

$$\frac{-\beta_i \cos i - r_0 \sin i}{\mu_1} = \frac{\alpha_i}{\mu_2} = \frac{\beta_i \sin i - r_0 \cos i}{\mu_3}$$

Trajectory of a Light Ray

$$\frac{-\beta_i \cos i - r_0 \sin i}{\mu_1} = \frac{\alpha_i}{\mu_2} = \frac{\beta_i \sin i - r_0 \cos i}{\mu_3}$$

$$\frac{-\beta_i \cos i - r_0 \sin i}{\sin i + r_0 \cos i \left. \frac{d\theta}{dr} \right|_{r_0}} = \frac{\alpha_i}{r_0 \sin i \left. \frac{d\phi}{dr} \right|_{r_0}} = \frac{\beta_i \sin i - r_0 \cos i}{\cos i - r_0 \sin i \left. \frac{d\theta}{dr} \right|_{r_0}}$$

Celestial Coordinates

$$\alpha_i = \lim_{r_0 \rightarrow \infty} -r_0^2 \sin^2 \theta_0 \left. \frac{d\varphi}{dr} \right|_{r_0}$$

$$\beta_i = \lim_{r_0 \rightarrow \infty} r_0^2 \left. \frac{d\theta}{dr} \right|_{r_0}$$

Celestial Coordinates

Using the equations of motion for the photons, we obtain

$$\alpha_i = -\xi \csc i$$

$$\beta_i = \sqrt{\eta + a^2 \cos^2 i - \xi^2 \cot^2 i}$$