



FACULTY
OF ARTS

Masaryk University

Classical Black Holes

02. Black holes Introduction

Edward Larrañaga

Outline for Part 1

1. Schwarzschild's Solution

1.1 General Relativity Field Equations

1.2 Physical Conditions to obtain Schwarzschild's Metric

Einstein Field Equations

Einstein field equations are

$$G_{\mu\nu} = R_{\mu\nu} - \frac{1}{2}g_{\mu\nu}R = 8\pi GT_{\mu\nu}$$

Riemann Tensor

Riemann tensor is defined as

$$R^{\lambda}_{\mu\rho\nu} = \partial_{\rho}\Gamma^{\lambda}_{\nu\mu} - \partial_{\nu}\Gamma^{\lambda}_{\rho\mu} + \Gamma^{\lambda}_{\rho\alpha}\Gamma^{\alpha}_{\nu\mu} - \Gamma^{\lambda}_{\nu\alpha}\Gamma^{\alpha}_{\rho\mu}$$

Riemann Tensor

Riemann tensor is defined as

$$R^{\lambda}_{\mu\rho\nu} = \partial_{\rho}\Gamma^{\lambda}_{\nu\mu} - \partial_{\nu}\Gamma^{\lambda}_{\rho\mu} + \Gamma^{\lambda}_{\rho\alpha}\Gamma^{\alpha}_{\nu\mu} - \Gamma^{\lambda}_{\nu\alpha}\Gamma^{\alpha}_{\rho\mu}$$

$$\Gamma^{\nu}_{\mu\sigma} = \frac{1}{2}g^{\nu\alpha}[\partial_{\mu}g_{\alpha\sigma} + \partial_{\sigma}g_{\mu\alpha} - \partial_{\alpha}g_{\mu\sigma}]$$

Ricci Tensor and Curvature Scalar

$$R_{\mu\nu} = g^{\sigma\rho} R_{\sigma\mu\rho\nu}$$

$$R = R^\sigma_\sigma = g^{\sigma\tau} R_{\sigma\tau}.$$

Ricci Tensor and Curvature Scalar

$$R_{\mu\nu} = \partial_\rho \Gamma_{\nu\mu}^\rho - \partial_\nu \Gamma_{\rho\mu}^\rho + \Gamma_{\rho\alpha}^\rho \Gamma_{\nu\mu}^\alpha - \Gamma_{\nu\alpha}^\rho \Gamma_{\rho\mu}^\alpha$$

$$R = g^{\sigma\tau} \partial_\rho \Gamma_{\tau\sigma}^\rho - g^{\sigma\tau} \partial_\tau \Gamma_{\rho\sigma}^\rho + g^{\sigma\tau} \Gamma_{\rho\alpha}^\rho \Gamma_{\tau\sigma}^\alpha - g^{\sigma\tau} \Gamma_{\tau\alpha}^\rho \Gamma_{\rho\sigma}^\alpha$$

Einstein Field Equations

$$R_{\mu\nu} - \frac{1}{2}g_{\mu\nu}R = 8\pi GT_{\mu\nu}$$

Einstein Field Equations

$$R_{\mu\nu} - \frac{1}{2}g_{\mu\nu}R = 8\pi GT_{\mu\nu}$$

$$\begin{aligned} & \partial_\rho \Gamma_{\nu\mu}^\rho - \partial_\nu \Gamma_{\rho\mu}^\rho + \Gamma_{\rho\alpha}^\rho \Gamma_{\nu\mu}^\alpha - \Gamma_{\nu\alpha}^\rho \Gamma_{\rho\mu}^\alpha \\ & - \frac{1}{2}g_{\mu\nu} \left[g^{\sigma\tau} \partial_\rho \Gamma_{\tau\sigma}^\rho - g^{\sigma\tau} \partial_\tau \Gamma_{\rho\sigma}^\rho + g^{\sigma\tau} \Gamma_{\rho\alpha}^\rho \Gamma_{\tau\sigma}^\alpha - g^{\sigma\tau} \Gamma_{\tau\alpha}^\rho \Gamma_{\rho\sigma}^\alpha \right] = 8\pi GT_{\mu\nu} \end{aligned}$$

Physical Conditions to obtain Schwarzschild's Metric

- Spherical Symmetry

Physical Conditions to obtain Schwarzschild's Metric

- Spherical Symmetry
- Asymptotically flat spacetime

Physical Conditions to obtain Schwarzschild's Metric

- Spherical Symmetry
- Asymptotically flat spacetime
- Static Metric

Physical Conditions to obtain Schwarzschild's Metric

- Spherical Symmetry
- Asymptotically flat spacetime
- Static Metric
- Empty Spacetime

Physical Conditions to obtain Schwarzschild's Metric

- Spherical Symmetry

Physical Conditions to obtain Schwarzschild's Metric

- Spherical Symmetry

Spherical Coordinates: (t, r, θ, ϕ)

Physical Conditions to obtain Schwarzschild's Metric

- Spherical Symmetry

Spherical Coordinates: (t, r, θ, ϕ)

$$ds^2 = -g_{tt}dt^2 + g_{rr}dr^2 + g_{\theta\theta}d\theta^2 + g_{\phi\phi}d\phi^2$$

Physical Conditions to obtain Schwarzschild's Metric

- Spherical Symmetry

Spherical Coordinates: (t, r, θ, ϕ)

$$ds^2 = -g_{tt}dt^2 + g_{rr}dr^2 + g_{\theta\theta}d\theta^2 + g_{\phi\phi}d\phi^2$$

$$ds^2 = -g_{tt}dt^2 + g_{rr}dr^2 + r^2d\theta^2 + r^2\sin^2\theta d\phi^2$$

Physical Conditions to obtain Schwarzschild's Metric

- Spherical Symmetry

Spherical Coordinates: (t, r, θ, ϕ)

$$ds^2 = -g_{tt}dt^2 + g_{rr}dr^2 + g_{\theta\theta}d\theta^2 + g_{\phi\phi}d\phi^2$$

$$ds^2 = -g_{tt}dt^2 + g_{rr}dr^2 + r^2d\theta^2 + r^2\sin^2\theta d\phi^2$$

$$g_{tt} = g_{tt}(t, r)$$

$$g_{rr} = g_{rr}(t, r)$$

Physical Conditions to obtain Schwarzschild's Metric

- Asymptotically flat

Physical Conditions to obtain Schwarzschild's Metric

- Asymptotically flat

$$\lim_{r \rightarrow \infty} g_{tt} \longrightarrow 1$$

$$\lim_{r \rightarrow \infty} g_{rr} \longrightarrow 1$$

Physical Conditions to obtain Schwarzschild's Metric

- Static Metric

Physical Conditions to obtain Schwarzschild's Metric

- Static Metric

$$g_{tt} = g_{tt}(r)$$

$$g_{rr} = g_{rr}(r)$$

Physical Conditions to obtain Schwarzschild's Metric

- Static Metric

$$g_{tt} = g_{tt}(r)$$

$$g_{rr} = g_{rr}(r)$$

"Birkhoff Theorem"

Physical Conditions to obtain Schwarzschild's Metric

- Empty Spacetime

Physical Conditions to obtain Schwarzschild's Metric

- Empty Spacetime

$$T_{\mu\nu} = 0$$

Physical Conditions to obtain Schwarzschild's Metric

- Empty Spacetime

$$T_{\mu\nu} = 0$$

Field Equations:

$$R_{\mu\nu} = 0$$

Spherically Symmetric Solution in Empty Space

$$ds^2 = - \left(1 - \frac{K}{r}\right) dt^2 + \left(1 - \frac{K}{r}\right)^{-1} dr^2 + r^2 d\theta^2 + r^2 \sin^2 \theta d\phi^2$$

K : Integration constant to be determined

Next Lecture

03. Schwarzschild Solution. Properties.