PROJECT COMPLETION REPORT

FALCON: Autonomous Aerial Biodiversity & Nature-Credit Auditing System

A Deep-Learning-Powered System for Aerial Micro-Fauna Auditing, Ecological Indexing, and Verifiable Green Credit Issuance

Overall Project Completion

100%

Repository:

https://github.com/ashcodes28/FALCON

Date of Submission

24 October 2026

Table of Contents

Table of Contents 2
1. Executive Summary 3
2. Introduction 3
  2.1 Background and Motivation 3
  2.2 Objectives 3
  2.3 Scope of This Final Report 4
3. Overall Project Implementation Snapshot 4
4. Scientific Grounding, Literature & Patent Landscape 3
  4.1 Key Findings from Academic Literature 5
  4.2 Patent Landscape & Novelty Gaps 5
  4.3 Patented Novelty Claims 5
5. Multi-Guild Dataset & Ecological Taxonomy 6
  5.1 Invertebrate Image Corpus 6
  5.2 Functional Ecological Guild Mapping 6
6. Deep Learning Detector Architecture & Performance Evaluation 6
  6.1 YOLOv8 & YOLOv11n Baseline Implementations 6
  6.2 Per-Class Performance Metrics 7
  6.3 Morphological Confusion Analysis 7
7. Platform & Hardware System Design 7
  7.1 UAV Airframe & Payload Bay CAD 7
  7.2 End-to-End Autonomous Data Pipeline 7
  7.3 Guild Isolation Testing Module 8
8. Mathematical Biodiversity Engine & Green Credit Formulation 8
  8.1 Shannon-Wiener Faunal Diversity Index 8
  8.2 Simpson's Diversity Index 8
  8.3 Pollinator-to-Pest Ratio 8
  8.4 Soil Bioturbation Proxy Score 9
  8.5 Composite Ecological Quality Score & Net Uplift 9
  8.6 Standardized Green Credit Units & Accreditation Tiers 10
9. Cryptographic Verification & Tamper-Evident Audit Ledger 3
  9.1 Sequential SHA-256 Chained Block Architecture 10
  9.2 Cryptographic Anti-Tamper Security Validation 11
  9.3 Audit Certificate & GeoJSON Vector Exports 3
10. System Validation, Simulation Flight Trials & Experimental Results 11
  10.1 Automated Self-Test & Unit Verification 3
  10.2 Autonomous Transect Simulation Flight Trial 11
  10.3 Real-Time Ecological & Green Credit HUD Streaming 12
11. Regulatory, Compliance & Environmental Market Impact 12
12. Conclusion 12
13. References 12

1. Executive Summary

This report documents the completed implementation of the FALCON autonomous unmanned aerial system (UAS). While originally designed for agricultural pest surveillance, the platform was expanded to audit terrestrial invertebrate biodiversity and issue verifiable nature credits. Emerging carbon and Biodiversity Net-Gain (BNG) markets often rely on satellite NDVI data, which tracks canopy density and chlorophyll content but cannot detect living micro-fauna. Ground entomology surveys measure species accurately, but manual sampling is too slow and labor-intensive to scale across commercial acreage.

FALCON resolves this limitation by pairing low-altitude aerial flight (3–8 m AGL) with edge computer vision (YOLOv11n at 1024×1024 resolution) and an automated ecological calculation engine. The system classifies 12 invertebrate taxa into four functional ecological guilds (pollinators, biocontrol predators, soil detritivores, and herbivores). These counts feed into the Shannon-Wiener Faunal Diversity Index (H'), the Pollinator-to-Pest Ratio (PPR), and an epigeic soil bioturbation score. The engine combines these metrics into a standardized Ecological Quality Score (Q), subtracts a regional degraded monoculture baseline ($Q_{baseline} = 0.20$) to calculate net ecological uplift (ΔQ), and converts the uplift into certified Green Credit Units (GCUs) scaled by surveyed acreage.

Each observation waypoint is recorded in a sequential SHA-256 hash-chained ledger, preventing retroactive modification of audit records. In a five-waypoint flight simulation covering a 15-hectare parcel, the pipeline processed 25 invertebrate observations across 12 taxa, measured a net uplift of +0.711 Delta-Q, and awarded 10.665 net Green Credit Units under a verified Tier-AAA rating.

---

2. Introduction

2.1 Background and Motivation

Statutory frameworks like the UK Biodiversity Net Gain (BNG) mandate and voluntary frameworks such as the Taskforce on Nature-related Financial Disclosures (TNFD) establish economic value for verified ecological restoration. Landowners and agricultural enterprises can generate nature credits and carbon premiums when they provide verifiable evidence of biological recovery on their land.

Monitoring, Reporting, and Verification (MRV) for terrestrial biodiversity has historically encountered practical obstacles. Satellite remote sensing cannot resolve insects or soil organisms, while manual entomological field surveys take substantial time, cost thousands of dollars per transect, and vary between observers. Paper-based reporting also leaves credit underwriters vulnerable to unverifiable claims. FALCON addresses these issues by operating an autonomous multirotor at low altitude (3–8 m AGL) to classify micro-fauna in near real time, logging every observation directly into a cryptographically sealed audit ledger.

2.2 Objectives

1. Deploy a lightweight deep learning detector (YOLOv11n) trained at 1024×1024 resolution to classify 12 ground and foliage invertebrate taxa.
2. Group detected taxa into four functional ecological guilds to assess pollination, natural pest predation, soil vitality, and herbivore pressure.
3. Formulate and implement mathematical indices for community diversity (H'), pollinator-to-pest balance (PPR), and upper-soil biological activity.
4. Construct an automated Green Credit Unit (GCU) calculation engine that converts verified ecological uplift over regional baselines into accredited units.
5. Engineer a sequential SHA-256 cryptographic audit ledger that exports signed underwriter certificates (`audit_certificate.json`) and GIS-ready vector files (`audit_transect.geojson`).
6. Validate the complete system architecture through unit test suites, anti-tamper verification checks, and a simulated 15-hectare aerial flight audit.

2.3 Scope of This Final Report

This document marks the conclusion of the FALCON engineering project, covering all hardware, machine learning, algorithmic, cryptographic, and validation deliverables.

---

3. Overall Project Implementation Snapshot

Every planned work package across the project timeline has reached completion.

Work Package	Deliverable	Status	% Weight
1. Literature Review & Regulatory Mapping	Academic state of the art, IP landscape, patent claim outlines	Completed	10%
2. Problem Definition & Architectural Pivot	Transition from pest spraying to autonomous biodiversity auditing	Completed	5%
3. Dataset Curation & Benchmark Linking	12-class micro-fauna corpus, Kaggle/IP102/ AIDN benchmark curation	Completed	10%
4. Baseline Model Training & Validation	YOLOv8 single-modality baseline (85.5% accuracy)	Completed	10%
5. Edge Model Development & Optimization	Lightweight YOLOv11n (5.56 MB) at 1024×1024 resolution	Completed	10%
6. Airframe & Payload Bay Mechanical CAD	3D CAD modeling of airframe, gimbal mount, and electronics bay	Completed	10%

7. Ecological Guild Classification Engine	4-guild functional mapping (Pollinators, Predators, Soil, Herbivores)	Completed	10%
8. Mathematical Biodiversity Engine	Shannon-Wiener H', Simpson's 1-D, PPR, and soil bioturbation scoring	Completed	10%
9. Green Credit Unit (GCU) Formulation	Quality score Q, net uplift Delta-Q, area scaling, 5 accreditation tiers	Completed	10%
10. Cryptographic Audit Ledger & Exports	SHA-256 sequential chaining, `audit_certificate.json`, GeoJSON exporter	Completed	10%
11. System Validation & Flight Simulation	Automated test suite (`biodiversity_engine.py`), flight trial (`simulate_audit.py`)	Completed	5%
Total	**Full System Implementation**	**100% Completed**	**100%**

---

4. Scientific Grounding, Literature & Patent Landscape

4.1 Key Findings from Academic Literature

- *Small-object aerial invertebrate localization:* Recent aerial entomology studies (Bai et al., 2023; AIDN, 2024) confirm that insects typically occupy only 10×10 to 30×30 pixels in low-altitude drone imagery. Training detector backbones on high-resolution 1024×1024 inputs preserves the edge gradients and fine texture cues needed to separate visually similar taxa.
- *Limitations of optical vegetation indices:* While NDVI and related multispectral reflectance metrics track canopy volume and chlorophyll density, they do not correlate reliably with invertebrate species richness or soil biological activity (Alsadik et al., 2024). Direct counts of micro-fauna provide the empirical baseline required by nature credit accreditation bodies.

4.2 Patent Landscape & Novelty Gaps

Prior agricultural drone patents (e.g., US 10,364,029 B2; US 2017/0231213 A1; KR 20230062713 A) focus on mechanical hardware: spray tanks, nozzle arrays, and chemical delivery valves. Algorithmic methods converting multi-guild micro-fauna aerial classifications into cryptographically sealed ecological credits were completely unaddressed in surveyed prior art.

4.3 Patented Novelty Claims

FALCON establishes novel system and method claims:

1. *Method Claim:* Autonomous capture of low-altitude aerial transects, real-time edge

classification into functional ecological guilds, spatial calculation of multi-guild diversity and pollinator ratios, and derivation of certified Green Credit Units based on verified uplift over baseline.

2. *System Claim:* An onboard companion processor cryptographically binding video frames, GPS waypoints, derived faunal metrics, and prior block hashes into an immutable SHA-256 ledger.

---

5. Multi-Guild Dataset & Ecological Taxonomy

5.1 Invertebrate Image Corpus

The model training dataset comprises 12 distinct invertebrate taxa commonly encountered in agricultural and terrestrial ecosystems, linked to the Agricultural Pests Image Dataset (Kaggle), supplemented by representation learning transfer from IP102 and AIDN.

5.2 Functional Ecological Guild Mapping

FALCON categorizes the 12 observed invertebrate classes into four functional ecological guilds based on their agroecological roles:

Ecological Guild	Included Classes	Ecological Function & Role
Pollinators	Bees	Floral resource indicator, angiosperm cross-pollination.
Biocontrol Predators	Wasps	Parasitoid regulation, natural suppression of herbivore larvae.
Soil & Detrital Fauna	Earthworms, Slugs, Snails	Soil bioturbation, topsoil moisture indicator, organic decomposition.
Herbivore Pressure	Caterpillars, Grasshoppers, Weevils, Beetles, Earwigs, Ants, Moths	Foliage defoliation pressure, primary consumer biomass.

---

6. Deep Learning Detector Architecture & Performance Evaluation

6.1 YOLOv8 & YOLOv11n Baseline Implementations

Two detector baselines were trained and benchmarked at 1024×1024 resolution:

- **YOLOv8 Baseline:** Reached 85.5% overall accuracy on validation splits.

- **YOLOv11n Edge Detector:** Compressed into a lightweight 5.56 MB model (`best.pt`), optimized for edge deployment on companion computers (such as NVIDIA Jetson Orin Nano).

6.2 Per-Class Performance Metrics

High classification accuracy was achieved across visually distinct invertebrate categories:

- **Snail:** 1.00 Precision
- **Weevil:** 1.00 Precision
- **Moth:** 0.99 Precision
- **Wasp:** 0.97 Precision
- **Ants:** 0.87 Precision

6.3 Morphological Confusion Analysis

Analysis of the normalized confusion matrix revealed specific inter-class confusion patterns:

- Small bounding contours produced cross-class overlap between Beetles ⇔ Wasps and Caterpillars ⇔ Bees/Beetles.
- Subterranean/surface detritivores exhibited expected visual overlap (Slugs ⇔ Earthworms at 0.29).
- Because both members of confused pairs often belong to the same functional guild (e.g., Slugs and Earthworms are both Detritivores), guild-level index derivation remains robust against individual species misclassifications.

7. Platform & Hardware System Design

7.1 UAV Airframe & Payload Bay CAD

The mechanical airframe (detailed in `CAD Images/`) incorporates:

- A rigid carbon-fiber multi-rotor frame with vibration-isolated motor arms.
- A downward-facing 2-axis brushless camera gimbal maintaining nadir optical alignment.
- A ventilated payload bay housing the companion single-board computer, GNSS module, and laser altimeter.

7.2 End-to-End Autonomous Data Pipeline

1. Downward gimbal camera captures 1024×1024 frames at 3–8 m AGL.
2. Edge YOLOv11n performs inference, outputting bounding coordinates and confidence scores.
3. The biodiversity calculation engine classifies detections into ecological guilds.
4. Mathematical routines compute H', PPR, and soil vitality.

5. The cryptographic ledger generates a sequentially hashed block.
6. Real-time metrics are streamed to the ground station HUD and written to persistent storage.

7.3 Guild Isolation Testing Module

`src/lab_test_detection.py` provides bench testing capabilities to isolate and verify specific guilds (`POLLINATORS`, `BIOCONTROL`, `SOIL_FAUNA`, `HERBIVORES`), allowing operators to validate guild-specific confidence thresholds before deployment.

---

8. Mathematical Biodiversity Engine & Green Credit Formulation

8.1 Shannon-Wiener Faunal Diversity Index (H')

Quantifies invertebrate species richness and relative abundance evenness across the surveyed parcel:

> **Equation 8.1 (Shannon-Wiener Diversity Index):**
> $$H' = -\sum [\, p_i \cdot \ln(p_i) \,]$$ *(summed across all observed taxa $i = 1$ to S)*
> where:
>
> - S = Total number of observed taxa ($S = 12$ in the FALCON ecological taxonomy).
> - n_i = Number of individuals recorded for taxon i.
> - N = Total number of individuals recorded across all taxa ($N = \sum n_i$).
> - p_i = Relative abundance proportion of taxon i ($p_i = n_i / N$).
>
> The theoretical maximum diversity for 12 equally distributed taxa is $\ln(12) \approx 2.4849$. A higher H' indicates an ecologically balanced invertebrate community with high evenness and species richness.

8.2 Simpson's Diversity Index (1 – D)

Measures the probability that two randomly sampled invertebrates from the surveyed transect belong to different species, quantifying community evenness and resistance to single-species dominance:

> **Equation 8.2 (Simpson's Diversity Index):**
> $$1 - D = 1 - \sum [\, p_i^2 \,]$$ *(summed across all observed taxa $i = 1$ to S)*
> where p_i is the relative abundance proportion of taxon i. Values approach 1.0 in highly diverse habitats and drop toward 0.0 under monoculture pest infestations.

8.3 Pollinator-to-Pest Ratio (PPR)

Directly quantifies the balance of ecosystem service providers (beneficial pollinators and parasitoid biocontrol regulators) against destructive foliage-feeding herbivores:

Equation 8.3 (Pollinator-to-Pest Ratio):
PPR = (N_Bees + N_Wasps) / (N_Weevils + N_Caterpillars + 10^{-6})
where:

- *N_Bees* and *N_Wasps* represent beneficial pollinating and parasitoid predatory guilds.
- *N_Weevils* and *N_Caterpillars* represent primary economic pest pressures.
- A smoothing constant of 10^{-6} in the denominator prevents division-by-zero artifacts in pest-free or heavily suppressed parcels.

An audited parcel with PPR $\geq$ 2.0 indicates an ecosystem where natural regulatory mechanisms outweigh herbivore pressure.

8.4 Soil Bioturbation Proxy Score

Assesses biological activity and structural aeration in the upper soil and litter layer through aerial observation of epigeic detrital macro-invertebrates:

Equation 8.4 (Soil Bioturbation Proxy Score):
Score_soil = 1.5 · N_Earthworms + 0.5 · (N_Slugs + N_Snails)
where Earthworms receive a 1.5× biological weighting reflecting their primary role in soil macro-pore creation, water infiltration, and organic matter incorporation, while Gastropods receive a 0.5× weighting as moisture and detrital decomposition indicators.

8.5 Composite Ecological Quality Score & Net Uplift

The composite Ecological Quality Score ($Q \in [0.0, 1.0]$) synthesizes the three biological pillars into a single standardized scalar metric:

Equation 8.5 (Composite Ecological Quality Score):
Q = 0.40 · (H' / ln 12) + 0.35 · min(PPR / 2.0, 1.0) + 0.25 · min(Score_soil / 10.0, 1.0)
where:

- 40% weight is assigned to overall community diversity and evenness (H' normalized by its theoretical maximum ln(12) $\approx$ 2.4849).
- 35% weight is assigned to the functional ecosystem balance between pollinators/predators and pests (PPR normalized to 2.0).
- 25% weight is assigned to epigeic soil biological activity (Score_soil normalized to 10.0).

Net ecological uplift (ΔQ) measures verified ecological gain above the degraded agricultural monoculture baseline (*Q_baseline* = 0.20):

Equation 8.6 (Net Ecological Uplift):
ΔQ = max(Q − Q_baseline, 0.0)
Ensuring that credits are only issued for verified ecological additionality above the regional degraded benchmark.

8.6 Standardized Green Credit Units & Accreditation Tiers

Credits are issued on a per-hectare uplift basis, directly tying financial issuance to physical land area and verifiable biodiversity additionality:

> **Equation 8.7 (Gross Green Credit Units):**
> **Gross GCU = Surveyed Area (ha) × Q**

> **Equation 8.8 (Net Certified Green Credit Units Awarded):**
> **Net Awarded GCU = Surveyed Area (ha) × ΔQ**

Audited parcels are classified into five standardized accreditation tiers based on their certified Ecological Quality Score (Q):

Quality Score (Q) Range	Accreditation Tier	Ecological Classification & Market Status
Q ≥ 0.80	**Tier-AAA**	Exemplary Biodiversity Net-Gain (Premium market issuance)
0.65 ≤ Q < 0.80	**Tier-AA**	High Ecological Integrity (Standard nature credit issuance)
0.50 ≤ Q < 0.65	**Tier-A**	Moderate Ecological Value (Transitional conservation credit)
0.35 ≤ Q < 0.50	**Tier-B**	Early Regenerative Transition (Monitoring credit)
Q < 0.35	**Tier-C**	Depleted / Sub-Baseline Monoculture (Ineligible for credit issuance)

---

9. Cryptographic Verification & Tamper-Evident Audit Ledger

9.1 Sequential SHA-256 Chained Block Architecture

Implemented in src/biodiversity_engine.py, the `BiodiversityAuditLedger` class records observations in sequential blocks. Each block contains:

- Block index, parcel ID, operator ID, UTC timestamp, and GNSS coordinates.
- Complete faunal counts and derived ecological indicators.
- Certified Green Credit calculations.
- The SHA-256 hash of the preceding block (`previous_hash`).

9.2 Cryptographic Anti-Tamper Security Validation

Modifying any historical block's insect count, coordinates, or credit units alters its SHA-256 digest, breaking the hash chain for all subsequent blocks. The `verify_integrity()` method performs end-to-end verification across the entire chain prior to issuing a certificate.

9.3 Audit Certificate & GeoJSON Vector Exports

Upon mission completion, the system serializes:

1. `audit_certificate.json`: Official underwriter certificate containing total counts, certified GCUs, accreditation tier, genesis/final hashes, and chain validity proofs.
2. `audit_transect.geojson`: Standard GIS vector file mapping localized diversity scores, PPR, and instant credits across geographic coordinates.

---

10. System Validation, Simulation Flight Trials & Experimental Results

10.1 Automated Self-Test & Unit Verification

Execution of `python src/biodiversity_engine.py` confirmed:

- Index calculation accuracy on multi-guild invertebrate sets.
- Mathematical convergence of the Green Credit formula.
- Invalidation of ledger integrity when simulated tampering was introduced (`Pre-tamper: True`, `Post-tamper: False`).

10.2 Autonomous Transect Simulation Flight Trial

Execution of `python src/simulate_audit.py` over a 15.0-hectare parcel across five waypoints produced the following verified results:

```
======================================================================
 OFFICIAL GREEN CREDIT AUDIT SUMMARY
======================================================================
 Target Parcel ID        : AGRI-PARCEL-GREEN-VALLEY-01
 Total Surveyed Area     : 15.00 Hectares
 Total Invertebrates     : 25 individuals across 12 taxa
 Shannon Diversity (H')  : 2.165 (Theoretical maximum: 2.485)
 Pollinator-to-Pest Ratio: 3.33 (Beneficials significantly outnumber pests)
 Soil Bioturbation Score : 8.5
 Certified Quality Score : 0.911 / 1.000
 Baseline Floor Score    : 0.200
 Net Ecological Uplift   : +0.711 Delta-Q
----------------------------------------------------------------------
 >>> CERTIFIED GREEN CREDIT UNITS (GCUs) AWARDED: 10.665 GCU <<<
```

```
------------------------------------------------------------------------
 Accredited Rating Tier  : Tier-AAA (Exemplary Biodiversity Net-Gain)
 Ledger Chain Integrity  : VALID (Zero Tampering Detected)
 Final Block SHA-256     : 6e5cc66bd414d9c5761f1fd2f43c8f757debd98b...
========================================================================
```

10.3 Real-Time Ecological & Green Credit HUD Streaming

src/detection.py integrates real-time video inference with the audit ledger, displaying a live telemetry HUD overlaying bounding boxes, running Shannon index, Pollinator-to-Pest Ratio, certified GCUs, and blockchain integrity status.

---

11. Regulatory, Compliance & Environmental Market Impact

The audit outputs produced by FALCON align directly with current environmental reporting frameworks and nature credit registries:

- *Verra Nature Framework and Plan Vivo:* The ledger supplies the empirical, individual-level invertebrate counts and spatial transects required to substantiate nature credit claims.
- *Taskforce on Nature-related Financial Disclosures (TNFD):* Corporate landowners can disclose localized biodiversity health metrics supported by immutable SHA-256 proofs.
- *UK Biodiversity Net Gain (BNG):* Delivers repeatable, field-level faunal condition measurements to satisfy statutory 10% ecological uplift requirements on development and agricultural parcels.
- *Regenerative Agriculture Certification:* Farmers and land stewards can document balanced pollinator-to-pest ratios (PPR $\geq$ 2.0) and soil biological activity without relying on subjective self-reporting.

---

12. Conclusion

The FALCON engineering project has reached 100% implementation completion. Expanding beyond agricultural pest scouting, the platform establishes that a lightweight drone running edge computer vision can survey ground-level micro-fauna, compute empirical diversity metrics, and issue cryptographically verifiable nature credits. Autonomous simulations and validation test suites confirmed the complete data flow: classifying 12 invertebrate taxa into functional ecological guilds, calculating Shannon-Wiener and pollinator-to-pest ratios, logging sequential SHA-256 blocks, and exporting GIS-compatible audit certificates. All CAD models, trained weights, calculation libraries, and verification routines are operational in the project repository.

---

13. References

- *Advanced Insect Detection Network (AIDN) for UAV-Based Biodiversity Monitoring*. Remote Sensing (MDPI), 2024. https://doi.org/10.3390/rs17060962
- Alsadik, B., Ellsäßer, F. J., Awawdeh, M., et al. (2024). *Remote Sensing Technologies Using UAVs for Monitoring*. Remote Sensing, 16(23), 4371.
- Bai, Y., Hou, F., Fan, X., et al. (2023). *A Lightweight Pest Detection Model for Drones Based on Transformer and Super-Resolution Sampling Techniques.* Agriculture, 13(9), 1812.
- Shannon, C. E. (1948). *A Mathematical Theory of Communication.* Bell System Technical Journal, 27(3), 379–423.
- Simpson, E. H. (1949). *Measurement of Diversity*. Nature, 163, 688.
- Subramanian, K. S., Pazhanivelan, S., Srinivasan, G., et al. (2021). *Drones in Insect Pest Management*. Frontiers in Agronomy.
- Wu, X., Zhan, C., Lai, Y. K., Cheng, M. M., & Yang, J. (2019). *IP102: A Large-Scale Benchmark Dataset for Insect Pest Recognition.* IEEE/CVF Conference on Computer Vision and Pattern Recognition (CVPR), 8788–8797. https://github.com/xpwu95/IP102
- *Agricultural Pests Image Dataset (12-class micro-fauna benchmark).* Kaggle Datasets. https://www.kaggle.com/datasets/fihmeh/agricultural-pests-image-dataset