

# Automated Assessment System for Embodied Cognition in Children


Computer Vision (6367) Project - Spring 2020

*Ashish Jaiswal (1001748734)*  
*Lalit Goski (1001728801)*

# Introduction

- A system that assesses ATEC tasks for measuring cognitive skills in children
- ATEC stands for Activate Test for Embodied Cognition
- Implements RNN on top of CNN layers on data recorded via a Kinect sensor
- Focuses only on Ball-Drop-to-the-Beat task from the ATEC tasks


# Ball-Drop-to-the-Beat Task



Green Light: Pass the ball

Red Light: No Pass

Yellow Light: Raise the hand that has the ball



# System Architecture I




Fig 1: Data Collection and Body Keypoints Extraction

# Body Keypoints from AlphaPose

{0, "Nose"}

{9, "LWrist"}

{1, "LEye"}

{10, "RWrist"}

{2, "REye"}

{11, "LHip"}

{3, "LEar"}

{12, "RHip"}

{4, "REar"}

{13, "LKnee"}

{5, "LShoulder"}

{14, "Rknee"}

{6, "RShoulder"}

{15, "LAnkle"}

{7, "LElbow"}

{16, "RAnkle"}

{8, "RElbow"}

The ones highlighted in **RED** are the ones that were selected as input for our Deep Neural Network

# System Architecture II




Fig 2: Model for Segment Classification

# Model Evaluation

- Training data size: 2908 and Testing data size: 300
- Test Accuracy: **87%**
- Confusion Matrix:

Pass	No Pass	Raise Hand
92	7	1
10	86	4
13	6	81

# Applications

- Helps in reducing manual labor in annotating and assessing video segments containing ATEC tasks
- Activity recognition for other tasks with a different dataset.
- Detection of early stages in cognitive disorders like developmental/motor skill disorders.
- Monitoring the progress of an individual in pre-post intervention studies of ATEC tasks.



# Conclusion and Future Enhancements

- Extension for the generalization of activity recognition for any task
- Discriminant validity will be determined comparing community samples with ADHD and ASD samples.
- Development of an adult version for use with mild to moderate TBI, Parkinson's Disease and other movement disorders.
- Pre-post intervention studies to determine ATEC sensitivity to interventions and to study course of illness.

# References

- [1] Alex Dillhoff, Konstantinos Tsiakas, Ashwin ramesh babu, Mohammad Zakizade-hghariehali, Benjamin Buchanan, Morris Bell, Vassilis Athitsos, and Fillia Makedon. An automated assessment system for embodied cognition in children: from motion data to executive functioning. pages 1–6, 09 2019.
- [2] Joel T Nigg Stephen V Faraone Erik G Willcutt, Alysa E Doyle and Bruce F Pennington. Validity of the executive function theory of attention-deficit/hyperactivity disorder: a meta-analytic review. pages 1336–1346, 11 2005.
- [3] Sepp Hochreiter and Jürgen Schmidhuber. Long short-term memory. *Neural Computation*, 9(8):1735–1780, 1997.
- [4] Z. Lai and W. Xie. Self-supervised learning for video correspondence flow. In *BMVC*, 2019.
- [5] Bastiaan Quast. rnn: a recurrent neural network in r. *Working Papers*, 2016.4
- [6] Yuliang Xiu, Jiefeng Li, Haoyu Wang, Yinghong Fang, and Cewu Lu. Pose Flow: Efficient online pose tracking. In *BMVC*, 2018.
- [7] Tomas Simon Shih-En Wei Zhe Cao, Gines Hidalgo and Yaser Sheikh. Openpose: real-time multi-person 2d pose estimation using part affinity field

THANK YOU