

Calculus III

MTH 331 | Section A | Fall 2012

Dr. James Ashe

Johnson C. Smith

beamer-ics

Calc III: 9.1 Sequences and Infinites Series

beamer-ics

Definition

A **sequence** $\{a_n\}$ is an ordered list of numbers of the form $\{a_0, a_1, a_2, \dots, a_n, \dots\}$.

Definition

A **sequence** $\{a_n\}$ is an ordered list of numbers of the form $\{a_0, a_1, a_2, \dots, a_n, \dots\}$.

The Fibonacci Sequence

$\{0, 1, 1, 2, 3, 5, 8, 13, 21, 34, \dots\}$ where $a_0 = 0$, $a_1 = 1$, and $a_n = a_{n-1} + a_{n-2}$ for $n > 2$.

Definition

A **sequence** $\{a_n\}$ is an ordered list of numbers of the form $\{a_0, a_1, a_2, \dots, a_n, \dots\}$.

The Fibonacci Sequence

$\{0, 1, 1, 2, 3, 5, 8, 13, 21, 34, \dots\}$ where $a_0 = 0$, $a_1 = 1$, and $a_n = a_{n-1} + a_{n-2}$ for $n > 2$.

- a_n is the n^{th} term in the sequence; above the 4^{th} term is $a_4 = 2$.

Definition

A **sequence** $\{a_n\}$ is an ordered list of numbers of the form $\{a_0, a_1, a_2, \dots, a_n, \dots\}$.

The Fibonacci Sequence

$\{0, 1, 1, 2, 3, 5, 8, 13, 21, 34, \dots\}$ where $a_0 = 0$, $a_1 = 1$, and $a_n = a_{n-1} + a_{n-2}$ for $n > 2$.

- a_n is the n^{th} term in the sequence; above the 4^{th} term is $a_4 = 2$.
- A sequence is generated by a **recurrence relation** if further terms are defined as a function of preceding terms (like in the Fibonacci sequence).

A sequence may also be defined by an **explicit formula**.

A sequence may also be defined by an **explicit formula**.

ex) $a_n = 1 + \sin(n\frac{\pi}{2}) \Rightarrow$

.

A sequence may also be defined by an **explicit formula**.

ex) $a_n = 1 + \sin(n\frac{\pi}{2}) \Rightarrow \{a_n\}_{n=1}^{\infty} = \{2, 1, 0, 1, 2, 1, 0, \dots\}$.

A sequence may also be defined by an **explicit formula**.

ex) $a_n = 1 + \sin(n\frac{\pi}{2}) \Rightarrow \{a_n\}_{n=1}^{\infty} = \{2, 1, 0, 1, 2, 1, 0, \dots\}$.

- There is an explicit formula for the Fibonacci sequence....

A sequence may also be defined by an **explicit formula**.

ex) $a_n = 1 + \sin(n\frac{\pi}{2}) \Rightarrow \{a_n\}_{n=1}^{\infty} = \{2, 1, 0, 1, 2, 1, 0, \dots\}$.

- There is an explicit formula for the Fibonacci sequence....

... $a_n = \frac{1}{\sqrt{5}} ((\varphi)^n - (1 - \varphi)^n)$ where $\varphi = \frac{1+\sqrt{5}}{2}$, the *Golden Ratio!*

A sequence may also be defined by an **explicit formula**.

ex) $a_n = 1 + \sin(n\frac{\pi}{2}) \Rightarrow \{a_n\}_{n=1}^{\infty} = \{2, 1, 0, 1, 2, 1, 0, \dots\}$.

- There is an explicit formula for the Fibonacci sequence....

... $a_n = \frac{1}{\sqrt{5}} ((\varphi)^n - (1 - \varphi)^n)$ where $\varphi = \frac{1+\sqrt{5}}{2}$, the *Golden Ratio*!

- More fun facts: $\lim_{n \rightarrow \infty} \frac{a_n}{a_{n-1}} = \left\{ \frac{a_n}{a_{n-1}} \right\}_{n=2}^{\infty} = \varphi$ where $\{a_n\}$ is the Fibonacci sequence!

A sequence may also be defined by an **explicit formula**.

ex) $a_n = 1 + \sin(n\frac{\pi}{2}) \Rightarrow \{a_n\}_{n=1}^{\infty} = \{2, 1, 0, 1, 2, 1, 0, \dots\}$.

- There is an explicit formula for the Fibonacci sequence....

... $a_n = \frac{1}{\sqrt{5}} ((\varphi)^n - (1 - \varphi)^n)$ where $\varphi = \frac{1+\sqrt{5}}{2}$, the *Golden Ratio*!

- More fun facts: $\lim_{n \rightarrow \infty} \frac{a_n}{a_{n-1}} = \left\{ \frac{a_n}{a_{n-1}} \right\}_{n=2}^{\infty} = \varphi$ where $\{a_n\}$ is the Fibonacci sequence!
- We say that if $\lim_{n \rightarrow \infty} a_n = L$ for a unique number L than a_n **converges**.

A sequence may also be defined by an **explicit formula**.

ex) $a_n = 1 + \sin(n\frac{\pi}{2}) \Rightarrow \{a_n\}_{n=1}^{\infty} = \{2, 1, 0, 1, 2, 1, 0, \dots\}$.

- There is an explicit formula for the Fibonacci sequence....

... $a_n = \frac{1}{\sqrt{5}} ((\varphi)^n - (1 - \varphi)^n)$ where $\varphi = \frac{1+\sqrt{5}}{2}$, the *Golden Ratio*!

- More fun facts: $\lim_{n \rightarrow \infty} \frac{a_n}{a_{n-1}} = \left\{ \frac{a_n}{a_{n-1}} \right\}_{n=2}^{\infty} = \varphi$ where $\{a_n\}$ is the Fibonacci sequence!
- We say that if $\lim_{n \rightarrow \infty} a_n = L$ for a unique number L than a_n **converges**. Otherwise we say that a_n **diverges**

A sequence may also be defined by an **explicit formula**.

ex) $a_n = 1 + \sin(n\frac{\pi}{2}) \Rightarrow \{a_n\}_{n=1}^{\infty} = \{2, 1, 0, 1, 2, 1, 0, \dots\}$.

- There is an explicit formula for the Fibonacci sequence....

... $a_n = \frac{1}{\sqrt{5}} ((\varphi)^n - (1 - \varphi)^n)$ where $\varphi = \frac{1+\sqrt{5}}{2}$, the *Golden Ratio*!

- More fun facts: $\lim_{n \rightarrow \infty} \frac{a_n}{a_{n-1}} = \left\{ \frac{a_n}{a_{n-1}} \right\}_{n=2}^{\infty} = \varphi$ where $\{a_n\}$ is the Fibonacci sequence!
- We say that if $\lim_{n \rightarrow \infty} a_n = L$ for a unique number L than a_n **converges**. Otherwise we say that a_n **diverges**

ex) $\{1, 2, 3, \dots\} \rightarrow$

A sequence may also be defined by an **explicit formula**.

ex) $a_n = 1 + \sin(n\frac{\pi}{2}) \Rightarrow \{a_n\}_{n=1}^{\infty} = \{2, 1, 0, 1, 2, 1, 0, \dots\}$.

- There is an explicit formula for the Fibonacci sequence....

... $a_n = \frac{1}{\sqrt{5}} ((\varphi)^n - (1 - \varphi)^n)$ where $\varphi = \frac{1+\sqrt{5}}{2}$, the *Golden Ratio*!

- More fun facts: $\lim_{n \rightarrow \infty} \frac{a_n}{a_{n-1}} = \left\{ \frac{a_n}{a_{n-1}} \right\}_{n=2}^{\infty} = \varphi$ where $\{a_n\}$ is the Fibonacci sequence!
- We say that if $\lim_{n \rightarrow \infty} a_n = L$ for a unique number L than a_n **converges**. Otherwise we say that a_n **diverges**
ex) $\{1, 2, 3, \dots\} \rightarrow$ diverges.

A sequence may also be defined by an **explicit formula**.

ex) $a_n = 1 + \sin(n\frac{\pi}{2}) \Rightarrow \{a_n\}_{n=1}^{\infty} = \{2, 1, 0, 1, 2, 1, 0, \dots\}$.

- There is an explicit formula for the Fibonacci sequence....

... $a_n = \frac{1}{\sqrt{5}} ((\varphi)^n - (1 - \varphi)^n)$ where $\varphi = \frac{1+\sqrt{5}}{2}$, the *Golden Ratio*!

- More fun facts: $\lim_{n \rightarrow \infty} \frac{a_n}{a_{n-1}} = \left\{ \frac{a_n}{a_{n-1}} \right\}_{n=2}^{\infty} = \varphi$ where $\{a_n\}$ is the Fibonacci sequence!
- We say that if $\lim_{n \rightarrow \infty} a_n = L$ for a unique number L than a_n **converges**. Otherwise we say that a_n **diverges**

ex) $\{1, 2, 3, \dots\} \rightarrow \text{diverges}$.

ex) $\{1, 2, 1, 2, 1, \dots\} \rightarrow$

A sequence may also be defined by an **explicit formula**.

ex) $a_n = 1 + \sin(n\frac{\pi}{2}) \Rightarrow \{a_n\}_{n=1}^{\infty} = \{2, 1, 0, 1, 2, 1, 0, \dots\}$.

- There is an explicit formula for the Fibonacci sequence....

... $a_n = \frac{1}{\sqrt{5}} ((\varphi)^n - (1 - \varphi)^n)$ where $\varphi = \frac{1+\sqrt{5}}{2}$, the *Golden Ratio*!

- More fun facts: $\lim_{n \rightarrow \infty} \frac{a_n}{a_{n-1}} = \left\{ \frac{a_n}{a_{n-1}} \right\}_{n=2}^{\infty} = \varphi$ where $\{a_n\}$ is the Fibonacci sequence!
- We say that if $\lim_{n \rightarrow \infty} a_n = L$ for a unique number L than a_n **converges**. Otherwise we say that a_n **diverges**

ex) $\{1, 2, 3, \dots\} \rightarrow$ diverges.

ex) $\{1, 2, 1, 2, 1, \dots\} \rightarrow$ diverges.

A sequence may also be defined by an **explicit formula**.

ex) $a_n = 1 + \sin(n\frac{\pi}{2}) \Rightarrow \{a_n\}_{n=1}^{\infty} = \{2, 1, 0, 1, 2, 1, 0, \dots\}$.

- There is an explicit formula for the Fibonacci sequence....

... $a_n = \frac{1}{\sqrt{5}} ((\varphi)^n - (1 - \varphi)^n)$ where $\varphi = \frac{1+\sqrt{5}}{2}$, the *Golden Ratio*!

- More fun facts: $\lim_{n \rightarrow \infty} \frac{a_n}{a_{n-1}} = \left\{ \frac{a_n}{a_{n-1}} \right\}_{n=2}^{\infty} = \varphi$ where $\{a_n\}$ is the Fibonacci sequence!
- We say that if $\lim_{n \rightarrow \infty} a_n = L$ for a unique number L than a_n **converges**. Otherwise we say that a_n **diverges**

ex) $\{1, 2, 3, \dots\} \rightarrow$ diverges.

ex) $\{1, 2, 1, 2, 1, \dots\} \rightarrow$ diverges.

- Fun question. Is $\{0, 1, 2, 3, \dots\} > \{1, 2, 3, \dots\}$?

Definition

An **infinite series** is the sum of an (infinite) sequence; It has the

form: $\sum_{k=1}^{\infty} a_n = a_1 + a_2 + \cdots + a_n \dots = S_n$. As before, if

$\lim_{n \rightarrow \infty} \sum_{k=1}^{\infty} a_n = \lim_{n \rightarrow \infty} S_n = L$ then the series **converges**; otherwise it **diverges**.

Definition

An **infinite series** is the sum of an (infinite) sequence; It has the

form: $\sum_{k=1}^{\infty} a_n = a_1 + a_2 + \cdots + a_n \dots = S_n$. As before, if

$\lim_{n \rightarrow \infty} \sum_{k=1}^{\infty} a_n = \lim_{n \rightarrow \infty} S_n = L$ then the series **converges**; otherwise it **diverges**.

ex) (see pg 52) Let $S_n = \sum_{k=1}^n \frac{1}{2^k}$.

Definition

An **infinite series** is the sum of an (infinite) sequence; It has the

form: $\sum_{k=1}^{\infty} a_n = a_1 + a_2 + \cdots + a_n \dots = S_n$. As before, if

$\lim_{n \rightarrow \infty} \sum_{k=1}^{\infty} a_n = \lim_{n \rightarrow \infty} S_n = L$ then the series **converges**; otherwise it **diverges**.

ex) (see pg 52) Let $S_n = \sum_{k=1}^n \frac{1}{2^k}$.

So then $S_3 =$

Definition

An **infinite series** is the sum of an (infinite) sequence; It has the

form: $\sum_{k=1}^{\infty} a_n = a_1 + a_2 + \cdots + a_n \dots = S_n$. As before, if

$\lim_{n \rightarrow \infty} \sum_{k=1}^{\infty} a_n = \lim_{n \rightarrow \infty} S_n = L$ then the series **converges**; otherwise it **diverges**.

ex) (see pg 52) Let $S_n = \sum_{k=1}^n \frac{1}{2^k}$.

So then $S_3 = \frac{1}{2^1} + \frac{1}{2^2} + \frac{1}{2^3} = \frac{7}{8}$.

Definition

An **infinite series** is the sum of an (infinite) sequence; It has the

form: $\sum_{k=1}^{\infty} a_n = a_1 + a_2 + \cdots + a_n \dots = S_n$. As before, if

$\lim_{n \rightarrow \infty} \sum_{k=1}^{\infty} a_n = \lim_{n \rightarrow \infty} S_n = L$ then the series **converges**; otherwise it **diverges**.

ex) (see pg 52) Let $S_n = \sum_{k=1}^n \frac{1}{2^k}$.

So then $S_3 = \frac{1}{2^1} + \frac{1}{2^2} + \frac{1}{2^3} = \frac{7}{8}$.

What is $\lim_{n \rightarrow \infty} S_n$?

HW 9.1: 21, 55, 60, 66, 68

Next time, 9.2: Sequences