

Section 4.4: Part C: More on Probability and the Standard Distribution

Finite Mathematics MTH 107 Fall 2014

Dr. James Ashe

Mars Hill University

Friday 14th November, 2014



Schedule

Wednesday 12 th November	worksheet 4.4
Friday 14 th November	finding values for given probability 4.4
Monday 17 th November	more section 4.4 stuff/Review
Wednesday 19 th November	Review
Friday 21 st November	Exam 4
Monday 24 th November	Review
Wednesday 26 th November	Thanksgiving Break
Friday 28 th November	Thanksgiving Break
Monday 1 st December	Review
	Final Exam

Finding the value from a probability


Example

A shoe company collects data and finds that male shoe size is normally distributed with a mean shoe size of 11 with a standard deviation of 1.5. If the company wants to make shoes that will fit 99.7% of the male population with sizes centered around the mean, what shoe sizes should they produce?

Finding the value from a probability

Example

A shoe company collects data and finds that male shoe size is normally distributed with a mean shoe size of 11 with a standard deviation of 1.5. If the company wants to make shoes that will fit 99.7% of the male population with sizes centered around the mean, what shoe sizes should they produce?




Finding the value from a probability

Example

A shoe company collects data and finds that male shoe size is normally distributed with a mean shoe size of 11 with a standard deviation of 1.5. If the company wants to make shoes that will fit 99.7% of the male population with sizes centered around the mean, what shoe sizes should they produce?

$$11 + 3 \cdot 1.5 = 15.5$$




Finding the value from a probability

Example

A shoe company collects data and finds that male shoe size is normally distributed with a mean shoe size of 11 with a standard deviation of 1.5. If the company wants to make shoes that will fit 99.7% of the male population with sizes centered around the mean, what shoe sizes should they produce?

$$11 + 3 \cdot 1.5 = 15.5$$

$$11 - 3 \cdot 1.5 = 6.5$$



Finding the value from a probability


Example

A shoe company collects data and finds that male shoe size is normally distributed with a mean shoe size of 11 with a standard deviation of 1.5. If the company wants to make shoes that will fit 99.7% of the male population with sizes centered around the mean, what shoe sizes should they produce?

$$11 + 3 \cdot 1.5 = 15.5$$

$$11 - 3 \cdot 1.5 = 6.5$$

$$\begin{aligned} p(6.5 < x < 15.5) \\ = p(-3 < z < 3) \approx 99.7\% \end{aligned}$$



z	.00	.01	.02	.03	.04	.05	.06	.07	.08	.09
-3.4	.0003	.0003	.0003	.0003	.0003	.0003	.0003	.0003	.0003	.0002
-3.3	.0005	.0005	.0005	.0004	.0004	.0004	.0004	.0004	.0004	.0003
-3.2	.0007	.0007	.0006	.0006	.0006	.0006	.0006	.0005	.0005	.0005
-3.1	.0010	.0009	.0009	.0009	.0008	.0008	.0008	.0008	.0007	.0007
-3.0	.0013	.0013	.0013	.0012	.0012	.0011	.0011	.0011	.0010	.0010
-2.9	.0019	.0018	.0018	.0017	.0016	.0016	.0015	.0015	.0014	.0014
-2.8	.0026	.0025	.0024	.0023	.0023	.0022	.0021	.0021	.0020	.0019
-2.7	.0035	.0034	.0033	.0032	.0031	.0030	.0029	.0028	.0027	.0026
-2.6	.0047	.0045	.0044	.0043	.0041	.0040	.0039	.0038	.0037	.0036
-2.5	.0062	.0060	.0059	.0057	.0055	.0054	.0052	.0051	.0049	.0048
-2.4	.0082	.0080	.0078	.0075	.0073	.0071	.0069	.0068	.0066	.0064
-2.3	.0107	.0104	.0102	.0099	.0096	.0094	.0091	.0089	.0087	.0084
-2.2	.0139	.0136	.0132	.0129	.0125	.0122	.0119	.0116	.0113	.0110
-2.1	.0179	.0174	.0170	.0166	.0162	.0158	.0154	.0150	.0146	.0143
-2.0	.0228	.0222	.0217	.0212	.0207	.0202	.0197	.0192	.0188	.0183
-1.9	.0287	.0281	.0274	.0268	.0262	.0256	.0250	.0244	.0239	.0233
-1.8	.0359	.0351	.0344	.0336	.0329	.0322	.0314	.0307	.0301	.0294
-1.7	.0446	.0436	.0427	.0418	.0409	.0401	.0392	.0384	.0375	.0367
-1.6	.0548	.0537	.0526	.0516	.0505	.0495	.0485	.0475	.0465	.0455
-1.5	.0668	.0655	.0643	.0630	.0618	.0606	.0594	.0582	.0571	.0559
-1.4	.0808	.0793	.0778	.0764	.0749	.0735	.0721	.0708	.0694	.0681
-1.3	.0968	.0951	.0934	.0918	.0901	.0885	.0869	.0853	.0838	.0823
-1.2	.1151	.1131	.1112	.1093	.1075	.1056	.1038	.1020	.1003	.0985
-1.1	.1357	.1335	.1314	.1292	.1271	.1251	.1230	.1210	.1190	.1170
-1.0	.1587	.1562	.1539	.1515	.1492	.1469	.1446	.1423	.1401	.1379
-0.9	.1841	.1814	.1788	.1762	.1736	.1711	.1685	.1660	.1635	.1611
-0.8	.2119	.2090	.2061	.2033	.2005	.1977	.1949	.1922	.1894	.1867
-0.7	.2420	.2389	.2358	.2327	.2296	.2266	.2236	.2206	.2177	.2148
-0.6	.2743	.2709	.2676	.2643	.2611	.2578	.2546	.2514	.2483	.2451
-0.5	.3085	.3050	.3015	.2981	.2946	.2912	.2877	.2843	.2810	.2776
-0.4	.3446	.3409	.3372	.3336	.3300	.3264	.3228	.3192	.3156	.3121
-0.3	.3821	.3783	.3745	.3707	.3669	.3632	.3594	.3557	.3520	.3483
-0.2	.4207	.4168	.4129	.4090	.4052	.4013	.3974	.3936	.3897	.3859
-0.1	.4602	.4562	.4522	.4483	.4443	.4404	.4364	.4325	.4286	.4247
-0.0	.5000	.4960	.4920	.4880	.4840	.4801	.4761	.4721	.4681	.4641

z	.00	.01	.02	.03	.04	.05	.06	.07	.08	.09
0.0	.5000	.5040	.5080	.5120	.5160	.5199	.5239	.5279	.5319	.5359
0.1	.5398	.5438	.5478	.5517	.5557	.5596	.5636	.5675	.5714	.5753
0.2	.5793	.5832	.5871	.5910	.5948	.5987	.6026	.6064	.6103	.6141
0.3	.6179	.6217	.6255	.6293	.6331	.6368	.6406	.6443	.6480	.6517
0.4	.6554	.6591	.6628	.6664	.6700	.6736	.6772	.6808	.6844	.6879
0.5	.6915	.6950	.6985	.7019	.7054	.7088	.7123	.7157	.7190	.7224
0.6	.7257	.7291	.7324	.7357	.7389	.7422	.7454	.7486	.7517	.7549
0.7	.7580	.7611	.7642	.7673	.7704	.7734	.7764	.7794	.7823	.7852
0.8	.7881	.7910	.7939	.7967	.7995	.8023	.8051	.8078	.8106	.8133
0.9	.8159	.8186	.8212	.8238	.8264	.8289	.8315	.8340	.8365	.8389
1.0	.8413	.8438	.8461	.8485	.8508	.8531	.8554	.8577	.8599	.8621
1.1	.8643	.8665	.8686	.8708	.8729	.8749	.8770	.8790	.8810	.8830
1.2	.8849	.8869	.8888	.8907	.8925	.8944	.8962	.8980	.8997	.9015
1.3	.9032	.9049	.9066	.9082	.9099	.9115	.9131	.9147	.9162	.9177
1.4	.9192	.9207	.9222	.9236	.9251	.9265	.9279	.9292	.9306	.9319
1.5	.9332	.9345	.9357	.9370	.9382	.9394	.9406	.9418	.9429	.9441
1.6	.9452	.9463	.9474	.9484	.9495	.9505	.9515	.9525	.9535	.9545
1.7	.9554	.9564	.9573	.9582	.9591	.9599	.9608	.9616	.9625	.9633
1.8	.9641	.9649	.9656	.9664	.9671	.9678	.9686	.9693	.9699	.9706
1.9	.9713	.9719	.9726	.9732	.9738	.9744	.9750	.9756	.9761	.9767
2.0	.9772	.9778	.9783	.9788	.9793	.9798	.9803	.9808	.9812	.9817
2.1	.9821	.9826	.9830	.9834	.9838	.9842	.9846	.9850	.9854	.9857
2.2	.9861	.9864	.9868	.9871	.9875	.9878	.9881	.9884	.9887	.9890
2.3	.9893	.9896	.9898	.9901	.9904	.9906	.9909	.9911	.9913	.9916
2.4	.9918	.9920	.9922	.9925	.9927	.9929	.9931	.9932	.9934	.9936
2.5	.9938	.9940	.9941	.9943	.9945	.9946	.9948	.9949	.9951	.9952
2.6	.9953	.9955	.9956	.9957	.9959	.9960	.9961	.9962	.9963	.9964
2.7	.9965	.9966	.9967	.9968	.9969	.9970	.9971	.9972	.9973	.9974
2.8	.9974	.9975	.9976	.9977	.9977	.9978	.9979	.9979	.9980	.9981
2.9	.9981	.9982	.9982	.9983	.9984	.9984	.9985	.9985	.9986	.9986
3.0	.9987	.9987	.9987	.9988	.9988	.9989	.9989	.9989	.9990	.9990
3.1	.9990	.9991	.9991	.9991	.9992	.9992	.9992	.9992	.9993	.9993
3.2	.9993	.9993	.9994	.9994	.9994	.9994	.9994	.9995	.9995	.9995
3.3	.9995	.9995	.9995	.9996	.9996	.9996	.9996	.9996	.9996	.9997
3.4	.9997	.9997	.9997	.9997	.9997	.9997	.9997	.9997	.9997	.9998

Finding the value for a probability

Example

A shoe company collects data and finds that male shoe size is normally distributed with a mean shoe size of 11 with a standard deviation of 1.5.

- a. How many *standard deviations* separate the bottom 74.72% from the remaining population.

Finding the value for a probability

Example

A shoe company collects data and finds that male shoe size is normally distributed with a mean shoe size of 11 with a standard deviation of 1.5.

- a. How many *standard deviations* separate the bottom 74.72% from the remaining population.

a. $p(z < c) = 74.2\%$

Finding the value for a probability

Example

A shoe company collects data and finds that male shoe size is normally distributed with a mean shoe size of 11 with a standard deviation of 1.5.

- a. How many *standard deviations* separate the bottom 74.72% from the remaining population.

a. $p(z < c) = 74.2\%$

z	.00	.01	.02	.03	.04	.05	.06
0.0	.5000	.5040	.5080	.5120	.5160	.5199	.5239
0.1	.5398	.5438	.5478	.5517	.5557	.5596	.5636
0.2	.5793	.5832	.5871	.5910	.5948	.5987	.6026
0.3	.6179	.6217	.6255	.6293	.6331	.6368	.6406
0.4	.6554	.6591	.6628	.6664	.6700	.6736	.6772
0.5	.6915	.6950	.6985	.7019	.7054	.7088	.7122
0.6	.7257	.7291	.7324	.7357	.7389	.7422	.7454
0.7	.7580	.7611	.7642	.7673	.7704	.7734	.7764
0.8	.7881	.7910	.7939	.7967	.7995	.8023	.8051
0.9	.8159	.8186	.8212	.8238	.8264	.8289	.8315
1.0	.8413	.8438	.8461	.8485	.8508	.8531	.8554
1.1	.8643	.8665	.8686	.8708	.8729	.8749	.8770
1.2	.8849	.8869	.8888	.8907	.8925	.8944	.8962
1.3	.9032	.9049	.9066	.9082	.9099	.9115	.9131

Finding the value for a probability

Example

A shoe company collects data and finds that male shoe size is normally distributed with a mean shoe size of 11 with a standard deviation of 1.5.

- a. How many *standard deviations* separate the bottom 74.72% from the remaining population.

a. $p(z < c) = 74.2\%$

z	.00	.01	.02	.03	.04	.05	.06
0.0	.5000	.5040	.5080	.5120	.5160	.5199	.5239
0.1	.5398	.5438	.5478	.5517	.5557	.5596	.5636
0.2	.5793	.5832	.5871	.5910	.5948	.5987	.6026
0.3	.6179	.6217	.6255	.6293	.6331	.6368	.6406
0.4	.6554	.6591	.6628	.6664	.6700	.6736	.6772
0.5	.6915	.6950	.6985	.7019	.7054	.7088	.7122
0.6	.7257	.7291	.7324	.7357	.7389	.7422	.7454
0.7	.7580	.7611	.7642	.7673	.7704	.7734	.7764
0.8	.7881	.7910	.7939	.7967	.7995	.8023	.8051
0.9	.8159	.8186	.8212	.8238	.8264	.8289	.8315
1.0	.8413	.8438	.8461	.8485	.8508	.8531	.8554
1.1	.8643	.8665	.8686	.8708	.8729	.8749	.8770
1.2	.8849	.8869	.8888	.8907	.8925	.8944	.8962
1.3	.9032	.9049	.9066	.9082	.9099	.9115	.9131

Finding the value for a probability

Example

A shoe company collects data and finds that male shoe size is normally distributed with a mean shoe size of 11 with a standard deviation of 1.5.

- a. How many *standard deviations* separate the bottom 74.72% from the remaining population.

a. $p(z < c) = 74.2\%$

<i>z</i>	.00	.01	.02	.03	.04	.05	.06
0.0	.5000	.5040	.5080	.5120	.5160	.5199	.5239
0.1	.5398	.5438	.5478	.5517	.5557	.5596	.5636
0.2	.5793	.5832	.5871	.5910	.5948	.5987	.6026
0.3	.6179	.6217	.6255	.6293	.6331	.6368	.6406
0.4	.6554	.6591	.6628	.6664	.6700	.6736	.6772
0.5	.6915	.6950	.6985	.7019	.7054	.7088	.7122
0.6	.7257	.7291	.7324	.7357	.7389	.7422	.7454
0.7	.7580	.7611	.7642	.7673	.7704	.7734	.7764
0.8	.7881	.7910	.7939	.7967	.7995	.8023	.8051
0.9	.8159	.8186	.8212	.8238	.8264	.8289	.8315
1.0	.8413	.8438	.8461	.8485	.8508	.8531	.8554
1.1	.8643	.8665	.8686	.8708	.8729	.8749	.8770
1.2	.8849	.8869	.8888	.8907	.8925	.8944	.8962
1.3	.9032	.9049	.9066	.9082	.9099	.9115	.9131

Finding the value for a probability

Example

A shoe company collects data and finds that male shoe size is normally distributed with a mean shoe size of 11 with a standard deviation of 1.5.

- a. How many *standard deviations* separate the bottom 74.72% from the remaining population.

a. $p(z < c) = 74.2\%$

<i>z</i>	.00	.01	.02	.03	.04	.05	.06
0.0	.5000	.5040	.5080	.5120	.5160	.5199	.5239
0.1	.5398	.5438	.5478	.5517	.5557	.5596	.5636
0.2	.5793	.5832	.5871	.5910	.5948	.5987	.6026
0.3	.6179	.6217	.6255	.6293	.6331	.6368	.6406
0.4	.6554	.6591	.6628	.6664	.6700	.6736	.6772
0.5	.6915	.6950	.6985	.7019	.7054	.7088	.7122
0.6	.7257	.7291	.7324	.7357	.7389	.7422	.7454
0.7	.7580	.7611	.7642	.7673	.7704	.7734	.7764
0.8	.7881	.7910	.7939	.7967	.7995	.8023	.8051
0.9	.8159	.8186	.8212	.8238	.8264	.8289	.8315
1.0	.8413	.8438	.8461	.8485	.8508	.8531	.8554
1.1	.8643	.8665	.8686	.8708	.8729	.8749	.8770
1.2	.8849	.8869	.8888	.8907	.8925	.8944	.8962
1.3	.9032	.9049	.9066	.9082	.9099	.9115	.9131

Finding the value for a probability

Example

A shoe company collects data and finds that male shoe size is normally distributed with a mean shoe size of 11 with a standard deviation of 1.5.

- a. How many *standard deviations* separate the bottom 74.72% from the remaining population.

a. $p(z < c) = 74.2\%$

<i>z</i>	.00	.01	.02	.03	.04	.05	.06
0.0	.5000	.5040	.5080	.5120	.5160	.5199	.5239
0.1	.5398	.5438	.5478	.5517	.5557	.5596	.5636
0.2	.5793	.5832	.5871	.5910	.5948	.5987	.6026
0.3	.6179	.6217	.6255	.6293	.6331	.6368	.6406
0.4	.6554	.6591	.6628	.6664	.6700	.6736	.6772
0.5	.6915	.6950	.6985	.7019	.7054	.7088	.7122
0.6	.7257	.7291	.7324	.7357	.7389	.7422	.7454
0.7	.7580	.7611	.7642	.7673	.7704	.7734	.7764
0.8	.7881	.7910	.7939	.7967	.7995	.8023	.8051
0.9	.8159	.8186	.8212	.8238	.8264	.8289	.8315
1.0	.8413	.8438	.8461	.8485	.8508	.8531	.8554
1.1	.8643	.8665	.8686	.8708	.8729	.8749	.8770
1.2	.8849	.8869	.8888	.8907	.8925	.8944	.8962
1.3	.9032	.9049	.9066	.9082	.9099	.9115	.9131

.742 → .67
area *z* — score

Finding the value for a probability

Example

A shoe company collects data and finds that male shoe size is normally distributed with a mean shoe size of 11 with a standard deviation of 1.5.

- a. How many *standard deviations* separate the bottom 74.72% from the remaining population.

a. $p(z < c) = 74.2\%$

$$\begin{array}{ccc} .742 & \rightarrow & .67 \\ \text{area} & & z - \text{score} \end{array}$$

$$p(z < .67) = 74.2$$

Finding the value for a probability

Example


A shoe company collects data and finds that male shoe size is normally distributed with a mean shoe size of 11 with a standard deviation of 1.5.

- a. How many *standard deviations* separate the bottom 74.72% from the remaining population.

a. $p(z < c) = 74.2\%$

$$\begin{array}{ccc} .742 & \rightarrow & .67 \\ \text{area} & & z\text{-score} \end{array}$$

$$p(z < .67) = 74.2$$



z	.00	.01	.02	.03	.04	.05	.06	.07	.08	.09
-3.4	.0003	.0003	.0003	.0003	.0003	.0003	.0003	.0003	.0003	.0002
-3.3	.0005	.0005	.0005	.0004	.0004	.0004	.0004	.0004	.0004	.0003
-3.2	.0007	.0007	.0006	.0006	.0006	.0006	.0006	.0005	.0005	.0005
-3.1	.0010	.0009	.0009	.0009	.0008	.0008	.0008	.0008	.0007	.0007
-3.0	.0013	.0013	.0013	.0012	.0012	.0011	.0011	.0011	.0010	.0010
-2.9	.0019	.0018	.0018	.0017	.0016	.0016	.0015	.0015	.0014	.0014
-2.8	.0026	.0025	.0024	.0023	.0023	.0022	.0021	.0021	.0020	.0019
-2.7	.0035	.0034	.0033	.0032	.0031	.0030	.0029	.0028	.0027	.0026
-2.6	.0047	.0045	.0044	.0043	.0041	.0040	.0039	.0038	.0037	.0036
-2.5	.0062	.0060	.0059	.0057	.0055	.0054	.0052	.0051	.0049	.0048
-2.4	.0082	.0080	.0078	.0075	.0073	.0071	.0069	.0068	.0066	.0064
-2.3	.0107	.0104	.0102	.0099	.0096	.0094	.0091	.0089	.0087	.0084
-2.2	.0139	.0136	.0132	.0129	.0125	.0122	.0119	.0116	.0113	.0110
-2.1	.0179	.0174	.0170	.0166	.0162	.0158	.0154	.0150	.0146	.0143
-2.0	.0228	.0222	.0217	.0212	.0207	.0202	.0197	.0192	.0188	.0183
-1.9	.0287	.0281	.0274	.0268	.0262	.0256	.0250	.0244	.0239	.0233
-1.8	.0359	.0351	.0344	.0336	.0329	.0322	.0314	.0307	.0301	.0294
-1.7	.0446	.0436	.0427	.0418	.0409	.0401	.0392	.0384	.0375	.0367
-1.6	.0548	.0537	.0526	.0516	.0505	.0495	.0485	.0475	.0465	.0455
-1.5	.0668	.0655	.0643	.0630	.0618	.0606	.0594	.0582	.0571	.0559
-1.4	.0808	.0793	.0778	.0764	.0749	.0735	.0721	.0708	.0694	.0681
-1.3	.0968	.0951	.0934	.0918	.0901	.0885	.0869	.0853	.0838	.0823
-1.2	.1151	.1131	.1112	.1093	.1075	.1056	.1038	.1020	.1003	.0985
-1.1	.1357	.1335	.1314	.1292	.1271	.1251	.1230	.1210	.1190	.1170
-1.0	.1587	.1562	.1539	.1515	.1492	.1469	.1446	.1423	.1401	.1379
-0.9	.1841	.1814	.1788	.1762	.1736	.1711	.1685	.1660	.1635	.1611
-0.8	.2119	.2090	.2061	.2033	.2005	.1977	.1949	.1922	.1894	.1867
-0.7	.2420	.2389	.2358	.2327	.2296	.2266	.2236	.2206	.2177	.2148
-0.6	.2743	.2709	.2676	.2643	.2611	.2578	.2546	.2514	.2483	.2451
-0.5	.3085	.3050	.3015	.2981	.2946	.2912	.2877	.2843	.2810	.2776
-0.4	.3446	.3409	.3372	.3336	.3300	.3264	.3228	.3192	.3156	.3121
-0.3	.3821	.3783	.3745	.3707	.3669	.3632	.3594	.3557	.3520	.3483
-0.2	.4207	.4168	.4129	.4090	.4052	.4013	.3974	.3936	.3897	.3859
-0.1	.4602	.4562	.4522	.4483	.4443	.4404	.4364	.4325	.4286	.4247
-0.0	.5000	.4960	.4920	.4880	.4840	.4801	.4761	.4721	.4681	.4641

z	.00	.01	.02	.03	.04	.05	.06	.07	.08	.09
0.0	.5000	.5040	.5080	.5120	.5160	.5199	.5239	.5279	.5319	.5359
0.1	.5398	.5438	.5478	.5517	.5557	.5596	.5636	.5675	.5714	.5753
0.2	.5793	.5832	.5871	.5910	.5948	.5987	.6026	.6064	.6103	.6141
0.3	.6179	.6217	.6255	.6293	.6331	.6368	.6406	.6443	.6480	.6517
0.4	.6554	.6591	.6628	.6664	.6700	.6736	.6772	.6808	.6844	.6879
0.5	.6915	.6950	.6985	.7019	.7054	.7088	.7123	.7157	.7190	.7224
0.6	.7257	.7291	.7324	.7357	.7389	.7422	.7454	.7486	.7517	.7549
0.7	.7580	.7611	.7642	.7673	.7704	.7734	.7764	.7794	.7823	.7852
0.8	.7881	.7910	.7939	.7967	.7995	.8023	.8051	.8078	.8106	.8133
0.9	.8159	.8186	.8212	.8238	.8264	.8289	.8315	.8340	.8365	.8389
1.0	.8413	.8438	.8461	.8485	.8508	.8531	.8554	.8577	.8599	.8621
1.1	.8643	.8665	.8686	.8708	.8729	.8749	.8770	.8790	.8810	.8830
1.2	.8849	.8869	.8888	.8907	.8925	.8944	.8962	.8980	.8997	.9015
1.3	.9032	.9049	.9066	.9082	.9099	.9115	.9131	.9147	.9162	.9177
1.4	.9192	.9207	.9222	.9236	.9251	.9265	.9279	.9292	.9306	.9319
1.5	.9332	.9345	.9357	.9370	.9382	.9394	.9406	.9418	.9429	.9441
1.6	.9452	.9463	.9474	.9484	.9495	.9505	.9515	.9525	.9535	.9545
1.7	.9554	.9564	.9573	.9582	.9591	.9599	.9608	.9616	.9625	.9633
1.8	.9641	.9649	.9656	.9664	.9671	.9678	.9686	.9693	.9699	.9706
1.9	.9713	.9719	.9726	.9732	.9738	.9744	.9750	.9756	.9761	.9767
2.0	.9772	.9778	.9783	.9788	.9793	.9798	.9803	.9808	.9812	.9817
2.1	.9821	.9826	.9830	.9834	.9838	.9842	.9846	.9850	.9854	.9857
2.2	.9861	.9864	.9868	.9871	.9875	.9878	.9881	.9884	.9887	.9890
2.3	.9893	.9896	.9898	.9901	.9904	.9906	.9909	.9911	.9913	.9916
2.4	.9918	.9920	.9922	.9925	.9927	.9929	.9931	.9932	.9934	.9936
2.5	.9938	.9940	.9941	.9943	.9945	.9946	.9948	.9949	.9951	.9952
2.6	.9953	.9955	.9956	.9957	.9959	.9960	.9961	.9962	.9963	.9964
2.7	.9965	.9966	.9967	.9968	.9969	.9970	.9971	.9972	.9973	.9974
2.8	.9974	.9975	.9976	.9977	.9977	.9978	.9979	.9979	.9980	.9981
2.9	.9981	.9982	.9982	.9983	.9984	.9984	.9985	.9985	.9986	.9986
3.0	.9987	.9987	.9987	.9988	.9988	.9989	.9989	.9989	.9990	.9990
3.1	.9990	.9991	.9991	.9991	.9992	.9992	.9992	.9992	.9993	.9993
3.2	.9993	.9993	.9994	.9994	.9994	.9994	.9994	.9995	.9995	.9995
3.3	.9995	.9995	.9995	.9996	.9996	.9996	.9996	.9996	.9996	.9997
3.4	.9997	.9997	.9997	.9997	.9997	.9997	.9997	.9997	.9997	.9998

Finding the value from a probability

Example

A shoe company collects data and finds that male shoe size is normally distributed with a mean shoe size of 11 with a standard deviation of 1.5.

- b. Find the shoe size that separates the bottom 15.15% from the rest of the population.

$$p(x < 9.455) = 15.15\%$$

Finding the value from a probability

Example

A shoe company collects data and finds that male shoe size is normally distributed with a mean shoe size of 11 with a standard deviation of 1.5.

- b. Find the shoe size that separates the bottom 15.15% from the rest of the population.

$$b. \quad p(x < c) = 15.15\%$$

$$p(x < 9.455) = 15.15\%$$

Finding the value from a probability

Example

A shoe company collects data and finds that male shoe size is normally distributed with a mean shoe size of 11 with a standard deviation of 1.5.

- b. Find the shoe size that separates the bottom 15.15% from the rest of the population.

$$\begin{array}{lll} \text{b. } p(x < c) = 15.15\% & & \\ .1515 & \rightarrow & -1.03 \\ \text{area} & & z\text{-score} \end{array}$$

$$p(x < 9.455) = 15.15\%$$

Finding the value from a probability

Example

A shoe company collects data and finds that male shoe size is normally distributed with a mean shoe size of 11 with a standard deviation of 1.5.

- b. Find the shoe size that separates the bottom 15.15% from the rest of the population.

Convert z-score to data value

$$c = z \cdot \sigma + \mu$$

$$\begin{array}{lll} \text{b. } p(x < c) = 15.15\% \\ .1515 & \rightarrow & -1.03 \\ \text{area} & & z\text{-score} \end{array}$$

$$p(x < 9.455) = 15.15\%$$

Finding the value from a probability

Example

A shoe company collects data and finds that male shoe size is normally distributed with a mean shoe size of 11 with a standard deviation of 1.5.


- b. Find the shoe size that separates the bottom 15.15% from the rest of the population.

Convert z-score to data value

$$c = z \cdot \sigma + \mu$$

b. $p(x < c) = 15.15\%$

$$\begin{array}{ccc} .1515 & \rightarrow & -1.03 \\ \text{area} & & z\text{-score} \end{array}$$



$$p(x < 9.455) = 15.15\%$$

Finding the value from a probability

Example

A shoe company collects data and finds that male shoe size is normally distributed with a mean shoe size of 11 with a standard deviation of 1.5.


- b. Find the shoe size that separates the bottom 15.15% from the rest of the population.

Convert z-score to data value

$$c = z \cdot \sigma + \mu$$

b. $p(x < c) = 15.15\%$

$$\begin{array}{ccc} .1515 & \rightarrow & -1.03 \\ \text{area} & & z\text{-score} \end{array}$$



$$c = z \cdot \sigma + \mu$$

$$p(x < 9.455) = 15.15\%$$

Finding the value from a probability

Example

A shoe company collects data and finds that male shoe size is normally distributed with a mean shoe size of 11 with a standard deviation of 1.5.


- b. Find the shoe size that separates the bottom 15.15% from the rest of the population.

Convert z-score to data value

$$c = z \cdot \sigma + \mu$$

b. $p(x < c) = 15.15\%$

$$\begin{array}{ccc} .1515 & \rightarrow & -1.03 \\ \text{area} & & z\text{-score} \end{array}$$



$$c = (-1.03) \cdot \sigma + \mu$$

$$p(x < 9.455) = 15.15\%$$

Finding the value from a probability

Example

A shoe company collects data and finds that male shoe size is normally distributed with a mean shoe size of 11 with a standard deviation of 1.5.


- b. Find the shoe size that separates the bottom 15.15% from the rest of the population.

Convert z-score to data value

$$c = z \cdot \sigma + \mu$$

b. $p(x < c) = 15.15\%$

$$\begin{array}{ccc} .1515 & \rightarrow & -1.03 \\ \text{area} & & z\text{-score} \end{array}$$



$$c = (-1.03) \cdot 1.5 + 11$$

$$p(x < 9.455) = 15.15\%$$

Finding the value from a probability

Example

A shoe company collects data and finds that male shoe size is normally distributed with a mean shoe size of 11 with a standard deviation of 1.5.


- b. Find the shoe size that separates the bottom 15.15% from the rest of the population.

Convert z-score to data value

$$c = z \cdot \sigma + \mu$$

b. $p(x < c) = 15.15\%$

$$\begin{array}{ccc} .1515 & \rightarrow & -1.03 \\ \text{area} & & z\text{-score} \end{array}$$



$$c = 9.455$$

$$p(x < 9.455) = 15.15\%$$

z	.00	.01	.02	.03	.04	.05	.06	.07	.08	.09
-3.4	.0003	.0003	.0003	.0003	.0003	.0003	.0003	.0003	.0003	.0002
-3.3	.0005	.0005	.0005	.0004	.0004	.0004	.0004	.0004	.0004	.0003
-3.2	.0007	.0007	.0006	.0006	.0006	.0006	.0006	.0005	.0005	.0005
-3.1	.0010	.0009	.0009	.0009	.0008	.0008	.0008	.0008	.0007	.0007
-3.0	.0013	.0013	.0013	.0012	.0012	.0011	.0011	.0011	.0010	.0010
-2.9	.0019	.0018	.0018	.0017	.0016	.0016	.0015	.0015	.0014	.0014
-2.8	.0026	.0025	.0024	.0023	.0023	.0022	.0021	.0021	.0020	.0019
-2.7	.0035	.0034	.0033	.0032	.0031	.0030	.0029	.0028	.0027	.0026
-2.6	.0047	.0045	.0044	.0043	.0041	.0040	.0039	.0038	.0037	.0036
-2.5	.0062	.0060	.0059	.0057	.0055	.0054	.0052	.0051	.0049	.0048
-2.4	.0082	.0080	.0078	.0075	.0073	.0071	.0069	.0068	.0066	.0064
-2.3	.0107	.0104	.0102	.0099	.0096	.0094	.0091	.0089	.0087	.0084
-2.2	.0139	.0136	.0132	.0129	.0125	.0122	.0119	.0116	.0113	.0110
-2.1	.0179	.0174	.0170	.0166	.0162	.0158	.0154	.0150	.0146	.0143
-2.0	.0228	.0222	.0217	.0212	.0207	.0202	.0197	.0192	.0188	.0183
-1.9	.0287	.0281	.0274	.0268	.0262	.0256	.0250	.0244	.0239	.0233
-1.8	.0359	.0351	.0344	.0336	.0329	.0322	.0314	.0307	.0301	.0294
-1.7	.0446	.0436	.0427	.0418	.0409	.0401	.0392	.0384	.0375	.0367
-1.6	.0548	.0537	.0526	.0516	.0505	.0495	.0485	.0475	.0465	.0455
-1.5	.0668	.0655	.0643	.0630	.0618	.0606	.0594	.0582	.0571	.0559
-1.4	.0808	.0793	.0778	.0764	.0749	.0735	.0721	.0708	.0694	.0681
-1.3	.0968	.0951	.0934	.0918	.0901	.0885	.0869	.0853	.0838	.0823
-1.2	.1151	.1131	.1112	.1093	.1075	.1056	.1038	.1020	.1003	.0985
-1.1	.1357	.1335	.1314	.1292	.1271	.1251	.1230	.1210	.1190	.1170
-1.0	.1587	.1562	.1539	.1515	.1492	.1469	.1446	.1423	.1401	.1379
-0.9	.1841	.1814	.1788	.1762	.1736	.1711	.1685	.1660	.1635	.1611
-0.8	.2119	.2090	.2061	.2033	.2005	.1977	.1949	.1922	.1894	.1867
-0.7	.2420	.2389	.2358	.2327	.2296	.2266	.2236	.2206	.2177	.2148
-0.6	.2743	.2709	.2676	.2643	.2611	.2578	.2546	.2514	.2483	.2451
-0.5	.3085	.3050	.3015	.2981	.2946	.2912	.2877	.2843	.2810	.2776
-0.4	.3446	.3409	.3372	.3336	.3300	.3264	.3228	.3192	.3156	.3121
-0.3	.3821	.3783	.3745	.3707	.3669	.3632	.3594	.3557	.3520	.3483
-0.2	.4207	.4168	.4129	.4090	.4052	.4013	.3974	.3936	.3897	.3859
-0.1	.4602	.4562	.4522	.4483	.4443	.4404	.4364	.4325	.4286	.4247
-0.0	.5000	.4960	.4920	.4880	.4840	.4801	.4761	.4721	.4681	.4641

z	.00	.01	.02	.03	.04	.05	.06	.07	.08	.09
0.0	.5000	.5040	.5080	.5120	.5160	.5199	.5239	.5279	.5319	.5359
0.1	.5398	.5438	.5478	.5517	.5557	.5596	.5636	.5675	.5714	.5753
0.2	.5793	.5832	.5871	.5910	.5948	.5987	.6026	.6064	.6103	.6141
0.3	.6179	.6217	.6255	.6293	.6331	.6368	.6406	.6443	.6480	.6517
0.4	.6554	.6591	.6628	.6664	.6700	.6736	.6772	.6808	.6844	.6879
0.5	.6915	.6950	.6985	.7019	.7054	.7088	.7123	.7157	.7190	.7224
0.6	.7257	.7291	.7324	.7357	.7389	.7422	.7454	.7486	.7517	.7549
0.7	.7580	.7611	.7642	.7673	.7704	.7734	.7764	.7794	.7823	.7852
0.8	.7881	.7910	.7939	.7967	.7995	.8023	.8051	.8078	.8106	.8133
0.9	.8159	.8186	.8212	.8238	.8264	.8289	.8315	.8340	.8365	.8389
1.0	.8413	.8438	.8461	.8485	.8508	.8531	.8554	.8577	.8599	.8621
1.1	.8643	.8665	.8686	.8708	.8729	.8749	.8770	.8790	.8810	.8830
1.2	.8849	.8869	.8888	.8907	.8925	.8944	.8962	.8980	.8997	.9015
1.3	.9032	.9049	.9066	.9082	.9099	.9115	.9131	.9147	.9162	.9177
1.4	.9192	.9207	.9222	.9236	.9251	.9265	.9279	.9292	.9306	.9319
1.5	.9332	.9345	.9357	.9370	.9382	.9394	.9406	.9418	.9429	.9441
1.6	.9452	.9463	.9474	.9484	.9495	.9505	.9515	.9525	.9535	.9545
1.7	.9554	.9564	.9573	.9582	.9591	.9599	.9608	.9616	.9625	.9633
1.8	.9641	.9649	.9656	.9664	.9671	.9678	.9686	.9693	.9699	.9706
1.9	.9713	.9719	.9726	.9732	.9738	.9744	.9750	.9756	.9761	.9767
2.0	.9772	.9778	.9783	.9788	.9793	.9798	.9803	.9808	.9812	.9817
2.1	.9821	.9826	.9830	.9834	.9838	.9842	.9846	.9850	.9854	.9857
2.2	.9861	.9864	.9868	.9871	.9875	.9878	.9881	.9884	.9887	.9890
2.3	.9893	.9896	.9898	.9901	.9904	.9906	.9909	.9911	.9913	.9916
2.4	.9918	.9920	.9922	.9925	.9927	.9929	.9931	.9932	.9934	.9936
2.5	.9938	.9940	.9941	.9943	.9945	.9946	.9948	.9949	.9951	.9952
2.6	.9953	.9955	.9956	.9957	.9959	.9960	.9961	.9962	.9963	.9964
2.7	.9965	.9966	.9967	.9968	.9969	.9970	.9971	.9972	.9973	.9974
2.8	.9974	.9975	.9976	.9977	.9977	.9978	.9979	.9979	.9980	.9981
2.9	.9981	.9982	.9982	.9983	.9984	.9984	.9985	.9985	.9986	.9986
3.0	.9987	.9987	.9987	.9988	.9988	.9989	.9989	.9989	.9990	.9990
3.1	.9990	.9991	.9991	.9991	.9992	.9992	.9992	.9992	.9993	.9993
3.2	.9993	.9993	.9994	.9994	.9994	.9994	.9994	.9995	.9995	.9995
3.3	.9995	.9995	.9995	.9996	.9996	.9996	.9996	.9996	.9996	.9997
3.4	.9997	.9997	.9997	.9997	.9997	.9997	.9997	.9997	.9997	.9998


Problem 4.4.1. The results of a statewide exam for assessing math skills of realtors was normally distributed with a mean of 72 and a standard deviation of 12. Those who score in the top 10% will receive a certificate, and those who score in the bottom 20% will be required to attend a remedial class.

- a. What score does a realtor need in order to receive the certificate?
- b. What score will dictate that a realtor must attend the remedial class?

Other Fun Stuff

Often plotted data lies *close* to a line.


Hours Studying	Exam Scores
2	55
4	70
5	64
6	80
9	87



Other Fun Stuff

Often plotted data lies *close* to a line.


Hours Studying	Exam Scores
2	55
4	70
5	64
6	80
9	87



Other Fun Stuff

Often plotted data lies *close* to a line.


Hours Studying	Exam Scores
2	55
4	70
5	64
6	80
9	87



Other Fun Stuff

Often plotted data lies *close* to a line.


Hours Studying	Exam Scores
2	55
4	70
5	64
6	80
9	87



Other Fun Stuff

Often plotted data lies *close* to a line.

Hours Studying	Exam Scores
2	55
4	70
5	64
6	80
9	87




Other Fun Stuff

Often plotted data lies *close* to a line.

Hours Studying	Exam Scores
2	55
4	70
5	64
6	80
9	87

- We call this a *linear relation*.




Other Fun Stuff

Often plotted data lies *close* to a line.

Hours Studying	Exam Scores
2	55
4	70
5	64
6	80
9	87

- We call this a *linear relation*.
- But which line?




Other Fun Stuff

Often plotted data lies *close* to a line.

Hours Studying	Exam Scores
2	55
4	70
5	64
6	80
9	87

- We call this a *linear relation*.
- But which line?
- We need a reasonable *approximation*.



The Least Squares Line (“best-fit line”)

Hours Studying	Exam Scores
2	55
4	70
5	64
6	80
9	87

The Least Squares Line (“best-fit line”)

Hours Studying	Exam Scores
2	55
4	70
5	64
6	80
9	87

Least Square Line *slope*

$$m = \frac{n(\sum xy) - (\sum x)(\sum y)}{n(\sum x^2) - (\sum x)^2}$$

The Least Squares Line (“best-fit line”)

Hours Studying	Exam Scores
2	55
4	70
5	64
6	80
9	87

Least Square Line *slope*

$$m = \frac{n(\sum xy) - (\sum x)(\sum y)}{n(\sum x^2) - (\sum x)^2}$$

Least Square Line *y-int*

$$b = \frac{\sum y - m(\sum x)}{n}$$

The Least Squares Line (“best-fit line”)

Hours Studying	Exam Scores
2	55
4	70
5	64
6	80
9	87

Least Square Line *slope*

$$m = \frac{n(\sum xy) - (\sum x)(\sum y)}{n(\sum x^2) - (\sum x)^2}$$

Least Square Line *y-int*

$$b = \frac{\sum y - m(\sum x)}{n}$$


Least Square Line

$$y = mx + b$$

Hours Studying	Exam Scores
2	55
4	70
5	64
6	80
9	87

Equation of the least squares line

$$y = (4.54)x + 47.57$$




Hours Studying	Exam Scores
2	55
4	70
5	64
6	80
9	87

Equation of the least squares line

$$y = (4.54)x + 47.57$$

Correlation Coefficient


$$r = \frac{n(\sum xy) - (\sum x)(\sum y)}{\sqrt{n(\sum x^2) - (\sum x)^2} \sqrt{n(\sum y^2) - (\sum y)^2}}$$



Problem 4.4.2. Find the equation of the least squares line.

The paired data below consist of the costs of advertising (in thousands) and the number of products sold (in thousands).


Cost(x)	9	2	3	4	2	5	9	10
Number(y)	85	52	55	68	67	86	83	73



Problem 4.4.2. Find the equation of the least squares line.

The paired data below consist of the costs of advertising (in thousands) and the number of products sold (in thousands).

Cost(x)	9	2	3	4	2	5	9	10
Number(y)	85	52	55	68	67	86	83	73




Problem 4.4.2. Find the equation of the least squares line.

The paired data below consist of the costs of advertising (in thousands) and the number of products sold (in thousands).

Cost(x)	9	2	3	4	2	5	9	10
Number(y)	85	52	55	68	67	86	83	73

$$y = 2.79x + 55.8$$




Problem 4.4.2. Find the equation of the least squares line.

The paired data below consist of the costs of advertising (in thousands) and the number of products sold (in thousands).

Cost(x)	9	2	3	4	2	5	9	10
Number(y)	85	52	55	68	67	86	83	73

$$y = 2.79x + 55.8$$

$$r \approx .708$$



Margin of Error

Margin of Error

- A poll of 30 people is taken.

Margin of Error

- A poll of 30 people is taken.
- There are many ways to select 30 people from the population.

Margin of Error

- A poll of 30 people is taken.
- There are many ways to select 30 people from the population.
- The set of all possible polls is also a population with its own μ and σ .

Margin of Error

- A poll of 30 people is taken.
- There are many ways to select 30 people from the population.
- The set of all possible polls is also a population with its own μ and σ .
- 5% chance a randomly selected poll is within $2 \cdot \sigma$ of the actual result.

Margin of Error

- A poll of 30 people is taken.
- There are many ways to select 30 people from the population.
- The set of all possible polls is also a population with its own μ and σ .
- 5% chance a randomly selected poll is within $2 \cdot \sigma$ of the actual result.

