### PROTOCOLE OPERATOIRE / OPERATIVE REPORT

Site: Ste-Agathe Hospital Date: 2025-06-26 14:02

Anesthésiste / Anesthetist: Dr. Omar Fahmy

Chirurgien / Surgeon: Dr. Aisha Patel Assistant(s): Dr. resident Leo Morel

Diagnostic préopératoire / Pre-operative diagnosis:

### APPENDICITIS WITH ABSCESS.

Diagnostic postopératoire / Post-operative diagnosis:

PERFORATED APPENDICITIS.

Opération / Operation:

#### OPEN APPENDECTOMY WITH PELVIC LAVAGE.

Tissu envoyé en pathologie / Tissue sent to pathology: Appendix with surrounding lymph nodes

Anesthésie / Anesthesia: General anesthesia and epidural block

# Historique et constatations opératoires / History and operative findings:

A 12-year-old male who presented with abdominal pain with rebound tenderness. Initially evaluated 4 days prior and diagnosed with IBD. Now has normal WBC, normal CRP, low-grade fever. Imaging: MRI demonstrating RLQ inflammation.

# Procédure(s) opératoire(s) / Operative procedure(s):

Patient in supine position. General anesthesia and epidural block administered. Time-out was performed and abdomen prepped in sterile fashion. A vertical infraumbilical incision is made and carried down to the fascia which is incised sharply. Two working ports are established in the right and left lower quadrants. No iatrogenic injuries occurred during trocar placement. Upon entering the abdominal cavity, significant adhesions noted. Operative findings included suppurative appendix and mild inflammation. A moderate amount of purulent material was present in the pelvis. The omentum was wrapped around the inflamed appendix. Surrounding omentum and bowel are separated from the inflammatory mass. The appendiceal artery is ligated and divided. Absorbable ligatures are applied prior to amputation. Specimen placed in EndoCatch bag for removal. Saline is used for thorough irrigation of all quadrants. Umbilical port site is closed with Vicryl 3-0 and skin with subcuticular Vicryl 4-0. No bowel injury noted.

The patient will be continued on IV antibiotics and advanced to diet as tolerated. The patient tolerated the procedure well with minimal blood loss.

Case ID: CASE-YTKH6O-13527