PROTOCOLE OPERATOIRE / OPERATIVE REPORT

Site: St. Mary's Hospital Date: 2023-10-17 07:44

Anesthésiste / Anesthetist: Dr. Lisa Garcia Chirurgien / Surgeon: Dr. Daniel Fortin Assistant(s): Dr. resident Leo Morel

Diagnostic préopératoire / Pre-operative diagnosis:

APPENDICITIS WITH FREE FLUID.

Diagnostic postopératoire / Post-operative diagnosis:

ACUTE APPENDICITIS.

Opération / Operation:

APPENDECTOMY WITH REMOVAL OF APPENDICOLITH.

Tissu envoyé en pathologie / Tissue sent to pathology: Appendix and inflamed tissue

Anesthésie / Anesthesia: General anesthesia

Historique et constatations opératoires / History and operative findings:

Patient (1 years, male) presented with abdominal pain with elevated WBC, markedly elevated WBC, normal CRP. Imaging: ultrasound reporting lymphadenopathy. Past medical history is otherwise unremarkable.

Procédure(s) opératoire(s) / Operative procedure(s):

Patient in supine position. General anesthesia and epidural block administered. Time-out was performed and abdomen prepped in sterile fashion. A small infraumbilical incision is made and the abdominal cavity is entered under direct vision. We place two additional trocars, one in the suprapubic region and one in the left lower quadrant, both under direct vision. No iatrogenic injuries occurred during trocar placement. Upon entering the abdominal cavity, significant adhesions noted. The surgical field demonstrated shrunken appendix with marked surrounding inflammation. Multiple small abscesses were encountered. Minimal adhesions were noted. No evidence of peritoneal carcinomatosis. Surrounding omentum and bowel are separated from the inflammatory mass. Mesenteric vessels to the appendix are secured prior to removal. Endoscopic stapling is used for appendiceal division. Specimen placed in EndoCatch bag for removal. Thorough irrigation of the abdominal cavity is performed, removing all purulent material. We close the fascia at the umbilical port with figure-of-eight sutures of Ethibond 2-0 and the skin with non-absorbable Prolene 4-0.

Patient will remain on IV antibiotics with gradual diet advancement as per surgery protocol. The patient tolerated the procedure well with minimal blood loss.

Case ID: CASE-UGTZWX-12124